



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Nurse-physician collaboration: a meta-analytical investigation of survey scores.

This is the peer reviewed version of the following article:

Original

Nurse-physician collaboration: a meta-analytical investigation of survey scores / Sollami, Alfonso; Caricati, Luca; Sarli, Leopoldo. - In: JOURNAL OF INTERPROFESSIONAL CARE. - ISSN 1356-1820. - 29:3(2015), pp. 223-229. [10.3109/13561820.2014.955912]

Availability:

This version is available at: 11381/2781734 since: 2022-01-21T11:18:35Z

Publisher:

Informa Healthcare

Published

DOI:10.3109/13561820.2014.955912

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

Abstract

This meta-analysis investigated differences between nurses and physicians in interprofessional collaboration (IPC) ratings. Fifty-one surveys, representing a total of 18,782 professionals and students (13,132 nurses and nursing students, and 5,650 physicians and medical students), were meta-analyzed, considering several moderating variables. Overall, nurses scored higher on IPC than physicians. Sensitivity analysis revealed that while physicians perceived more existing collaboration than nurses, nurses had a more positive attitude toward collaboration than physicians. Moreover, IPC ratings of nursing and medical students did not differ from those of practitioners. Finally, it appeared that interprofessional education interventions were able to reduce the difference in IPC between nurses and physicians.

Introduction

Nurse-physician **interprofessional** collaboration (**IPC**) can be defined as the joint decision process in which nurses and physicians share objectives and responsibility of results. This process **should be** characterized by mutual trust, open communication, respect and knowledge about one's role and autonomy (Weiss & Davis, 1985). Physicians and nurses **should** work together in a non-hierarchical way, and **should** contribute **equally** to decisions regarding patient care (Taylor, 1996).

For two decades, nurse-physician collaboration has been advocated as an important strategy for improving both clinical care deliveries and organizational outcomes (**WHO, 2010**). Accordingly, good **IPC** has been shown to improve several aspects of the care process, such as quality of patient care, health outcomes (e.g., Baggs et al., 1999) and decreased mortality (Schraeder, **Shelton, & Sager, 2001**). **Contrariwise**, poor **IPC** negatively impacts patient outcomes and quality of delivered care (Callahan et al., 2006; Rosenstein & O'Daniel, 2005), **which leads** to a high level of dissatisfaction among professionals' (e.g., Lim, **Bogossian, & Ahern, 2010**), thus increasing the rates of resignation (Nelson, **King, & Brodine, 2008**) and turnover (Hughes & Fizpatrick, 2010).

Despite the fact that **IPC** is advantageous for **professionals and patients**, there are still several barriers to the effectiveness of nurse-physician collaboration (Rice et al., 2010). Several evidences demonstrate that nurses and physicians have a different understanding of collaboration (Kripalani et al., 2007), work together in a conflictual way (Robinson et al., 2010), and experience tense and even hostile relationships (e.g., Papathanassolou et al., 2012). These barriers are rooted in historical differences between the roles of nurses and physicians, which traditionally are separated by status and power in hospitals (Schmalenberg & Kramer, 2009) **as well as by** educational **and training program** differences (Skjoshammer, 2001).

Other studies, **however**, demonstrate that the nurse-physician relationship is improving, becoming more collegial (Schmalenberg & Kramer, 2009) and characterized by mutual trust, respect, autonomy and participation regarding patient care (Tang, **Chan, Zhou, & Liaw, 2013**).

Some studies indicate that common interprofessional education programs (IPE) **are one of the factors that** can improve **IPC**, and that nurses and physicians are increasingly understanding the importance of working together for healthcare outcomes (Reeves, **Perrier, Goldman, Freeth, & Zwarenstein, 2013**).

In summary, **notwithstanding some** evidence of improved nurse-physician collaboration, differences between nurses and physicians still remain, and require further effort to enhance their professional relationship. Thus, knowing the extent to which nurses and physicians differ in their predisposition toward **IPC** is crucial to the process of improving IPC.

Objectives

The present meta-analysis aims to **investigate the extent to which nurses and physicians differ in their ratings of IPC**, and to **evaluate** potential moderators of **any observed differences**. The specific questions are:

- 1) Do nurses and physicians differ in their *attitude* toward **IPC**?
- 2) Do nurses and physicians differ in their *perception* of **IPC**?
- 3) What factors can moderate nurse-physician difference **in IPC rating**?

Before proceeding, questions 1 and 2 merit some clarifications. Several scales have been developed for measuring different dimensions of IPC that can, nevertheless, be **assumed** in the **aforementioned** general definition of **IPC**. However, in our opinion, these scales do not measure precisely the same thing, because some scales measure professionals' *attitude* toward **IPC**, asking professionals to express their agreement toward some statement related to **IPC**. Other scales measure the *perception* of existing **IPC**, asking practitioners to indicate the extent to which certain collaborative behaviors occur in their care units. **Attitudes and perceptions influence each other (Eagly & Chaiken, 1993), but are very different psychosocial constructs, since** attitudes refer to an expression of favor or disfavor toward a particular social object (in this case, **IPC**), **thus reflecting** the individual's desired aspect of the social context. Perceptions, instead, refer to individuals' experience **of** particular behaviors in a certain situation. Thus, perception pertains to the existing

situation, rather than the desired situation. Unfortunately, researchers tend to not state explicitly which kind of measure they use, and to overlap attitude and perception of IPC. According to Ødegård and Bjørkly (2012), the lack of clarity regarding the IPC measures developed and used (i.e., the IPC operationalization) could in part explain the motive for which “the meaning of IPC is somewhat inconclusive” (p. 284). For this reason, in this meta-analysis, the measure (attitudes vs. perception) was coded in order to control for the moderating effect.

The next sections follow the PRISMA statement for reporting systematic reviews and meta-analyses (Moher, Liberati, Tetzlaff, & Altman, 2009).

Method

Eligibility criteria

Reports were included if the following criteria were met: (a) published between 1999 and 2013, (b) quantitatively analyzed the nurse-physician collaboration, (c) included results from both nurses and physicians, (d) used a validated instrument to assess IPC, (e) reported sufficient statistics for computing effect size. Usually, meta-analyses are limited to English publications, and this has been advocated as a limit for systematic inquiries (Card, 2012). The present meta-analysis also considered articles in Italian, given that Italian is the first language of the authors.

Literature search

Search of studies comprised both scientific and gray literature. Computerized searches were conducted using Medline, CINAHL, SSCI, Psych INFO and PUBMED, using OVID and EBSCO software. For gray literature, Google and Google Scholar were used. Searching keywords were *collaboration, interdisciplinary, interprofessional, relations, nurses and physicians*, using Boolean operators AND and OR in several combinations of the search.

Study selection

Excluding duplicated records and those not in English or Italian, the search yielded a total of 890 records, which were screened for inclusion in the meta-analysis. Figure 1 shows the flowchart for inclusion/exclusion of records. This screening procedure left a total of 35 studies supplying 51

unique measures of **IPC**.

Figure 1 here

Data collection

Two authors (**A.S. and L.C.**) independently read all considered studies, **coding several items (complete description of surveys available on request from the first author)**. In order to compute effect size, the mean, standard deviations and sample size of both nurses and physicians, as well as statistical tests of differences, were collected. When the total score¹ was not reported, weighted means and pooled standard deviations for both nurses and physicians were computed from disaggregated data.

Quality rating of surveys

Quality assessment is fundamental for the evaluation of consistency and validity of findings in each meta-analyzed survey. **Given that** studies were primarily correlational, it was impossible to apply standardized instruments to evaluate quality. Thus, a set of nine dichotomous (**yes/no**) *ad-hoc* indicators relevant to correlational designs were realized in accordance with Zangaro and Soeken (2007): 1) research questions are clearly stated, 2) participants in sample are described, 3) sub-samples of nurses and physicians are described, 4) the setting is described, 5) methods of data collection are described, 6) response rate is indicated, 7) operational definition of **IPC** is clearly stated, 8) instrument used to measure **IPC** are clearly indicated, and 9) internal consistency of the instruments are reported. A rating score was computed, summing the responses to questions 1-9. A total score lower or equal to 4 indicated poor quality, a score of 5 to 7 indicated medium quality, and a score of 8 to 9 indicated high quality.

Three judges (**C.F., G.M., A.S.**) independently evaluated each survey with an inter-rater agreement (Krippendorff's alpha, see Hayes & Krippendorff, 2007) ranging between 0.82 and 1, indicating strong agreement among judges. The few disagreements were discussed and a consensus

was quickly reached.

Data analysis

The mean difference between nurses' and physicians' IPC ratings was measured by Cohen's d , and negative values of d indicate a higher rating for nurses. The standardized mean difference, SDs and weights were corrected for small sample size bias (Lipsey & Wilson, 2001).

Given that meta-analyzed studies were very different in many aspects, the random effect model was chosen. Heterogeneity among studies was assessed with I -squared statistic, which represents the percentage of variance in meta-analysis that is attributable to heterogeneity.

Moderation with sub-group analysis (in case of categorical moderators), and meta-regression with restricted maximum likelihood estimation (in case of continuous moderators), were analyzed. For sub-group analysis, difference in effect size has been statistically tested by comparing Q for each sub-group (Lipsey & Wilson, 2001). This analysis compares between and within group heterogeneity, supplying statistical tests for the difference between levels of the moderator.

Finally, publication bias was assessed using both Begg's and Egger's tests.

Results

Characteristics of study samples

The total number of participants was 18,782. Thirty-nine out of 51 surveys concerned professional nurses ($N = 7,898$; 65.9%) and physicians ($N = 4,084$; 34.1%), for a total of 11,982 practitioners (63.8%). The remaining 12 surveys covered nursing ($N = 5,234$; 77.0%) and medicine ($N = 1566$; 23.0%) students for a total of 6,800 students (36.2%).

Mean age of the participants (number of studies $k = 25$) was 41.29 years ($SD = 7.14$), and average tenure ($k = 18$) was 16.99 years ($SD = 9.69$).

Characteristics of studies

Three studies were longitudinal and measured collaboration before and after an IPE. Other studies did not include a pre-test, but collected data after an IPE intervention. Surveys without ($k = 36$) or before ($k = 3$) an IPE intervention were coded in the same way ("single or pre-IPE," $k = 39$).

Surveys after an IPE intervention were coded as “post-IPE” ($k = 12$) regardless of whether they had a pre-IPE collection ($k = 4$) or not ($k = 8$).

Thirty-six surveys used scales measuring attitude toward IPC while 15 surveys measured perception of IPC. Forty-one surveys (80.4%) were from scientific literature, while ten surveys (19.6%) came from gray literature. Twenty-eight surveys (54.9%) were conducted in the U.S., 12 (23.5%) in Europe and the remaining 11 (21.6%) in other countries. Only 25 surveys reported the facility: eleven (44.0%) were from an intensive care facility, 13 (52.0%) from an ordinary care facility and one (4.0%) from a community care facility.

Finally, 30 surveys (58.8%) were rated as medium quality, 20 (39.2%) were rated as high quality and only one was rated as poor quality ($M = 7.10$, $SD = 1.28$).

Pooled results

The overall effect size was significant ($d = -.34$, 95% C.I. = $-.51$ to $-.17$, $Z = 3.93$, $p < .001$), indicating a moderately higher rating for nurses (Figure 2).

No single survey influenced the overall result, given that overall d showed very little change (range = $-.37$ to $-.32$) after omitting each survey in turn.

The overall effect, however, had great heterogeneity ($Q(50) = 1196.70$, $p < .001$, $I^2 = 95.8\%$), indicating effect size variability among surveys.

Figure 2 here

Sensitivity analysis

Results of moderation analysis are reported in Table 1. Type of measurement yielded different results ($Q(1) = 46.42$, $p < .001$): when attitude was considered, a strong effect favoring nurses emerged ($d = -.64$, $Z = 7.76$, $p < .001$), while when perception was considered, a moderate effect favoring physicians appeared ($d = .45$, $Z = 3.44$, $p = .001$).

Scores for students versus professionals showed no significant difference ($Q(1) = .04$, $p =$

.84), because both students ($d = -.32, Z = 2.52, p = .01$) and professionals ($d = -.35, Z = 3.12, p = .002$) showed moderated effects favoring nurses or nursing students.

The type of design revealed **no significant difference** ($Q(1) = 3.36, p = .07$), although studies with unique measure, or pre-IPE intervention showed a higher score for nurses ($d = -.43, Z = 4.00, p < .001$), while studies realized after an IPE intervention evidenced no differences between professions ($d = -.06, Z = .44, p = .66$).

Facility significantly moderated the difference between nurses and physicians ($Q(1) = 4.00, p = .045$), **showing** that, in ordinary care units, nurses tended to have a higher mean than physicians ($d = -.42, Z = 1.86, p = .06$), while the opposite tended to occur in ICU ($d = .23, Z = 1.00, p = .32$).

Country did not significantly moderate the difference between nurses and physicians ($Q(2) = 1.01, p = .60$).

Also, publication type **was not significant** ($Q(1) = .04, p = .84$) as surveys from both scientific ($d = -.33, Z = 3.51, p < .001$) and gray literature ($d = -.37, Z = 1.63, p = .10$) showed effects favoring nurses over physicians.

Finally, meta-regression showed no significant effect of quality ratings ($b = .11, SE = .08, t = 1.32, p = .19$), but a significant effect of publication year ($b = .06, SE = .03, t = 2.28, p = .02$), indicating that difference between nurses and physicians regarding **IPC** has reduced in recent years.

Table 1 here

Further analysis

The specific scales **measuring IPC** deserve further attention. Thus, a sensitive analysis considering the scale was conducted (Table 2).

Only CSACDS and NPCCS yielded no significant differences between nurses and physicians. However, analysis of the influence of a single survey revealed that by excluding one survey (Nair et al., 2012), the pooled effect of NPCCS **was** significant ($d = .53, Z = 3.52, p < .001, 95\% CI = .23$ to $.82$).

Table 2 here

Publication bias

Begg's and Egger's tests indicated no publication bias (Begg's test: $Z = 1.50, p = .13$; Egger's test: $t = .84, p = .40$).

Discussion

This is one of the first meta-analyses on differences in nurse-physician ratings of IPC, and factors which moderate these differences.

Results indicated that overall, nurses showed a greater predisposition towards IPC than did physicians, independently by the country in which they worked. This finding, albeit of moderate intensity, indicates that physicians would be somewhat reluctant to engage in an effective collaborative practice (e.g., Makary, Sexton, Freischlag, Holzmueller, & Millman, 2006).

Meta-analysis also showed considerable variability in the differences between physicians and nurses across studies. A powerful moderator was the considered measure of IPC. Indeed, if scales measuring IPC attitude were considered, a systematic and strong effect favoring nurses emerged, indicating that nurses hold more positive attitudes toward IPC than do physicians. Contrariwise, considering scales measuring IPC perception, physicians perceived more existing collaboration than did nurses. This evidence suggests that nurses and physicians have a different understanding of what collaboration is, and recognize IPC to different extents (Lingard et al., 2012; Makary et al., 2006). Indeed, on one hand, nurses seem to ask for more collaboration (e.g., more professional autonomy, or "emancipation" in Haddara & Lingard's (2013) terms), while on the other hand, physicians seem to affirm that a good collaboration already exists. Given that attitudes and perceptions about IPC refer to two different aspects of the work environment (desired vs. interpretation of the existing context), the difference between nurses and physicians indicates not simply a difference between professionals, but rather a different understanding of the job context,

and different interest implied in the **IPC**.

These results also raise some important concerns with respect to the use of instruments to measure **IPC**. Analysis suggests that researchers should be well aware of the conceptual definition of **IPC** they are considering, and cautious when they choose the instrument to measure nurse-physician collaboration. Indeed, different kinds of measures may supply biased results favoring either nurses or physicians (see also Ødegård & Bjørkly, 2012).

Another **moderator** was IPE intervention. Studies collecting data after an IPE intervention indicated no difference between nurses and physicians regarding **IPC**, while studies collecting data first **or** without an IPE showed an effect favoring nurses over physicians. This suggests that interdisciplinary trainings can effectively improve interprofessional skills and cooperative behavior, especially for physicians. However, the difference between studies with and without IPE **was** not statistically significant. This imposes caution about the strength of the positive effect of interdisciplinary training, and confirms the urgent need to implement strategies for **improving** collaborative work (Mann et al., 2009).

Also, facility type appeared to significantly affect **IPC** between nurses and physicians. Results show that in ordinary wards, nurses have higher scores than physicians, while in **ICUs**, physicians showed a higher score. This seems to strengthen the idea **that work organization of ICUs could be used as a lesson for improving IPC in other areas of care** (Stein-Parbury & Liaschenko, 2007).

An important, and somewhat surprising, result **is** that both **practitioners** and students showed the same pattern, that is, nurses and nursing students have higher scores in **IPC** than do physicians and medical students. Given the strong emphasis on common education in healthcare, raised in the last decade, **this result seems to indicate that further work is needed to train students to go beyond the confines of their own discipline** (e.g., Abbot, Watson, & Townsley, 2005). **This seems** to be more urgent for medical students, who are still trained to be independent and autonomous (Atwal & Caldwell, 2005).

A final **observation** regards **the significant decrease of nurse-physician difference in IPC** in the

last years. Although substantial work is needed to overcome the remaining barriers to collaboration, the reduction of differences in attitude and behavior linked to nurse-physician collaboration may be due to strong efforts made by healthcare organizations, in order to enhance professionals' awareness of the importance of IPC for improving the quality of patient care. This indicates that organizations are effectively answering the principal challenges for healthcare industries, which, nowadays, are organizational rather than clinical (Ramanujam & Rousseau, 2006).

Limits

This meta-analysis has several limitations. Firstly, despite an attempt to consider the gray literature and not limit the analysis to English papers, studies in other languages were omitted. However, the relatively wide geographical provenience of surveys permits some optimism regarding the generalizability of results. Another limitation concerns the fact that in moderation analysis, some surveys were not actually independent. This may affect in particular the estimation of the IPE effect. However, the small number of longitudinal surveys (3 over 51), seems to restrict this threat. A further limitation is the use of the total score of each scale instead of the score of each sub-dimension. This may obscure the differences between nurses and physicians on particular aspects of IPC. However, this choice was necessary for comparing ratings of both groups of professionals. Moreover, although scales may be composed of several subscales, they normally refer to one main, general construct of IPC. This increases the possibility of comparing overall scores of different scales. Finally, further moderation analyses on each scale or kind of measure would be addressed in future work.

Conclusion

The most significant implication of this study concerns the difference in the understanding and meaning of collaboration between nurses and physicians. Given that healthcare organizations are spending considerable money in order to improve IPC, it is crucial to be aware that nurses and physicians view collaboration in a different way. This, in turn, is reflected in the choice of the instrument used to measure IPC.

Declaration of Interest

The authors report no conflicts of interest.

Reference (asterisks indicate studies included in the meta-analysis)

- Abbot, D., Watson, D., & Townsley, R. (2005). The proof of the pudding: what difference does multi-agency working make to families with disabled children with complex health care needs? *Child and Family Social Work*, 10, 229–238.
- *Ardahan, M., Akçasu, B., & Engin, E. (2010). Professional collaboration in students of medicine faculty and school of nursing. *Nurse Education Today*, 30(4), 350-354.
- Atwal, A., & Caldwell, K. (2005). Do all health and social care professionals interact equally: A study of interactions in multidisciplinary teams in the United Kingdom. *Scandinavian Journal of Caring Sciences*, 19(3), 268–273.
- Baggs, J.G., Schmitt, H.M., Mushlin, A.I., Mitchell, P.H., Eldredge, D.H., Oakes, D. & Hutson, A.D. (1999). Association between nurse-physician collaboration and patient outcomes in three intensive care units. *Critical Care Medicine*, 27, 1991–1998.
- Callahan, C.M., Boustani, M.A., Unverzagt, F.W., Austrom, M.G., Damush, T.M., Perkins, A.J., ... Hendrie, H.C. (2006). Effectiveness of collaborative care for older adults with Alzheimer disease in primary care: a randomized controlled trial. *JAMA*, 295(18), 2148–2157.
- Card, N.A. (2012). *Applied Meta-Analysis for Social Science Research*. New York: Guilford press.**
- *Caricati, L., Mancini, T., Sollami, A., Guidi, C., Prandi, C., Bianconcini, M., ... Artioli, G. (2013a). Nurse-Physician Collaborative Scale: A contribution to the Italian validation. *TPM - Testing, Psychometrics, Methodology in Applied Psychology*, 20(3), 263-276
- *Caricati, L., Mancini, T., Bianconcini, M., Guidi, C., Prandi, C., Silvano, R., ... Artioli, G. (2013b). Nurse-physician collaboration in hospital: A social identity approach. *Unpublished Manuscript*.
- *Cicchetti, A., Lo Scalzo, A., Tangucci, M., Amicosante, A.M.V., & Liva, C. (2002). *La relazione medico-paziente e la relazione medico-infermiere presso il Policlinico Agostino Gemelli – Agenzia per i Servizi Sanitari Regione Lazio*.

- *Curran, V., Sharpe, D., Flynn, K., & Button, P. (2010). A longitudinal study of the effect of an interprofessional education curriculum on student satisfaction and attitudes towards interprofessional teamwork and education. *Journal of Interprofessional Care*, 24(1), 41-52.
- *Dillon, P., Noble, K.A., & Kaplan, L. (2009). Collaborative Interdisciplinary Education Nursing. *Education Research*, 30(2), 87-90.
- *Donald, F., Mohide, E.A., Dicenso, A., Brazil, K., Stephenson, M., & Akhtar-Danesh, N. (2009). Nurse practitioner and physician collaboration in long-term care homes: survey results. *Canadian Journal of Aging*, 28(1), 77-87.
- Eagly, A., & Chaiken, S. (1993). *Psychology of Attitudes*. New York, NY: Harcourt, Brace Jovanovich.
- *El Sayed, K.A., & Sleem, W.F. (2011). Nurse–physician collaboration: A comparative study of the attitudes of nurses and physicians at Mansoura University Hospital. *Life Science Journal*, 8(2), 140-146.
- *Garber, J. S., Madigan, E., Click, E.R., & Fitzpatrick J.J.(2009). Attitudes towards collaboration and servant leadership among nurses, physicians and residents. *Journal of Interprofessional Care*, 23(4), 331-340.
- *Gerard, C.J. (2011). *The effect of a communication protocol implementation on nurse/physician collaboration and communication*. University of Louisville: Louisville, Kentucky
- *Gillen, K. (2007). *Understanding Attitude toward nurse/physician Collaboration in Practicing Nurses and Physicians*. Master Thesis, University of Nevada, Reno
- Haddara, W. & Lingard, L. (2013). Are we all on the same page? A discourse analysis of interprofessional collaboration. *Academic Medicine*, 88(10), 1509-1515.
- *Hamric, A.B., & Blackhall, L.J. (2007). Nurse-physician perspectives on the care of dying patients in intensive care units: Collaboration, moral distress, and ethical climate. *Critical Care Medicine*, 35(2), 422-429.

- *Hansson, A., Avremon, T., Marklund, B., Gedda, B., & Mattson B. (2010). Working together-- primary care doctors' and nurses' attitudes to collaboration. *Scandinavian Journal of Public Health*, 38, 78-85.
- Hayes, A. F., & Krippendorff, K. (2007). Answering the call for a standard reliability measure for coding data. *Communication Methods and Measures*, 1, 77-89.
- *Hind, M., Norman, I., Cooper, S., Gill E., Hilton, R., Judd P., & Jones S.C. (2003). Interprofessional perceptions of health care students. *Journal of Interprofessional Care*, 17(1), 21-34.
- *Hojat, M., Fields, S., Veloski, J., Griffiths, M., Cohen, M., & Plumb, J. (1999). Psychometric properties of an attitude scale measuring physician nurse collaboration. *Evaluation & the Health Professions*, 22(2), 208- 220.
- *Hojat, M., Nasca, T.J., Cohen, M.J., Fields, S.K., Rattner, S.L., Griffiths, M., ... Garcia, A. (2001). Attitudes toward physician-nurse collaboration: a cross-cultural study of male and female physicians and nurses in the United States and Mexico. *Nursing Research*, 50(2), 123-128.
- *Hojat, M., Gonnella, J., Nacsa, T., Fields, S., Cicchetti, A., & Scalzo, A. (2003). Comparison of American, Israeli, Italian and Mexican physician and nurse on the total and factor scores of Jefferson Scale of attitudes toward physician nurse collaborative relationships. *International Journal of Nursing Studies*, (40), 427-435.
- *Hughes, B., & Fitzpatrick, J. J. (2010). Nurse-physician collaboration in an acute care community hospital. *Journal of Interprofessional Care*, 24(6), 625-632.
- Kripalani, S., LeFevre, F., Philips, C.O., Williams, M.V., Basaviah, P., & Baker, D.W. (2007). Deficit in communication and information transfer between hospital-based and primary care physician: implication for patient safety and continuity of care. *JAMA*, 297, 831-841.
- *Jones, T.S., & Fitzpatrick, J.J. (2009). CRNA-physician collaboration in anesthesia. *AANA Journal*, 77(6), 431-437.

- Lim, J., Bogossian, F. & Ahern, K. (2010). Stress and coping in Singaporean nurses: a literature review. *Nursing and Health Sciences*, 12, 251–258.
- Lingard, L., Vanstone, M., Durrant, M., Fleming-Carroll, B., Lowe, M., Rashotte, J., ... & Tallett, S. (2012). Conflicting messages: examining the dynamics of leadership on interprofessional teams. *Academic Medicine*, 87(12), 1762-1767.
- Lipsey, M., & Wilson, D. (2001). The role of method in treatment effectiveness research: Evidence from meta-analysis. *Psychological Methods*, 6(4), 413-429.
- Mann, K.V., Mcfetridge-Durdle, J., Martin-Misener, R., Clovis, J., Rowe, R., Beanlands, H., & Sarria, M. (2009). Interprofessional education for students of the health professions: the "Seamless Care" mode. *Journal of Interprofessional Care*, 23(3), 224-233.
- Makary, M. A., Sexton, J. B., Freischlag, J. A., Holzmueller, C. G., & Millman, E. A. (2006). Operating room teamwork among physician and nurses: teamwork in the eye of the beholder. *Journal of American College Surgery*, 2(202), 746-752.
- *McCaffrey, R., Hayes, R.M., Cassell, A., Miller-Reyes, S., Donaldson, A. & Ferrell, C. (2012). The effect of an educational programme on attitudes of nurse and medical resident towards the benefits of positive communication and collaboration *Journal of Advanced Nursing*, 68(2), 293-301.
- *Miller, P.A. (2001). Nurse-Physician Collaboration in an Intensive care-Unit. *American Journal of Critical Care*, 10(5), 341-350.
- Moher, D., Liberati, A., Tetzlaff, J., & Altman, D.G. (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Annual Internal Medicine*, 151(4), 264-269.
- *Nair, D.M., Fitzpatrick, J.J., McNulty, R., Click, E.R., & Glembocki, M.M., (2012). Frequency of nurse-physician collaborative behaviors in an acute care hospital. *Journal of Interprofessional Care*, 26(2), 115-120.

- *Nathanson, B.H., Henneman E.A., Blonaisz, E.R., Doubleday N.D., Lusardi, P., & Jodka, P.G. (2011). How much teamwork exists between nurses and junior doctors in the intensive care unit? *Journal of Advanced Nursing* 67(8), 1817-1823.
- *Nelson, G. A., King, M.L., & Brodine, S. (2008). Nurse-Physician Collaboration on Medical-Surgical Units. *Medical Surgical Nursing Journal*, 17 (1), 35-40.
- Ødegård, A., & Bjørkly, S. (2012). A mixed method approach to clarify the construct validity of interprofessional collaboration: An empirical research illustration. *Journal of Interprofessional Care*, 26, 283-288.
- Papathanassoglou, E.D., Karanikola, M.N., Kalafati, M., Giannakopoulou, M., Lemoindou, C., & Albarran, J.W. (2012). Professional autonomy, collaboration with physicians, and moral distress among European intensive care nurses. *American Journal of Critical Care*, 21, 41–52.
- *Pomari, C., (2009). Studio descrittivo sugli atteggiamenti riguardanti la collaborazione medico-infermiere in un'Azienda Sanitaria Locale veneta. *Professioni Infermieristiche* 62 (2), 109-119.
- *Pevida, A. (2009). *Measurement of attitude towards nurse/physician collaboration in the health care corporation of St. John's*. Master Thesis: Memorial University of Newfoundland.
- Ramanujam, R., & Rousseau, D.M., (2006). The challenges are organizational not just clinical. *Journal of Organizational Behavior*, 27 (7), 811-827.
- Reeves, S., Perrier, L., Goldman, J., Freeth, D., & Zwarenstein, M. (2013). Interprofessional education: effect on professional practice and health care outcomes (update). *The Cochrane Database of Systematic Reviews*, 28(3), CD002213.
- *Refatti, C., & Bevilacqua, A. (2007). L'integrazione tra infermiere e medico nelle medicine. Risultati di un'indagine. *Assistenza Infermieristica e Ricerca*, 26 (1), 32-37.
- Rice, K., Zwarenstein, M., Conn, G.L., Kenaszchuck, C., Russel, A., & Reeves, S. (2010). An intervention to improve interprofessional collaboration and communications: A comparative qualitative study, *Journal of Interprofessional Care*, 24(4), 350-361.

- Rosenstein, A.H., & O'Daniel, M. (2005). Disruptive behavior and clinical outcomes: perceptions of nurses and physicians. *The American Journal of Nursing*, 105(1), 54–64.
- *Russel, L., Nyohf-Young, J., Abosh, B., & Robinson, S. (2006). An exploratory analysis of an interprofessional learning environment in two hospital clinical teaching units. *Journal of Interprofessional Care*, 20(1), 29-39.
- Schmalenberg, C., & Kramer, M. (2009). Nurse-physician relationships in hospitals: 20000 nurses tell their story. *Critical Care Nurse*, 29 (1), 74–83.
- Schraeder, C., Shelton, P., & Sager, M. (2001). The effects of a collaborative model of primary care on the mortality and hospital use of community-dwelling older adults. *The Journals of Gerontology Series A*, 56(2), 106–12.
- Skjorshammer, M. (2001). Cooperation and conflict in a hospital: interprofessional differences in perception and management of conflict. *Journal of Interprofessional Care*, 15, 7-18.
- Stein-Parbury, J. & Liaschenko, J. (2007). Understanding collaboration between nurses and physicians as knowledge at work. *American Journal of Critical Care*, 16 (5), 470–478.
- *Sterchi, S. (2007). Perception that affect physician-nurse collaboration in the perioperative setting. *AORN Journal*, 86(1), 46-57.
- *Taylor, C. (2009). Attitudes toward physician-nurse collaboration in anesthesia. *American Association of Nurse Anesthetist Journal*, 77(5), 343-348.
- Taylor, J.S. (1996). Collaborative practice within the intensive care unit: a deconstruction. *Intensive and Critical Care Nursing*, 12, 64–70.
- Tang, C.J., Chan, S.W., Zhou, W.T. & Liaw, S.Y. (2013). Collaboration between hospital physicians and nurses: An integrated literature review. *International Nursing Review*, 60, 291–302.
- *Tschannen, D. (2004). The effect of individual characteristics on perceptions of collaboration on the work environment. *Medsurg Nursing*, 13(5), 312- 318.
- *Ushiro, R., (2009). Nurse-Physician Collaboration Scale: development and psychometric testing. *Journal of Advanced Nursing*, 65(7), 1497-1508.

*Yasemin, E., & Yldirim A. (2011). *Attitudes towards physician-nurse collaboration in intensive care unit*. Paper presented at the 4th EfCCNa & FSAIO Congress, Session 18, Copenhagen.

*Yildirim, A., Atesa, M., Akincib, F., Rossb,T., Selimenc, D., Isseverd, H., ... Akgün, M. (2005). Physician–nurse attitudes toward collaboration in Istanbul’s public hospitals. *International Journal of Nursing Studies*, 42, 429–437.

*Yildirim, A, Akinci, F., Ates, M., Ross, T., Issever, H., Isci, E., & Selimen, D. (2006). Turkish version of the Jefferson Scale of Attitudes toward Physician-Nurse Collaboration: a preliminary study. *Contemporary Nurse*, 23(1), 38-45.

Weiss, S., & Davis, H. (1985). Validity and reliability of the collaborative practice scales. *Nursing Research*, 34(5), 299-304.

World Health Organization. (2010). *Framework for action on interprofessional education and collaborative practice*. Geneva: Author. Retrieved from http://whqlibdoc.who.int/hq/2010/WHO_HRH_HP_N_10.3_eng.pdf

Zangaro, G.A., & Soeken, K.L. (2007). A Meta-Analysis of Studies of Nurses’ Job Satisfaction. *Research in Nursing & Health*, 30, 445–458.

Footnotes

¹ Several scales have more than one factor. However, in this meta-analysis, the total score was considered in order to compare differences between nurses and physicians on a global assessment of IPC. Only 5 studies did not report an overall score.

Table 1. Sensitivity analysis for considered moderators

	<i>Q</i> between	<i>k</i>	<i>d</i>	95% C.I.	<i>Q</i> within
Type of measure	46.42**				703.28
Attitude		36	-.64**	-.80 to -.48	617.10
Perception		15	.45**	.19 to .70	86.18
Sample	.04				1191.61
Students		12	-.32*	-.58 to -.07	168.04
Professionals		39	-.35**	-.57 to -.13	1023.57
Design	3.36				1173.62
Unique or Pre IPE		39	-.43**	-.64 to -.22	974.58
Post IPE		12	-.06	-.35 to .22	199.03
Facility type	4.00*				646.07
Intensive care		11	.23	-.22 to .69	248.21
Ordinary care		13	-.42^	-.86 to .02	397.86
Country	1.01				1086.29
Europe		12	-.50*	-.92 to -.08	310.48
USA		28	-.30**	-.52 to -.08	483.01
Other		11	-.28	-.64 to .07	292.80
Publication	.04				1164.90
Scientific		41	-.33**	-.52 to -.15	987.03
Gray		10	-.37	-.81 to .07	177.87

^ $p = .06$; * $p < .05$, ** $p < .01$

Table 2. Sensitivity analysis for scales measuring IPC.

Scale	Q between	k	d	95% C.I.	Q within
	54.71**				589.62
Perception					
CPS		4	.87**	.64 to 1.11	1.60
CSACDS		3	.20	-.69 to 1.09	8.48
ICU NPQ		2	.47**	.13 to .80	0.58
NPCS		5	.23	-.25 to .72	55.81
Attitude					
IHCT		3	-.20**	-.29 to -.11	1.53
JSAPNC		29	-.68**	-.88 to -.48	514.13
RIPLS		4	-.70**	-.84 to -.55	7.48

** $p < .01$

CPS = Collaboration Practice Scale, CSACDS = Collaboration and Satisfaction About Care Decision

Scale, ICU NPQ = ICU Nurse-Physician Questionnaire, IHCT = Interprofessional HealthCare Teams;

JSAPNC = Jefferson Scale of Attitude towards Physician-Nurse Collaboration, NPCS = Nurse-Physician

Collaboration Scale, RIPLS = Readiness for Interprofessional Learning Scale

Figure 1. Flowchart for records inclusion/exclusion




Figure 2. Effect size, variance and forest plot of the analyzed surveys

