



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Influence of milk somatic cell content on Parmigiano-Reggiano cheese yield

This is the peer reviewed version of the following article:

Original

Influence of milk somatic cell content on Parmigiano-Reggiano cheese yield / Summer, Andrea; Franceschi, Piero; Formaggioni, Paolo; Malacarne, Massimo. - In: THE JOURNAL OF DAIRY RESEARCH. - ISSN 0022-0299. - 82:2(2015), pp. 222-227. [10.1017/S0022029915000102]

Availability:

This version is available at: 11381/2787871 since: 2021-10-05T10:33:11Z

Publisher:

Cambridge University Press

Published

DOI:10.1017/S0022029915000102

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

02 May 2026

Proof Delivery Form
Journal of Dairy Research

Date of delivery: 02/05

Journal and vol/article ref: dar /F5/10 JDR no.: 3947

Number of pages (not including this page): 7

This proof is sent to you on behalf of Cambridge University Press. Please check the proofs carefully. Make any corrections necessary on a hardcopy and answer queries on each page of the proofs

Please return the **marked proof** within **2** days of receipt to:

Eric Needs, Editor
Journal of Dairy Research
171 Whitleywood Road
Reading RG2 8LB

Authors are strongly advised to read these proofs thoroughly because any errors missed may appear in the final published paper. This will be your ONLY chance to correct your proof. Once published, either online or in print, no further changes can be made.

To avoid delay from overseas, please send the proof by airmail or courier.

If you have **no corrections** to make, please email **ericneeds@jdrcredit.co.uk** to save having to return your paper proof. If corrections are light, you can also send them by email, quoting both page and line number.

- The proof is sent to you for correction of typographical errors only. Revision of the substance of the text is not permitted, unless discussed with the editor of the journal. Only **one** set of corrections are permitted.
- Please answer carefully any author queries.
- Corrections which do NOT follow journal style will not be accepted.
- A new copy of a figure must be provided if correction of anything other than a typographical error introduced by the typesetter is required.

Copyright: if you have not already done so, please download a copyright form from: <http://journals.cambridge.org/action/displayMoreInfo?jid=DAR&type=tcr> Please sign the form by hand and return by mail to the address shown on the form. Failure to send this form will delay the publication of your article.

- If you have problems with the file please contact **sperkins@cambridge.org**

Please note that this pdf is for proof checking purposes only. It should not be distributed to third parties and may not represent the final published version.

Important: you must return any forms included with your proof. We cannot publish your article if you have not returned your signed copyright form.

NOTE - for further information about **Journals Production** please consult our **FAQs** at http://journals.cambridge.org/production_faqs

Offprint order form



CAMBRIDGE
UNIVERSITY PRESS

PLEASE COMPLETE AND RETURN THIS FORM. WE WILL BE UNABLE TO SEND OFFPRINTS UNLESS A RETURN ADDRESS AND ARTICLE DETAILS ARE PROVIDED.

VAT REG NO. GB 823 8476 09

Journal of Dairy Research (DAR)

Volume: no:

If you require paid offprints please complete this form and send it to **Sue Perkins (address below)**. Please give the address to which your offprints should be sent. They will be dispatched by surface mail within one month of publication. For an article by **more than one author this form is sent to you as the first named. All offprints should be ordered by you in consultation with your co-authors.**

Number of paid offprints required

Email:

Offprints to be sent to (print in BLOCK CAPITALS):

.....

Post/Zip Code:
Telephone: Date (dd/mm/yy): / /

Author(s):

Article Title:

All enquiries about offprints should be addressed to **Sue Perkins: Journals Production Department, Cambridge University Press, The Edinburgh Building, Shaftesbury Road, Cambridge CB2 2RU, UK.**

Charges for extra offprints (excluding VAT) Please circle the appropriate charge:

Number of copies	25	50	100	150	200	per 50 extra
1-4 pages	£68	£109	£174	£239	£309	£68
5-8 pages	£109	£163	£239	£321	£399	£109
9-16 pages	£120	£181	£285	£381	£494	£120
17-24 pages	£131	£201	£331	£451	£599	£131
Each Additional 1-8 pages	£20	£31	£50	£70	£104	£20

Methods of payment

If you live in Belgium, France, Germany, Ireland, Italy, Portugal, Spain or Sweden and are not registered for VAT we are required to charge VAT at the rate applicable in your country of residence. If you live in any other country in the EU and are not registered for VAT you will be charged VAT at the UK rate. If registered, please quote your VAT number, or the VAT number of any agency paying on your behalf if it is registered. VAT Number:

Payment **must** be included with your order, please tick which method you are using:

- Cheques should be made out to Cambridge University Press.
- Payment by someone else. Please enclose the official order when returning this form and ensure that when the order is sent it mentions the name of the journal and the article title.
- Payment may be made by any credit card bearing the Interbank Symbol.

Card Number:

Expiry Date (mm/yy): / Card Verification Number:

The card verification number is a 3 digit number printed on the **back** of your **Visa** or **Master card**, it appears after and to the right of your card number. For **American Express** the verification number is 4 digits, and printed on the **front** of your card, after and to the right of your card number.

Signature of card holder: Amount (Including VAT if appropriate): £

Please advise if address registered with card company is different from above

Author Queries

Journal: DAR (Journal of Dairy Research)

Manuscript: S0022029915000102jra

- Q1** The distinction between surnames can be ambiguous, therefore to ensure accurate tagging for indexing purposes online (eg for PubMed entries), please check that the highlighted surnames have been correctly identified, that all names are in the correct order and spelt correctly.

Influence of milk somatic cell content on Parmigiano-Reggiano cheese yield

Andrea Summer^{1,2}, Piero Franceschi^{1*}, Paolo Formaggioni¹ and Massimo Malacarne^{1,2}

¹ Department of Food Science, University of Parma, Via del Taglio 10, I-43126 Parma, Italy

² MILC Center, University of Parma, Parco Area delle Scienze, 59/A, I-43124 Parma, Italy

Received 25 November 2013; accepted for publication 14 January 2015

The aim of this study was to determine the influence of the somatic cell content (SCC) of milk on Parmigiano-Reggiano cheese yield, produced in commercial cheese factories under field conditions. The study was carried out following the production of 56 batches of Parmigiano-Reggiano in 13 commercial cheese factories by processing milk collected from Italian Friesian cattle herds. The vat-milk (V-milk) used for making each cheese batch was obtained by mixing evening milk (partially skimmed following spontaneous separation of fat overnight, natural creaming) and morning milk. The batches of cheese produced were divided into 5 classes according to the SCC value of the evening milk determined prior to natural creaming (class 1, from 0 to 200 000; 2, 201 000–300 000; 3, 301 000–400 000; 4, 401 000–500 000; 5, over 501 000 cells/ml). The cheese yield was calculated as the amount of 24-h cheese, expressed in kilograms, obtained from 100 kg of V-milk (24 h ACY). The values of fat, crude protein, true protein, casein and 24 h ACY of V-milk were negatively correlated with the somatic cell score (SCS) of the evening milk. Conversely, a positive correlation was observed between chloride and SCS. Fat, protein fractions (crude protein, casein and whey proteins), P and titratable acidity of V-milk were positively correlated with its 24 h ACY, while chloride, pH and SCS showed a negative correlation. A significant drop in 24 h ACY was observed in classes 3, 4 and 5, therefore when the SCC of the evening milk exceeded 300 000 cells/ml. Finally a lower recovery of milk fat in cheese was observed as SCC of evening milk increase.

Keywords: Bulk milk, cheese-making losses, mastitis, milk composition.

The mammary gland inflammation process is characterised by an increased number of the somatic cells (mainly macrophages, leucocytes and PMN) in milk (Mazal et al. 2007; Viguier et al. 2009). Inflammatory response is also characterised by a transfer of some blood components to the milk (Urech et al. 1999) and by a decrease in secretory activity resulting in reduced milk production (Lescourret & Coulon, 1994; Klei et al. 1998; de los Campos et al. 2006; Le Maréchal et al. 2011). Moreover, alterations are observed in the chemical composition of the milk secreted (Dang et al. 2008; Franceschi et al. 2009; Le Maréchal et al. 2011) and its physicochemical properties (Summer et al. 2003; Franceschi et al. 2009; Le Maréchal et al. 2011).

Many studies have been carried out on the relationships between the characteristics of milk and its cheese yielding capacity. The quantity of cheese produced per unit of milk

mainly depends on the total amount of solid contents, in particular fat and casein, and on the cheese-making technology (Verdier-Metz et al. 2001; Auldust et al. 2004). The influence of the rennet-coagulation aptitude, milk protein composition and genetic variants of milk proteins on the efficiency of the cheese-making process were also considered (Hallén et al. 2010). Milk with better rennet-coagulation properties (RCP) generally produces a larger cheese yield and recoveries of milk casein and fat in the cheese (Aleandri et al. 1989; Malacarne et al. 2006; De Marchi et al. 2008) although no significant differences were also observed in cheese yield between well- or poorly-coagulating milks (Ikonen et al. 1999; Wedholm et al. 2006; Bonfatti et al. 2014). The presence of κ -CN B in milk is associated with a higher casein content with a higher κ -CN proportion, resulting in smaller micelles, which in turn leads to an improvement in the curdling properties of the milk (Pearse et al. 1986) and a greater cheese yield (Mariani et al. 1976; Walsh et al. 1998a, b). There is general consensus concerning the negative effect of milk on the somatic cell

*For correspondence; e-mail: piero.franceschi@unipr.it

content (SCC), cheese yield and quality (Geary et al. 2013). The same effects were also observed in goat and sheep milk (Leitner et al. 2008). In general, a decrease of fat and protein recoveries in cheese was observed as milk SCC increases (Politis & Ng-Kwai-Hang, 1988a, b; Barbano et al. 1991; Klei et al. 1998). This was due to impaired rennet coagulation properties and/or increased proteolysis and lipolysis in high SCC milk. Mazal et al. (2007) found no effect of milk SCC on Prato cheese yield, although cheese produced with high-SCC had higher moisture content and a more intense proteolysis than low-SCC milk cheese. Marino et al. (2005) reported the influence of somatic cell enzymes on cheese proteolysis.

Most of the studies on the yielding capacity of milk cheese were carried out in model cheese produced at laboratory- or pilot-scale level, using individual milk or bulk milk obtained from few selected cows (Cipolat-Gotet et al. 2013). However, the dairy industry would benefit from the being able to investigate the effect of milk characteristics on cheese yield under field conditions. Pretto et al. (2013) recently analysed the relationships between milk composition, coagulation properties and Grana Padano cheese yield under field conditions but they did not consider the influence of milk SCC. Grana Padano cheese is a very hard Protected Designated of Origin (PDO) cheese whose cheese-making technology is very similar to that used for making Parmigiano-Reggiano cheese. The aim of this study was to determine the influence of milk SCC on Parmigiano-Reggiano cheese yield, produced in commercial cheese factories under field conditions. Moreover, an attempt was made to establish a threshold for herd bulk milk SCC above which a significant decrease in Parmigiano-Reggiano cheese yield was observed.

Materials and methods

Experimental design, sampling procedure and classification of cheese batches

The study was carried out following the production of 56 Parmigiano-Reggiano cheese batches in 13 commercial cheese factories under field conditions. The raw milk used for cheese-making was collected from Italian Friesian cattle herds. A brief description of the origin of the raw milk used for cheese-making (known as vat milk, V-milk) is required in order to understand the sampling procedures and how the cheese batches were grouped into SCC classes. In short, the V-milk used for each batch of Parmigiano-Reggiano cheese (about 1100 kg) was composed (1 : 1) of the milk of two consecutive milkings collected from the same herd: the partially skimmed evening milk (obtained through the overnight gravity separation of fat, a process called natural creaming) and the full-cream morning milk. For each batch of cheese made in this study, a sample of full-cream evening milk (E-milk) was collected at its arrival at the cheese factory before proceeding

with the natural creaming process. The following morning, the V-milk was weighed using a volumetric pump and a sample was collected directly from the vat before adding the natural whey-starter. At the end of the cheese-making procedure, a sample of the residual cheese whey was collected from the vat. Sodium merthiolate (0.02 g/100 ml) was added to each type of sample (E-milk, V-milk and cheese whey), cooled to 5 °C and transferred to the laboratory within 30 min where the analyses were carried out within the following 24 h. The 2 twin wheels obtained from each batch of cheese were weighed 24 h after extraction from the vat.

The batches of cheese were divided into 5 classes according to the SCC of the E-milk: class 1, from 0 to 200 000 cells/ml; class 2, from 201 000 to 300 000 cells/ml; class 3, from 301 000 to 400 000 cells/ml; class 4, from 401 000 to 500 000 cells/ml; class 5, over 501 000 cells/ml. The SCC value of the V-milk was not reliable enough for dividing the cheese batches into classes as half of it had undergone the natural creaming process, which caused a significant reduction of somatic cells which only partially depends on the original value and is mainly influenced by technological and environmental factors (Malacarne et al. 2008). Therefore, classification can be only be carried out by measuring the SCC of E-milk or morning milk (the full cream milk). Since the E-milk and morning milk were obtained from two consecutive milkings, it was assumed that they had the same SCC value and we therefore decided to only take the SCC value of the E-milk.

Parmigiano-Reggiano cheese-making

The cheese was produced with the method approved by the Consortium (Council Regulation, 1992).

A natural whey-starter culture (about 3 l for every 100 kg V-milk), obtained by means of the spontaneous acidification of the previous day's cheese whey, was added to the V-milk before coagulation. The cheese milk was then heated to 33 °C and clotted in 10–12 min with calf rennet (2.5 g/100 kg V-milk, calf activity 1 : 125 000, Bellucci Srl, 41123 Modena, Italy). The curd was broken up into small granules (approximately the size of a rice grain) and cooked. During this step, the temperature was increased to 55 °C in 10–15 min and the curd granules were stirred continuously. After cooking the broken-up curd particles settled at the bottom of the vat where they aggregated and blended together spontaneously during the so-called resting phase. During this step, which lasted 45–60 min, the temperature was 55–53 °C. The result was a consolidated cheese mass, which was removed from the vat, divided into two parts and placed in special moulds called 'fascere' for 2 d. During this period the cheese wheels were naturally cooled and periodically turned over in order to dry them homogeneously. The cheese wheels were then submerged in saturated brine for a period of 20–25 d. The cheese finally entered the ripening phase, which lasts about 24 months. At the end of the ripening

stage, the cheese wheels were cylindrical in shape with a slightly convex side, 22–24 cm high, 40–45 cm diam., and weighed 35–36 kg.

Analytical methods

The following standard analyses were performed on the V-milk samples: somatic cells (this parameter was also measured on E-milk); the fat and lactose contents were assessed with mid-infrared readings (Biggs, 1978) with Milko-Scan 134 A/B (Foss Electric); total N (TN), non casein N (NCN) and non protein N (NPN) on milk, acid whey at pH 4.6 and TCA 12% filtered whey, respectively, were determined with the Kjeldahl method (Aschaffenburg & Drewry, 1959), from which the values of crude protein ($TN \times 6.38/1000$), casein ($(TN - NCN) \times 6.38/1000$), casein number ($(TN - NCN) \times 100/TN$), NPN $\times 6.38$ ($NPN \times 6.38/1000$), true protein ($(TN - NPN) \times 6.38/100$) were calculated. Digestion was carried out using DK 6 macro Kjeldahl heating block (Velp Scientifica, Usmate, MB, Italy) and distillation with 126 UDK distillation unit (Velp Scientifica, Usmate, MB, Italy). The content of phosphorous was measured with the colorimetric method (Allen, 1940) while the calcium and magnesium contents were determined with the complexometric method (Ntalianas & Whitney, 1964) (V-milk, cheese whey); chloride concentration was measured by titration according to Savini (1946) (V-milk); pH and titratable acidity were measured in V-milk with a potentiometer and by titration of 50 ml of milk with 0.25 N sodium hydroxide according to the Soxhlet-Henkel method (Anon., 1963).

The content of TN, Ca, P and Mg were determined with the same methods described above also for cheese whey. The fat in the cheese whey was determined with the volumetric Gerber method (Savini, 1946).

The SCC in E-milk was measured by means of the fluoro-opto-electronic method (Schmidt-Madsen, 1975) with Fossomatic 250 (Foss Electric, DK-3400 Hillerød Denmark). The somatic cell score (SCS) was calculated as follows: $SCS = (\log_2(SCC/100) + 3)$ (Shook & Schutz, 1994).

The 24-h actual cheese yield (kg of cheese /100 kg of milk) was calculated as follows:

$$Y = [cw] \times 100/mw$$

where: Y=cheese yield; cw=cheese weight measured 24 h after removing it from the vat, kg; mw=weight of V-milk measured before adding the natural whey starter, kg.

Statistical analysis

The data were analysed with ANOVA univariate using the following model:

$$y_{ijk} = \mu + \alpha_i + \beta_j + \epsilon_{ijk}$$

where: y_{ijk} =dependent variable; μ =overall mean; α_i =fixed effect SCC class ($i = 1, \dots, 5$); β_j =fixed effect of the cheese factory ($j = 1, \dots, 13$, one level for each cheese factory); ϵ_{ijk} =residual error.

ANOVA and Pearson correlation coefficients were calculated with IBM SPSS statistics software, version 21 (Armonk, NY, USA).

Results and discussion

The mean values of the V-milk characteristics and their correlation coefficients with both E-milk SCS and 24 h actual cheese yield (24 h ACY) are shown in Table 1. The contents of the protein fractions and the value of titratable acidity are higher and lower, respectively, than those reported for Parmigiano-Reggiano V-milk collected from Italian Friesian cattle herds by Malacarne et al. (2006). The increase in milk protein is actually due to breeding programs aimed at increasing the amount of casein in milk. Conversely, the values of titratable acidity observed confirm the decreasing trend for this parameter over the years (Formaggioni et al. 2005; Franceschi et al. 2012; Summer et al. 2014).

The values of fat, crude protein, true protein, casein and 24 h ACY were negatively correlated with SCS. Conversely, a positive correlation was observed between chloride and SCS. Fat, protein fractions (crude protein, casein and whey proteins), P and titratable acidity were positively correlated with 24 h ACY while a negative correlation was observed for chloride, pH and SCS. The negative correlation between E-milk SCS and fat content in V-milk could be due to the fact that half of V-milk underwent a natural creaming process in which somatic cells are partially removed from within the surfaced fat globules (Malacarne et al. 2008). In order to determine a possible effect of SCC on milk fat and/or the natural creaming process, the fat content values of the E-milk and the partially skimmed evening are required. Unfortunately, these parameters were not calculated in this study. An interesting result is the decrease of crude protein as the SCS increase observed in this study since there is generally an increase of crude protein in milk with high SCC (Politis & Ng-Kwai-Hang, 1988a, b). However, Le Marechal et al. (2011) obtained conflicting results concerning the impact of mastitis on total protein. In fact, the rupture of the tight junction between epithelial cells within the inflamed quarters increases the transfer of haematic whey proteins into the alveolar lumen via the paracellular route thus causing an increase in milk crude protein and a decrease in casein number (Kitchen, 1981; Shennan & Peaker, 2000). No significant variations of both whey protein and casein number were found in milk with different SCC in this study even if there was a decreasing trend in the casein number in V-milk as SCS increased. The relationships between these parameters and SCC may no longer be detectable at bulk milk level due to the mixing of the milk from inflamed and healthy quarters. Conversely, the decrease in casein and the increase in chloride as the SCC increases are in agreement with previous observations. Hydrolysis of casein fractions by plasmin and somatic

Table 1. Characteristics of vat milk (V-milk) and their correlations with both somatic cell score (SCS) of the full cream evening milk (E-milk) and the actual cheese yield measured after 24 h (24 h ACY)

	Descriptive statistics			Pearson correlation coefficient	
	Mean	SD	<i>n</i>	SCS	24 h ACY
Lactose, g/100 g	4.94	0.16	50	—	—
Fat, g/100 g	2.84	0.20	56	-0.49	0.67
Crude protein, g/100 g	3.26	0.15	56	-0.36	0.71
Wheyprotein, g/100 g	0.57	0.04	56	—	0.37
Casein, g/100 g	2.52	0.12	56	-0.36	0.71
Casein number, %	77.24	0.76	56	—	—
NPN × 6.38, g/100 g	0.18	0.03	56	—	—
True protein, g/100 g	3.09	0.15	56	-0.34	0.70
Ca, mg/100 g	114.15	3.48	56	—	—
P, mg/100 g	90.05	3.69	56	—	0.36
Mg, mg/100 g	10.51	0.67	56	—	—
Cl, mg/100 g	100.28	8.02	56	0.67	-0.35
pH	6.74	0.04	50	—	-0.38
Titrate acidity, °SH/50 ml	3.17	0.15	56	—	0.48
SCS (E-milk)	4.37	0.67	56	—	-0.42

Only significant correlations ($P < 0.05$) are reported




Fig. 1 - BW online, BW in print

Fig. 1. Parmigiano-Reggiano cheese yield, calculated as the kg of 24-h cheese obtained from 100 kg of processed vat milk (24 h ACY), according to somatic content of evening milk, measured prior natural creaming: class 1, from 0 to 200 000; 2, 201 000–300 000; 3, 301 000–400 000; 4, 401 000–500 000; 5, over 501 000 cells/ml. ^{a, b}Are different at $P < 0.05$.

cells proteases are the main reasons for the lower values of casein observed in high SCC milk (Urech et al. 1999; Somers et al. 2003). The increase in chlorides is a typical feature of milk obtained from inflamed quarters and is related to the increased permeability of the epithelial cells layer (Rogers et al. 1989).

As rennet-cheese consists in the formation and consequent dehydration of a paracasein reticulum in which fat droplets

are entrapped, the strong positive correlations between 24 h ACY and the contents of casein and fat in milk were expected and the values were similar to those reported for Grana Padano cheese-making by Pretto et al. (2013). The significant correlations observed between milk P and 24 h ACY mainly depends on the high content of P in milk with elevated casein content since approximately half of the phosphorus contained in milk is associated with casein (Bijl et al. 2013). The negative correlation between 24 h ACY and milk pH could rely on the relationships between milk pH and its rennet coagulation properties (RCP). Although Pretto et al. (2013) did not find a significant correlation between milk pH and its 24 h ACY, these authors report that a higher Grana Padano cheese yield was obtained using milk with better RCP. In fact, there is an improvement in milk RCP as pH decrease is observed (Ikonen et al. 2004; Cassandro et al. 2008). Therefore, a decrease of milk pH should improve its RCP and consequently its 24 h ACY. However, there is not general consensus regarding the relationships between milk RCP in its cheese yielding capacity since several studies carried out at pilot- or laboratory-scale level did not find a significant association (Ikonen et al. 1999; Mazal et al. 2007; Bonfatti et al. 2014). The positive correlation between milk titrate acidity and 24 h ACY was also reported by Pretto et al. (2013). The casein content of milk probably mediates this correlation since milk with high casein levels is expected to have elevated values of titrate acidity. The increase in the proportion of milk produced from inflamed quarters in the bulk tank should be followed by a contextual increase of SCS and chloride. Therefore, the negative correlations observed between chloride and 24 h ACY confirm the reduced cheese yielding capacity of milk produced from inflamed quarters.

The cheese batches were divided into five classes according to the SCC contained in the E-milk. The 24 h ACY was

Table 2. Cheese-making losses† and their correlations with both somatic cell score (SCS) of the full cream evening milk (E-milk) and the actual cheese yield measured after 24 h (24 h ACY)

	Descriptive statistics			Simple correlation	
	Mean	SD	<i>n</i>	SCS	24 h ACY
Protein losses %	27.33	1.06	56	—	—
Fat losses, %	14.95	3.50	56	0.53	—
P losses, %	50.23	2.06	56	—	—
Ca losses, %	35.42	2.17	56	—	—
Mg losses, %	77.52	3.14	56	—	—

Only significant correlations ($P < 0.05$) are reported

†Mean values expressed as follow: concentration in the residual cheese whey $\times 100$ / concentration in V-milk

higher in the batches of cheese in classes 1 and 2 than in the other classes, where no significant differences were observed (Fig. 1). On the other hand, no differences were observed among the classes (data not shown) when the cheese yield was corrected according to the fat and casein content in V-milk. The values of 24 h ACY observed in this study are similar to those previously observed for Parmigiano-Reggiano cheese (Malacarne et al. 2006), and other hard Italian PDO very hard cheeses, namely Grana Padano and Grana Trentino (De Marchi et al. 2008; Pretto et al. 2013). The negative influence of milk somatic cells on its cheese yielding ability confirms results observed for other types of cheeses (Politis & Ng-Kwai-Hang, 1988a, b; Barbano et al. 1991; Klei et al. 1998). However, it seems that variations among classes are mainly due to the difference in milk fat and casein contents as no differences were found when adjusted cheese yield was considered.

The mean values of estimated cheese-making losses (expressed as concentration in cheese whey $\times 100$ /concentration in vat milk) and their correlations with E-milk SCS are reported in Table 2. The values are comparable to those reported for Parmigiano-Reggiano cheese-making by Malacarne et al. (2006). The only significant correlation with SCS was observed for estimated fat loss (14.95%). The lower recovery of fat in cheese is attributed to increased lipolysis in high SCC milk (Barbano et al. 1991)

Conclusions

The increase of milk somatic cell content is associated with a reduction of its cheese yielding ability. The reduced cheese yield is the consequence of a decrease of both milk casein and recovery of milk fat in cheese, as SCC increases. The data reported suggest that there was a significant decrease in 24 h ACY when the value of SCC was above 300 000 cells/ml, therefore before milk SCC reaches the legal limit of 400 000 cell/ml. The difference observed may be relevant for the profit of the cheese factory. Assuming standard Parmigiano-Reggiano cheese-making conditions (1100 kg of V-milk processed, 15% loss of cheese weight throughout a ripening of 24 months) and a price of 9.76 euro/kg for 24 month-old-cheese, according to ACY observed in this study, it is possible to estimate a

decrease in profit of 46 euro for a cheese batches made with milk SCC within the range 300 000–400 000 cells/ml compared to batches made with milk SCC below 300 000 cells/ml.

Further research is required to analyse the relationships between milk somatic cell content and quality of cheese.

The authors would like to thank Prof. Primo Mariani for the important contributions in the writing of this paper.

References

- Aleandri R, Schneider JC & Buttazzoni LG 1989 Evaluation of milk for cheese production based on milk characteristics and formagraph measures. *Journal of Dairy Science* **72** 1967–1975
- Allen R 1940 The estimation of phosphorus. *Biochemical Journal* **34** 858–865
- Anon. 1963 Titratable acidity evaluation with the Soxhlet-Henkel (SH) method. *Milchwissenschaft* **18** 520
- Aschaffenburg R & Drewry J 1959 New procedure for the routine determination of the various non-casein proteins of milk. *Proceedings, 15th International Dairy Congress* **3** 1631–1637
- Auldist MJ, Johnston KA, White NJ, Fitzsimons WP & Boland MJ 2004 A comparison of the composition, coagulation characteristics and cheese-making capacity of milk from Friesian and Jersey dairy cows. *Journal of Dairy Research* **71** 51–57
- Barbano DM, Rasmussen RR & Lynch JM 1991 Influence of milk somatic cell count and milk age on cheese yield. *Journal of Dairy Science* **74** 369–388
- Biggs DA 1978 Instrumental infrared estimation of fat, protein and lactose in milk: collaborative study. *Journal of the Association of Official Analytical Chemists* **61** 1015–1034
- Bijl E, van Valenberg HJF, Huppertz T & van Hooijdonk ACM 2013 Protein, casein, and micellar salts in milk: current content and historical perspectives. *Journal of Dairy Science* **96** 5455–5464
- Bonfatti V, Tuzzato M, Chiarot G & Carnier P 2014 Variation in milk coagulation properties does not affect cheese yield and composition of model cheese. *International Dairy Journal* **39** 139–145
- Cassandro M, Comin A, Ojala M, Dal Zotto R, De Marchi M, Gallo L, Carnier P & Bittante G 2008 Genetic parameters of milk coagulation properties and their relationships with milk yield and quality traits in Italian Holstein cows. *Journal of Dairy Science* **91** 371–376
- Cipolat-Gotet C, Cecchinato A, De Marchi M & Bittante G 2013 Factors affecting variation of different measures of cheese yield and milk nutrient recovery from an individual model cheese-manufacturing process. *Journal of Dairy Science* **96** 7952–7965
- Council Regulation 1992 (EEC) No 2081/92 of 14 July 1992 on the protection of geographical indications and designations of origin for

- agricultural products and foodstuffs. *Official Journal of the European Union L* **208** 1–8
- Dang AK, Suman K, Charan S & Sehgal JP** 2008 Milk differential cell counts and compositional changes in cows during different physiological stages. *Milchwissenschaft* **63** 239–242
- de los Campos G, Gianola D & Heringstad B** 2006 A structural equation model for describing relationships between somatic cell score and milk yield in first-lactation dairy cows. *Journal of Dairy Science* **89** 4445–4455
- De Marchi M, Bittante G, Dal Zotto R, Dalvit C & Cassandro M** 2008 Effect of Holstein Friesian and Brown Swiss breeds on quality of milk and cheese. *Journal of Dairy Science* **91** 4092–4102
- Formaggioni P, Sandri S, Franceschi P, Malacarne M & Mariani P** 2005 Milk acidity, curd firming time, curd firmness and protein and fat losses in the Parmigiano-Reggiano cheesemaking. *Italian Journal of Animal Science* **4** (Suppl. 2) 239–241
- Franceschi P, Summer A, Sandri S, Formaggioni P, Malacarne M & Mariani P** 2009 Effects of the full cream milk somatic cell content on the characteristics of vat milk in the manufacture of Parmigiano-Reggiano cheese. *Veterinary Research Communications* **33**(Suppl. 1) 281–283
- Franceschi P, Malacarne M, Nordera B, Vergnani M, Ghizzoni PG, Formaggioni P, Mariani P & Summer A** 2012 Effect of pedoclimatic conditions on the physico-chemical characteristics, hygiene and coagulation properties of milk for Parmigiano-Reggiano cheese. *Milchwissenschaft* **67** 166–169
- Geary U, Lopez-Villalobos N, O'Brien B, Garrick DJ & Shalloo L** 2013 Meta-analysis to investigate relationships between somatic cell count and raw milk composition, Cheddar cheese processing characteristics and cheese composition. *Irish Journal of Agricultural and Food Research* **52** 119–133
- Hallén E, Lundén A, Allmere T & Andrén A** 2010 Casein retention in curd and loss of casein into whey at chymosin-induced coagulation of milk. *Journal of Dairy Research* **77** 71–76
- Ikonen T, Ruottinen O, Syväoja EL, Saarinen K, Pakkala E & Ojala M** 1999 Effect of milk coagulation properties of herd bulk milk on yield and composition of Emmental cheese. *Agricultural and Food Science in Finland* **8** 411–422
- Ikonen T, Morri S, Tyrisevä AM, Ruottinen O & Ojala M** 2004 Genetic and phenotypic correlations between milk coagulation properties, milk production traits, somatic cell count, casein content, and pH of milk. *Journal of Dairy Science* **87** 458–467
- Kitchen BJ** 1981 Bovine mastitis: milks compositional change and related diagnostic test. *Journal of Dairy Research* **48** 167–188
- Klei L, Yun J, Sapru A, Lynch J, Barbano DM, Sears P & Galton D** 1998 Effects of milk somatic cell count on Cottage cheese yield and quality. *Journal Dairy Science* **81** 1205–1213
- Leitner G, Silanikove N & Merin U** 2008 Estimate of milk and curd yield loss of sheep and goats with intramammary infection and its relation to somatic cell count. *Small Ruminant Research* **74** 221–225
- Le Maréchal C, Thiéry R, Vautor E & Le Loir Y** 2011 Mastitis impact on technological properties of milk and quality of milk products – a review. *Dairy Science and Technology* **91** 247–282
- Lescouret F & Coulon JB** 1994 Modelling the impact of mastitis on milk production by dairy cows. *Journal of Dairy Science* **77** 2289–2301
- Malacarne M, Summer A, Fossa E, Formaggioni P, Franceschi P, Pecorari M & Mariani P** 2006 Composition, coagulation properties and Parmigiano-Reggiano cheese yield of Italian Brown and Italian Friesian herd milks. *Journal of Dairy Research* **73** 171–177
- Malacarne M, Summer A, Formaggioni P, Franceschi P, Sandri S, Pecorari M, Vecchia P & Mariani P** 2008 Dairy maturation of milk used in the manufacture of Parmigiano-Reggiano cheese: effects on chemical characteristics, rennet coagulation aptitude and rheological properties. *Journal of Dairy Research* **75** 218–224
- Mariani P, Losi G, Russo V, Castagnetti GB, Grazia L, Morini D & Fossa E** 1976 Cheesemaking tests made with milk characterized by variants A and B of k-casein in the production of Parmigiano-Reggiano cheese. *Scienza e Tecnica Lattiero-Casearia* **27** 208–227
- Marino R, Considine T, Sevi A, McSweeney PLH & Kelly AL** 2005 Contribution of proteolytic activity associated with somatic cell sin milk to cheese ripening. *International Dairy Journal* **15** 1026–1033
- Mazal G, Vianna PCB, Santos MV & Gigante ML** 2007 Effect of somatic cell count on Prato cheese composition. *Journal of Dairy Science* **90** 630–636
- Ntalianas HA & Whitney RMCL** 1964 Calcein as an indicator for the determination of total calcium and magnesium and calcium alone in the same aliquot of milk. *Journal of Dairy Science* **47** 19–27
- Pearse MJ, Linklater PM, Hall RJ & MacKinlay AG** 1986 Effect of casein micelle composition and casein dephosphorylation on coagulation and syneresis. *Journal of Dairy Research* **53** 381–390
- Politis I & Ng-Kwai-Hang KF** 1988a Association between somatic cell count of milk and cheese-yielding capacity. *Journal of Dairy Science* **71** 1720–1727
- Politis I & Ng-Kwai-Hang KF** 1988b Effects of somatic cell count and milk composition on cheese composition and cheese making efficiency. *Journal of Dairy Science* **71** 1711–1719
- Pretto D, De Marchi M, Penasa M & Cassandro M** 2013 Effect of milk composition and coagulation traits on Grana Padano cheese yield under field conditions. *Journal of Dairy Research* **80** 1–5
- Rogers SA, Mitchell GE & Bartley JP** 1989 The relationship between somatic cell count, composition and manufacturing properties of bulk milk 4. Non-protein constituents. *Australian Journal of Dairy Technology* **44** 53–56
- Savini E** 1946 *Analysis of Milk and Dairy Products*. Milano: Hoepli
- Schmidt-Madsen P** 1975 Fluoro-opto-electronic cell-counting on milk. *Journal of Dairy Research* **42** 227–239
- Shennan DB & Peaker M** 2000 Transport of milk constituents by the mammary gland. *Physiological Reviews* **80** 925–951
- Shook GE & Schutz MM** 1994 Selection on somatic cell score to improve resistance to mastitis in the United States. *Journal of Dairy Science* **77** 648–658
- Somers J, O'Brien B, Meany W & Kelly AL** 2003 Heterogeneity of proteolytic enzyme activities in milk samples of different somatic cell count. *Journal of Dairy Research* **70** 45–50
- Summer A, Formaggioni P, Franceschi P, Malacarne M & Mariani P** 2003 Proteose-peptone content in the milk of Italian Friesian cows with moderate and high somatic cell values. *Italian Journal of Animal Science* **2** (Suppl. 1) 266–268
- Summer A, Franceschi P, Formaggioni P & Malacarne M** 2014 Characteristics of raw milk produced by free-stall or tie-stall cattle herds in the Parmigiano-Reggiano cheese production area. *Dairy Science and Technology* **94** 581–590
- Urech E, Puhán Z & Schällibaum M** 1999 Changes in milk protein fraction as affected by subclinical mastitis. *Journal of Dairy Science* **82** 2402–2411
- Verdier-Metz I, Coulon JB & Pradel P** 2001 Relationship between milk fat and protein contents and cheese yield. *Animal Research* **50** 365–371
- Viguiet C, Arora S, Gilmartin N, Welbeck K & O'Kennedy R** 2009 Mastitis detection: current trends and future perspectives. *Trends Biotechnology* **27** 486–493
- Walsh CD, Guinee TP, Harrington D, Mehra R, Murphy J & Fitzgerald RJ** 1998a Cheesemaking, compositional and functional characteristics of low-moisture part-skim Mozzarella cheese from bovine milks containing k-casein AA, AB or BB genetic variants. *Journal of Dairy Research* **65** 307–315
- Walsh CD, Guinee TP, Reville WD, Harrington D, Murphy JJ, O'Kennedy BT & FitzGerald RJ** 1998b Influence of k-Casein genetic variant on rennet gel microstructure, Cheddar cheesemaking properties and casein micelle size. *International Dairy Journal* **8** 707–714
- Wedholm A, Larsen LB, Lindmark-Månsson H, Karlsson AH & Andrén A** 2006 Effect of protein composition on the cheese-making properties of milk from individual dairy cows. *Journal of Dairy Science* **89** 3296–3305