

University of Parma Research Repository

FGFR as potential target in the treatment of squamous non small cell lung cancer

This is the peer reviewd version of the followng article:

Original

FGFR as potential target in the treatment of squamous non small cell lung cancer / Tiseo, Marcello; Gelsomino, Francesco; Alfieri, Roberta; Cavazzoni, Andrea; Bozzetti, Cecilia; De Giorgi, Anna Maria; Petronini, Pier Giorgio; Ardizzoni, Andrea. - In: CANCER TREATMENT REVIEWS. - ISSN 0305-7372. -41:6(2015), pp. 527-539-539. [10.1016/j.ctrv.2015.04.011]

Availability: This version is available at: 11381/2794251 since: 2021-10-20T15:23:44Z

Publisher: W.B. Saunders Ltd

Published DOI:10.1016/j.ctrv.2015.04.011

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

Dear Author,

Please, note that changes made to the HTML content will be added to the article before publication, but are not reflected in this PDF.

Note also that this file should not be used for submitting corrections.

Cancer Treatment Reviews xxx (2015) xxx-xxx

Contents lists available at ScienceDirect

Cancer Treatment Reviews

journal homepage: www.elsevierhealth.com/journals/ctrv

Anti-Tumour Treatment

FGFR as potential target in the treatment of squamous non small cell lung cancer

Marcello Tiseo ^{a,*,1}, Francesco Gelsomino ^{b,1}, Roberta R. Alfieri ^{c,1}, Andrea Cavazzoni ^c, Cecilia Bozzetti ^a,
 Anna Maria De Giorgi ^a, Pier Giorgio Petronini ^c, Andrea Ardizzoni ^d

10 a Division of Medical Oncology, University Hospital of Parma, Parma, Italy

11 ^b Division of Medical Oncology, Istituto Nazionale Tumori, Milano, Italy

¹² ^c Department of Clinical and Experimental Medicine, University of Parma, Parma, Italy

13 ^d Division of Medical Oncology, Sant'Orsola-Malpighi University Hospital, Bologna, Italy

ARTICLE INFO

20 Received in revised form 12 April 2015 21 Accepted 25 April 2015

- 21 Accepted 25 April 201522 Available online xxxx
- Available online
- 23 Keywords:
- 24 NSCLC 25 Squamous cell carcing
- 5 Squamous cell carcinoma
- 26 FGF 27 FGFR
- 27 FGF 28 TKI
- 29

46

59

60

61

5

14

ABSTRACT

To date therapeutic options for squamous cell lung cancer patients remain scarce because no druggable targets have been identified so far. Aberrant signaling by FGFs (fibroblast growth factors) and FGFRs (fibroblast growth factors receptors) has been implicated in several human cancers and, particularly, in squamous non-small cell lung cancer (NSCLC).

FGFR gene amplifications, somatic missense mutations, chromosomal translocations are the most frequent mechanisms able to induce aberrant activation of this pathway. Data from literature have established that the presence of an aberrant FGFR signaling has to be considered a possible negative prognostic factor but predictive of potential sensitivity to FGFR inhibitors.

In the last years, clinical research efforts allowed to identify and evaluate promising FGFR inhibitors, such as monoclonal antibodies, ligand traps, non-selective or selective tyrosine kinase inhibitors. This review summarizes the current knowledge about FGFR alterations in NSCLC and the relative inhibitors in development, in particular in squamous NSCLC.

© 2015 Published by Elsevier Ltd.

47 Introduction

Lung cancer is classified into two main histologic types: nonsmall cell lung cancer (NSCLC) and small cell lung cancer (SCLC), accounting for 87% and 13% of all cases, respectively. The main histologic subtypes of NSCLC are adenocarcinoma (Ad-NSCLC, 50– 60%), squamous cell carcinoma (Sq-NSCLC, 30–35%) and large-cell carcinoma (LCC, 5–10%) [1].

In the last years, potentially targetable oncogene products have been recognized in approximately 60% of Ad-NSCLC [2–5]; conversely, Sq-NSCLC remains an "orphan" tumor to date; in fact, targeted agents have not yet been developed and chemotherapy continues to be the standard of care in this histotype [6].

By using high-throughput molecular technologies, ever-growing information about distinct genomic alterations in Sq-NSCLC are becoming available [7,8]. Recently, the Cancer Genome Atlas

¹ Equal contributors.

http://dx.doi.org/10.1016/j.ctrv.2015.04.011 0305-7372/© 2015 Published by Elsevier Ltd. Research Network reported an integrated analysis based on DNA copy number, exonic mutations, mRNA sequencing and expression and epigenetic alterations in 178 Sq-NSCLC samples [8]. A mean of 360 exonic mutations, 323 altered copy number segments and 165 genomic rearrangements per tumor was identified. SOX2 amplification, NFE2L2, KEAP1, discoidin domain receptor 2 (DDR2) mutations, phosphatidyl-inositol 3-kinase (PI3K) pathway changes and fibroblast growth factor receptor 1 (FGFR1) amplification, rare in Ad-NSCLC, confirmed the distinct molecular features of Sq-NSCLC. Potential druggable alterations were identified in 64% of cases.

Among them, DDR2 mutation was reported in 1–4% of Sq-NSCLC and its sensitivity to dasatinib inhibition was demonstrated both *in vitro* and *in vivo* [9,10]. Alterations in PI3K/AKT/mTOR pathway have shown to be mutually exclusive with EGFR ones (in contrast to that reported in Ad-NSCLC) and include PIK3CA mutations (3–10%) or amplification (25–40%), loss of PTEN (8–59%) or PTEN mutation (3–10%) and AKT1 or AKT2 overexpression (19% and 32%, respectively) [11]. Together with PI3K pathway, fibroblast growth factor receptor (FGFR) turns out as one of the most promising druggable target in Sq-NSCLC [12,13].

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

31

32

33

Please cite this article in press as: Tiseo M et al. FGFR as potential target in the treatment of squamous non small cell lung cancer. Cancer Treat Rev (2015), http://dx.doi.org/10.1016/j.ctrv.2015.04.011



^{*} Corresponding author at: Division of Medical Oncology, University Hospital of Parma, Via Gramsci 14, Parma 43125, Italy. Tel.: +39 0521 702316; fax: +39 0521 995448.

E-mail address: mtiseo@ao.pr.it (M. Tiseo).

M. Tiseo et al./Cancer Treatment Reviews xxx (2015) xxx-xxx

2

83 FGF-FGFR: structure, signaling and functions

84 FGFs and FGFRs

In humans 22 members of FGFR ligands (FGFs) have been 85 identified. The majority of FGFs contains a high sequence 86 87 homology region, necessary for the ligand-receptor interaction. 88 Furthermore, most of the FGFs contain a heparin sulfate proteogly-89 can (HSPG) region; the binding with HSPG protects the ligands 90 from degradation and it is also involved in the complex formation 91 between the FGFs and the FGFRs. [Fig. 1A] [14,15].

92 The FGFRs are 4 single-pass, transmembrane, tyrosine kinase 93 (TK) receptors (FGFR1-4) consisting of an extracellular portion, a transmembrane region and an intracellular domain. The extracel-94 95 lular region is composed of three immunoglobulin-like domains 96 (Ig-I, Ig-II and Ig-III). Ig-I is responsible for binding affinity; Ig-II 97 constitutes the binding site for FGF and the latter is crucial for 98 the ligand binding selectivity [Fig. 1B]. Due to alternative splicing 99 in Ig-III domain of FGFR1-3, there are several isoforms with different FGF-binding specificity; among them, FGFR IIIb and IIIc are 100 epithelial and mesenchymal isoforms, respectively [16]. The extra-101 102 cellular domain contains also the acidic box necessary for the 103 interaction between HSPGs and FGFRs, the HSPG binding domain 104 and another region responsible for the interaction with cellular 105 adhesion molecules and extracellular matrix. The transmembrane domain acts as stabilizer of receptor conformation, essential for 106 107 its ligand-dependent activation. The juxtamembrane region con-108 tains the binding sites for signaling effectors/modulators, such as 109 PKC and FRS2α. Finally, the TK domains have the catalytic activity 110 and recruit specific effectors able to activate different downstream 111 signaling pathways [14,15]. In addition, there is a fifth FGFR 112 (FGFRL1) without TK activity [17].

113 FGF/FGFR signaling and functions

Upon ligand-receptor binding, FGFR dimerizes and, in turn, it 114 115 phosphorylates FRS2a, leading to GRB2 recruitment and, ultimately, to the activation of different pathways. The main down-116 stream signaling pathways are RAS/MAPK and PI3K/AKT/mTOR 117 cascades. The activation of these signaling pathways, together with 118 119 PLC γ /Ca²⁺, promotes cell survival, motility and invasiveness, cell 120 proliferation, epithelial-to-mesenchymal transition (EMT), angio-121 genesis [Fig. 2]. After activation, the receptor complex is internal-122 ized by endocytosis, degraded by lysosomes or inactivated by 123 polyubiquitination process [18].

FGFs and FGFRs are expressed in several normal cells and tis-124 125 sues of epithelial and mesenchymal origin. In embryonic life, they have a crucial role in angiogenesis, formation and development of 126 different organs and specific FGFR2 and FGFR3 point mutations 127 128 have been associated with congenital skeletal disorders [19–21]. In adults, FGF/FGFR signaling mediates tissue homeostasis and 129 130 plays a fundamental role in inflammation processes, in neoangiogenesis and vessel maturation during wound healing and tissue 131 repair [14,18,22,23]. 132

FGFR alterations in NSCLC 133

134 In human cancers, FGF/FGFR signaling can be aberrantly acti-135 vated as a result of ligand-dependent or ligand-independent mech-136 anisms. FGFR gene amplifications, somatic missense mutations and 137 chromosomal translocations, leading to receptor overexpression 138 and/or constitutive FGFR or FGFR-TK activation; alternative splic-139 ing, leading to altered ligand-receptor specificity; paracrine/au-140 tocrine signaling, due to FGF upregulation, and FGFR germline 141 single nucleotide polymorphisms (SNPs) are the most frequent

mechanisms of activation [24]. Table 1 and Fig. 3 summarize the 142 principal FGF/FGFR alterations in NSCLC cancers. In particular, in 143 Sq-NSCLC, FGF/FGFR pathway has been recognized as one of the hallmark alterations with relevant clinical implications.

FGFR gene amplifications

The amplification of FGFR1 gene, located on chromosome 8p12, is a common potentially druggable alteration in NSCLC. Several studies have demonstrated that FGFR1 gene amplification incidence is significantly higher in Sq-NSCLC (22%) than in Ad-NSCLC (3%) [25–33]. Moreover, FGFR1 amplification tends to correlate with smoking status, with higher frequency in current smokers than in former and never-smokers [25,32]. No other specific clinico-demographic features correlate with FGFR1 amplification.

Recently, several groups have investigated the potential role of FGFR1 gene amplification as prognostic factor in resected NSCLC patients [29,31,32,34]. Kim and colleagues [32] showed that resected patients whose tumors harbored high FGFR1 gene amplification (34/262, 13%) had significant shorter disease-free survival (DFS) and overall survival (OS) than those without FGFR1 gene amplification (26.9 vs. 94.6 months, p < 0.001; 51.2 vs. 115.0 months, p = 0.002, respectively). A multivariate analysis confirmed FGFR1 gene amplification as an independent negative prognostic factor, as well as other studies [31,34,35], whereas a Canadian report did not [29]. A meta-analysis of 13 published studies in Sq-NSCLC showed a FGFR1 amplification rate of 19%, significantly correlated with smoking status and lymph node metastasis, not able to influence OS [25].

More recently, it has been shown that tumorigenic potential of FGFR1 amplification depends on multiple factors. A recent study demonstrated a genomic heterogeneity of FGFR1-amplified Sq-NSCLC that could affect its sensitivity to FGFR inhibition. It has been shown that the 8p12 (including FGFR1) amplicon clustered together with the 11q13 (including CCND1, FGF4 and FGF19). This co-occurred event seems to be related to a higher sensitivity to FGFR inhibition. Furthermore, the co-expression of high levels of c-MYC, as well as FGFs, induced in FGFR1-amplified cells higher oncogenic transformation, cell-autonomous signaling and higher FGFR inhibitor sensitivity both in vitro and in vivo [30]. Hence, these data suggest an evidence to refine patients who will be likely benefit from FGFR inhibitor treatment.

FGFR gene mutations

Somatic FGFR mutations in lung tumors occur at the same positions to germline FGFR2 and FGFR3 mutations and may contribute to lung cancer development [36]. No correlation between FGFR mutations and histology subtype has been reported.

Recently, through a whole-genome sequencing platform, 210 human cancers were screened and more than one thousand somatic mutations were identified. In this cohort, 88 lung tumors were included and the highest prevalence (4.21 per Mb) of somatic mutations was seen among them. Overall, FGFR mutations were detected in 10% of lung tumor samples. FGFR2 mutations (3.4% in lung tumors) showed a high selection pressure with an excess of non-synonymous mutations, providing evidence for its role as oncogenic driver [37].

Another study evaluated FGFR2 and FGFR3 mutations in Sq-NSCLC. Overall, 12 (6.7%) mutations in the FGFR2 (6 cases) and FGFR3 (6 cases) genes were detected. The presence of specific mutations in the extracellular domains of FGFR2 (W290C and S320C) and FGFR3 (R248C and S249C) was responsible for cellular transformation, as did K660E and K660N mutations in the kinase domains of FGFR2, both in vitro and in xenograft models [38]. It is noteworthy that W290C, S320C and K660N mutations showed

144 145

146

147

148

149

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

179

Please cite this article in press as: Tiseo M et al. FGFR as potential target in the treatment of squamous non small cell lung cancer. Cancer Treat Rev (2015), http://dx.doi.org/10.1016/j.ctrv.2015.04.011





Fig. 1. Structure of the majority of FGFs (A) and FGF receptors (B). (A) FGF is composed of a signal peptide (SP), an amino-terminal coil, a conserved core (a high-sequence homology region), including both FGFR- and heparin (HPSG)-binding sites, and the carboxy-terminal coil. (B) FGF receptor includes three immunoglobulin (Ig) domains, an acidic box, a trans-membrane (TM) domain, a juxta-membrane (JM) region, and two tyrosine kinase (TK) domains.



Fig. 2. The FGF/FGFR signaling cascade. The FGF/FGFR network regulates cell survival, proliferation and motility. The main downstream signaling pathways are RAS/MAPK and PI3K/AKT/mTOR cascades.

210

211

204

different oncogenic potential in xenograft models while other FGFR2 (E471Q and T787K) or FGFR3 (S435C and K717M) mutations were not oncogenic when tested in an anchorage-independent growth assay. However, *in vivo* oncogenic activity of many other variants has not been validated so far, and further studies are needed. In any case, the sensitivity to pan-FGFR and multi-kinase inhibitors provided preclinical evidence for their role as potential oncogenic driver and druggable targets [38]. Sporadic cases of non-synonymous FGFR4 mutations (P672T and E681K) in current smoker Ad-NSCLC patients have also been reported [36,39].

FGFR gene fusions

FGFR1/3 fusions occurred in about 1% of patients with NSCLC and 2–3.5% of patients with Sq-NSCLC [40,41].

By using an integrative whole-exome and transcriptome sequencing platform, Wu et al. [40], isolated from Sq-NSCLCs six rearrangements (1 BAG4-FGFR1, 1 FGFR2-KIAA1967 and 4 FGFR3-TACC3) biologically active and sensitive to FGFR inhibitors, such as PD173074 and pazopanib. It was noteworthy that FGFR3 gene fusion-positive tumors had enhanced susceptibility to FGFR inhibition over FGFR3-mutant cell lines [40].

Furthermore, it has been shown that compared with the FGFR fusion-negative group, patients with FGFR fusions were more likely to be smokers, significantly associated with larger tumor and with a tendency to be more poorly differentiated. No difference in terms of progression free survival (PFS) and OS was seen between FGFR1/3 fusion positive and negative tumor patients [42].

FGFR3-TACC3 fusion transcripts were also detected in 0.5% of Ad-NSCLC and predominantly in never/light smoker patients [41].

Table 1

Principal genomic FGFR alterations in NSCLC.

_					
	Tumor type	Gene	Alteration	Incidence (%)	Refs.
	Sq-NSCLC	FGFR1 FGFR1 FGFR2 FGFR3 FGFR4 FGFR3-TACC3 BAG4-FGFR1	Amplification Mutation Mutation Mutation Translocation Translocation	11-22 2 3-5 2-3 2 1-3.5 0.3-0.6	[25–33] [37] [37,38] [37,38] [37] [40,42] [40,42]
		FGFR2-KIAA1967	Translocation	0.3	[40]
	Ad-NSCLC	FGFR1 FGFR3-TACC3	Amplification Translocation	2.2–3.4 0.5	[31,35] [41,42]

Sq-NSCLC, squamous non small cell lung cancer; Ad-NSCLC, adenocarcinoma non small cell lung cancer.

Please cite this article in press as: Tiseo M et al. FGFR as potential target in the treatment of squamous non small cell lung cancer. Cancer Treat Rev (2015), http://dx.doi.org/10.1016/j.ctrv.2015.04.011

214

212

213

215 216

217

223

224

225

226

227

228

229

230

4

M. Tiseo et al./Cancer Treatment Reviews xxx (2015) xxx-xxx



Fig. 3. FGF-FGFR alterations. Potential targetable FGF/FGFR biomarkers include overexpression of FGF family members, FGFR1 overexpression (by gene amplification or increased mRNA FGFR1 transcription), FGFR2 mutations in the extracellular and in the kinase domain (W290C, S320C and K660E/N respectively), FGFR3 mutation in the extracellular domain (R248C, S249C) and FGFR gene fusions (BAG4-FGFR1, FGFR2-KIA1967, FGFR3-TACC3).

232 Autocrine/paracrine signaling

Multiple FGFs, such as FGF2 (also called basic FGF, b-FGF) and 233 FGF9, are upregulated in NSCLC cell lines and involved in EMT 234 235 and cancer progression. Furthermore, the existence of an additive synergism between FGFs and other angiogenic pathways, such as 236 237 VEGFR and PDGFR, has been shown [43]. The use of a FGF2-specific 238 shRNA or a multiple anti-angiogenic TK inhibitor blocked cell proliferation and tumor growth by reduction of FGF2 expression [43]. 239 240 Finally, high levels of FGF2 were predictive of resistance to EGFR-TK inhibitors [43], as also of resistance to VEGF and VEGFR-TK inhi-241 bitors in colorectal [44] and glioblastoma [45] models, 242 243 respectively.

244 Different studies have evaluated b-FGF as prognostic factor in 245 NSCLC [46–53]; some of them reported a significant correlation 246 between tumor b-FGF expression and poor outcome [46,49]; con-247 versely, others showed opposite results [47,50].

FGFR germline polymorphisms and new insights 248

Genetic variations, such as germline SNPs, have been associated 249 with enhanced tumor risk. A correlation between specific variants 250 251 in the FGFR2 gene and increased susceptibility to breast cancer has been confirmed [54]. Conversely, a specific polymorphism in the 252 253 FGFR4 gene (Gly388Arg) has been associated with tumor progression rather than tumor risk in different cancer types [55,56], 254 255 including also lung adenocarcinoma [57]. The analysis of the 256 Gly388Arg polymorphism has revealed a significant association 257 with an earlier age at cancer onset (p = 0.002), advanced stage (p = 0.002) and poor survival (p = 0.007) [57]. 258

More recently, a large-scale RNAi-based mouse tumorigenesis 259 260 screen identified twenty-four tumor suppressor genes (TSGs) that 261 were significantly down-regulated in human Sq-NSCLCs. A part 262 of these TSGs encoded repressors of FGFR signaling. The downregulation of these suppressors led to activation of FGFR signaling, 263 as a consequence of increased levels of phosphorylated FRS2-Y436, 264 and rendered in most cases tumor cells sensitive to FGFR inhibi-265 tors. These data support the hypothesis that aberrant FGFR signal-266 ing pathway may drive tumorigenesis in human Sq-NSCLC even in 267 the absence of FGFR amplification, translocation or activating 268 mutation [58]. 269

FGF/FGFR pathway as predictive factor

Several studies have identified FGFR1 amplification as the major 271 predictive factor of response to FGFR inhibitors [26,27]. However, 272 in a recent study on lung cancer cell lines with ponatinib, FGFR1 273 mRNA and protein expression, together with FGF2 and FGF9 274 mRNA, were better predictive biomarkers of sensitivity than 275 FGFR1 gene copy number [59]. These data were evaluated also in 276 resected lung tumors. Approximately 50% of Sq-NSCLC with high 277 FGFR1 gene copy number expressed high FGFR1 mRNA levels. At 278 once, increased FGFR1 gene copy number included a significant 279 fraction of Sq-NSCLC not expressing FGFR1 mRNA or protein 280 (18%), providing evidence for considering that FGFR1 amplification 281 could be occurred in both sensitive and resistant cell lines [59]. 282

Furthermore, hot topics on the role of FGF/FGFR pathway as 283 predictor of resistance to other targeted agents are emerging. 284 FGFR signaling seems to be involved in tumor growth by an auto-285 crine mechanism and may be considered a specific pathway of 286 insensitivity to EGFR-TK inhibitors. FGF2 and FGF9 mRNA levels 287 were significantly higher in gefitinib-insensitive cells and the use 288 of FGF2-specific shRNAs or FGFR TK inhibitors induced cell growth 289 arrest [43,60,61]. Furthermore, also FGFR2 and FGFR3 were de-re-290 pressed after gefitinib exposure, leading to a rapid mechanism of 291 acquired resistance to EGFR-TK inhibitor [62]. A treatment with 292 gefitinib and AZD4547 allowed to prevent the outgrowth of drug 293 resistant cell clones, supporting the use of a combinatorial therapy 294

270

Please cite this article in press as: Tiseo M et al. FGFR as potential target in the treatment of squamous non small cell lung cancer. Cancer Treat Rev (2015), http://dx.doi.org/10.1016/j.ctrv.2015.04.011

Please cite this article in press as: Tiseo M et al. FGFR as potential target in the treatment of squamous non small cell lung cancer. Cancer Treat Rev (2015),

295 to prevent or delay acquired resistance in EGFR-addicted NSCLC 296 [63]. Another study showed that the concomitant use of FGFR-TK 297 inhibitors (SU5402 or PD166866) and EGFR-TK inhibitors (erlotinib 298 or lapatinib) was responsible for a markedly reduced cell proliferation over FGFR-TK inhibitor alone. Conversely, the concomitant 299 administration of FGFR inhibitors and chemotherapeutics was 300 301 detrimental [64]. Unfortunately, when this approach was applied 302 in clinical trials it failed because of toxicity (see phase I trial, NCT01515969, combining erlotinib and dovitinib in advanced 303 NSCLC). 304

More recently, a potential role of FGFR3 as predictive marker for MET-targeted therapy has been reported. FGFR3 seems to be highly expressed in specific MET-amplified cells and its suppression enhanced the antiproliferative effects of a MET-monoclonal antibody, providing evidence for the presence of a specific crosstalk between FGFR and MET pathways [65].

Furthermore, a growing body of research have suggested that FGF/FGFR signaling pathway may also play a role in the resistance to anti-VEGF therapy after the evidence of high FGF2 levels in different cell models at the time of disease progression [22].

315 From preclinical evidences to clinical data

The deregulation of FGF/FGFR pathway can be considered as a mechanism of oncogene-addiction and, in some cases, oncogeneexpedience. Therefore, different treatment strategies including the administration of a FGFR inhibitor alone or in combination with other target drugs or chemotherapy (concomitantly or sequentially) have to be investigated.

Several preclinical data have shown that selective FGFR-TK inhibitors [26,27], multi-TK inhibitors [38,40,41,43], FGFR1-specific
 shRNAs [27] or FGF2-specific shRNAs [43] were capable of blocking
 tumor growth in different FGFR-addicted NSCLC cell lines.

In the last years, clinical research efforts allowed to identify and
 evaluate promising FGFR inhibitors and some of them are under
 development in clinical trials.

To block FGF/FGFR pathway, different targeted molecules have been developed: (1) monoclonal antibodies (mAbs); (2) ligand traps; (3) non-selective TK inhibitors and (4) selective TK inhibitors. Ongoing clinical trials in lung cancer are reported in Table 2.

333 FGFR mAbs

The clinical development of several mAbs directed against specific FGFRs may be an attractive strategy for treating different FGFR-addicted solid tumors because of their high selectivity and low toxicity when compared with pan-FGFR inhibitors.

338 MFGR1877S is the most promising agent of this class. It specifi-339 cally acts on isoforms IIIb/IIIc of FGFR3. A phase I dose escalation 340 study showed that MFGR1877S was well tolerated in 26 patients with advanced solid malignancies. A maximum tolerated dose 341 (MTD) was not reached. There was a dose limiting toxicity (DLT) 342 343 of grade 4 thrombocytopenia and recommended phase 2 dose 344 (RP2D) was set at 30 mg/kg. The most frequent adverse events 345 (AEs) were nausea and fatigue. In terms of efficacy, 9 out of 26 346 patients (35%) experienced a disease stabilization [66]. No clinical 347 trial of MFGR1877S is currently ongoing in lung cancer.

348 FGF-ligand traps

This class of compounds includes some specific proteins (FGFtraps) that are capable of blocking FGF-FGFR binding, by sequestering endogenous FGF ligands released by tumor cells or tumor stroma.

http://dx.doi.org/10.1016/j.ctrv.2015.04.011

FP-1039/CSK3052230 is the first-in-class ligand trap, currently investigated in phase I/II clinical trials. It is a soluble decoy receptor containing the extracellular domains of human FGFR1c isoform linked to the Fc regions of human immunoglobulin G1.

In preclinical models, this compound demonstrated its activity by inhibiting tumor growth both *in vitro* and in xenograft models of several tumor types, including NSCLC [67].

A phase I dose escalation study reported preliminary data on safety. This study enrolled 33 patients with advanced solid tumors, including 3 with lung cancer. The main AEs were gastrointestinal disorders, fatigue, and peripheral edema. No DLTs were observed in the 15 patients dosed from 2.0 to 16.0 mg/kg and the maximum feasible dose (MFD) was set at 20 mg/kg weekly. Pharmacokinetics (PKs) supported weekly dosing and pharmacodynamics demonstrated a significant decrease of serum FGF2 levels during treatment period. Disease control rate (DCR) was 42% [68]. This compound is currently being evaluated in a three-arm, non-randomized, open-label phase Ib trial. The objective of this study is to evaluate safety and efficacy of FP1039 in combination with chemotherapy in FGFR1-amplified Sq-NSCLC and in malignant pleural mesothelioma [NCT01868022] [Table 2].

Non-selective FGFR-TK inhibitors

This class of compounds includes several ATP-competitive small molecules directed against FGFR TK domain, but also other structurally related tyrosine kinase domains, including VEGFR, RET, PDGFR, KIT, FLT3, BCR-ABL. These multitarget inhibitors have demonstrated to be active in the context of FGFR-addicted tumors but, because of the lack of kinase selectivity, their use is limited by a variety of side effects. Among them, the molecules at more advanced clinical development are dovitinib, nintedanib, cediranib, ponatinib, lucitanib and pazopanib. Other compounds, such as brivanib, lenvatinib, orantinib are under investigation in other tumor types and have demonstrated to be more active against VEGFR or other TKs than against FGFRs. The IC₅₀ versus the different targets are shown in Table 3. The non-selective FGFR TK inhibitors have a specific toxicity profile primarily dependent on VEGF/VEGFR inhibition. Treatment with the VEGF inhibitors, such as bevacizumab, has shown a significant higher risk of cardiovascular complications and bleeding, particularly in Sq-NSCLC patients [69]. This risk led to approval bevacizumab in combination with chemotherapy as first-line treatment in NSCLC, except squamous histology. For these reasons, exclusion criteria in ongoing trials, particularly with nonselective FGFR TK inhibitors, do not allow to enroll patients with a history of clinically significant bleeding disorder or baseline hemoptysis, therapeutic anticoagulation and central tumors with cavitation.

Dovitinib (TKI258) is an inhibitor of VEGFR1-3, FGFR1/3, FLT3, KIT, RET and PDGFR-β. It has a potent anti-angiogenic activity through the inhibition of PDGFR, VEGFR and FGFR [70,71]. In a phase I dose escalation study enrolling 35 patients with advanced solid tumors a partial response (PR) and 2 long stable disease (SD) with a tolerable toxicity profile were observed. The most frequent AEs were diarrhea and fatigue; 14% of patients experienced toxicities related to VEGFR pathway inhibition, such as hypertension and left ventricular ejection fraction decrease [72]. Currently, this compound is being evaluated in different solid malignancies both in phase II and III trials [73,74]. A phase I trial enrolling advanced NSCLC patients for a combined therapy with dovitinib and erlotinib was suspended because of toxicity data at the interim analysis. Two phase II studies are currently evaluating dovitinib as monotherapy in NSCLC [NCT01676714] and in FGFR1-amplified Sq-NSCLC patients [NCT01861197] [Table 2].

Nintedanib (BIBF1120) is a multitarget small tyrosine kinase inhibitor directed against FGFR1–4, VEGFR1–3, PDGFRA-B and

y n

5

364

365

353

354 355

356

357

358

359

366 367 368

369

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

|--|

Table 2Ongoing trials on FGFR inhibitors in NSCLC.

Drug	Target (s)	Tumor type (s)	Inclusion criteria	Phase	Arm (s)	Objectives	Status
FGF traps FP-1039/GSK3052230 Non-selective FCFR inhib	FGF	Metastatic recurrent Sq-NSCLC with FGFR1 amplification and recurrent/unresectable MPM [NCT01868022]	Sq-NSCLC • FGFR1-ampl • 1st line (Arm A) • 2nd line (Arm B) MPM • FGF2 expression • 1st line (Arm C) • ECOG PS 0-1 (Arm A) • ECOG PS 0-2 (Arms B, C)	Ib	A. FP1039 5 mg/kg i.v. weekly + paclitaxel 200 mg/mq and CBDCA AUC 6 d1 q21 B. FP1039 5 mg/kg i.v. weekly + docetaxel 75 mg/mq d1 q21 C. FP1039 5 mg/kg i.v. weekly + pemetrexed 500 mg/ mq and CDDP 75 mg/mq d1 q21	AES, DLTS, MTD, ORR; PFS, PK	Currently recruiting
Dovitinib/TKI258	VEGFR1–3, FGFR1/3, FLT3, KIT, RET, PDGFRB	Advanced NSCLC or colorectal cancer (CRC) previously treated with anti-VEGF therapy [NCT01676714]	 Immediate prior treatment regimen with a anti-VEGF agent (bevacizumab, sunitinib, sorafenib) No potentially curative treatment options 	Ш	Dovitinib 500 mg once daily for 5 continuous day weekly	ORR; DCR, PFS, toxicities	Currently recruiting
		Previously treated FGFR1-amplified Sq-NSCLC patients [NCT01861197]	 Sq-NSCLC 1-2 prior therapies FGFR-ampl (FISH > 5 gene copies) 	II	Dovitinib 500 mg daily for 5 days on/2 day off	ORR	Currently recruiting
Nintedanib/BIBF1120	VEGFR1–3 PDGFRα-β FGFR1–3	Phase I dose escalation trial in elderly patients with stage IV NSCLC [NCT01684111]	 Age > 70 years ECOG PS < 2 No previous therapies 	Ι	Nintedanib 150 mg BID (starting dose level) + Vinorelbine25 mg/ mq i.v. dd1,8 q21	MTD; AEs, PK, ORR	Currently recruiting
		Phase I safety run-in trial in Japanese patients with advanced/metastatic Ad-NSCLC [NCT02300298]	 Ad-NSCLC Japanese population Prior 1st line platinum-based ECOG PS < 1 	Ι	Nintedanib 200 mg BID q28 + docetaxel 75 mg/mq iv q21	DLTs cycle 1; AEs, PK	Currently recruiting
		First-line treatment in Sq-NSCLC [NCT01346540]	 Stage IIIB/IV or recurrent Sq-NSCLC ECOG PS ≤ 1 No PRO after 2 cycles of GEM/CDDP No previous therapy with other antivascular drugs 	1/11	Phase I (open label): nintedanib + gemcitabine and cisplatin Phase II: ≥SD after 2 courses of gemcitabine + cisplatin: A. Nintedanib + gemcitabine and cisplatin B. Placebo + gemcitabine and cisplatin	MTD (Phase I), PFS (Phase II); ORR, OS, AEs	Ongoing, not recruiting
		Recurrent FGFR1-amplified and wild-type Sq- NSCLC after failure of 1 or 2 chemotherapy regimens [NCT01948141]	 Sq-NSCLC Prior 2 chemoterapies ECOG PS ≤ 1 FGFR1 ampl/non ampl 	ΙΙ	Nintedanib 200 mg BID q28	6-month PFS in FRGF1-amplified tumor group; 6-month PFS in FGFR1-wild type and FGFR1- amplified groups; PFS, OS, ORR by FGFR1 group; FGFR polymorphism, toxicity and efficacy	Currently recruiting
Ponatinib/AP24534	Pan-FGFR PDGFRA/B VEGFR1/3 RET	Locally-advanced or metastatic lung cancer preselected using different candidate predictive biomarkers [NCT01935336]	 Any histology (except carcinoid) NSCLC (IIIA-IV) SCLC (LD-ED) EGFR wild type No ALK fusions FGFR1 [SISH+/ISH+, SISH+/ISH-, SISH-/ISH+, SISH-/ISH-] RET FISH positive ECOG PS ≤ 2 	ΙΙ	Ponatinib 45 mg orally once or twice daily q28	Biomarkers, ORR; AEs	Currently recruiting
		Open-label study in advanced NSCLC with RET translocations [NCT01813734]	 RET translocation No restriction on number of prior therapies 	II	Ponatinib 45 mg orally once a day q28	ORR; DCR, PFS, 1-year OS rate, AEs, safety and tolerability	Currently recruiting

Drug	Target (s)	Tumor type (s)	Inclusion criteria	Phase	Arm (s)	Objectives	Status
Lucitanib/E3810	FGFR1/2 VEGFR1-3	FGFR1-amplified Sq-NSCLC [NCT02109016]	 Sq-NSCLC FGFR1 amplification At least one prior therapy ECOG PS ≤ 1 	II	Lucitanib 15 mg once daily q28	ORR; CBR, PFS, DOR, OS, AEs, PK, PD	Currently recruitin
Pazopanib/GW786034	FGFR1/3 VEGFR PDGFRc-KIT	First-line treatment for advanced NSCLC patients not eligible for front-line therapy with a platinum doublet [NCT01179269]	 No previous therapy ECOG PS ≤ 2 Unfit for doublet-chemotherapy regimen 	Π	Pazopanib daily and weekly paclitaxel i.v.	ORR	Currentl recruitin
Selective FGFR TK inhibit	ors						
AZD4547	FGFR1-3	Recurrent Sq-NSCLC [NCT01824901]	 Sq-NSCLC 2nd line treatment ECOG PS ≤ 1 FGFR1-ampl (Phase II) 	I/II	Phase I: docetaxel d1 + AZD4547 d2-15 q21 Phase II: A. Docetaxel d1 q21 \times 6 \rightarrow PRO \rightarrow AZD4547 d1- 14 q21 B. Docetaxel d1 q21 \times 6 + AZD4547 d1-14 q21 \times 6 + AZD4547 d1-14 q21 \rightarrow AZD4547 d1-14	RP2D (Ph I), PFS (Ph II); PK, safety/toxicity, clinical activity, ORR, OS	Ongoing not recruitir
		FGFR1-amplified mBC, advanced Sq-NSCLC and FGFR2-amplified gastric cancer progressed following previous chemotherapy [NCT01795768]	Sq-NSCLC • FGFR1-ampl • 1-2 previous therapies Breast • FGFR1-ampl • Her2 negative • At least 1 prior hormone therapy • 1-3 previous chemotherapy Gastro-oesophageal	Π	AZD4547 80 mg orally twice daily 2 weeks on/1 week off q21	Antitumor activity; ORR, safety, DCR, PFS	Currentl recruitir
		Metastatic NSCLC according to genomic profiles, including FGFR [NCT02117167]	 FGFR2-allpl 1-2 previous chemotherapy ECOG PS ≤ 1 Squamous Non-squamous No previous chemotherapy Suitable for 1st line platinumbased ECOG PS ≤ 1 	П	Induction platinum-based chemotherapy for 4 cycles with SD or PR → Maintenance: A. AZD4547 (FGFR) 80 mg BID; AZD2014 (mTOR) 50 mg BID; AZD5363 (AKT) 480 mg BID	PFS; OS, ORR, toxicities	Currentl recruitir
			 No PRO after current line EGFR wild type No ALK fusions Targeted therapy according to specific genomic profile (FGFR, mTOR, AKT, HER2/EGFR, MEK, VEGF/EGFR) 		4 days on/3 days off; AZD8931 (HER2, EGFR) 40 mg BlD; Selumetinib (MEK) 75 mg BlD; Vandetanib (VEGF, EGFR) 300 mg B. Pemetrexed 500 mg/mq eV d1 q21 (NSq-NSCLC); Erlotinib 150 mg (Sq-NSCLC)		
		Recurrent IIIB/IV Sq-NSCLC [NCT02154490]	 Sq-NSCLC 2nd line treatment ECOG PS ≤ 2 EGFR wild type No ALK fusions No drug-biomarkers (Arm A) PIK3CA positive (Arm B) CDK4/6, CCND1, CCND2 and CCND3 positive (Arm C) FGFR1-3 positive (Arm D) HGF-cMET positive (Arm E) 	11/111	A. MEDI4736 (anti-B7H1) eV d1 q14 vs docetaxel d1 q21 B. GDC-0032 once daily q21 vs docetaxel d1 q21 C. Palbociclib once daily d1–21 q28 vs Docetaxel d1 q21 D. AZD4547 BID d1–21 vs docetaxel d1 q21 E. Rilotumumab eV d1 + erlotinib once daily q21 vs erlotinib once daily q21	PFS (Ph II), OS (Ph III); ORR, toxicities	Currentl recruitir
					• •		

No. of Pages 13, Model 5G

 $\overline{}$

4 May 2015 **YCTRV 1402**

Drug	Target (s)	Tumor type (s)	Inclusion criteria	Phase	Arm (s)	Objectives	Status
BGJ398	Pan-FGFR	Advanced solid tumors with FGFR1/2 amplifications or FGFR3 mutations [NCT01004224]	 FGFR1-3 gene alterations FGFR1-ampl (Sq-NSCLC) FGFR3 mutation/fusion (bladder 	I	BGJ398 orally	MTD, RP2D, schedule; safety, tolerability, PK, PD, preliminary activity	Currently recruiting
		Advanced solid tumors with FGFR alteration in Asian patients [NCT01697605]	 FGFR gene alteration Asian ethnicity ECOG PS ≤ 2 	Ι	BGJ398 orally once daily or twice daily	DLT; AEs, ORR, PFS	Currently recruitin
		Advanced solid tumors (ASTs) expressing PIK3CA mutations with or without FGFR alterations [NCT01928459]	 PIK3CA mutations (dose escalation + expansion) FGFR gene alteration (expansion cohort) No CRC (expansion cohort) ECOG PS ≤ 2 	Ιb	BGJ398 orally once daily d1–21 q28 + BYL719 orally once daily die 1–28 Dose escalation part: A. ASTs PIK3CA mut Expansion part: B. mBC (PIK3CA mut + FGFR pos) C. ASTs (PIK3CA mut) D. ASTs (PIK3CA mut + FGFR pos)	DLT; safety and tolerability, ORR, PFS, PK	Currentl recruitir
		Advanced solid tumors, including NSCLC, and haematological malignancies with FGFR genetic alterations [NCT02160041]	 FGFR gene alteration ECOG PS ≤ 1 	II	BGJ398 125 mg once daily d1–21 q28	CBR; ORR, PFS, OS, DOR, safety and tolerability	Currentl recruitin
BAY1163877	Pan-FGFR	Advanced solid tumors (dose escalation), including Ad-NSCLC and Sq-NSCLC according to FGFR profile [NCT01976741]	 Ad-NSCLC Sq-NSCLC High FGFR expression FGFR mutation 	Ι	BAY1163877 BID	MTD, PK, PD, safety, tolerability; tumor response	Currentl recruitin
JNJ-42756493	Pan-FGFR	Advanced solid tumors, including NSCLC, or lymphoma [NCT01703481]	 Pre- and post-treatment biopsies Sq-NSCLC ECOG PS ≤ 1 	Ι	JNJ-42756493 once daily q21 Part 1: ASTs or lymphoma Part 2: ASTs Part 3: •NSCLC (cohort A) •SCLC (cohort B) •mBC (cohort C) •ASTs (cohort D)	MTD; PK, PD, ORR, AEs, DOR, PFS	Currentl recruitir
GSK3052230	FGFR1	Advanced solid tumors and deregulated FGF pathway signaling [NCT01868022]	Sq-NSCLC • FGFR1-ampl • 1st line (Arm A) • 2nd line (Arm B) MPM • FGF2 expression • 1st line (Arm C) • ECOG PS ≤ 1 (Arm A) • ECOG PS ≤ 2 (Arms B C)	lb	A. GSK3052230 eV d1,8,15 + paclitaxel d1 + CBDCA d1 q21 B. GSK3052230 eV d1,8,15 + docetaxel d1 q21 C. GSK3052230 eV d1,8,15 + pemetrexed d1 + cisplatin d1 q21	AEs, safety and tolerability, DLT, MTD, ORR; PFS, PK	Currentl recruitin

RTICLE

z

PRESS

Table 2 (continued)

ø

9

M. Tiseo et al./Cancer Treatment Reviews xxx (2015) xxx-xxx

Table	3
-------	---

Targets and IC₅₀ values (nM).

	FGFR1	FGFR2	FGFR3	FGFR4	VEGFR1	VEGFR2	VEGFR3	PDGFR-a	PDGFR-β
Non-selective FGFR-TK inhibitors Dovitinib/TKI258									
Ref. [70]*	8	-	9	-	10	13	8	200	27
Ref. [71]*	16	50	53	341					
Nintedanib/BIBF1	120								
Ref. [75]*	69	37	108	610	34	21	13	59	65
Ref. [71]*	47	63	122	451					
Cediranib/AZD217	71								
Ref. [81]*	26	-	-	-	5.0	1.0	3.0	36	5
Ref. [71]*	5	33	36	697					
Ponatinib/AP2453	4								
Ref. [86]*	2.2	1.6	18.2	7.7	3.7	1.5	2.3	1.1	7.7
Ref. [71]*	2	2	18	8					
Lucitanib/E3810									
Ref. [88]*	17	82	237	>1000	7	25	10	175	525
Ref. [89]**	7	-	-	-	12	4	-	13	8
Pazopanib/GW78	5034								
Ref. [91]*	140	-	130	800	10	30	47	71	84
Selective FGFR-TK inhibitors									
Ref. [94]*	2.8	2.6	6.4	6.0	-	7		-	-
AZD4547 Ref. [95]*	0.2	2.5	1.8	165	-	24	-	_	_
BGI398									
Ref. [100]*	0.9	1.4	1	60	-		-	-	-

Please cite this article in press as: Tiseo M et al. FGFR as potential target in the treatment of squamous non small cell lung cancer. Cancer Treat Rev (2015),

* Biochemical kinase inhibition assay.

** Elisa assay.

FLT-3 [71,75]. Phase I studies evaluated this agent as single agent
[76], in combination with pemetrexed in pretreated NSCLC
patients [77] and with carboplatin and paclitaxel as first-line treatment of NSCLC [78], showing gastrointestinal disorders as the most
frequent AEs and a response rate of 5–27%.

A phase I/II study evaluating nintedanib with cisplatin-gemcitabine as first-line treatment in advanced Sq-NSCLC patients is ongoing [NCT01346540] [Table 2]. A phase II trial confirmed its promising anti-tumor activity as monotherapy in 73 previously treated advanced NSCLC patients, with 1 PR and 35 SD and median PFS and OS of 6.9 and 21.9 weeks, respectively. The toxicity profile was similar to that seen in phase I trials [79].

Considering its activity and tolerable toxicity profile, nintedanib 429 430 has been evaluated in two phase III trials (LUME-Lung 1 and 2) per-431 formed in advanced NSCLC patients after failure of first-line stan-432 dard treatment. LUME-Lung 1 was designed to assess the efficacy 433 and safety of docetaxel plus nintedanib as second-line treatment with PFS as primary end-point both in adenocarcinoma and in 434 435 intent-to-treat (ITT) population. Patients were assigned to receive docetaxel plus nintedanib (n = 655) or plus placebo (n = 659). 436 Nintedanib significantly improved PFS in ITT (3.4 vs. 2.7 months; 437 p = 0.0019), adenocarcinoma (p = 0.0193) and in squamous 438 (p = 0.02) populations. A significant improvement in OS was seen 439 in adenocarcinoma group (12.6 vs. 10.3 months; p = 0.0359), but 440 not in squamous one (8.6 vs. 8.7 months; p = 0.8907). The most 441 442 common AEs in the experimental arm were diarrhea (42.3% vs. 21.8%), increased alanine aminotransferase (ALT) (28.5% vs. 8.4%) 443 444 and aspartate aminotransferase (AST) (22.5% vs. 6.6%) [80].

LUME-Lung 2 trial enrolled only pretreated patients with nonsquamous histology and was stopped after an interim futility analysis. Considering the LUME-Lung 1 results, EMA approved nintedanib in combination with docetaxel for the treatment of locally
advanced, metastatic or locally recurrent NSCLC of only adenocarcinoma histology after first-line chemotherapy.

http://dx.doi.org/10.1016/j.ctrv.2015.04.011

Other trials evaluating nintedanib in NSCLC are reported in Table 2.

Cediranib (AZD2171) is a multitarget tyrosine kinase inhibitor primarily directed against VEGFR2, but it has also demonstrated some inhibitory activity against FGFR1 and FGFR2 [71,81].

In a phase I trial 20 advanced NSCLC patients were treated with cediranib in combination with paclitaxel and carboplatin and the reported ORR was 45% [82].

Two randomized phase II/III studies assessed the safety and efficacy of cediranib in association with carboplatin and paclitaxel [83] or gemcitabine [84] as first-line treatment in advanced NSCLC patients. Both studies provided a starting cediranib dose of 45 mg orally once daily; however, this dose was not tolerated and it was set at 30 mg.

The first study [83] enrolled 296 patients and showed a significant higher response rate (38% vs. 16%; p > 0.0001) regardless of histology in oral cediranib arm over placebo, but no significant advantage in survival was seen.

In the phase II portion of the N0528 trial [84], 91 patients were randomized, but only 20% of all population had tumors with squamous cell histology. The study did not meet its primary end-point (ORR: 19% versus 20%, p = 1.0); furthermore, patients in the cediranib arm experienced significantly higher rates of non-hematological toxicities of grade ≥ 3 (71% versus 45%, p = 0.01). Interestingly, an exploratory pharmacogenetic study focused on SNPs of target genes, such as VEGFR1–3, FGFR1–3, showed that genetic variants of FGFR2 (rs17542768, p = 0.019; rs2071616, p = 0.010) were significantly associated with lower toxicity, whereas other variants of FGFR1 (rs7012413, p = 0.007) and FGFR2 (rs2912791, p = 0.0002) were associated with survival.

Another phase II trial evaluated cediranib in combination with pemetrexed in previously treated NSCLC patients, regardless of histology. The combination was tolerable and promising, but it appeared ineffective in recurrent squamous cell patients [85]. A

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

458

10

489

490

491

492

493

494

495

496

497

498

499

500

501

M. Tiseo et al./Cancer Treatment Reviews xxx (2015) xxx-xxx

randomized phase III trial of cediranib plus carboplatin and paclitaxel in patients with recurrent or metastatic NSCLC patients has
been completed. Its primary endpoint was OS and results are
awaited [NCT00795340].

Ponatinib (AP24534) is an oral multi-TK inhibitor that primarily inhibits BCR-ABL. It has been demonstrated that it is highly active in most solid tumor cell lines harboring different FGFR alterations, including NSCLCs, both *in vitro* and *in vivo*. Furthermore, its inhibitory capacity was more potent than that of other anti-FGFR compounds [71,86]. These data have suggested the investigation of ponatinib in clinical trials for patients with FGFR-addicted solid tumors. A phase II study of ponatinib in advanced lung and head and neck cancers with FGFR kinase alterations has suspended recruitment due to increase risk of blood clot [NCT01761747]. Another phase II trial evaluating the safety and effectiveness of ponatinib in lung cancer patients preselected for specific predictive biomarkers is currently ongoing [NCT01935336] [Table 2].

502 Lucitanib (E3810) is a potent oral dual TK inhibitor directed 503 against VEGFR1-3 and FGFR1-2. Lucitanib showed its antitumor activity as single agent in FGFR1-amplified lung cancers both 504 505 in vitro and in xenograft models [87-89]. Recently, the results of 506 an open-label phase I/IIa study have been published. Seventy-six 507 patients with advanced and potentially sensitive solid tumors 508 (tumors harboring FGF-aberrant pathway or sensitive to angio-509 genic inhibition) were enrolled. In overall population, 7 patients 510 (9%) had NSCLC. In the dose expansion phase, based on the 511 emerged safety data, lucitanib dose was set at 15 mg daily. The 512 most frequent AEs were hypertension, proteinuria, thrombotic 513 microangiopathy, asthenia, hypothyroidism, anorexia, diarrhea, 514 nausea, weight decrease, thrombocytopenia. Treatment-related 515 AEs of grade 4 were reported in two patients and consisted of a 516 case of increased lipase and a case of depressed level of conscious-517 ness. Sixty-four out of 76 patients were evaluable for response and 518 58 had misurable disease; among these, 4 patients with NSCLC (3 519 with FGFR1+ and 1 with 11q+) obtained a SD as the best response. 520 In the overall population the DCR reached 80% with several durable 521 responses and long-lasting SD [90]. To date, a phase II trial is cur-522 rently ongoing for evaluating the efficacy of lucitanib in FGFR1-am-523 plified Sq-NSCLC [NCT02109016] [Table 2].

524 Pazopanib (GW786034) is a multi-TK inhibitor including FGFR 525 family [91]. A phase II study comparing first-line combination 526 pazopanib-pemetrexed with the standard cisplatin-pemetrexed in NSCLC was terminated early due to unacceptable levels of toxi-527 528 city [92]. Results from a phase II study of pazopanib in 15 patients with stage IV non-squamous NSCLC progressed to first line therapy 529 530 containing bevacizumab, demonstrated limited activity to justify 531 additional accrual [93]. A phase II trial is currently ongoing evalu-532 ating the efficacy of first-line treatment of pazopanib in combina-533 tion with paclitaxel for advanced NSCLC patients not eligible for 534 front-line therapy with a platinum doublet [NCT01179269] 535 [Table 2].

536 Selective FGFR-TK inhibitors

This class of compounds includes several selective small molecules specifically directed against one or more FGFR tyrosine kinase domains. The IC_{50} versus the different targets are shown in Table 3.

LY2874455, an ATP-competitive pan-FGFR inhibitor, has demon-541 542 strated its potent and selective antitumor activity both in cell lines 543 and in tumor xenografts. Its inhibitory activity was particularly 544 evident in cancer cell lines with elevated FGF or FGFR levels, 545 whereas its activity against VEGFR2 was significantly lower. In 546 xenograft models derived from diverse cancer cell lines (including 547 NSCLCs), its administration promoted tumor growth arrest, 548 through inhibition of FRS2 phosphorylation [94]. This compound

has been evaluated in a phase I trial [NCT01212107] and its results are awaited.

AZD4547, a novel and selective FGFR1–3 inhibitor [95], was investigated both in FGFR1-amplified NSCLC cell lines and patient-derived tumor xenograft models. Its potent antitumor activity was shown in FGFR1-addicted NSCLCs and correlated with FGFR1 gene copies (FISH score 6) and protein (IHC 3+) expression. Inhibition of p-ERK and p-S6 correlated with its efficacy, suggesting a possible role as pharmacodynamic biomarkers [96]. Recently, the results of a dose escalation phase I trial of AZD4547 have been presented. The most common AEs were alopecia, fatigue, different gastrointestinal disturbances, nail disorders, dry skin, hyperphosphatemia and retinal pigment epithelial detachment. Encouraging evidence of antitumor activity was seen in some patients; clinical benefit was observed in 5 out of 20 patients with FGFR alterations (two of whom with Sq-NSCLC) [97]. The responder patient with Sq-NSCLC (prolonged SD) had high FGFR1 amplification together with amplification of 11q13 (containing FGF3/4/ 19 genes) and CCND1 [98]. Recently, the results of phase I expansion of AZD4547 in patients with previously treated metastatic FGFR1-amplified Sq-NSCLC have been updated. Among 15 patients, fourteen were evaluable for tumor response with 1 PR and 4 SD. The PR was observed in a patient with high FGFR1 amplification (FISH ratio > 2.8). AZD4547 was well tolerated but did not meet its efficacy endpoint (ORR) for continuation. Moreover, the increase in serum phosphatase levels could be a circulating biomarker of pharmacologic target inhibition [99].

AZD4547 is currently being evaluated in phase II/III trials across a range of solid tumors, including patients with FGFR1-amplified Sq-NSCLC [Table 2].

BGJ398 is a novel and selective FGFR1–3 inhibitor that demonstrated its dose-dependent activity in different xenograft tumor models [100]. Preliminary results on the first 26 patients enrolled in a phase I trial have been reported [NCT01004224]. The patients had advanced solid tumors harboring FGFR1/2 amplifications or FGFR3 mutations and among them 3 had FGFR1-amplified Sq-NSCLC. DLT was related to grade 3 elevation in AST/ALT levels and grade 2 corneal events at the dose of 100 mg. However, AEs were generally of mild/moderate grade and included fatigue (37%), diarrhea (37%), nausea (32%) and hyperphosphatemia (30%). A patient with FGFR1-amplified Sq-NSCLC obtained a confirmed PR, providing early evidence that inhibition of the FGFR pathway was effective in patients with FGFR-addicted tumors [101]. This trial is currently recruiting together with others, particularly in genetically pre-selected solid tumors [Table 2].

BAY1163877 and JNJ42756493 are selective FGFR family inhibitors with nanomolar affinity [102,103] currently under evaluation in phase I trials [NCT01976741-NCT01703481, Table 2]. JNJ-42756493, an orally bioavailable pan-FGFR inhibitor, has been evaluated in a phase I study [NCT01962532] in different solid tumors (including squamous NSCLC) with FGFR gene amplification, mutations or translocations. A total of 41 patients have been enrolled. The investigational drug was safe and well tolerated; 9 mg daily was the first RP2D, but safety evaluation at higher doses is ongoing. Hyperphosphatemia was the most common but manageable on-target AE (58.5% of any grade in ITT population). Two FGFR-positive NSCLC patients had a clinical benefit (1 PR and 1 SD) at a dose of 12 mg daily. Pharmacodynamic serum (e.g., increased phosphate, FGF23 and calcium levels and decreased PTH levels) and tissue biomarkers (e.g., decreased phospho-ERK levels after drug exposure), capable to predict tumor response are being studied [104].

Irreversible FGRF inhibitors: FIIN-1 is the first covalent FGRF irreversible inhibitor targeting a cysteine residue conserved in all four FGFR kinases. *In vitro* the inhibitor inhibited numerous FGFR-dependent cancer cell lines with a potency in the nM range [105].

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

Please cite this article in press as: Tiseo M et al. FGFR as potential target in the treatment of squamous non small cell lung cancer. Cancer Treat Rev (2015), http://dx.doi.org/10.1016/j.ctrv.2015.04.011 M. Tiseo et al./Cancer Treatment Reviews xxx (2015) xxx-xxx

Very recently, a second generation of irreversible FGFR inhibitors
(FIIN-2 and FIIN-3) with improved affinity for wild type and gatekeeper mutants has been developed and characterized [106]. These
new inhibitors can overcome clinical resistance to first-generation
FGFR inhibitors.

620 **Conclusions: open issues and future perspectives**

FGF/FGFR signaling pathway has been recognized as one of the hallmark alterations with relevant clinical implications, particularly in Sq-NSCLC. Data from literature have established that the presence of an aberrant FGFR pathway has to be considered as possible prognostic factor of poor outcome and predictive of potential sensitivity to FGFR inhibitors.

627 Among the several DNA alterations involved in aberrant FGFR 628 pathway, FGFR1 amplification, identified particularly in Sq-629 NSCLC, is the most frequent. Several techniques are being used to detect specific genomic aberrations involving FGFR pathway, such 630 as in situ hybridization assays (FISH and CISH), quantitative real-631 time PCR and RNA sequencing. Among them, FISH represents the 632 633 standard assay to screen tumor samples, but various studies have 634 considered different cut-off values. Therefore, no standardized cri-635 teria have been defined to consider test as positive to date. 636 Moreover, traditional FISH analysis is not able to detect some vari-637 ants of FGFR gene fusions. By using this test, at least 20% of patients 638 potentially suitable for FGFR inhibition, such as those with FGF/ 639 FGFR tumor upregulation or FGFR mutations, may be missed. 640 Therefore, this issue remains a relevant question because of impli-641 cation in screening patients amenable for FGFR-targeted therapy 642 and in drug clinical development.

Standardization of different molecular diagnostic procedures,
together with the use of high throughput sequencing technologies,
will allow us to better select patients suitable for FGFR targeted
therapy and, at the same time, to increase the number of significant and potentially predictive alterations in the FGFR pathway.

648 Initially, FGFR inhibitors were tested primarily as anti-angio-649 genics and in molecularly unselected patients. In these popula-650 tions, the results of different clinical trials have shown a 651 potential benefit from adding non-selective FGFR inhibitors to 652 standard chemotherapy, due to their synergistic antiangiogenic 653 activity. Both preclinical and clinical data have demonstrated a biologic rationale to use an FGFR inhibitor in this setting, although we 654 are not able to establish when we have to consider this combina-655 656 tion treatment.

Furthermore, it seems that a possible explanation of disease 657 658 progression during a VEGFR inhibitor-based therapy would be 659 related to high levels of FGFs; at this time, the use of an FGFR inhi-660 bitor may be able to overcome resistance. Nowadays, we are not 661 able to recognize the best treatment strategy (up-front versus after 662 anti-VEGF treatment failure) because of the lack of comparison tri-663 als. Probably, there are no significant differences in terms of efficacy between the two strategies, although a patients' selection 664 according to serum FGF levels might allow optimizing the treat-665 666 ment strategy.

667 Both nonselective and selective FGFR inhibitors demonstrated 668 their activity in FGFR-addicted NSCLC. Efficacy data with multitar-669 get inhibitors are more robust, but selective FGFR inhibitors might 670 be more potent. The time for obtaining these data is not mature, 671 because direct comparison trials lack, and poses the fascinating 672 question of whether the patients' selection can be influenced and 673 enriched by these different molecules.

Furthermore, there is a potential role of FGFR inhibitors in combination with other targeted agents. It is possible that a combined treatment might be a feasible strategy to delay or overcome acquired resistance to EGFR inhibitor. Unfortunately, when applied in clinical trial, it failed because of toxicity. At this regard, several data have been reported on different toxicity profiles between selective and nonselective anti-FGFR drugs. The prevalent side effects of non-selective FGFR inhibitors are related to VEGF inhibition (e.g., hypertension, proteinuria, hypotiroidism and cardiovascular events); in addition, other common (off-target) adverse events are gastrointestinal disorders and skin reaction. This broad toxicity profile has limited the use of these compounds in combination with other anticancer drugs, particularly with anti-EGFR agents. Conversely, selective FGFR inhibitors showed a limited toxicity profile with hyperphosphatemia as the principal on-target toxicity and no events related to VEGF inhibition. In addition, new FGF inhibitors (FGF-ligand traps) have a better tolerability, also avoiding the onset of hyperphosphatemia, due to their little or no affinity for the hormonal FGFs (19, 21 and 23). Therefore, selective FGFR inhibitors lend themselves to combination with other drugs, according to their favorable toxicity profile, or alone after progression to another targeted agent.

The development of new covalent, irreversible FGFR inhibitors may overcome resistance to first-generation inhibitor even if at this time, no clinical data are available about mechanisms of resistance to anti-FGFR inhibitors.

Finally, in absence of clinical data on different activity and longterm safety of FGFR inhibition, in our opinion the toxicity profile, host and tumor selection (FGFR status and predictive biomarkers) should drive the drug development and the design of clinical trials.

In conclusion, it is plausible to hope that in the next years the research efforts in preclinical and clinical fields allow to establish an optimal treatment strategy also in FGFR-addicted NSCLC population.

Conflict of interest	
----------------------	--

All authors of this paper have not conflict of interest to disclose.

Acknowledgement

We thank Associazione Italiana per la Ricerca sul Cancro (AIRC) Milan grant IG 14214.

References

- Travis WD, Brambilla E, Muller-Hermelink HK, Harris CC. World Health Organization CLassification of tumors. Pathology and genetics of tumours of the lung, pleura, thymus and heart. Lyon, France: IARCC Press; 2004.
- [2] Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, et al. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. Lancet Oncol 2012;13:239–46.
- [3] Kwak EL, Bang YJ, Camidge DR, Shaw AT, Solomon B, Maki RG, et al. Anaplastic lymphoma kinase inhibition in non-small-cell lung cancer. N Engl J Med 2010;363:1693–703.
- [4] Shaw AT, Ou SH, Bang YJ, Camidge DR, Solomon BJ, Salgia R, et al. Crizotinib in ROS1-rearranged non-small-cell lung cancer. N Engl J Med 2014;371:1963–71.
- [5] Oxnard GR, Binder A, Janne PA. New targetable oncogenes in non-small-cell lung cancer. J Clin Oncol 2013;31:1097–104.
- [6] Scagliotti GV, Parikh P, von Pawel J, Biesma B, Vansteenkiste J, Manegold C, et al. Phase III study comparing cisplatin plus gemcitabine with cisplatin plus pemetrexed in chemotherapy-naive patients with advanced-stage non-smallcell lung cancer. J Clin Oncol 2008;26:3543–51.
- [7] Drilon A, Rekhtman N, Ladanyi M, Paik P. Squamous-cell carcinomas of the lung: emerging biology, controversies, and the promise of targeted therapy. Lancet Oncol 2012;13:e418–26.
- [8] Comprehensive genomic characterization of squamous cell lung cancers. Nature 2012;489:519–25.
- [9] Hammerman PS, Sos ML, Ramos AH, Xu C, Dutt A, Zhou W, et al. Mutations in the DDR2 kinase gene identify a novel therapeutic target in squamous cell lung cancer. Cancer Discov 2011;1:78–89.
- [10] Payne LS, Huang PH. Discoidin domain receptor 2 signaling networks and therapy in lung cancer. J Thorac Oncol 2014;9:900–4.

Please cite this article in press as: Tiseo M et al. FGFR as potential target in the treatment of squamous non small cell lung cancer. Cancer Treat Rev (2015), http://dx.doi.org/10.1016/j.ctrv.2015.04.011

679

680

681

682

683

684

685

686 687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704 705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734 735

736

737

738

739

740

741

742

YCTRV 1402 4 May 2015

ARTICLE IN PRESS

12

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770 771

772

773

778 779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

M. Tiseo et al./Cancer Treatment Reviews xxx (2015) xxx-xxx

- [11] Fumarola C, Bonelli MA, Petronini PG, Alfieri RR. Targeting PI3K/AKT/mTOR pathway in non small cell lung cancer. Biochem Pharmacol 2014;90:197–207.
 - [12] Heist RS, Sequist LV, Engelman JA. Genetic changes in squamous cell lung cancer: a review. J Thorac Oncol 2012;7:924–33.
 - [13] Gold KA, Wistuba II, Kim ES. New strategies in squamous cell carcinoma of the lung: identification of tumor drivers to personalize therapy. Clin Cancer Res 2012;18:3002–7.
- [14] Wesche J, Haglund K, Haugsten EM. Fibroblast growth factors and their receptors in cancer. Biochem J 2011;437:199–213.
- [15] Beenken A, Mohammadi M. The FGF family: biology, pathophysiology and therapy. Nat Rev Drug Discov 2009;8:235–53.
- [16] Holzmann K, Grunt T, Heinzle C, Sampl S, Steinhoff H, Reichmann N, et al. Alternative splicing of fibroblast growth factor receptor IgIII loops in cancer. J Nucleic Acids 2012;2012:950508.
- [17] Trueb B. Biology of FGFRL1, the fifth fibroblast growth factor receptor. Cell Mol Life Sci 2011;68:951–64.
- [18] Turner N, Grose R. Fibroblast growth factor signalling: from development to cancer. Nat Rev Cancer 2010;10:116–29.
- [19] Jabs EW, Li X, Scott AF, Meyers G, Chen W, Eccles M, et al. Jackson-Weiss and Crouzon syndromes are allelic with mutations in fibroblast growth factor receptor 2. Nat Genet 1994;8:275–9.
- [20] Przylepa KA, Paznekas W, Zhang M, Golabi M, Bias W, Bamshad MJ, et al. Fibroblast growth factor receptor 2 mutations in Beare-Stevenson cutis gyrata syndrome. Nat Genet 1996;13:492–4.
- [21] Paznekas WA, Cunningham ML, Howard TD, Korf BR, Lipson MH, Grix AW, et al. Genetic heterogeneity of Saethre-Chotzen syndrome, due to TWIST and FGFR mutations. Am J Hum Genet 1998;62:1370–80.
- [22] Lieu C, Heymach J, Overman M, Tran H, Kopetz S. Beyond VEGF: inhibition of the fibroblast growth factor pathway and antiangiogenesis. Clin Cancer Res 2011;17:6130–9.
- [23] Powers CJ, McLeskey SW, Wellstein A. Fibroblast growth factors, their receptors and signaling. Endocr Relat Cancer 2000;7:165–97.
- [24] Dienstmann R, Rodon J, Prat A, Perez-Garcia J, Adamo B, Felip E, et al. Genomic aberrations in the FGFR pathway: opportunities for targeted therapies in solid tumors. Ann Oncol 2014;25:552–63.
- [25] Jiang T, Gao G, Fan G, Li M, Zhou C. FGFR1 amplification in lung squamous cell carcinoma: a systematic review with meta-analysis. Lung Cancer 2015;87:1–7.
- [26] Weiss J, Sos ML, Seidel D, Peifer M, Zander T, Heuckmann JM, et al. Frequent and focal FGFR1 amplification associates with therapeutically tractable FGFR1 dependency in squamous cell lung cancer. Sci Transl Med 2010;2:62ra93.
- [27] Dutt A, Ramos AH, Hammerman PS, Mermel C, Cho J, Sharifnia T, et al. Inhibitor-sensitive FGFR1 amplification in human non-small cell lung cancer. PLoS One 2011;6:e20351.
- [28] Kohler LH, Mireskandari M, Knosel T, Altendorf-Hofmann A, Kunze A, Schmidt A, et al. FGFR1 expression and gene copy numbers in human lung cancer. Virchows Arch 2012;461:49–57.
- [29] Craddock KJ, Ludkovski O, Sykes J, Shepherd FA, Tsao MS. Prognostic value of fibroblast growth factor receptor 1 gene locus amplification in resected lung squamous cell carcinoma. | Thorac Oncol 2013;8:1371–7.
- [30] Malchers F, Dietlein F, Schottle J, Lu X, Nogova L, Albus K, et al. Cellautonomous and non-cell-autonomous mechanisms of transformation by amplified FGFR1 in lung cancer. Cancer Discov 2014;4:246–57.
- [31] Cihoric N, Savic S, Schneider S, Ackermann I, Bichsel-Naef M, Schmid RA, et al. Prognostic role of FGFR1 amplification in early-stage non-small cell lung cancer. Br J Cancer 2014;110:2914–22.
- [32] Kim HR, Kim DJ, Kang DR, Lee JG, Lim SM, Lee CY, et al. Fibroblast growth factor receptor 1 gene amplification is associated with poor survival and cigarette smoking dosage in patients with resected squamous cell lung cancer. J Clin Oncol 2013;31:731–7.
- [33] Heist RS, Mino-Kenudson M, Sequist LV, Tammireddy S, Morrissey L, Christiani DC, et al. FGFR1 amplification in squamous cell carcinoma of the lung. J Thorac Oncol 2012;7:1775–80.
- [34] Seo AN, Jin Y, Lee HJ, Sun PL, Kim H, Jheon S, et al. FGFR1 amplification is associated with poor prognosis and smoking in non-small-cell lung cancer. Virchows Arch 2014;465:547–58.
- [35] Gadgeel SM, Chen W, Cote ML, Bollig-Fischer A, Land S, Schwartz AG, et al. Fibroblast growth factor receptor 1 amplification in non-small cell lung cancer by quantitative real-time PCR. PLoS One 2013;8:e79820.
- [36] Davies H, Hunter C, Smith R, Stephens P, Greenman C, Bignell G, et al. Somatic mutations of the protein kinase gene family in human lung cancer. Cancer Res 2005;65:7591–5.
- [37] Greenman C, Stephens P, Smith R, Dalgliesh GL, Hunter C, Bignell G, et al. Patterns of somatic mutation in human cancer genomes. Nature 2007;446:153–8.
- [38] Liao RG, Jung J, Tchaicha J, Wilkerson MD, Sivachenko A, Beauchamp EM, et al. Inhibitor-sensitive FGFR2 and FGFR3 mutations in lung squamous cell carcinoma. Cancer Res 2013;73:5195–205.
- [39] Marks JL, McLellan MD, Zakowski MF, Lash AE, Kasai Y, Broderick S, et al. Mutational analysis of EGFR and related signaling pathway genes in lung adenocarcinomas identifies a novel somatic kinase domain mutation in FGFR4. PLoS One 2007;2:e426.

- [40] Wu YM, Su F, Kalyana-Sundaram S, Khazanov N, Ateeq B, Cao X, et al. Identification of targetable FGFR gene fusions in diverse cancers. Cancer Discov 2013;3:636–47.
- [41] Capelletti M, Dodge ME, Ercan D, Hammerman PS, Park SI, Kim J, et al. Identification of recurrent FGFR3-TACC3 fusion oncogenes from lung adenocarcinoma. Clin Cancer Res 2014;20:6551–8.
- [42] Wang R, Wang L, Li Y, Hu H, Shen L, Shen X, et al. FGFR1/3 tyrosine kinase fusions define a unique molecular subtype of non-small cell lung cancer. Clin Cancer Res 2014;20:4107–14.
- [43] Marek L, Ware KE, Fritzsche A, Hercule P, Helton WR, Smith JE, et al. Fibroblast growth factor (FGF) and FGF receptor-mediated autocrine signaling in non-small-cell lung cancer cells. Mol Pharmacol 2009;75:196–207.
- [44] Kopetz S, Hoff PM, Morris JS, Wolff RA, Eng C, Glover KY, et al. Phase II trial of infusional fluorouracil, irinotecan, and bevacizumab for metastatic colorectal cancer: efficacy and circulating angiogenic biomarkers associated with therapeutic resistance. J Clin Oncol 2010;28:453–9.
- [45] Batchelor TT, Sorensen AG, di Tomaso E, Zhang WT, Duda DG, Cohen KS, et al. AZD2171, a pan-VEGF receptor tyrosine kinase inhibitor, normalizes tumor vasculature and alleviates edema in glioblastoma patients. Cancer Cell 2007;11:83–95.
- [46] Shou Y, Hirano T, Gong Y, Kato Y, Yoshida K, Ohira T, et al. Influence of angiogenetic factors and matrix metalloproteinases upon tumour progression in non-small-cell lung cancer. Br J Cancer 2001;85:1706–12.
- [47] Volm M, Koomagi R, Mattern J, Stammler G. Prognostic value of basic fibroblast growth factor and its receptor (FGFR-1) in patients with non-small cell lung carcinomas. Eur J Cancer 1997;33:691–3.
- [48] Bremnes RM, Camps C, Sirera R. Angiogenesis in non-small cell lung cancer: the prognostic impact of neoangiogenesis and the cytokines VEGF and bFGF in tumours and blood. Lung Cancer 2006;51:143–58.
- [49] Takanami I, Tanaka F, Hashizume T, Kikuchi K, Yamamoto Y, Yamamoto T, et al. The basic fibroblast growth factor and its receptor in pulmonary adenocarcinomas: an investigation of their expression as prognostic markers. Eur J Cancer 1996;32A:1504–9.
- [50] Brattstrom D, Bergqvist M, Hesselius P, Larsson A, Lamberg K, Wernlund J, et al. Elevated preoperative serum levels of angiogenic cytokines correlate to larger primary tumours and poorer survival in non-small cell lung cancer patients. Lung Cancer 2002;37:57–63.
- [51] Brattstrom D, Bergqvist M, Hesselius P, Larsson A, Wagenius G, Brodin O. Serum VEGF and bFGF adds prognostic information in patients with normal platelet counts when sampled before, during and after treatment for locally advanced non-small cell lung cancer. Lung Cancer 2004;43:55–62.
- [52] Iwasaki A, Kuwahara M, Yoshinaga Y, Shirakusa T. Basic fibroblast growth factor (bFGF) and vascular endothelial growth factor (VEGF) levels, as prognostic indicators in NSCLC. Eur J Cardiothorac Surg 2004;25:443–8.
- [53] Donnem T, Al-Shibli K, Al-Saad S, Busund LT, Bremnes RM. Prognostic impact of fibroblast growth factor 2 in non-small cell lung cancer: coexpression with VEGFR-3 and PDGF-B predicts poor survival. J Thorac Oncol 2009;4:578–85.
- [54] Easton DF, Pooley KA, Dunning AM, Pharoah PD, Thompson D, Ballinger DG, et al. Genome-wide association study identifies novel breast cancer susceptibility loci. Nature 2007;447:1087–93.
- [55] Bange J, Prechtl D, Cheburkin Y, Specht K, Harbeck N, Schmitt M, et al. Cancer progression and tumor cell motility are associated with the FGFR4 Arg(388) allele. Cancer Res 2002;62:840–7.
- [56] Wang J, Stockton DW, Ittmann M. The fibroblast growth factor receptor-4 Arg388 allele is associated with prostate cancer initiation and progression. Clin Cancer Res 2004;10:6169–78.
- [57] Spinola M, Leoni V, Pignatiello C, Conti B, Ravagnani F, Pastorino U, et al. Functional FGFR4 Gly388Arg polymorphism predicts prognosis in lung adenocarcinoma patients. J Clin Oncol 2005;23:7307–11.
- [58] Lin L, Chamberlain L, Pak ML, Nagarajan A, Gupta R, Zhu LJ, et al. A large-scale RNAi-based mouse tumorigenesis screen identifies new lung cancer tumor suppressors that repress FGFR signaling. Cancer Discov 2014;4:1168–81.
- [59] Wynes MW, Hinz TK, Gao D, Martini M, Marek LA, Ware KE, et al. FGFR1 mRNA and protein expression, not gene copy number, predict FGFR TKI sensitivity across all lung cancer histologies. Clin Cancer Res 2014;20:3299–309.
- [60] Terai H, Soejima K, Yasuda H, Nakayama S, Hamamoto J, Arai D, et al. Activation of the FGF2-FGFR1 autocrine pathway: a novel mechanism of acquired resistance to gefitinib in NSCLC. Mol Cancer Res 2013;11:759–67.
- [61] Azuma K, Kawahara A, Sonoda K, Nakashima K, Tashiro K, Watari K, et al. FGFR1 activation is an escape mechanism in human lung cancer cells resistant to afatinib, a pan-EGFR family kinase inhibitor. Oncotarget 2014;5:5908–19.
- [62] Ware KE, Marshall ME, Heasley LR, Marek L, Hinz TK, Hercule P, et al. Rapidly acquired resistance to EGFR tyrosine kinase inhibitors in NSCLC cell lines through de-repression of FGFR2 and FGFR3 expression. PLoS One 2010;5:e14117.
- [63] Ware KE, Hinz TK, Kleczko E, Singleton KR, Marek LA, Helfrich BA, et al. A mechanism of resistance to gefitinib mediated by cellular reprogramming and the acquisition of an FGF2-FGFR1 autocrine growth loop. Oncogenesis 2013;2:e39.
- [64] Fischer H, Taylor N, Allerstorfer S, Grusch M, Sonvilla G, Holzmann K, et al. Fibroblast growth factor receptor-mediated signals contribute to the malignant phenotype of non-small cell lung cancer cells: therapeutic

910

911

912

913

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

914

915

916

917

918

919

920

921

922

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

13

or can remail cell lung or xenograft models. Clin oil R, Stockman PK, et al. of fibroblast growth factor olid tumors. Cancer Res oney C, Smith NR, et al. of AZD4547, an inhibitor of ents with advanced solid ol. patients with advanced antitumor activity and suppl abst 8035]. Ito M, Barys L, et al. FGFR 2-BCI308 a selective nan-

1038

1039

1040

1041

1042

1043

1044

1045 1046

1047

1077

implications and synergism with epidermal growth factor receptor inhibition. Mol Cancer Ther 2008;7:3408-19.

- [65] Kim B, Wang S, Lee JM, Jeong Y, Ahn T, Son DS, et al. Synthetic lethal screening reveals FGFR as one of the combinatorial targets to overcome resistance to Met-targeted therapy. Oncogene 2014;34:1083–93.
- [66] O'Donnell P, Goldman JW, Gordon MS, Shih K, Choi D, Lu O, et al. A phase I dose escalation study of MFGR1877S, a human monoclonal anti-fibroblast growth factor 3 (FGFR3) antibody, in patients (pts) with advanced solid tumors. Eur J Cancer 2012;48(Suppl. 6) [abst 621].
- [67] Harding TC, Long L, Palencia S, Zhang H, Sadra A, Hestir K, et al. Blockade of nonhormonal fibroblast growth factors by FP-1039 inhibits growth of multiple types of cancer. Sci Transl Med 2013;5:178ra39.
 [68] Tolcher A, Pandopoulos K, Pathiak A, Heath F, Weise A, Prokop T, et al.
 - [68] Tolcher A, Papadopoulos K, Patniak A, Heath E, Weise A, Prokop T, et al. Preliminary results of a dose escalation study of the fibroblast growth factor (FGF) "trap" FP-1039 (FGFR1:Fc) in patients with advanced malignancies. Eur J Cancer 2010;8 [8 issue 7 abst 381].
 - [69] Johnson DH, Fehrenbacher L, Novotny WF, Herbst RS, Nemunaitis JJ, Jablons DM, et al. Randomized phase II trial comparing bevacizumab plus carboplatin and paclitaxel with carboplatin and paclitaxel alone in previously untreated locally advanced or metastatic non-small-cell lung cancer. J Clin Oncol 2004;22:2184–91.
 - [70] Lee SH, Lopes de Menezes D, Vora J, Harris A, Ye H, Nordahl L, et al. In vivo target modulation and biological activity of CHIR-258, a multitargeted growth factor receptor kinase inhibitor, in colon cancer models. Clin Cancer Res 2005;11:3633–41.
 - [71] Gozgit JM, Wong MJ, Moran L, Wardwell S, Mohemmad QK, Narasimhan NI, et al. Ponatinib (AP24534), a multitargeted pan-FGFR inhibitor with activity in multiple FGFR-amplified or mutated cancer models. Mol Cancer Ther 2012;11:690–9.
 - [72] Sarker D, Molife R, Evans TR, Hardie M, Marriott C, Butzberger-Zimmerli P, et al. A phase I pharmacokinetic and pharmacodynamic study of TKI258, an oral, multitargeted receptor tyrosine kinase inhibitor in patients with advanced solid tumors. Clin Cancer Res 2008;14:2075–81.
 - [73] Andre F, Bachelot D, Campone M, Dalenc F, Perez-Garcia JM, Hurvitz SA, et al. A multicenter, open-label phase II trial of dovitinib, an FGFR1 inhibitor, in FGFR1-amplified and non-amplified metastatic breast cancer. J Clin Oncol 2011;29 [suppl abst 508].
 - [74] Argevin E, Grunwald V, Ravaud A, Castellano DE, Lin CC, Gschwend JE, et al. A phase I study of dovitinib (TKI258), an FGFR-and VEGFR-inhibitor, in patients with advanced or metastatic renal cell carcinoma. J Clin Oncol 2014;29 [suppl abst 4551].
 - [75] Hilberg F, Roth GJ, Krssak M, Kautschitsch S, Sommergruber W, Tontsch-Grunt U, et al. BIBF 1120: triple angiokinase inhibitor with sustained receptor blockade and good antitumor efficacy. Cancer Res 2008;68:4774–82.
 - [76] Mross K, Stefanic M, Gmehling D, Frost A, Baas F, Unger C, et al. Phase I study of the angiogenesis inhibitor BIBF 1120 in patients with advanced solid tumors. Clin Cancer Res 2010;16:311–9.
 - [77] Ellis PM, Kaiser R, Zhao Y, Stopfer P, Gyorffy S, Hanna N. Phase I open-label study of continuous treatment with BIBF 1120, a triple angiokinase inhibitor, and pemetrexed in pretreated non-small cell lung cancer patients. Clin Cancer Res 2010;16:2881–9.
 - [78] Doebele RC, Conkling P, Traynor AM, Otterson GA, Zhao Y, Wind S, et al. A phase I, open-label dose-escalation study of continuous treatment with BIBF 1120 in combination with paclitaxel and carboplatin as first-line treatment in patients with advanced non-small-cell lung cancer. Ann Oncol 2012;23:2094–102.
 - [79] Reck M, Kaiser R, Eschbach C, Stefanic M, Love J, Gatzemeier U, et al. A phase II double-blind study to investigate efficacy and safety of two doses of the triple angiokinase inhibitor BIBF 1120 in patients with relapsed advanced nonsmall-cell lung cancer. Ann Oncol 2011;22:1374–81.
 - [80] Reck M, Kaiser R, Mellemgaard A, Douillard JY, Orlov S, Krzakowski M, et al. Docetaxel plus nintedanib versus docetaxel plus placebo in patients with previously treated non-small-cell lung cancer (LUME-Lung 1): a phase 3, double-blind, randomised controlled trial. Lancet Oncol 2014;15:143–55.
 - [81] Wedge SR, Kendrew J, Hennequin LF, Valentine PJ, Barry ST, Brave SR, et al. AZD2171: a highly potent, orally bioavailable, vascular endothelial growth factor receptor-2 tyrosine kinase inhibitor for the treatment of cancer. Cancer Res 2005;65:4389–400.
 - [82] Laurie SA, Gauthier I, Arnold A, Shepherd FA, Ellis PM, Chen E, et al. Phase I and pharmacokinetic study of daily oral AZD2171, an inhibitor of vascular endothelial growth factor tyrosine kinases, in combination with carboplatin and paclitaxel in patients with advanced non-small-cell lung cancer: the National Cancer Institute of Canada clinical trials group. J Clin Oncol 2008;26:1871–8.
 - [83] Goss GD, Arnold A, Shepherd FA, Dediu M, Ciuleanu TE, Fenton D, et al. Randomized, double-blind trial of carboplatin and paclitaxel with either daily oral cediranib or placebo in advanced non-small-cell lung cancer: NCIC clinical trials group BR24 study. J Clin Oncol 2010;28:49–55.
 - [84] Dy GK, Mandrekar SJ, Nelson GD, Meyers JP, Adjei AA, Ross HJ, et al. A randomized phase II study of gemcitabine and carboplatin with or without cediranib as first-line therapy in advanced non-small-cell lung cancer: North Central Cancer Treatment Group Study N0528. J Thorac Oncol 2013;8:79–88.

- [85] Gadgeel SM, Ruckdeschel JC, Wozniak A, Chen W, Hackstock D, Galasso C, et al. Cediranib, a VEGF receptor 1, 2, and 3 inhibitor, and pemetrexed in patients (pts) with recurrent non-small cell lung cancer (NSCLC). J Clin Oncol 2011;29 [suppl abst 7564].
- [86] O'Hare T, Shakespeare WC, Zhu X, Eide CA, Rivera VM, Wang F, et al. AP24534, a pan-BCR-ABL inhibitor for chronic myeloid leukemia, potently inhibits the T3151 mutant and overcomes mutation-based resistance. Cancer Cell 2009;16:401–12.
- [87] Colella G, Damia G, D' Incalci M, Cereda R, Peitz I, Kelter G, et al. E-3810 antitumor activity in human xenografts expressing different levels of FGF receptor 1. Cancer Res 2011;71(Suppl. 8) [abst 595].
- [88] Bello E, Colella G, Scarlato V, Oliva P, Berndt A, Valbusa G, et al. E-3810 is a potent dual inhibitor of VEGFR and FGFR that exerts antitumor activity in multiple preclinical models. Cancer Res 2011;71:1396–405.
- [89] Zhou Y, Chen Y, Tong L, Xie H, Wen W, Zhang J, et al. AL3810, a multi-tyrosine kinase inhibitor, exhibits potent anti-angiogenic and anti-tumour activity via targeting VEGFR, FGFR and PDGFR. J Cell Mol Med 2012;16:2321–30.
- [90] Soria JC, DeBraud F, Bahleda R, Adamo B, Andre F, Dientsmann R, et al. Phase I/ Ila study evaluating the safety, efficacy, pharmacokinetics, and pharmacodynamics of lucitanib in advanced solid tumors. Ann Oncol 2014;25:2244–51.
- [91] Kumar R, Knick VB, Rudolph SK, Johnson JH, Crosby RM, Crouthamel MC, et al. Pharmacokinetic-pharmacodynamic correlation from mouse to human with pazopanib, a multikinase angiogenesis inhibitor with potent antitumor and antiangiogenic activity. Mol Cancer Ther 2007;6:2012–21.
- [92] Scagliotti GV, Felip E, Besse B, von Pawel J, Mellemgaard A, Reck M, et al. An open-label, multicenter, randomized, phase II study of pazopanib in combination with pemetrexed in first-line treatment of patients with advanced-stage non-small-cell lung cancer. J Thorac Oncol 2013;8:1529–37.
- [93] Weiss JM, Villaruz LC, Socinski MA, Ivanova A, Grilley-Olson J, Dhruva N, et al. A single-arm phase II trial of pazopanib in patients with advanced non-small cell lung cancer with non-squamous histology with disease progression on bevacizumab containing therapy. Lung Cancer 2014;86:288–90.
- [94] Zhao G, Li WY, Chen D, Henry JR, Li HY, Chen Z, et al. A novel, selective inhibitor of fibroblast growth factor receptors that shows a potent broad spectrum of antitumor activity in several tumor xenograft models. Mol Cancer Ther 2011;10:2200–10.
- [95] Gavine PR, Mooney L, Kilgour E, Thomas AP, Al-Kadhimi K, Beck S, et al. AZD4547: an orally bioavailable, potent, and selective inhibitor of the fibroblast growth factor receptor tyrosine kinase family. Cancer Res 2012;72:2045–56.
- [96] Zhang J, Zhang L, Su X, Li M, Xie L, Malchers F, et al. Translating the therapeutic potential of AZD4547 in FGFR1-amplified non-small cell lung cancer through the use of patient-derived tumor xenograft models. Clin Cancer Res 2012;18:6658–67.
- [97] Andre F, Ranson M, Dean E, Varga A, Van der Noil R, Stockman PK, et al. Results of a phase I study of AZD4547, an inhibitor of fibroblast growth factor receptor (FGFR) in patients with advanced solid tumors. Cancer Res 2013;73(Suppl. 1) [abst LB-145].
- [98] Kilgour E, Ferry D, Saggese M, Arkenau HT, Rooney C, Smith NR, et al. Exploratory biomarker analysis of a phase I study of AZD4547, an inhibitor of fibroblast growth factor receptor (FGFR), in patients with advanced solid tumors. J Clin Oncol 2014;32 [5s Suppl abst 11010].
- [99] Paik PK, Shen R, Ferry D, Soria JC, Mathewson A, Kilgour E, et al. A phase 1b open-label multicenter study of AZD4547 in patients with advanced squamous cell lung cancer: preliminary antitumor activity and pharmacodynamics data. J Clin Oncol 2014;32 [5s suppl abst 8035].
- [100] Guagnano V, Kauffmann A, Wohrle S, Stamm C, Ito M, Barys L, et al. FGFR genetic alterations predict for sensitivity to NVP-BGJ398, a selective pan-FGFR inhibitor. Cancer Discov 2012;2:1118–33.
- [101] Wolf J, LoRusso PM, Camidge RD, Perez JM, Tabernero J, Hidalgo M, et al. A phase I dose escalation study of NVP-BGJ398, a selective pan FGFR inhibitor in genetically preselected advanced solid tumors. Cancer Res 2012;72(Suppl. 8) [abst LB-122].
- [102] Angibaud PR, Mavellec L, Saxty G, Adelinet C, Akkari R, Berdini V, et al. Discovery of JNJ-42756493, a potent fibroblast growth factor receptor (FGFR) inhibitor using a fragment based approach. Cancer Res 2014;74(Suppl. 19) [abst 4748].
- [103] Heroult M, Ellinghaus P, Sieg C, Brohm D, Gruenewald S, Collin MP, et al. Preclinical profile of BAY 1163877 – a selective pan-FGFR inhibitor in phase 1 clinical trial. Cancer Res 2014;74(Suppl. 19) [abst 1739].
- [104] Dienstmann R, Bahleda R, Adamo B, Rodon J, Varga A, Gazzah A, et al. First in human study of JNJ-42756493, a potent pan fibroblast growth factor receptor (FGFR) inhibitor in patients with advanced solid tumors. Cancer Res 2014;74(Suppl. 19) [abst CT325].
- [105] Zhou W, Hur W, McDermott U, Dutt A, Xian W, Ficarro SB, et al. A structureguided approach to creating covalent FGFR inhibitors. Chem Biol 2010;17:285–95.
- [106] Tan L, Wang J, Tanizaki J, Huang Z, Aref AR, Rusan M, et al. Development of covalent inhibitors that can overcome resistance to first-generation FGFR kinase inhibitors. Proc Natl Acad Sci USA 2014;111:E4869–77.