



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Survey of prevalence and seasonal variability of *Listeria monocytogenes* in raw cow milk from Northern Italy

This is a pre print version of the following article:

Original

Survey of prevalence and seasonal variability of *Listeria monocytogenes* in raw cow milk from Northern Italy / Dalzini, Elena; Bernini, Valentina; Barbara, Bertasi; Paolo, Daminelli; Nadia, Losio Marina; Giorgio, Varisco. - In: FOOD CONTROL. - ISSN 0956-7135. - 60:(2016), pp. 466-470.
[10.1016/j.foodcont.2015.08.019]

Availability:

This version is available at: 11381/2795453 since: 2021-11-02T11:08:42Z

Publisher:

Elsevier Ltd

Published

DOI:10.1016/j.foodcont.2015.08.019

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

Survey of prevalence and seasonal variation of *Listeria monocytogenes* in raw cow milk from Northern Italy

Elena Dalzini ^{a,*}, Valentina Bernini ^b, Barbara Bertasi ^c, Paolo Daminelli ^a, Marina-Nadia Losio ^c and Giorgio Varisco ^a

^a National Reference Centre for Emerging Risk in Food Safety, Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna "B. Ubertini", Brescia, Italy.

^b Department of Food Science, University of Parma, Parco Area delle Scienze 49/A, 43124, Parma, Italy.

^c Department of Food Microbiology, Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna "B. Ubertini", Brescia, Italy.

* **Corresponding author:** Tel.: +39 0302290533; fax: +39 2290629; *E-mail address:* elena.dalzini@izsler.it (Elena Dalzini).

Abstract

Listeria monocytogenes is an important food-borne pathogen causing meningitis, meningo-encephalitis and abortion. Both sporadic and epidemic human listeriosis cases are associated with the consumption of contaminated foods. To assess the potential risk to consumer health, the presence of *L. monocytogenes* was investigated using qualitative and quantitative methods in raw milk (bulk tank milk and milk for vending machine) collected from 2010 to 2013 in Northern Italy (Lombardy and Emilia-Romagna regions). Overall, *L. monocytogenes* was detected in 145 on 8716 of raw milk samples, with a prevalence of 1.66% (95% C.I. 1.4%-1.7%). The prevalence ranged from 0.52% (95% C.I. 0.3%-0.9%) in 2012 to 2.7% (95% C.I. 2.0%-3.8%) in 2013, but no trend of increase was observed in four-years of investigation. The pathogen was detected from 2.2% (95% C.I. 1.9%-2.6%) of bulk tank milk and from 0.5% (95% C.I. 0.3%-0.8%) of milk for vending machine. A significative difference ($p < 0.05$) of the prevalence data was observed between data collected in two different regions of Northern Italy with an higher prevalence in Lombardy. In addition to the geographical area, the *L. monocytogenes* presence was influenced also by the seasonal period of collection samples, with peaks in spring and autumn. These results confirm the raw milk can be a source of foodborne illness outbreaks if consumed without sanitizing treatments, but the low prevalence and the low contamination levels (more than 80% of the contaminated samples contained <10 cfu ml⁻¹ of *L. monocytogenes*) proving the hygienic quality of the milk produced in Northern Italy.

Key words: Raw milk; incidence; *Listeria monocytogenes*; risk analysis.

1. Introduction

Milk and dairy products are assumed to be implicated in 2-6% of the bacterial foodborne diseases (De Buyser et al., 2001) and, in particular, the presence of foodborne pathogens in raw milk and bulk tank milk has been widely reported (D'Amico and Donnelly, 2010; De Reu et al., 2004; Gaya et al., 1996; Hassan et al., 2000; Jayarao and Henning, 2001; Kousta et al., 2010; Moshtaghi and Mohamadpour, 2007; Van Kessel et al., 2004; Waak et al., 2002). The prevalence of pathogens in milk is influenced by numerous factors such as farm size, number of animals on the farm, hygienic conditions, farm management practices, variation in sampling and types of samples evaluated, differences in detection methodologies used, geographical location, and season. However, in spite of the variation, all of these surveys clearly demonstrated that milk can be a major source of foodborne pathogens of human health significance (Oliver et al., 2005).

In this context, *Listeria monocytogenes*, responsible of listeriosis, represents one of the most serious food safety concerns. It has been isolated from many foods (Gianfranceschi et al., 2003), among which milk and cheeses (Bernini et al., 2013; Dalmaso and Jordan, 2014; Manfreda et al., 2005; Pintado et al., 2005; Torres-Vitela et al., 2012) and it has been involved in numerous outbreaks occurring after consumption of contaminated milk and milk products throughout the world (Linnan et al., 1988; Lyytikainen et al., 2000; Donnelly, 2001; Lunde'n et al., 2004). In particular, dairy products contaminated with *L. monocytogenes* have been implicated at almost the half of the reported listeriosis outbreaks in Europe (Lunden, Tolvanen & Korkeala, 2004). The pathogen is widespread in nature and lives naturally in plants and soil environments. Its ability to survive and grow over a wide range of environmental conditions, including refrigeration temperatures, high salt concentration and low pH, makes it a potential hazard in foods (Ryser, 2007). Moreover, *L. monocytogenes* into food processing plants results in reservoirs that are difficult to eradicate: this is the case of

biofilms that are a constant issue in food processing environments (Oliver et al., 2005). So, in addition to the risk associated to direct consumption, the introduction of raw contaminated milk into dairy processing plants represents a risk to human health if milk is used unpasteurized for cheese making or in case of cross contamination (Kousta et al., 2010). Considering the threat represented by the pathogen in raw milk, a survey was conducted from January 2010 to September 2013, involving a large number of samples collected in different geographical area in the North of Italy. This research aimed to give a considerable overview of *L. monocytogenes* presence in raw cow milk, intended both for cheese making and for direct consumption, by evaluating the prevalence of the pathogen at farm level also in relation to the seasonality.

2. Materials and methods

2.1. Samples collection

A total of 8716 raw cow milk samples were collected from January 2010 to September 2013 in 942 farms located in Lombardy and Emilia-Romagna regions, Northern Italy. All the samples, consisting in 5897 samples of bulk tank milk intended for cheese making and 2819 samples of milk intended for sale in automatic vending machines, were collected into sterile containers, kept below 4 °C during transportation and analysed within 2 h after receipt. Samples were collected in the frame of Food Business Operator's self-control programs or in the frame of monitoring surveys officers of the Regional Veterinary Authority.

*2.2. Detection and enumeration of *L. monocytogenes**

All samples were tested for the presence of *L. monocytogenes* on 25 ml of raw cow milk by means of qualitative methods.

The samples collected from 2010 to 2011 were examined qualitatively according to ISO 11290-1 (ISO, 1996). The samples collected from 2012 to 2013 were examined by a biomolecular method (real-time PCR) (Biorad AFNOR BRD 07/ 10-04/05) (AFNOR, 2004) to detect *L. monocytogenes* DNA. Samples testing positive were retested under a microbiological protocol according to ISO 11290-1 (ISO, 1996). Typical colonies (n = 5) presumed to be *Listeria* spp. were streaked from Agar Listeria acc. to Ottaviani & Agosti (ALOA) (Biolife Italiana, Teramo, Italy) supplemented with ALOA enrichment-selective supplements (Biolife Italiana) onto Tryptone Soya Yeast Extract Agar, TSYEA (Oxoid, Basingstoke, UK) and plates were incubated at 37 °C for 24 h. By following the Gram's staining, catalase reaction and tumbling motility were performed using the pure cultures obtained from TSYEA. The isolates resulted positive to phenotypic tests were inoculated on 5% sheep blood agar (Oxoid) to determine the Beta haemolytic reaction. For following confirmation, carbohydrate utilization and CAMP tests were performed.

On samples found to be positive, the enumeration of *L. monocytogenes* was carried out according to the method described by ISO 11290-2 (ISO, 1998) to evaluate the prevalence of contamination level.

2.3. Data analyses

The prevalence of *L. monocytogenes* in raw milk, calculated as proportion between positive samples on total sample, was expressed in percentage values. Statistical analysis was performed by Epi tools (<http://epitools.ausvet.com.au>): the confidence intervals (C.I.) of proportions were calculated with using the binomial exact method and the statistical significance of differences between proportions was evaluated by Chi-square (χ^2) test.

3. Results and discussion

From 2010 to 2013, 8716 raw milk samples, intended both for cheese making and for vending machine, were collected in Northern Italy. Samples were taken from local farms within self-control sampling programs and by the official veterinarians within state surveillance programs and were investigated for the presence of *L. monocytogenes*.

The results are summarized in Table 1. The prevalence values in raw milk ranged from 0.52% (95% C.I. 0.3%-0.9%) in 2012 to 2.73% (95% C.I. 2.0%-3.8%) in 2013, but no trend of increase was observed in four-year investigation. Overall, *L. monocytogenes* was detected in 145 raw milk samples out of 8716, with a prevalence of 1.66% (95% C.I. 1.4%-2.0%). This result is mainly due ($p > 0.05$) to bulk tank milk contamination rather than to raw milk intended for vending machine. In fact, concerning bulk tank milk, *L. monocytogenes* was found in 2.22% (95% C.I. 1.9%-2.6%) of samples (131/5897) in four years of survey (Table 2).

These findings were in agreement with those reported in several studies carried out internationally and recently published, in which the prevalence of *L. monocytogenes* in raw milk has ranged from ‘not detected’ to 7.1%. In particular, the pathogen was “not detected” in Norway in 2011 (Jakobsen et al., 2011), and to the extent of 0.68% in New Zealand in 2012 (Hill et al., 2012), 2.12% in Turkey in 2006 (Aygün & Pehlivanlar, 2006), 2.61% in Algeria in 2007 (Hamdi et al., 2007), 5.5% in Finnish in 2013 (Ruusunen et al., 2013), 6.1% in North-West Spain in 2007 (Vilar et al., 2007) and 7.1% in USA in 2011 (Van Kessel et al., 2011).

Previously, in 2005, the prevalence of *L. monocytogenes* in bulk tank milk has been reported to range from 1 to 12% (Oliver et al., 2005), therefore, a reduction of contamination samples seems to have been monitored in the last years. Anyway, the above data collected in different world areas underline the wide variability of the *L. monocytogenes* prevalence, which can be due to several factors, as geographical area, size farm, types of housings for the cattle. As further confirmation of this variability, the overall data we collected in this study show that even in the same geographical area there may be a different prevalence of *L. monocytogenes*.

In Lombardy region, the pathogen was detected in 96 (2.6%; 95% C.I. 2.1%-3.1%) of 3721 samples analyzed and the prevalence was statistically highest ($p < 0.05$) compared to that found in Emilia-Romagna region, where *L. monocytogenes* was detected in 26 (1.2%; 95% C.I. 0.8%-1.8%) of 2176 milk samples (data not shown). The different prevalence values can be due to the different treatment of the cattle usually practiced in the two Italian regions. In Lombardy, the most of the cattle is fed with silages, the housing of cattle is indoors and these practices, when combined with poor hygiene on the farm, may contribute to contamination of milk, in agreement with Husu et al. (1990) and Sanaa et al. (1993). In contrast with this, many researchers have identified the raw milk as a source of *L. monocytogenes*, but environmental and fecal contamination during the transportation and storage of milk have also been reported (Frece et al., 2010). Moreover, *L. monocytogenes* may also directly contaminate milk from animals with mastitis (Hird & Genigeorgis, 1990).

A low prevalence of *L. monocytogenes* was observed over the four years survey in raw milk samples indented for vending machine (Table 2). The pathogen was detected in 0.50% (95% C.I. 0.03%-0.8%) of samples (14/2819), but the prevalence ranged from 0% (95% C.I. 0%-0.7%) in 2010 to 1.57% (95% C.I. 0.7%-3.1%) in 2011. A similar result was obtained by Bianchi et al. (2013) who detected the *L. monocytogenes* in 1.6% of the milk samples for vending machine during the monitoring survey from 2009 to 2011 in another Italian region (Piedmont region, near Lombardy).

L. monocytogenes is widely extended throughout the environment (Fenlon et al., 1996). This pathogen has the ability to survive in stress conditions and it is able to grow at low temperatures in several food types as cooked meat (Daminelli et al., 2014), vegetables (Sant'Ana et al., 2012), cold-smoked salmon (Beaufort et al., 2007), milk and cheese (Schvartzman et al., 2010). To prevent the growth of the pathogen in raw milk, in Italy, where the sale and the distribution of unpacked raw milk via automatic self-service vending

175 machines was authorized since 2007, the product must be maintained at constant temperature
176 between 0 °C and 4 °C, and the customers are instructed to boil the milk before consumption.
177 Even if the most important aspect remains the hygienic quality of the product, these practices
178 can contribute to improve the safety of the raw milk, as shown the low prevalence of *L.*
179 *monocytogenes* reported in the present study.

180 To evaluate the seasonal influence on the *L. monocytogenes* presence in bulk tank milk, in
181 Table 3 were reported the prevalence data for each year (from 2010 to 2013), broken down by
182 each month. In four years, only in January on a total of 188 samples analyzed, milk samples
183 positive for *L. monocytogenes* have never been found. Considering the seasonal variability,
184 the *L. monocytogenes* prevalence was statistically lower ($p < 0.05$) during the winter season,
185 with a prevalence of 0.8% (95% C.I. 0.3%-1.7%), in contrast with the spring prevalence of
186 3.04% (95% C.I. 2.3%-3.9%), the summer prevalence of 1.91% (95% C.I. 1.4%-2.6%) and
187 the autumn prevalence of 2.33% (95% C.I. 1.6%-3.3%) (Table 3). Previous studies on
188 *Listeria* spp. prevalence in raw milk reported some evidence of seasonal variation. Atil et al.
189 (2011) in eastern Turkey observed a high prevalence in spring and winter; in France, Meyer-
190 Broseta et al. (2003) reported peaks in winter. Ryser (1999) reported that seasonal variations
191 in *Listeria* prevalence may be related to silage feeding, with higher prevalence in months
192 when silage is fed to animals. In fact, *L. monocytogenes* could be present in silage, in which
193 the pathogen can multiply if the silage has been inadequately fermented (pH above 5.0 to 5.5)
194 (Husu, 1990). Seasonal differences in *L. monocytogenes* prevalence were observed in our
195 monitoring, but more studies are needed to determine the validity and the causes of these
196 differences. In fact, in the present study, the reduced number of samples found positive to *L.*
197 *monocytogenes* in the winter season may be due to the lower number of samples analyzed
198 (746 samples against over 1000 samples analyzed in each of the other seasons), or to other

variables such as the type of cattle feed (fresh grass or silage), the hygienic conditions of breeding or the presence of cattle infected with mastitis. The contamination levels of *L. monocytogenes* in bulk tank milk samples are reported in Figure 1 as percentage of samples with pathogen concentration variable between <1 and >1000 cfu ml⁻¹. Overall, more than 80% of the contaminated samples contained <10 cfu ml⁻¹ of *L. monocytogenes*. This trend is reflected from 2010 to 2012, while in 2013 the majority of the positive samples contained <1 cfu ml⁻¹ of the pathogen, as shown in Figure 1. It is difficult to compare the results from this study with those conducted in other countries, because most studies express the results only qualitatively. However, a study by Meyer-Broseta et al. (2003), shows that even in France, the level of contamination of bulk tank milk was very low, generally less than 1 cfu ml⁻¹ of the *L. monocytogenes*.

4. Conclusions

During 2007-2009 in Italy there was an increase of notifications of listeriosis with the most cases are reported in the Centre-North of Italy. This is probably attributable both to a real increase of listeriosis in Italy and to surveillance implementation (Pontello et al., 2012). However, statistically significant increasing trends in listeriosis notification rates from 2005 to 2009 were noted in Italy as elsewhere in Europe (EFSA/ECDC, 2011). In the present study the raw milk was found positive to the presence of *L. monocytogenes*, confirming the milk as potential source of the food borne disease, but the low prevalence and the reduced level of pathogen concentration, when present, can highlight the hygienic quality of the milk produced in Northern Italy. Furthermore, the collected data in the present study can be a useful tool for the quantitative risk assessment study for human listeriosis linked to the consumption of raw milk and cheese made from raw milk in Italy. Furthermore, the collected data in the present

study can be a useful tool for the quantitative risk assessment study for human listeriosis linked to the consumption of raw milk and cheese made from raw milk in Italy.

Disclosure

Authors declare that no conflict of interests exists.

Acknowledgements

We sincerely thank Dr. Silvia Todeschi (Food Microbiology Department, IZSLER, Brescia, Italy) for the technical assistance and for the informatics support development to the data management.

References

- AFNOR. (2004). Biorad AFNOR BRD 07/6-07/04. <http://www.afnor.validation.org/pdf/AFNOR-Validation2008.pdf>. Accessed 26 June 2014.
- Atil, E., Ertas, H. B., & Ozbey, G. (2011). Isolation and molecular characterization of *Listeria* spp. from animals, food and environmental samples. *Veterinari Medicina*, 56(8), 386-394.
- Aygun, O., & Pehlivanlar, S. (2006). *Listeria* spp. in the raw milk and dairy products in Antakya, Turkey. *Food Control*, 17, 676-679.
- Beaufort, A., Rudelle, S., Gnanou-Besse, N., Toquin, M. T., Kerouanton, A., Bergis, H., ... & Cornu, M. (2007). Prevalence and growth of *Listeria monocytogenes* in naturally contaminated cold-smoked salmon. *Letters in Applied Microbiology*, 44, 406-411.
- Bernini, V., Bottari, B., Dalzini, E., Sgarbi, E., Lazzi, C., Neviani, E., & Gatti, M. (2013). The presence, genetic diversity and behaviour of *Listeria monocytogenes* in blue-veined cheese rinds during the shelf life. *Food Control*, 34, 323-330.

247 Bianchi, D. M., Barbaro, A., Gallina, S., Vitale, N., Chiavacci, L., Caramelli, M., &
 248 Decastelli, L. (2013). Monitoring of foodborne pathogenic bacteria in vending machine
 249 raw milk in Piedmont, Italy. *Food Control*, 32, 435-439.

250 Dalmasso, M. & Jordan, K. (2014). Absence of growth of *Listeria monocytogenes* in naturally
 251 contaminated Cheddar cheese. *Journal of Dairy Research*, 81, 46-53.

252 D'Amico, D. J., & Donnelly, C. W. (2010). Microbiological quality of raw milk used for
 253 small-scale artisan cheese production in Vermont: effect of farm characteristics and
 254 practices. *Journal of Dairy Science*, 93, 134-147.

255 Daminelli, P., Dalzini E., Cosciani-Cunico, E., Finazzi, G., D'Amico, S. & Losio, M.N.
 256 (2014). Prediction of the maximal growth rate of *Listeria monocytogenes* in sliced
 257 mortadella by the square root type model. *Italian Journal of Food Science*, 26, 261-267.

258 De Buyser, M. L., Dufour, B., Maire, M., Lafarge, V. (2001). Implication of milk and milk
 259 products in food-borne diseases in France and in different industrialised countries.
 260 *International Journal of Food Microbiology*, 67, 1-17.

261 De Reu, K., Grijspeerdt, K., & Herman, L. (2004). A Belgian survey of hygiene indicator
 262 bacteria and pathogenic bacteria in raw milk and direct marketing of raw milk farm
 263 products. *Journal of Food Safety*, 24, 17-36.

264 Donnelly, C. (2001). Factors associated with hygienic control and quality of cheeses prepared
 265 from raw milk: a review. *Bulletin of the International Dairy Federation*, 369, 16-27.

266 European Food Safety Authority/European Centre for Disease Prevention and Control
 267 (EFSA/ECDC). The European Union summary report on trends and sources of zoonoses,
 268 zoonotic agents and food-borne outbreaks in 2009. *EFSA Journal* 2011;9(3):2090.
 269 <http://dx.doi.org/10.2903/j.efsa.2011.2090>

270 Fenlon, D.R., Wilson, J., Donachie, W., 1996. The incidence and level of *Listeria*
 271 *monocytogenes* contamination of food sources at primary production and initial processing.
 272 *Journal of Applied Bacteriology*, 81, 641-650.

273 Frece J, Markov K, Cvek D, Kolarec K, Delas F (2010): Comparison of conventional and
 274 molecular methods for the routine confirmation of *Listeria monocytogenes* in milk
 275 products produced domestically in Croatia. *Journal of Dairy Research*, 77, 112–116.

276 Gaya, P., Saralegui, C., Medina, M., & Nunez, M. (1996). Occurrence of *Listeria*
 277 *monocytogenes* and other *Listeria* spp. in raw caprine milk. *Journal of Dairy Science*, 79,
 278 1936-1941.

279 Gianfranceschi, M., Gattuso, A., Tartaro, S., & Aureli, P. (2003). Incidence of *Listeria*
 280 *monocytogenes* in food and environmental samples in Italy between 1990 and 1999:
 281 Serotype distribution in food, environmental and clinical samples. *European Journal of*
 282 *Epidemiology*, 18, 1001-1006.

283 Hamdi, T. M., Naïm, M., Martin, P., & Jacquet, C. (2007). Identification and molecular
 284 characterization of *Listeria monocytogenes* isolated in raw milk in the region of Algiers
 285 (Algeria). *International Journal of Food Microbiology*, 116, 190-193.

286 Hassan, L., Mohammed H. O., McDonough P. L., & Gonzalez, R. N. (2000). A cross-
 287 sectional study on the prevalence of *Listeria monocytogenes* and Salmonella in New York
 288 dairy herds. *Journal of Dairy Science*, 83, 2441-2447.

289 Hill, B., Smythe, B., Lindsay, D., & Shepherd, J. (2012). Microbiology of raw milk in New
 290 Zealand. *International Journal of Food Microbiology*, 157, 305-308.

291 Hird, D. W., & Genigeorgis, C. (1990). Listeriosis in food animals: clinical signs and
 292 livestock as a potential source of direct (non-foodborne) infection for humans. In A. J.
 293 Miller, J. L. Smith, & G. A. Somkuti (Eds.), Foodborne Listeriosis (pp. 31–39).
 294 Amsterdam: Elsevier Science Publishers.

295 Husu, J. R. (1990). Epidemiological studies on the occurrence of *Listeria monocytogenes* in
 296 the feces of dairy cattle. *Journal of Veterinary Medicine*, 37, 276-282.

297 Husu, J.R., Seppänen, J.T., Sivelä, S.K., Rauramaa, A.L., 1990. Contamination of raw milk
 298 by *Listeria monocytogenes* on dairy farms. *Journal of Veterinary Medicine*, 37, 268–275.

299 ISO 11290-1 (International Standards Organization), 1996/Amd. 1:2004 (E). Microbiology of
 300 food and animal feeding stuffs — Horizontal method for the detection and enumeration of
 301 *Listeria monocytogenes* — Part 1: Detection method. International Organization for
 302 Standardization, Geneva, Switzerland.

303 ISO 11290-2 (International Standards Organization), 1998/Amd. 1:2004 (E) Microbiology of
 304 Food and Animal Feeding Stuffs — Horizontal Method for the Detection and Enumeration
 305 of *Listeria monocytogenes* — Part 2: Enumeration Method. International Organization for
 306 Standardization, Geneva, Switzerland.

307 Jakobsen, R. A., Heggebø, R., Sunde, E. B., & Skjervheim, M. (2011). *Staphylococcus aureus*
 308 and *Listeria monocytogenes* in Norwegian raw milk cheese production. *Food*
 309 *Microbiology*, 28, 492-496.

310 Jayarao, B. M., & Henning D. R. (2001). Prevalence of foodborne pathogens in bulk tank
 311 milk. *Journal of Dairy Science*, 84, 2157-2162.

312 Kousta, M., mataragas, M., Skandamis, P., & Drosinos, E. H. (2010). Prevalence and sources
 313 of cheese contamination with pathogens at farm and processing levels. *Food Control*, 21,
 314 805-815.

315 Linnan, M. J., Nascola, L., Lou, X. D., Goulet, V., May, S., Salminen, C., et al. (1988).
 316 Epidemic listeriosis associated with Mexican style cheese. *New England Journal of*
 317 *Medicine*, 319, 823-824.

318 Lundén J, Tolvanen R, & Korkeala H. (2004). Human listeriosis outbreaks linked to dairy
 319 products in Europe. *Journal of Dairy Science*, 87, E6-E11.

320 Lyytikäinen, O., Autio, T., Maijala, R., Ruutu, P., Honkanen-Buzalski, T., Miettinen, M., et
321 al. (2000). An outbreak of *Listeria monocytogenes* serotype 3a infections from butter in
322 Finland. *The Journal of Infectious Diseases*, 181, 1838-1841.

323 Manfreda, G., De Cesare, A., Stella, S., Cozzi, M., & Cantoni, C. (2005). Occurrence and
324 ribotypes of *Listeria monocytogenes* in Gorgonzola cheeses. *International Journal of*
325 *Food Microbiology*, 102, 287-293.

326 Meyer-Broseta, S., Diot, A., Bastian, S., Rivière, J., & Cerf, O. (2003). Estimation of low
327 bacterial concentration: *Listeria monocytogenes* in raw milk. *International journal of food*
328 *microbiology*, 80, 1-15.

329 Moshtaghi, H., & Mohamadpour, A. A. (2007). Incidence of *Listeria* spp. in raw milk in
330 Shahrekord, Iran. *Foodborne Pathogens Disease*, 4, 107-110.

331 Oliver, S. P., Jayarao, B. M., & Almeida, R. A. (2005). Foodborne pathogens in milk and the
332 dairy farm environment: food safety and public health implications. *Foodborne Pathogens*
333 *& Disease*, 2, 115-129.

334 Pintado, C. M. B. S., Oliveira, A., Pampulha, M. E., & Ferriera, M. A. S. S. (2005).
335 Prevalence and characterization of *Listeria monocytogenes* isolated from soft cheese.
336 *Food Microbiology*, 22, 79-85.

337 Pontello, M., Guaita, A., Sala, G., Cipolla, M., Gattuso, A., Sonnessa, M., & Gianfranceschi,
338 M. V. (2012). *Listeria monocytogenes* serotypes in human infections (Italy, 2000-2010).
339 *Annali dell'Istituto Superiore di Sanità*, 48, 146-150.

340 Ruusunen, M., Salonen, M., Pulkkinen, H., Huuskonen, M., Hellström, S., Revez, J., ... &
341 Lindström, M. (2013). Pathogenic bacteria in Finnish bulk tank milk. *Foodborne*
342 *Pathogens and Disease*, 10, 99-106.

343 Ryser, E. T. (2007). Incidence and behavior of *Listeria monocytogenes* in cheese and other
 344 fermented dairy products. In Ryser, E. T., & Marth, E. H. (Eds) *Listeria*, Listeriosis and
 345 Food Safety. New York: Marcel Dekker Inc., pp. 405-453.

346 Ryser, E.T. (1999) Incidence and behaviour of *Listeria monocytogenes* in unfermented dairy
 347 products. In *Listeria*, Listeriosis and Food Safety ed. Ryser, E.T. and Marth, E.H. pp.
 348 359–409. New York: Marcel Dekker, Inc.

349 Sanaa, M., Poutrel, B., Menard, J.L., Serieys, F., 1993. Risk factors associated with
 350 contamination of raw milk by *Listeria monocytogenes* in dairy farms. *Journal of Dairy*
 351 *Science*, 76, 2891–2898.

352 Sant'Ana, A. S., Barbosa, M. S., Destro, M. T., Landgraf, M., & Franco, B. D. (2012). Growth
 353 potential of *Salmonella* spp. and *Listeria monocytogenes* in nine types of ready-to-eat
 354 vegetables stored at variable temperature conditions during shelf-life. *International*
 355 *Journal of Food Microbiology*, 157, 52-58.

356 Schwartzman, M. S., Belessi, X., Butler, F., Skandamis, P., & Jordan, K. (2010). Comparison
 357 of growth limits of *Listeria monocytogenes* in milk, broth and cheese. *Journal of applied*
 358 *microbiology*, 109, 1790-1799.

359 Torres-Vitela, M. R., Mendoza-Bernardo, M., Castro-Rosas, J., Gomez-Aldapa, C. A., Garay-
 360 Martinez, L. E., Navarro-Hidalgo, V., & Villarruel-López, A. (2012). Incidence of
 361 *Salmonella*, *Listeria monocytogenes*, *Escherichia coli* O157:H7, and Staphylococcal
 362 enterotoxin in two types of Mexican fresh cheeses. *Journal of Food Protection*, 75, 79-84.

363 Van Kessel, J. A. S., Karns, J. S., Lombard, J. E., & Kopral, C. A. (2011). Prevalence of
 364 *Salmonella enterica*, *Listeria monocytogenes*, and *Escherichia coli* virulence factors in
 365 bulk tank milk and in-line filters from US dairies. *Journal of Food Protection*, 74, 759-
 366 768.

367 Van Kessel, J. S., Karns, J. S., Gorski, L., McCluskey, B. J., & Perdue, M. L. (2004).
368 Prevalence of Salmonellae, *Listeria monocytogenes*, and fecal coliforms in bulk tank milk
369 on US dairies. *Journal of Dairy Science*, 87, 2822-2830.

370 Vilar, M. J., Yus, E., Sanjuán, M. L., Diéguez, F. J., & Rodríguez-Otero, J. L. (2007).
371 Prevalence of and Risk Factors for Listeria Species on Dairy Farms. *Journal of dairy*
372 *science*, 90, 5083-5088.

373 Waak, E., Tham, W., & Danielsson-Tham M. L. (2002). Prevalence of *Listeria*
374 *monocytogenes* strains isolated from raw whole milk in farm bulk tanks and dairy plants
375 receiving tanks. *Applied and Environmental Microbiology*, 68, 3366-3370.

376

Table 1. Detection and prevalence of *L. monocytogenes* in raw milk collected in Northern Italy according to the year of sampling.

Year	Samples	Positive for <i>L. monocytogenes</i> (%)
2010	1728	20 (1.16%)
2011	3150	76 (2.41%)
2012	2519	13 (0.52%)
2013	1319	36 (2.73%)
Total	8716	145 (1.66%)

Table 2. Detection and prevalence of *L. monocytogenes* in different categories of raw milk collected in Northern Italy according to the year of sampling.

Year	Bulk Tank Milk		Raw Milk for vending machine	
	Samples	Positive for <i>L. monocytogenes</i> (%)	Samples	Positive for <i>L. monocytogenes</i> (%)
2010	1176	20 (1.70%)	552	0
2011	2639	68 (2.58%)	511	8 (1.57%)
2012	1317	9 (0.68%)	1202	4 (0.33%)
2013	765	34 (4.44%)	554	2 (0.36%)
Total	5897	131 (2.22%)	2819	14 (0.50%)

401 **Table 3.** Seasonal detection and prevalence of *L. monocytogenes* in bulk tank milk collected in Northern Italy according to the year of sampling.


402

Season	Month	Sample positive / Sample analyzed (%)					
		2010	2011	2012	2013	Total for Month	Total for Season
Winter	December	0/58 (0%)	2/106 (1.89%)	0/45 (0%)	nc	2/209 (0.96%)	6/746 (0.8%)
	January	0/47 (0%)	0/49 (0%)	0/70 (0%)	0/22 (0%)	0/188 (0%)	
	February	0/108 (0%)	1/51 (1.96%)	3/158 (1.9%)	0/32 (0%)	4/349 (1.15%)	
Spring	March	2/174 (1.15%)	7/233 (3%)	0/215 (0%)	21/212 (9.91%)	30/834 (3.6%)	56/1844 (3.04%)
	April	0/74 (0%)	8/225 (3.56%)	0/79 (0%)	12/167 (7.19%)	20/545 (3.67%)	
	May	1/112 (0.89%)	5/218 (2.29%)	0/76 (0%)	0/59 (0%)	6/465 (1.29%)	
Summer	June	4/84 (4.76%)	5/297 (1.68%)	0/80 (0%)	0/73 (0%)	9/534 (1.69%)	37/1934 (1.91%)
	July	2/116 (1.72%)	8/463 (1.73%)	5/212 (2.36%)	1/148 (0.68%)	16/939 (1.7%)	
	August	2/129 (1.55%)	10/121 (8.26%)	0/162 (0%)	0/49 (0%)	12/461 (2.6%)	
Autumn	September	6/119 (5.04%)	12/485 (2.47%)	1/154 (0.65%)	0/3 (0%)	19/761 (2.5%)	32/1373 (2.33%)
	October	3/97 (3.09%)	5/273 (1.83%)	0/35 (0%)	nc	8/405 (1.98%)	
	November	0/58 (0%)	5/118 (4.24%)	0/31 (0%)	nc	5/207 (2.42%)	

nc: data not collected.

403

Figure 1. Counts of *L. monocytogenes* in bulk tank milk according the year of sampling, expressed as percentages of samples containing colony numbers ranging from < 1 to > 1000 cfu ml⁻¹.



Highlights

1. Survey study of *Listeria monocytogenes* in raw cow mil in Northern Italy.
2. Low level of milk contamination with *L. monocytogenes*.
3. Seasonal differences in *L. monocytogenes* prevalence (lower in winter).
4. Overall, more than 80% of the contaminated samples contained low *L. monocytogenes* concentration (<10 cfu ml⁻¹).