

University of Parma Research Repository

Palaeontological characterisation and analysis of the AND-2A core, ANDRILL southern McMurdo sound project, Antarctica

This is the peer reviewd version of the followng article:

Original

Palaeontological characterisation and analysis of the AND-2A core, ANDRILL southern McMurdo sound project, Antarctica / Taviani, M; Hannah, M; Harwood, D. M; Ishman, S. E; Johnson, K; Olney, M; Riesselman, C; Tuzzi, E; Askin, R; Beu, A. G; Blair, S; Cantarelli, V; Ceregato, A; Corrado, S; Mohr, B; Nielsen, S. H. H; Persico, Davide; Petrushak, S; Raine, J. I; Warny, S.. - In: TERRA ANTARTICA. - ISSN 1122-8628. - 15:1-2(2008), pp. 113-146.

Availability: This version is available at: 11381/2819991 since: 2016-12-01T12:12:18Z

Publisher:

Published DOI:

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

### University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

ANDRILL Research and Publications

Antarctic Drilling Program

2008

## Palaeontological Characterisation and Analysis of the AND-2A Core, ANDRILL Southern McMurdo Sound Project, Antarctica

M. Taviani Istituto di Scienze Marine (ISMAR)-CNR, marco.taviani@bo.ismar.cnr.it

M. Hannah Victoria University of Wellington

D. M. Harwood University of Nebraska at Lincoln, dharwood1@unl.edu

S. E. Ishman Southern Illinois University

K. Johnson Ohio State University

See next page for additional authors

Follow this and additional works at: http://digitalcommons.unl.edu/andrillrespub Part of the <u>Environmental Indicators and Impact Assessment Commons</u>

Taviani, M.; Hannah, M.; Harwood, D. M.; Ishman, S. E.; Johnson, K.; Olney, M.; Riesselman, C.; Tuzzi, E.; Beu, A. G.; Blair, S.; Cantarelli, V.; Ceregato, A.; Corrado, S.; Mohr, B.; Nielson, S. H.H.; Persico, D.; Petrushak, S.; Raine, J. I.; Warny, S.; and ANDRILL-SMS Science Team, "Palaeontological Characterisation and Analysis of the AND-2A Core, ANDRILL Southern McMurdo Sound Project, Antarctica" (2008). *ANDRILL Research and Publications*. Paper 41. http://digitalcommons.unl.edu/andrillrespub/41

This Article is brought to you for free and open access by the Antarctic Drilling Program at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in ANDRILL Research and Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

#### Authors

M. Taviani, M. Hannah, D. M. Harwood, S. E. Ishman, K. Johnson, M. Olney, C. Riesselman, E. Tuzzi, A. G. Beu, S. Blair, V. Cantarelli, A. Ceregato, S. Corrado, B. Mohr, S. H.H. Nielson, D. Persico, S. Petrushak, J. I. Raine, S. Warny, and ANDRILL-SMS Science Team

## Palaeontological Characterisation and Analysis of the AND-2A Core, ANDRILL Southern McMurdo Sound Project, Antarctica

M. TAVIANI<sup>1\*</sup>, M. HANNAH<sup>2</sup>, D.M. HARWOOD<sup>3</sup>, S.E. ISHMAN<sup>4</sup>, K. JOHNSON<sup>5</sup>, M. OLNEY<sup>6</sup>,
C. RIESSELMAN<sup>7</sup>, E. TUZZI<sup>3</sup>, R. ASKIN<sup>8</sup>, A.G. BEU<sup>9</sup>, S. BLAIR<sup>10</sup>, V. CANTARELLI<sup>11</sup>,
A. CEREGATO<sup>1</sup>, S. CORRADO<sup>11</sup>, B. MOHR<sup>12</sup>, S.H.H. NIELSEN<sup>10</sup>, D. PERSICO<sup>13</sup>,
S. PETRUSHAK<sup>10</sup>, J.I. RAINE<sup>9</sup>, S. WARNY<sup>14</sup> & THE ANDRILL-SMS SCIENCE TEAM<sup>15</sup>

<sup>1</sup>Istituto di Scienze Marine (ISMAR)-CNR, v. Gobetti 101, I-40129 Bologna – Italy
<sup>2</sup>Antarctic Research Centre, Victoria University of Wellington, P.O. Box 600, Wellington – New Zealand
<sup>3</sup>Dept. of Geosciences, ANDRILL SMO, 214 Bessey Hall, University of Nebraska, Lincoln, NE 68588-0340 – USA
<sup>4</sup>Dept. of Geology, Southern Illinois University, 1259 Lincoln Drive, Carbondale, IL 62901-6899 – USA
<sup>5</sup>School of Earth Sciences, The Ohio State University, 125 South Oval Mall, Columbus OH 43210 – USA
<sup>6</sup>University of South Florida, Dept. of Geology, 4202 E. Fowler Ave., SCA 528 Tampa, FL 33620-8100 – USA
<sup>7</sup>School of Earth Sciences, Stanford University, Braun Hall, Bldg. 320, Stanford, CA 94305-2115 – USA
<sup>8</sup>1930 Bunkhouse Dr., Jackson, WY 83001 – USA
<sup>9</sup>Paleontology & Environmental Change Section, GNS Science, 1 Fairway Drive, Lower Hutt 5040 – New Zealand

<sup>19</sup>Aleontology & Environmental Change Section, GNS Science, 1 Fairway Drive, Lower Hutt 5040 – New Zealand
<sup>10</sup>Antarctic Marine Geological Research Facility, Florida State University, Tallahassee, FL 32306-4100 – USA
<sup>11</sup>Dip. di Scienze Geologiche, Univ. degli Studi Roma Tre, Largo S. Leonardo Murialdo, 1, I-0014 Rome – Italy
<sup>12</sup>Naturhistorisches Forschungsinstitut, invalidenstraße 43 D-10115, Berlin – Germany
<sup>13</sup>Dip. di Scienze della Terra, Università degli Studi di Parma, V.le G.P. Usberti 157A, I-40100 Parma – Italy
<sup>14</sup>Dept. of Geology and Geophysics, Louisiana State University, Baton Rouge, LA 70803 – USA
<sup>15</sup> http://andrill.org/projects/sms/team.htm<sup>1</sup>

\*Corresponding author (marco.taviani@bo.ismar.cnr.it)

Abstract – The palaeontological yield of the 1138.54 metre-long AND-2A sedimentary rock core provides unique documentation of Neogene environments in the Ross Sea region of Antarctica. Especially important is the biological legacy of the climatically crucial 'mild' middle Miocene phase. Diatom-bearing units provide key information for stratigraphic intervals never previously recovered from locations proximal to the Antarctic continent and constrain the age model for the AND-2A core. Benthic calcareous (and agglutinated) foraminifera were present at many levels; remarkable is the occurrence of planktonic taxa only seldom found in the Neogene nearshore record of the Ross Sea region. The sporadic occurrence of calcareous dinoflagellate remains (thoracosphaerids) is consistent with warmer-than-present seawater during the Miocene. Marine palynomorphs are almost ubiquitous, although their abundance and diversity are variable. Pollen and spores from the middle Miocene section suggest a mossy tundra vegetation and represent the first stratigraphically-constrained record of terrestrial vegetation in Victoria Land during this time. Fragments of lignin-rich organic matter (huminite-vitrinite and inertinite groups) are particularly predominant during the Miocene climatic optimum, and continue into the Pliocene. Macrofossils are reasonably common throughout the core. Polychaete worm tubes were almost ubiquitous. Especially remarkable is the bivalve record (mainly pectinids), with 4-5 different taxa pointing out a mild climatic situation in the Miocene nearshore.

#### INTRODUCTION

Drilling operations for the Southern McMurdo Sound (SMS) Project were conducted in the austral spring 2007 to recover the AND-2A core at 77°44.27′.76″S and 165°17′18.24″E. This core comprises 1138.54 metres of various types of sedimentary rocks with some volcanics encompassing lower Miocene to Quaternary successions (Fielding et al., Fig. 1, this volume; Acton et al., this volume, b) (Fig. 1).

The Core Characterisation Phase of the SMS Project was conducted by an on-ice Palaeontology Team that included seven specialists of taxonomic groups of recognised biostratigraphic and environmental utility in the Antarctic Cenozoic, and an off-ice team who produced results reported herein after the drilling phase was completed. The palaeontologic record of the AND-2A core is highly remarkable in terms of quality and quantity of fossils, with over 1200 samples collected on-ice, including those for off-ice specialists. The on-ice treatment of such a large set of samples was possible through the collaborative work of additional SMS on-ice personnel (J. Carnes, R. Frisch-Gleason, J. Hamre, L. Jovane, K. Mankhoff, K. Pound, J. Reed, E. Strada, P.-N. Webb, and R. Williams). It is worth mentioning the important role provided by using thin-sections (Fig. 2) to identify fossils in the AND-2A core, that were otherwise undetectable through visual inspection or washing; these on-ice thin sections were provided by S. Petrushak. Sample aliquots from intervals of presumptive critical palaeoclimatic significance were

# © Terra Antartica Publication 2008-2009