



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

GPNA inhibits the sodium-independent transport system I for neutral amino acids

This is the peer reviewed version of the following article:

Original

GPNA inhibits the sodium-independent transport system I for neutral amino acids / Chiu, Martina; Sabino, Cosimo; Taurino, Giuseppe; Bianchi, Massimiliano; Andreoli, Roberta; Giuliani, Nicola; Bussolati, Ovidio. - In: AMINO ACIDS. - ISSN 0939-4451. - 49:8(2017), pp. 1365-1372. [10.1007/s00726-017-2436-z]

Availability:

This version is available at: 11381/2827725 since: 2021-10-26T10:13:42Z

Publisher:

Springer-Verlag Wien

Published

DOI:10.1007/s00726-017-2436-z

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

Amino Acids

GPNA inhibits the sodium-independent transport system L for neutral amino acids

--Manuscript Draft--

Manuscript Number:	AMAC-D-17-00013R2
Full Title:	GPNA inhibits the sodium-independent transport system L for neutral amino acids
Article Type:	Original Article
Corresponding Author:	Ovidio Bussolati, M.D., Ph.D. Universita degli Studi di Parma Parma, ITALY
Corresponding Author E-Mail:	ovidio.bussolati@unipr.it
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Universita degli Studi di Parma
Corresponding Author's Secondary Institution:	
First Author:	Martina Chiu, PhD
First Author Secondary Information:	
Order of Authors:	Martina Chiu, PhD Cosimo Sabino, BS Giuseppe Taurino, MS Massimiliano Giovanni Bianchi, PhD Roberta Andreoli, PhD Nicola Giuliani, MD, PhD Ovidio Bussolati, MD, PhD
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	<p>L-γ-glutamyl-p-nitroanilide (GPNA) is widely used to inhibit the glutamine transporter ASCT2, although it is known that it also inhibits other sodium-dependent amino acid transporters. In a panel of human cancer cell lines, which express the system L transporters LAT1 and LAT2, GPNA inhibits the sodium-independent influx of leucine and glutamine. The kinetics of the effect suggests that GPNA is a low affinity, competitive inhibitor of system L transporters. In Hs683 human oligodendroglioma cells, the incubation in the presence of GPNA, but not ASCT2 silencing, lowers the cell content of leucine. Under the same conditions the activity of mTORC1 is inhibited. Decreased cell content of branched chain amino acids and mTORC1 inhibition are observed in most of the other cell lines upon incubation with GPNA. It is concluded that GPNA hinders the uptake of essential amino acids through system L transporters and lowers their cell content.</p>
Response to Reviewers:	<p>Referee 1 Reviewer's comment: Overall, the authors have addressed each of the questions from my initial review, however this reviewer maintains that the manuscript would be far more beneficial to the research community with the simple addition of oocyte experiments to identify the transporter(s) involved. As it currently stands, the following line should be altered on page 3, since there are no data showing specific inhibition of LAT1 and LAT2: "Here we show that GPNA inhibits the sodium-independent influx of leucine and lowers its cell content, indicating that the inhibitor hinders the activity of system L transporters LAT1 and LAT2." Reply: We agree with the Reviewer but we feel that the identification of the sensitivity</p>

of each L transporter to GPNA, although very interesting, goes far beyond the main aim of our work and would significantly delay the publication of our results. We also agree that the sentence, in its present form, is somewhat overreaching and have modified it.

"Here we show that GPNA inhibits the sodium-independent influx of leucine and lowers its cell content, indicating that the inhibitor hinders the activity of system L."

Reviewer's comment: Glutamine uptake inhibition by siASCT2 should also be shown in Figure 6.

Reply: Glutamine uptake inhibition by siASCT2 is now shown in Figure 6, c. The results indicate that ASCT2 silencing significantly inhibits glutamine uptake and that the inhibition is exclusively attributable to the decrease of the sodium-dependent transport.

[Click here to view linked References](#)

1 Martina Chiu, Cosimo Sabino, Giuseppe Taurino, Massimiliano G. Bianchi, Roberta Andreoli, Nicola Giuliani, Ovidio
2 Bussolati
3

4 **GPNA inhibits the sodium-independent transport system L for neutral amino acids**
5
6

7
8 Martina Chiu, Cosimo Sabino, Giuseppe Taurino, Massimiliano G. Bianchi, Ovidio Bussolati[✉]

9 Laboratory of General Pathology

10 Department of Medicine and Surgery, University of Parma

11 Via Volturmo 39

12 43125 Parma

13 Italy
14
15
16

17
18 Roberta Andreoli

19 Unit of Occupational Medicine

20 Department of Medicine and Surgery, University of Parma
21
22
23

24 Nicola Giuliani

25 Myeloma Unit

26 Department of Medicine and Surgery, University of Parma

27 Hematology, BMT Center and CoreLab

28 Azienda Ospedaliero-Universitaria di Parma
29
30
31

32 Via Gramsci, 14

33 43121, Parma

34 Italy
35
36
37

38 [✉]Ovidio Bussolati, MD PhD

39 Laboratory of General Pathology

40 Department of Medicine and Surgery

41 University of Parma

42 Via Volturmo 39

43 43125 Parma
44
45
46

47 Italy

48 ovidio.bussolati@unipr.it

49 Phone: +39 0521 033783

50 Fax: +39 0521 033742
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

L- γ -glutamyl-p-nitroanilide (GPNA) is widely used to inhibit the glutamine transporter ASCT2, although it is known that it also inhibits other sodium-dependent amino acid transporters. In a panel of human cancer cell lines, which express the system L transporters LAT1 and LAT2, GPNA inhibits the sodium-independent influx of leucine and glutamine. The kinetics of the effect suggests that GPNA is a low affinity, competitive inhibitor of system L transporters. In Hs683 human oligodendroglioma cells, the incubation in the presence of GPNA, but not ASCT2 silencing, lowers the cell content of leucine. Under the same conditions the activity of mTORC1 is inhibited. Decreased cell content of branched chain amino acids and mTORC1 inhibition are observed in most of the other cell lines upon incubation with GPNA. It is concluded that GPNA hinders the uptake of essential amino acids through system L transporters and lowers their cell content.

Keywords: GPNA – Leucine – System L – LAT1 –ASCT2– Glutamine

Introduction

1
2 Glutamine uptake through the transporter ASCT2 has been found stimulated in many cancer models (Fuchs and Bode
3 2005). This metabolic feature has prompted various attempts to inhibit glutamine entry as a device to hinder cancer cell
4 proliferation. L- γ -Glutamyl-*p*-nitroanilide (GPNA) was proposed several years ago as an ASCT2 inhibitor (Esslinger et
5 al. 2005) and has been subsequently widely used to this purpose (Hassanein et al. 2015; Ren et al. 2015; Wang et al.
6 2015; Bolzoni et al. 2016; van Geldermalsen et al. 2016). However, GPNA selectivity for ASCT2 had never been
7 assessed in depth until, most recently, Broer et al. (Broer et al. 2016) definitely demonstrated that the compound
8 inhibits, besides ASCT2, other Na⁺-dependent carriers, such as several members of the SNAT family.
9

10 Besides ASCT2, also other transporters have been found overexpressed in human cancer cells *in vitro*, as well as in
11 primary and metastatic human tumors *in vivo*. In particular, two of these carriers, LAT1 (coded by *SLC7A5*) and LAT2
12 (coded by *SLC7A8*) are highly expressed in a wide array of human tumors (Fuchs and Bode 2005; Kaira et al. 2008;
13 Wang and Holst 2015; Barollo et al. 2016). LAT1 and LAT2, once complexed with the chaperone 4F2hc, account in
14 many tissues for the activity of System L, a sodium-independent, non electrogenic, exchange transport mechanism that
15 operates the transmembrane fluxes of most essential amino acids. LAT1 and LAT2 have comparable operational
16 features, both efficiently transport leucine, although LAT2 is endowed with a lower affinity for substrates (del Amo et
17 al. 2008), and, through leucine transport, have an important regulatory role in the stimulation of mTORC1 activity
18 (Nicklin et al. 2009; Chen et al. 2014; Milkereit et al. 2015).
19

20 Here we show that GPNA inhibits the sodium-independent influx of leucine and lowers its cell content, indicating that
21 the inhibitor hinders the activity of system L.
22
23
24
25
26
27
28
29
30
31
32
33

Materials and methods

34 Human oligodendroglioma Hs683 cells, provided by Prof. R. Kiss, University of Bruxelles, were grown in low-
35 glucose (1 g/L) Dulbecco's modified medium, DMEM (Euroclone), supplemented with 10 % FBS (Lonza, Basel,
36 Switzerland), 4 mM Gln, 25 mM HEPES, and antibiotics (100 U/ml penicillin, and 100 μ g/ml streptomycin). Human
37 cervix carcinoma HeLa cells, obtained from ATCC, human breast adenocarcinoma MCF7 cells, purchased from the
38 IZSLER Cell Bank (Brescia, Italy), and human hepatocellular carcinoma Huh7 cells, a gift of Prof. G. Raimondo,
39 University of Messina, were grown in high-glucose (4.5 g/L) DMEM supplemented with 10 % FBS, 4 mM Gln, and
40 antibiotics. Human lung alveolar carcinoma A549 cells, provided by Prof. L. Migliore, University of Pisa, were grown
41 in Ham's F12 medium supplemented with 10 % FBS, 1 mM Gln, and antibiotics. Cells were incubated at 37°C at 5%
42 CO₂; after thawing, all cells were used for less than 10 passages.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Total cell RNA (1 µg) was isolated and reverse transcribed, and cDNA analyzed as previously described (Chiu et
2 al. 2014). Primers were: 5'-GTGGAC TTCGGGA ACTATCACC (*SLC7A5*, for), 5'-GAACAGGGACCCATTGACGG
3 (*SLC7A5*, rev); 5'-AGGCTGGA ACTTTCTGAAT (*SLC7A8*, for), 5'-ACATAAGCGACATTGGCAA (*SLC7A8* rev);
4 5'-TGGTCTCCTGGATCATGTGG (*SLC1A5* for), 5'-TTTGCGGGTGAAGAGGAAGT (*SLC1A5* rev); 5'-
5 CACCACAGGGAAGTTCGTATTC (*SLC38A1* for), 5'-CGTACCAGGCTGAAAATGTCTC (*SLC38A1* rev); 5'-
6 ATGAAGAAGGCCGAAATGGGA (*SLC38A2* for), 5'-TGCTTGGTGGGGTAGGAGTAG (*SLC38A2* rev); 5'-
7 GCAGCCATCAGGTAAGCCAAG (*RPL-15*, for), 5'-AGCGGACCCTCAGAAGAAAGC (*RPL-15*, rev). Data
8 analysis was made according to the Relative Standard Curve Method (Bustin 2000).
9
10
11
12
13
14
15

16 Immunoblotting was performed as previously described (Chiu et al. 2012) using anti-LAT1 (rabbit, polyclonal,
17 1:1000, Cell Signaling Technology), anti-S6K1 phospho T389 (rabbit, monoclonal, 1:1000, Cell Signaling
18 Technology), anti-S6K1 total (rabbit, monoclonal, 1:1000, Cell Signaling Technology), anti-ASCT2 (rabbit,
19 monoclonal, 1:4000; Cell Signaling Technology) and anti-β-actin (rabbit, polyclonal, 1:1000, Sigma).
20
21
22
23

24 Intracellular leucine, isoleucine and glutamine were extracted with ice-cold absolute ethanol and determined with
25 liquid chromatography coupled with mass spectrometry as previously described (Bolzoni et al. 2016).
26
27

28 For ASCT2 gene silencing, Hs683 cells were transfected with a scrambled (ON-TARGETplus Non-targeting
29 Pool) or with a siRNA targeting ASCT2 (ON-TARGETplus SMARTpool, *SLC1A5*, Thermo Scientific DharmaFECT).
30 72 h after transfection, cells were rinsed in PBS and fresh medium was added. After 9 h, intracellular leucine was
31 extracted, and its content determined.
32
33
34
35

36 For transport experiments, cells were seeded on 96-well plates at a density of 15×10^3 cells/well in normal growth
37 medium. The initial influx of L-[3,4-³H]-glutamine (Amersham Biosciences) and L-[4,5-³H]-leucine (Amersham
38 Biosciences) was measured following the method previously described (Bianchi et al. 2012). For L-glutamine transport,
39 cells were rapidly washed with an Earle's Balanced Salt Solution (EBSS, composition in mM: NaCl 117, KCl 5.3,
40 CaCl₂ 1.8, MgSO₄·7H₂O 0.81, choline phosphate 0.9, glucose 5.5, supplemented with 0.02 % Phenol Red, kept at pH
41 7.4 with 26 mM Tris-HCl) and transport assay (30 s) was performed in the same solution. For L-leucine , before
42 transport determination, Na⁺-free EBSS, where N-methyl-D-glucamine chloride was used to replace NaCl, was used
43 for the washing and the transport assay
44
45
46
47
48
49
50
51

52 The non-saturable component of leucine influx was estimated measuring leucine uptake in the presence of 2 mM
53 leucine.
54
55

56 For the kinetic analysis, L-leucine influx data, obtained at different concentrations of the amino acid, were fit to

57 the equation: $v = \frac{v_{\max} \cdot [\text{Leu}]}{K_m + [\text{Leu}]} + K_D \cdot [\text{Leu}]$ (Eq. 1)
58
59
60
61
62
63
64
65

1 For the kinetic analysis of GPNA inhibition activity, L-leucine influx data, obtained at different concentrations of
2 the inhibitor, were fit to the equation for competitive inhibition: $v = v_0 - \frac{I_{\max} \cdot [\text{GPNA}]}{[I]_{0.5} + [\text{GPNA}]}$ (Eq. 2)
3

4 where v_0 is the influx in the absence of inhibitor, I_{\max} the maximal inhibition and $[I]_{0.5}$ the GPNA concentration at
5 which the inhibition is half-maximal.
6

7
8 GraphPad Prism 5.0™ was used for all the statistical analyses, and p values <0.05 were considered statistically
9 significant. Unless otherwise stated, Sigma-Aldrich was the source of all the chemicals, included GPNA.
10
11
12
13

14 Results

15
16 In a panel of human cancer cell lines, L- γ -glutamyl-*p*-nitroanilide (GPNA), an inhibitor of the sodium-dependent carrier
17 ASCT2, inhibited most of glutamine influx in the presence of sodium (Fig. 1a). However, GPNA significantly inhibited
18 glutamine transport also in the absence of sodium (Fig. 1b). To identify the sodium-independent transport system
19 inhibited by GPNA, the saturable influx of leucine was determined in the same cell models in the absence of sodium.
20 Leucine influx was different in the lines tested, with HeLa cells exhibiting the fastest influx and Huh7 cells the slowest
21 (Fig. 1c). In all the cell lines GPNA significantly inhibited the sodium-independent leucine influx, with inhibitions
22 ranging from almost 40% for Huh7 cells to over the 50% for Hs683 and MCF7 cells.
23
24

25 LAT1 and LAT2 system L transporters were consistently expressed, although at a variable degree, in all the cancer cell
26 lines (Fig. 2a, b). Human oligodendrogloma Hs683 cells showed the highest relative LAT1 and the lowest LAT2
27 mRNA expression, while MCF7 breast cancer cells had the highest LAT2 but relatively low LAT1 mRNA expression,
28 and hepatocellular carcinoma Huh7 cells had a low relative expression of both transporters.
29

30 The kinetic analysis of the Na⁺-independent Leu influx, performed in Hs683 cells (Fig. 3a), indicated that GPNA
31 increased the K_m for leucine from $56 \pm 2 \mu\text{M}$ to $104 \pm 15 \mu\text{M}$, while the V_{\max} was not significantly modified ($15.1 \pm$
32 2.10 nmol/mg/min , GPNA absent, versus $14.1 \pm 1.33 \text{ nmol/mg/min}$, GPNA present). The diffusion constant K_D was
33 also comparable in the absence and in the presence of GPNA ($49.5 \pm 2.42 \text{ min}^{-1}$, GPNA absent, versus $50.5 \pm 3.33 \text{ min}^{-1}$,
34 GPNA present). The inhibition pattern was satisfactorily fitted with an equation for a competitive inhibition (Fig. 3b).
35

36 At $10 \mu\text{M}$ [Leu], the maximal inhibition by GPNA was more than 65% of the uninhibited total influx with a half-
37 maximal inhibitory concentration of $807 \pm 70 \mu\text{M}$. The inhibitory effects on the sodium-independent leucine influx of
38 GPNA and of BCH, an amino acid analogue that preferentially inhibits system L, are compared in Fig. 3c. BCH
39 inhibited leucine influx by more than 80%, while GPNA-dependent inhibition was roughly 50% of the total leucine
40 uptake.
41

42 We next investigated the intracellular content of the two system L substrates leucine and isoleucine upon 9 hours of
43 incubation with 3 mM GPNA. Under control conditions, the intracellular content of both leucine and isoleucine varied
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 among the cell lines (Fig. 4a, b). Hs683 cells had the highest content of either leucine (14.4 ± 0.37 nmol/mg prot) or
2 isoleucine (14.3 ± 0.85 nmol/mg prot), while Huh7 cells had the lowest (Leu = 4.3 ± 0.09 nmol/mg prot; Ile = $3.9 \pm$
3 0.59 nmol/mg prot). In all but Huh7 cells, the intracellular content the two amino acids significantly decreased in the
4 presence of GPNA, with the highest inhibition (more than 40%) in Hs683 cells. The cell content of glutamine, measured
5 in the same cells (Fig. 4c) also varied among the various cell lines, with MCF7 showing the highest intracellular levels
6 and A549 cells the lowest. Only in Hs683 cells, GPNA significantly decreased intracellular glutamine.
7

8 Under the same conditions, the abundance of the phosphorylated form of S6K1, an indicator of the activity of the kinase
9 mTORC1, which is stimulated by intracellular leucine, was markedly lowered in all the cell lines (Fig. 5a). In all cells,
10 but Huh7, changes in pS6K1 were not paralleled by changes in total S6K1 and, therefore, could be attributed to an
11 effective decrease in mTORC1 activity. Interestingly, GPNA had inconsistent effects on LAT1 expression, with Hs683
12 and HeLa exhibiting sizable increases, while A549 and MCF7 showed a decreased expression of the transporter. As
13 expected, rapamycin suppressed S6K1 phosphorylation in all the cell lines, confirming its absolute dependence upon
14 mTORC1 activity (Fig. 4b).
15

16 To verify if the inhibition of ASCT2 by GPNA could be involved in the effects of the analog on the cell content of
17 leucine, SLC1A5 was silenced in Hs683 cells, causing an almost complete suppression of ASCT2 expression both at
18 mRNA (Fig. 6a) and protein level (Fig. 6b), and a significant inhibition of glutamine uptake that is exclusively detected
19 in the presence of sodium (Fig. 6c). No evidence of a compensatory increase in the expression of the other sodium-
20 dependent transporters SNAT1 (encoded by *SLC38A1*, Fig. 6e) or SNAT2 (encoded by *SLC38A2*, Fig. 6f) was detected.
21 However, ASCT2 silencing did not change the cell content of leucine (Fig. 6d).
22
23
24
25

26 Discussion

27 This report demonstrates that GPNA is a competitive inhibitor of system L transport activity. This sodium-independent
28 transport agency accounts for the cell uptake of most essential amino acids and is stimulated in many tumors (Kanai et
29 al. 1998). Although the kinetics of the inhibitory effect on leucine influx have been studied in Hs683 cells, which are
30 endowed with the highest expression of the system L transporter LAT1 when compared to the other cell lines tested,
31 both LAT1 and LAT2 transporters are likely inhibited by GPNA. Indeed, the percentage inhibition of the Na⁺-
32 independent leucine influx is substantially comparable in Hs683 and MCF7 cells, although the two cell lines exhibit
33 specular differences in LAT1 and LAT2 mRNA expression. However, the approach used in this study is not sufficient
34 to definitely discriminate if LAT1 or LAT2 are equally sensitive to GPNA or to exclude a possible sensitivity of the
35 other system L transporters LAT3 and LAT4.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In the past few years, GPNA has been used as an experimental device to inhibit glutamine transport through the Na⁺-dependent transporter ASCT2 in several models of cancer cells (Hassanein et al. 2013; Indo et al. 2013; Ren et al. 2015; Takahashi et al. 2015; Wang et al. 2015; Bolzoni et al. 2016; van Geldermalsen et al. 2016). Therefore, the inhibition of cell growth by GPNA has been taken as an evidence for the essential metabolic role of the ASCT2 transporter in those models. Inhibition by GPNA has been also used to demonstrate that glutamine analogues, synthesized as potential probes of ASCT2 transport function for Positron Emission Tomography, effectively interact with the transporter (Lieberman et al. 2011; Tang et al. 2016). At the light of the data presented here, these interpretations should be taken with caution. In particular, system L inhibition may contribute to the suppression of glutamine uptake by GPNA. Moreover, since the inhibition of system L by GPNA could directly hamper, besides glutamine uptake, the uptake of essential amino acids needed for protein synthesis and cell growth, the effects of GPNA on cell growth cannot be attributed to the sole inhibition of ASCT2. .

According to the model of Nicklin (Nicklin et al. 2009), leucine influx may be inhibited as an indirect effect of GPNA inhibition of the glutamine influx mediated by the sodium-dependent ASCT2 transporter that would lower the amount of intracellular glutamine available for promoting the influx of leucine through system L exchange transporters LAT1 and LAT2. The involvement of this mechanism in the GPNA-mediated inhibition of leucine influx is highly unlikely, since, in the transport experiments, the inhibitor is only present during the assay (30s), which, moreover, occurs in the absence of sodium. With the same argument, it is possible to exclude the involvement of other sodium-dependent transport systems, such as SNAT1 and SNAT2, recently described to be sensitive to GPNA inhibition (Broer et al. 2016).

On the contrary, it is possible that the significant depletion of the intracellular pool of leucine and isoleucine, observed in cells incubated for 9h with GPNA, may involve the decrease in cell glutamine attributable to ASCT2 inhibition. Actually, in human oligodendroglioma Hs683 cells long term GPNA treatment significantly lowered also the cell content of glutamine (Fig. 4c). However, GPNA decreased leucine influx and lowered the cell content of the essential amino acid in all the cell lines tested, with the only exception of Huh7 cells. In three of these cell lines (HeLa, A549, and MCF7) cell leucine decreased in the absence of a significant depletion of intracellular glutamine. Even in Hs683 cells a contribution of ASCT2 inhibition to leucine depletion seems unlikely, since ASCT2 silencing does not significantly affect cell leucine while markedly inhibits total and sodium-dependent glutamine uptake (Fig. 6). Although a role for other GPNA-sensitive sodium-dependent transporters, such as SNAT1 and SNAT2, cannot be excluded, it should be noted that no compensatory induction of these transporters is detected in ASCT2-silenced Hs683 cells (Fig. 6d,e), at variance with the results reported in other cell models (Broer et al. 2016).

1 Interestingly, in all the cell lines tested, the long term incubation with GPNA markedly decreases, although at a variable
2 degree, mTORC1 activity. Under the same conditions, the expression of LAT1 exhibited inconsistent changes, being
3 increased in Hs683 and in HeLa cells and decreased in A549 and MCF7 lines. Therefore, the inhibition of mTORC1
4 was not associated d to changes in LAT1 expression and should be attributable to the inhibition of leucine influx by
5 GPNA and/or to the partial depletion of the intracellular amino acid. Consistently, mTORC1 inhibition had been also
6 observed in cells incubated with the system L inhibitor BCH (Ishizuka et al. 2008). However, given that also glutamine
7 activates mTORC1 independently of leucine (Chiu et al. 2012; Jewell et al. 2015), GPNA-dependent inhibition of
8 ASCT2-mediated glutamine influx may also contribute to kinase inhibition.
9

10 In conclusion, this report demonstrates that the ASCT2-inhibitor GPNA also inhibits the influx of essential amino acids
11 through LAT1/2 and that this inhibition, depending on the expression level of LAT1/2, may affect the composition of
12 intracellular amino acid pool and the activity of mTORC1. The demonstration of the metabolic relevance of ASCT2 in a
13 cancer model should, therefore, rely on the genetic suppression of the transporter rather than on GPNA effects.
14
15

16 **Authors' contribution**

17 MC, CS, GT, and MGB performed the experiments. RA performed LC/MS-MS analysis. MC analyzed the data. MC
18 and OB designed the study and wrote the manuscript. NG discussed the results and revised the text. All Authors have
19 approved the final version.
20
21

22 **Funding Information**

23 MC is supported by a research fellowship of the University of Parma.
24
25

26 **Conflict of Interest**

27 The authors declare that they have no conflict of interest.
28
29

30 **Research involving Human Participants and/or Animals**

31 This article does not contain any study with human participants or animals performed by any of the authors.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 **References**
3
4
5

- 6 Barollo S, Bertazza L, Watutantrige-Fernando S et al (2016) Overexpression of L-Type Amino Acid Transporter 1
7 (LAT1) and 2 (LAT2): Novel Markers of Neuroendocrine Tumors. PLoS One 11:e0156044
8
9
10 Bianchi MG, Franchi-Gazzola R, Reia L et al (2012) Valproic acid induces the glutamate transporter excitatory amino
11 acid transporter-3 in human oligodendrogloma cells. Neuroscience 227:260-70
12
13 Bolzoni M, Chiu M, Accardi F et al (2016) Dependence on glutamine uptake and glutamine addiction characterize
14 myeloma cells: a new attractive target. Blood 128:667-79
15
16
17 Broer A, Rahimi F, Broer S (2016) Deletion of Amino Acid Transporter ASCT2 (SLC1A5) Reveals an Essential Role
18 for Transporters SNAT1 (SLC38A1) and SNAT2 (SLC38A2) to Sustain Glutaminolysis in Cancer Cells. J
19 Biol Chem 291:13194-205
20
21
22 Bustin SA (2000) Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction
23 assays. J Mol Endocrinol 25:169-93
24
25
26 Chen R, Zou Y, Mao D et al (2014) The general amino acid control pathway regulates mTOR and autophagy during
27 serum/glutamine starvation. J Cell Biol 206:173-82
28
29
30 Chiu M, Tardito S, Barilli A et al (2012) Glutamine stimulates mTORC1 independent of the cell content of essential
31 amino acids. Amino Acids 43:2561-7
32
33
34 Chiu M, Tardito S, Pillozzi S et al (2014) Glutamine depletion by crisantaspase hinders the growth of human
35 hepatocellular carcinoma xenografts. Br J Cancer 111:1159-67
36
37
38 del Amo EM, Urtti A, Yliperttula M (2008) Pharmacokinetic role of L-type amino acid transporters LAT1 and LAT2.
39 Eur J Pharm Sci 35:161-74
40
41
42 Esslinger CS, Cybulski KA, Rhoderick JF (2005) Ngamma-aryl glutamine analogues as probes of the ASCT2 neutral
43 amino acid transporter binding site. Bioorg Med Chem 13:1111-8
44
45
46 Fuchs BC, Bode BP (2005) Amino acid transporters ASCT2 and LAT1 in cancer: partners in crime? Semin Cancer Biol
47 15:254-66
48
49
50 Hassanein M, Hoeksema MD, Shiota M et al (2013) SLC1A5 mediates glutamine transport required for lung cancer cell
51 growth and survival. Clin Cancer Res 19:560-70
52
53
54 Hassanein M, Qian J, Hoeksema MD et al (2015) Targeting SLC1a5-mediated glutamine dependence in non-small cell
55 lung cancer. Int J Cancer 137:1587-97
56
57
58
59
60
61
62
63
64
65

1
2 Indo Y, Takeshita S, Ishii KA et al (2013) Metabolic regulation of osteoclast differentiation and function. *J Bone Miner Res* 28:2392-9
3
4 Ishizuka Y, Kakiya N, Nawa H et al (2008) Leucine induces phosphorylation and activation of p70S6K in cortical
5 neurons via the system L amino acid transporter. *J Neurochem* 106:934-42
6
7 Jewell JL, Kim YC, Russell RC et al (2015) Differential regulation of mTORC1 by leucine and glutamine. *Science*
8 347:194-8
9
10
11 Kaira K, Oriuchi N, Imai H et al (2008) L-type amino acid transporter 1 and CD98 expression in primary and metastatic
12 sites of human neoplasms. *Cancer Sci* 99:2380-6
13
14
15 Kanai Y, Segawa H, Miyamoto K et al (1998) Expression cloning and characterization of a transporter for large neutral
16 amino acids activated by the heavy chain of 4F2 antigen (CD98). *J Biol Chem* 273:23629-32
17
18
19 Lieberman BP, Ploessl K, Wang L et al (2011) PET imaging of glutaminolysis in tumors by 18F-(2S,4R)4-
20 fluoroglutamine. *J Nucl Med* 52:1947-55
21
22
23 Milkereit R, Persaud A, Vanoaica L et al (2015) LAPT4b recruits the LAT1-4F2hc Leu transporter to lysosomes and
24 promotes mTORC1 activation. *Nat Commun* 6:7250
25
26
27 Nicklin P, Bergman P, Zhang B et al (2009) Bidirectional transport of amino acids regulates mTOR and autophagy. *Cell*
28 136:521-34
29
30
31 Ren P, Yue M, Xiao D et al (2015) ATF4 and N-Myc coordinate glutamine metabolism in MYCN-amplified
32 neuroblastoma cells through ASCT2 activation. *J Pathol* 235:90-100
33
34
35 Takahashi K, Uchida N, Kitanaka C et al (2015) Inhibition of ASCT2 is essential in all-trans retinoic acid-induced
36 reduction of adipogenesis in 3T3-L1 cells. *FEBS Open Bio* 5:571-8
37
38
39 Tang C, Tang G, Gao S et al (2016) Radiosynthesis and preliminary biological evaluation of N-(2-
40 [18F]fluoropropionyl)-L-glutamine as a PET tracer for tumor imaging. *Oncotarget* 7:34100-11
41
42
43 van Geldermalsen M, Wang Q, Nagarajah R et al (2016) ASCT2/SLC1A5 controls glutamine uptake and tumour
44 growth in triple-negative basal-like breast cancer. *Oncogene* 35:3201-8
45
46
47 Wang Q, Hardie RA, Hoy AJ et al (2015) Targeting ASCT2-mediated glutamine uptake blocks prostate cancer growth
48 and tumour development. *J Pathol* 236:278-89
49
50
51 Wang Q, Holst J (2015) L-type amino acid transport and cancer: targeting the mTORC1 pathway to inhibit neoplasia.
52 *Am J Cancer Res* 5:1281-94
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legends

Fig. 1a,b L-[3,4-³H(N)]-Glutamine influx (25 μM; 5 μCi/mL; transport time, 30 s) was assayed in EBSS (a) or in Na⁺-free EBSS in the absence (Control) or in the presence of L-γ-glutamyl-*p*-nitroanilide (GPNA, 3 mM) in Hs683, HeLa, A549, MCF7 and Huh7 cells. **c** The determination of Na⁺-independent influx of L-[4,,5-³H(N)]-leucine (10 μM; 2 μCi/mL; transport time, 30 s) was performed in the same cells in the absence (Control) or in the presence of GPNA, 3 mM. The saturable influx was obtained subtracting the non-saturable influx (see Methods), determined in parallel under the same conditions, from the total influx. Data are expressed as nmol/mg prot/min. * *p* < 0.05, *** *p* < 0.001 versus control assessed with Student *t* test for unpaired data.

Fig. 2a *SLC7A5*, encoding for LAT1 (left), and *SLC7A8*, encoding for LAT2 (right), expression was assessed by RT-PCR in the indicated cells incubated in growth medium. Data were normalized to the expression of RPL-15. **b** LAT1 protein expression was assessed by Western Blot in Hs683, HeLa, A549, MCF7 and Huh7 cells. β-actin was used for loading control. A representative experiment, performed twice with comparable results, is shown.

Fig. 3a Kinetic analysis of Na⁺-independent leucine uptake by Hs683 cells. Cells were incubated for 30 s in Na⁺-free EBSS with L-[3,4-³H(N)]-Leu (2 μCi/mL; 1, 5, 10, 25, 50, 500 μM, left panel) in the absence (Control) or in the presence of 3 mM GPNA. Grey lines represent the saturable component of leucine influx for Control (dashed line) and GPNA (dotted line). Lines represent the best fit to Eq. 1 (*R*² 0.999 for control, 0.994 for GPNA inhibited influx). **b** Analysis of Leu uptake inhibition by GPNA. Cells were incubated for 30 s in Na⁺-free EBSS with L-[3,4-³H(N)]-Leu (2 μCi/mL; 10 μM) in the absence or in the presence of GPNA (0.3, 1, 3 mM). The line represents the best fit to Eq. 2 (*R*² 0.980). **c** Inhibition of Na⁺-independent leucine (10 μM) uptake by GPNA (3 mM) or BCH (1 mM). For **a-c**, data represent means ± SD of five independent determinations each. * *p* < 0.05; ** *p* < 0.01; *** *p* < 0.001 vs. control, assessed with Student *t* test for unpaired data.

Fig. 4 The cell content of Leu (**a**), Ile (**b**) and Gln (**c**) in Hs683, HeLa, A549, MCF7 and Huh7 cells incubated in growth medium without (Control) or with 3 mM GPNA for 9 h. Means ± SD of three experiments are shown. * *p* < 0.05, ** *p* < 0.01, *** *p* < 0.001 versus control; ns, not significant, as assessed with Student *t* test for unpaired data.

Fig. 5a Western blot of S6K1 (phosphoT389 and total) and LAT1 in Hs683, HeLa, A549, MCF7 and Huh7 cells incubated in growth medium (Control) or with 3 mM GPNA for 9 h. β-actin was used for loading control. **b** Cells were

1 incubated in the absence or in the presence of rapamycin (100 nm) for 9h. At the end of the incubation, the Western
2 Blot of S6K1 (phosphoT389 and total) was performed. Representative experiments, performed twice with comparable
3 results, are shown.
4
5
6

7 **Fig. 6** The expression of *SLCIA5* (for ASCT2, **a**) *SLC38A1* (for SNAT1, **e**) and *SLC38A2* mRNA (for SNAT2, **f**) was
8 assessed by RT-PCR in scramble- or anti-ASCT2-siRNA-transfected Hs683 cells 72h after transfection. Data were
9 normalized to the expression of RPL-15. **b** Western blot of ASCT2 in scramble- and anti-ASCT2-siRNA-transfected
10 Hs683 cells. **c** 72h after transfection, L-glutamine uptake (25 μ M, 10 μ Ci/ml) was measured, in the presence and in the
11 absence of sodium, in Hs683 cells transfected with scramble- or anti-ASCT2-siRNA as described under Materials and
12 Methods. **d** 72h after transfection, Hs683 cells, transfected with scramble- or anti-ASCT2-siRNA, were incubated for 9
13 h in fresh growth medium, and intracellular leucine was determined at the end of the incubation. For **a**, **e** and **f**, n = 4;
14 for **c**, n = 5; for **d**, n = 3; for **b**, a representative experiment, performed twice, is shown. * p < 0.05, ns, not significant, as
15 assessed with a Student t test for unpaired data.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65











