



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Above and belowground biodiversity in adjacent and distinct serpentine soils

This is the peer reviewed version of the following article:

Original

Above and belowground biodiversity in adjacent and distinct serpentine soils / Visioli, Giovanna; Sanangelantoni, Anna Maria; Conti a, Federica D.; A, Beatrice Bonati; B, Ciro Gardi; Menta, Cristina. - In: APPLIED SOIL ECOLOGY. - ISSN 0929-1393. - 133:(2019), pp. 98-103. [10.1016/j.apsoil.2018.09.013]

Availability:

This version is available at: 11381/2850549 since: 2021-10-01T09:52:30Z

Publisher:

Elsevier B.V.

Published

DOI:10.1016/j.apsoil.2018.09.013

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

02 May 2026

Manuscript Details

Manuscript number	APSOIL_2018_697_R1
Title	Above and below ground biodiversity in adjacent and distinct serpentine soils
Article type	Short Communication

Abstract

In this study we analysed the diversity of vascular plants, soil microorganisms and microarthropods on two serpentine outcrops in the Northern Apennines (Italy). The soils of the two sites are at different stages of evolution, as shown by soil depth and soil organic matter (SOM) content: the first site is a serpentine grassland clearing within sub-montane vegetation, while the second is a serpentinite scree rich in specialised flora only, including rare hyperaccumulator plant species such as *Noccaea caerulescens* and *Alyssum bertolonii*. The aboveground biodiversity was analysed via floristic relevés. The bacterial diversity was estimated through 16S rDNA profiling of the Ni hyperaccumulator *N. caerulescens* at the rhizosphere level, which thrives in both sites. Microarthropod communities were characterized by extracting and identifying organisms from the soil. The number of individuals per taxon, Acari/Collembola ratio, biodiversity indices and QBS-ar index were calculated. The two sites showed a clear difference as regards plant community: only three plant species were present in both sites, namely *Euphorbia cyparissias*, belonging to Euphorbiaceae, *Galium* spp., belonging to Rubiaceae, and *N. caerulescens*, belonging to Brassicaceae. Belowground diversity was positively correlated with vegetation cover and SOM content. In particular, among the bacteria colonising the rhizosphere of *N. caerulescens* (present in both sites), Actinobacteria, which exhibit K-strategist attributes and are more successful in resource-limited, crowded environments, appeared to be more abundant in the site with lower SOM and higher content of Ni. Bacteroidetes, which are specialised in degradation of cellulose, chitin and plant detritus, were instead more abundant in the site with higher SOM. As for soil microarthropod communities, the site with higher concentration and bioavailability of heavy metals and lower SOM showed poorer and less structured community. In conclusion, the data observed with plant, microbial and microarthropod communities identify different micro-habitats, in spite of proximal geographical position and the common metal-rich substrate.

Keywords	Serpentinite vegetation; <i>Noccaea caerulescens</i> ; Rhizosphere; Soil biodiversity; Soil fauna
Taxonomy	Biological Sciences, Environmental Science
Manuscript category	Invertebrate-related Submissions
Corresponding Author	Giovanna Visioli
Corresponding Author's Institution	University of Parma
Order of Authors	Giovanna Visioli, Anna Maria Sanangelantoni, Federica D. Conti, Beatrice Bonati, Ciro Gardi, Cristina Menta
Suggested reviewers	Arthurita Venter, Sara Branco, Nishanta Rajakaruna

Submission Files Included in this PDF

File Name [File Type]

Cover letter.doc [Response to Reviewers]

Highlights.docx [Highlights]

ManuscriptR1.doc [Manuscript File]

Fig1.tif [Figure]

Fig2.tif [Figure]

Figures SI1,2,3.docx [Figure]

Table SI1.docx [Table]

Table SI2.docx [Table]

Table SI3.docx [Table]

Submission Files Not Included in this PDF

File Name [File Type]

Supplementary data SI4.xlsx [Table]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Dr Andrey Zaitsev

Editor-in-Chief

Applied Soil Ecology

Parma 7 September, 2018

Dear Dr. Zaitsev,

Enclosed please find the revised manuscript “Above- and belowground biodiversity in adjacent and distinct serpentine soils”, by Giovanna Visioli, Anna Maria Sanangelantoni, Federica D. Conti, Beatrice Bonati, Ciro Gardi and Cristina Menta for publication in *Applied Soil Ecology*. We have followed the Editor’s and the Reviewers’ suggestions carefully, and moved most of the Tables and Figures to the Supplementary Materials section. The MS has also been thoroughly proofread by an English language instructor at our University Language Centre, Dr. Anila R. Scott-Monkhouse. As for the issue of pseudoreplication, we have added a sentence in the Conclusion, as suggested by Referee 2 too. Unfortunately, we found no other serpentine outcrop either nearby or elsewhere in the Apennines displaying the same conditions and plant species to perform comparative experiments. We have chosen to leave the Results and Discussion together since there are distinct subsets of data that were easily discussed independently.

We hope that this R1 version of the manuscript meets the standards required to be deemed suitable for publication in *Applied Soil Ecology*, as either a Research Article or a Short Communication.

Yours sincerely,

Giovanna Visioli

Corresponding author

The answers to the Editor’s and the reviewers’ comments are presented in bold:

Editor: Dear Authors, I received two reviews from the two independent Reviewers for your submission.

As you can see from their comments, they were very much satisfied with the editions you made. Please see below some further minor suggestions from the Reviewers.

However, after careful reading of your manuscript I still see that it needs more work before it could be considered for publication.

An important issue is language. Your text needs quite some editing. I tried to correct it but got discouraged after reading the abstract. I herewith kindly request you to let a native-speaker read and correct your manuscript.

I also suggest you to separate Results and Discussion. You have a full-size research paper and it is hard to follow the story with the current structure. Putting results and discussion together is more common in Short Communications. Alternatively, and this is my suggestion to convert this manuscript into a short communication as still the dataset is quite modest and the statistical analysis is very simple. Most of the results can be reported in supplementary materials.

Finally, there is a problem with pseudoreplication of your study. It is understandable, that it is hard to find truly replicated sites on serpentine, but this problem should be treated.

ANSWER TO THE GENERAL EDITOR'S COMMENTS: please see the cover letter.

Editor: Additionally, I ask you to make further corrections as described below.

Highlight 1 is uninformative. Please bring it closer to the context of your study.

Add one more highlight (make it Highlight 1) with some sort of rationale of your study. Considering its novelty this should not be too difficult.

Current highlight 2: how is affected.

Current highlight 3: shorten to fit 85 symbols. Alternatively, you may split it into two.

ANSWER: We have rewritten all the highlights, and hope that this amended version will be suitable for the journal.

L15 ...we analysed vascular plant, soil microorganism and microarthropod diversity...**DONE**

L19 mention at least one or two species here for example here. **DONE**

L21 ...bacterial and microarthropod diversity... This is easy to remember. We say e.g. "tree diversity", similarly we should say "microarthropod diversity". Please correct elsewhere. **DONE**

L23 ...thriving in both sites...**DONE**

L24 "Abundances" sounds strange. What kind of abundances? Microarthropod taxa separately, total, per species? **DONE - We have specified in the MS**

L27 ...and *N. caerulescens* belonging...**DONE**

L28 ...Belowground diversity...**DONE**

L33 ...abundant in the site with higher OC content in soil...**DONE**

L37 ...highest concentration and bioavailability of heavy metals...**DONE**

L38 Some sort of conclusion is needed (1 sentence). **DONE**

L45 ...of a stressful environment...**DONE**

NOTE! From now on I will not make any further corrections of language and will focus only on the scientific side of your work. A thorough check of the text by a native-speaker is required.

L46 correct Reference format. See <https://www.elsevier.com/journals/applied-soil-ecology/0929-1393/guide-for-authors> **DONE**

L80-84 The sentence is too long and hard to follow. Consider revision and splitting. **DONE**

L81 ...and nutrient elements cycling... sounds weird. **DONE**

L90 Single plant species? If yes, I would call it here "a model plant species" **DONE**

L97-98 "mountain range ridge" sounds strange- **DONE we eliminated the phrase**

L101-102 do these two sites have some proper names? If MP1 and MP2 have already been used in some earlier published work, it would be nice to have a reference here. **DONE**

L104 Specify horizons.:**DONE**

L106 Specify metals analyzed. **DONE**

L127 Format reference as required by the Guide for Authors. **DONE**

L131 Please explain, what is a "uclust".

ANSWER: A uclust algorithm divides a set of sequences into clusters.

L132 Explain, what is "QIIME".

ANSWER: Quantitative Insight Into Microbial Ecology is the bioinformatic pipeline used to analyse the microbial communities sampled through marker gene (e.g. 16S rRNA gene) amplicon sequencing.

L135 Please briefly explain, what is a "UniFrac".

ANSWER: Unifrac is a distance metric used to compare biological communities. It differs from dissimilarity measures such as Bray-Curtis dissimilarity in that it incorporates information on the relative relatedness of community members by incorporating phylogenetic distances between observed organisms in the computation. Both weighted (quantitative) and unweighted (qualitative) variants of UniFrac are widely used in microbial ecology.

L139 Did you split your monolith into horizontal layers during extraction? If not there is a fair chance that your extraction was not efficient. Please provide more details on that.

ANSWER: We collected a 10x10x10 cm soil core and distributed the sample in a 22 cm sieve of splitting the monolith into two parts only when the soil remained compact.

L142-143 It is very uncommon to place Myriapoda, all Insecta, all Chelicerata and Crustacea among microarthropods. For example, all myriapods and crustaceans (isopods actually?) are macroarthropods. Please clarify this issue and make changes accordingly throughout the text.

ANSWER: We agree with the Editor's comment and have understood the reason for this comment. However, we use the term "microarthropod" because we extracted and analysed only arthropods having 2 mm body size, in relation to the sieve mesh (we specified this aspect in the reviewed version). If the Editor believes that this use is wrong, we will replace the term with "arthropods".

Collembolans are often put separately from insects. You should be careful with your macrosystem. Please provide references.

ANSWER: We agree with the Editor's comment, and we know that Collembola are included in superclass Hexapoda, class Entognata (like Protura and Diplura). Considering the Editor's suggestion, we have replaced Insect with Hexapoda in the text.

L144-147 It is quite not correct to calculate diversity indices basing on different taxonomic resolution levels for different soil animal taxa. How did you deal with this?

ANSWER: We have understood the Editor's comment. We adopted this approach several times in previously published papers and the Reviewers accepted it. If the Editor considers this approach as being not entirely correct, we can calculate the indices using the orders only.

L148 Please explain, what is a "soil adaptation level"?

ANSWER: This concept is in relation to the QBS-ar index approach and the EMI (EcoMorfological score) assignment used for the QBS-ar computation. We did not explain this concept in the text because it was explained in several previous scientific papers, two of which are reported in the references.

L151 Obviously those are packages of R. Please explicitly state that and include the version of the software and a proper reference into the Reference list. **DONE**

L154 observed number of taxa? **ANSWER: J was calculated dividing H' by the logarithm of the number of taxa. We have clarified this in the text.**

L159 Results are always reported in the past tense. **DONE**

L162-163 Move to site description. **DONE**

L180-181 Are those SDs or SEs? Explain in Methods **DONE.**

L185 You mean "weakly?" **DONE**

L192-194 This belongs to Discussion.

ANSWER: We have considered the Editor's suggestion of shortening the manuscript and left the Results and discussion section together.

L199 ...*N. caerulescens*... At first mentioning of a species in the text you write it in full, then you abbreviate the genus name. Same is applicable for the Abstract as a standalone text. **DONE**

L210 ...Venter et al. (2018). **DONE.**

L215-220 The sentence is too long and unintelligible. Should be split into several sentences. The second part should go to Discussion. **ANSWER: We have modified the sentence. As explained above, we have considered the Editor's suggestion of shortening the manuscript and left the Results and discussion section together.**

L231 You refer to Fig. 2a, but not to 2b. Later you refer to Fig. 2 as a whole. Do you really need "a" and "b" then? **DONE**

L238 in the rhizosphere? **DONE**

L242-243 to Discussion and L248-250 to Discussion. **ANSWER:**

As explained above, we have considered the Editor's suggestion of shortening the manuscript and left the Results and discussion section together.

L252-255, L257-262 to Discussion. Further, please make very clear separation between Results and Discussion. I will not make further comments on that but anticipate, that this is fixed. **ANSWER:**

We have considered the Editor's suggestion of shortening the manuscript and left the Results and discussion section together.

L263 "degree of biodiversity" is not a common term. **DONE**

L276 From what I read here, there is no way to call all taxa collected as microarthropods. You have a mixture of macro- and mesofauna. Please adjust elsewhere that you were studying soil arthropods.

ANSWER: Please refer to the explanation above (L 142-143).

L305 The story ends nowhere and makes a paper rather confirmatory. A deeper discussion on the correlation of plant, microbial and arthropod diversity is needed.

Conclusions are also rather confirmatory and can be or replaced by a take-home message. The only interesting statement on the importance of OC content and Ni content is actually not based on any kind of statistical analysis or extensive literature search.

ANSWER: A final sentence was added in the Conclusion, following the suggestions of Referee 2 too.

Table 1 to Supplementary Material with most important data to methods.

ANSWER: DONE

Table 2 Are those means and SEs? In fact those are not replicates but pseudoreplicates. How have you treated this in your design? A star marking significance should be not after the mean value, but after the SE. **ANSWER: DONE**

L492 Organic matter or Organic carbon? If the former, how have you recalculated this? **ANSWER: This was indicated in the Materials and Methods section.**

L499 Space after "x"**DONE**

Table 3 can go to Supplementary Material.**DONE**

Figures.

First, it is hard to understand which Figure is which one. They are not signed.

Further I will call them in accordance to their appearance.

Figure 1 - to Supplementary material **DONE**

Figure 2 - What do percent refer to? Frequencies? Y-axis should be signed. Also in the text to refer to Figure 2a. Here there are no any "a's" and b's"...**DONE. This figure is now Figure 1. Percentagae of number of sequences of the phyla respect to the total number of sequences**

Figure 3 is not readable. Increase fonts. Overall it can be replaced with a Table and go to Supplementary Material. You have "a" and "b" in the Figure, but nothing in the figure caption about it. Also no separate references for Fig. 3a and Fig. 3b in the text. **ANSWER: We put Figure 3 in supplementary materials increasing font.**

Figure 4 - How can you prove significance of differences? This figure does not bring too much, particularly if you had multiple taxonomic resolution levels in your dataset. Consider skipping.

ANSWER: After carefully considering the Reviewer's suggestion, we have chosen to maintain the figure in the MS because the statistical analysis was made by comparing the couple of data (MP1 vs MP2) for each taxon. In our opinion this figure adds important explanation and would be best kept it in the Results discussion section.

References.

Please see attached file with the results of the references check. The color coding should be pretty much self-explanatory: Red - reference missing in text or vice-versa. Purple - problems with formatting.

ANSWER: All the references were checked according to the indications. Where there are open assess articles we put the doi

REVIEWER 1 The revisions have greatly improved the paper. Look forward to seeing the paper in print.

ANSWER: We thank Reviewer1 for their positive evaluation of the manuscript in its present form.

REVIEWER 2

The study characterizes the diversity of two different serpentine sites, having different environmental features, particularly regarding microclimate and soil structure. The manuscript is interesting, really well written, pleasant to read and rich in details and analyses. The only concern I have is about the only two sites where the authors made the experiment. Although they temporary and spatially replicate the sampling, only one site per factor level risks being not sufficient to make general conclusions about the aim of the study. However, if the sampling area provides spatial constraints as declared by the authors, this problem cannot be actually solved, so I think they have done a really good job anyway. Nonetheless, I suggest the authors to add a sentence in the

conclusion section where they declare that the study may suffer from this problem, and that it should be repeated in different contexts to ensure a general reliability of results. Adding a figure showing the outcrop and the relative position of the two sites, may help to better understand this issue.

However, I think that the manuscript is suitable for publication. I have only a few minor comments listed below.

Lines 54-68: I suggest the authors to consider these two studies:

Mengoni et al (2004). Genetic diversity of bacterial communities of serpentine soil and of rhizosphere of the nickel-hyperaccumulator plant *Alyssum bertolonii*. *Microbial ecology*, 48(2), 209-217.

Mengoni et al. (2001). Characterization of nickel-resistant bacteria isolated from serpentine soil. *Environmental Microbiology*, 3(11), 691-698.

ANSWER: We thank Reviewer 2 for their positive evaluation of the manuscript. We have followed their suggestions by adding a comment in the Conclusion section concerning the problem of the small number of samples. We have also added the studies suggested. The figure showing the outcrop and the relative position of the two sites is reported as supplementary figure 1.

Lines 84-91: In my opinion, the aim as described by the authors is a little bit weak if compared to the in-depth study they performed. Given that the study provides a comparative approach between two different sites, I think that the aim has to be rephrased also focusing this aspect, not only the 'simple' characterization of the two site. The authors should also list their expectations on the basis of the available literature, and answer to each expectation in the discussion section. **DONE**

Lines 173-174: Please add the type of test used and the value of the statistical parameter. **DONE**

Bullet points

- Above and belowground biodiversity in serpentine soils are still little known.
- Vascular plant, soil microorganism and microarthropod diversity were studied.
- Different micro-habitats on serpentinite showed high vascular plant diversity.
- *Noccaea caerulescens* rhizosphere is modulated by soil characteristics.
- Microarthropods are richer in lower Ni and higher soil organic matter contents.

1 **Above and belowground biodiversity in adjacent and distinct serpentine soils**

2 Giovanna Visioli^{a*}, Anna Maria Sanangelantoni^a, Federica D. Conti^a, Beatrice Bonati^a, Ciro Gardi^b,
3 Cristina Menta^a

4 ^a Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma,
5 Parco Area delle Scienze 11/A, 43124, Parma, Italy

6 ^bAnimal and Plant Health (Alpha), European Food Safety Authority (EFSA), Via Carlo Magno 1A,
7 43126 Parma – Italy.

8

9 * Corresponding author: Giovanna Visioli, Department of Chemistry, Life Sciences and
10 Environmental Sustainability, University of Parma, Parco Area delle Scienze 11/A, 43124, Parma,
11 Italy.

12 *E-mail address:* giovanna.visioli@unipr.it

14 **Abstract**

15 In this study we analysed the diversity of vascular plants, soil microorganisms and microarthropods
16 on two serpentine outcrops in the Northern Apennines (Italy). The soils of the two sites are at
17 different stages of evolution, as shown by soil depth and soil organic matter (SOM) content: the
18 first site is a serpentine grassland clearing within sub-montane vegetation, while the second is a
19 serpentinite scree rich in specialised flora only, including rare hyperaccumulator plant species such
20 as *Noccaea caerulescens* and *Alyssum bertolonii*. The aboveground biodiversity was analysed via
21 floristic relevés. The bacterial diversity was estimated through 16S rDNA profiling of the Ni
22 hyperaccumulator *N. caerulescens* at the rhizosphere level, which thrives in both sites.
23 Microarthropod communities were characterized by extracting and identifying organisms from the
24 soil. The number of individuals per taxon, Acari/Collembola ratio, biodiversity indices and QBS-ar
25 index were calculated. The two sites showed a clear difference as regards plant community: only
26 three plant species were present in both sites, namely *Euphorbia cyparissias*, belonging to
27 Euphorbiaceae, *Galium* spp., belonging to Rubiaceae, and *N. caerulescens*, belonging to
28 Brassicaceae. Belowground diversity was positively correlated with vegetation cover and SOM
29 content. In particular, among the bacteria colonising the rhizosphere of *N. caerulescens* (present in
30 both sites), Actinobacteria, which exhibit *K-strategist* attributes and are more successful in
31 resource-limited, crowded environments, appeared to be more abundant in the site with lower SOM
32 and higher content of Ni. Bacterioidetes, which are specialised in degradation of cellulose, chitin
33 and plant detritus, were instead more abundant in the site with higher SOM. As for soil
34 microarthropod communities, the site with higher concentration and bioavailability of heavy metals
35 and lower SOM showed poorer and less structured community. In conclusion, the data observed
36 with plant, microbial and microarthropod communities identify different micro-habitats, in spite of
37 proximal geographical position and the common metal-rich substrate.

38

39 *Keywords:* Serpentinite vegetation; *Noccaea caerulescens*; Rhizosphere; Soil biodiversity; Soil
40 fauna

41

42 **1. Introduction**

43 Serpentinites are widespread and represent a well-known example of a stressful environment
44 (Harrison and Rajakaruna, 2011), being constituted by tectonic blocks and intrusion of ultramafic
45 rock and by sedimentary and metamorphosed derivatives (Coleman and Jove, 1992). These soils are
46 generally characterised by low levels of macronutrients such as N, K, P, Fe. In addition, the high
47 Mg causes a low Ca/Mg molar ratio, inhibiting Ca uptake, and the high concentration of heavy
48 metals, particularly Ni, exerts toxic effects on plants and soil biota (Chiarucci et al., 2003; Lombini
49 et al., 1998). Dwarf plant communities often inhabit serpentine areas and are remarkably different
50 from the ones growing in adjacent soils (Lombini et al., 1998). Even microbial communities are an
51 important, highly specialised component in belowground serpentine environments (Branco, 2010;
52 Mengoni et al., 2004; 2001; Venter et al., 2018; 2015). Several studies based on culture-dependent
53 techniques have revealed that serpentine microorganisms show a very high degree of multi-metal
54 resistance. Moreover, bacteria present in serpentine soils surrounding plant roots (rhizobiome) are
55 known to enhance plant nutrient and water uptake and protect plants against toxic conditions, thus
56 greatly contributing to plant growth and fitness under harsh edaphic conditions (Seneviratne et al.,
57 2016). The culturable bacteria isolated from serpentine soils benefit plants by producing
58 siderophores and carboxylic acids, and help solubilisation of phosphates, which increase the
59 mobility of macro- and micronutrients in the rhizosphere (Rajkumar and Freitas, 2008) and produce
60 phytohormones (Visioli et al., 2014). Isolation-based microbiological techniques are limited
61 because culturable microorganisms represent only a tiny percentage of the soil's microbial
62 community. Metagenomics and metaproteomics are ideal approaches when investigating the

63 diversity and ecology of microbiome in serpentine soils, though very little information is currently
64 available (Mattarozzi et al., 2017).

65 Even the soil fauna community is affected by high metal concentrations, but to date
66 knowledge about soil animal communities in serpentine soils is scanty. There are examples of high-
67 Ni herbivorous insects which are well adapted to these severe habitats (Boyd, 2009). Frizzi et al.
68 (2017) reported an interesting pattern of mortality in the *Crematogaster scutellaris* ant in long-term
69 metal exposure in serpentine soils, while ants from intermediate contaminated sites are more
70 tolerant to acute exposure to nickel. Moreover, the authors found no difference in genetic diversity
71 between ant colonies from different sites, and hypothesised that this was due to queen mediated
72 gene flow during nuptial flights across contaminated and uncontaminated areas of limited
73 geographical extent. Visioli et al. (2013) reported significant differences in terms of microarthropod
74 communities between four serpentine soils characterized by different metal content. The site
75 characterized by lower metal content showed the richer community. Soil fauna is involved in
76 important processes such as organic matter decomposition, humus formation, nutrient cycles
77 (nitrogen, sulphur, and carbon), and bioturbation (Bird et al., 2004). For these reasons, soil animals
78 play an important role in the distribution and bioavailability of the trace elements generally present
79 in high concentration in serpentine soils.

80 The aim of this study was to compare the above- and belowground biodiversity in two micro-
81 habitats on the serpentine outcrop of Mt. Prinzera, an ophiolite mountain in the Italian Northern
82 Apennines, characterised by sharp changes in geomorphology and significant edaphic and
83 microclimatic variations due to exposure (Lombini et al., 1998). The study adopts a
84 multidisciplinary approach, from aboveground plant cover to soil microbial and fauna community.
85 In both sites we compared: i) vegetation cover; ii) bacterial diversity via 16S rDNA profiling within
86 the rhizosphere of a model plant species present in both sites; iii) composition and abundance of
87 microarthropod community.

88

89 **2. Materials and Methods**

90

91 *2.1 Study area, soil and aboveground classification*

92 The area studied is located on Mt. Prinzera (MP) (736 m a.s.l.), one of the most extreme ophiolitic
93 outcrops of the Italian Northern Apennines (Lombini et al., 1998). The landscape and ecological
94 features of Mt. Prinzera and its surroundings (237 ha) are protected as a nature reserve of the
95 Emilia–Romagna region. For this reason, the collection of few samples only was allowed, and only
96 two previously identified different micro-habitats were analysed in this study: MP1 (GPS
97 44.65138°N – 10.08330°E) and MP2 (GPS 44.64282°N – 10.07951°E) (Visioli et al., 2013, 2012).
98 The soil profile description and sampling were carried out in the two sites by collecting three soil
99 samples per horizon (horizon A and horizon A/C). The soils were classified according to the World
100 Reference Base for soil resources 2014 (IUSS Working Group, 2015) as Skeletic Mollic Leptosol
101 and Orthoskeletal Leptosol for site MP1 and site MP2 respectively (Figure SI1, Table SI1). The
102 water content, pH value and total Ca, K, Mg, Fe, Ni, Zn, Co and DTPA extractable Ni, Zn, Co
103 contents in sites MP1 and MP2 were analysed according to Visioli et al. (2012). SOM was
104 determined by “weight loss on ignition method. Two-gram aliquots of each soil were placed in
105 ceramic crucibles at 450°C for 3h. The SOM was then calculated with the equation: $SOM \% =$
106 $(\text{initial weight} - \text{final weight}) \times 100$.

107 Plant diversity was analysed by recording vascular flora in the sites being studied. Vascular plant
108 species were recorded as present during field surveys from May to October 2015. The aboveground
109 biomass was collected within three plots (0.25 x 0.25 m) for each sampling site, oven dried at 105°
110 C, and weighted, and standard deviations were calculated. The nomenclature of taxa follows
111 Pignatti (1982).

112

113 *2.2 Rhizosphere biodiversity*

114 2.2.1 Plant samples and extraction of total DNA from rhizosphere soil

115 *N. caerulea* plants (which accumulate up to 2,000 mg kg⁻¹ DW) were collected in May
116 2015 from the previously studied sites, MP1 and MP2 (Visioli et al., 2012). A total of 6 *N.*
117 *caerulea* plants per site, at the same phenological stage were uprooted manually (i.e. with steles
118 full of immature inflorescences). *N. caerulea* rhizospheric soils were separated from the roots
119 and collected. The DNA was extracted from 500 mg of soil with the FastDNA[®] Spin Kit for Soil
120 (MP Biomedicals, Santa Ana, CA) and visualised by electrophoresis on 0.8% (w/v) agarose gels to
121 test the DNA integrity.

122

123 2.2.2 16S rRNA gene-based community analysis

124 Partial 16S rDNA sequences were amplified from the DNA extracted by using primer pair
125 Probio_Uni and /Probio_Rev, which target region V3 of the 16S rRNA gene sequence, and 16S
126 rRNA gene amplification and amplicon checks were carried out as described (Mancabelli et al.,
127 2016). The 16S rDNA sequencing was performed via MiSeq (Illumina) at the facility of GenProbio
128 srl (www.genprobio.com) following the protocol previously reported (Mancabelli et al., 2016). In
129 order to calculate downstream diversity measures (alpha and beta diversity indices, Unifrac
130 analysis), 16S rRNA Operational Taxonomic Units (OTUs) were defined at ≥ 97 % sequence
131 identity by using uclust algorithm (Edgar, 2010). All reads were classified at the lowest possible
132 taxonomic rank by using QIIME (Quantitative Insights into Microbial Ecology) and a reference
133 dataset from the SILVA database (Quast et al., 2013). The biodiversity of the samples (alpha-
134 diversity) was calculated with Chao1 and Shannon indices. Similarities between samples (beta-
135 diversity) were calculated by unweighted UniFrac distance metric (Lozupone and Knight, 2005).
136 The range of similarities was calculated between the values 0 and 1.

137

138 2.3 Microarthropod community

139 Three soil cores (100 cm² and 10 cm in depth) per site were collected in May and October
140 2015, and put on the Berlese-Tüllgren funnel (2 mm sieve mesh) for microarthropods extraction.
141 The specimens extracted were collected in a 2 alcohol/1 glycerine (v/v) solution and identified via a
142 stereomicroscope. We considered class taxonomic level for Myriapoda, and order level for
143 Hexapoda, Chelicerata and Crustacea (<http://taxonomicon.taxonomy.nl/>), counted the number of
144 specimens present per taxon and calculated their abundance (ind/m²). Based on this classification
145 level, we assessed taxa richness and diversity using Margalef index (D; Margalef, 1958), Shannon
146 diversity index (H'; Shannon and Weaver, 1963), Pielou's evenness index (J; Pielou, 1975) and
147 Simpson's index 1-D (Simpson, 1949). We applied QBS-ar index in order to describe the soil
148 microarthropods community in terms of soil adaptation level (Menta et al., 2018).

149 *2.4 Statistical analyses*

150 Richness and biodiversity indices (Shannon diversity - H'-, Pielou's evenness -J- and Simpson's
151 dominance -1-D-) were calculated by using the *vegan* package (Oksanen et al., 2017), in particular
152 applying the *diversity* function to the absolute abundances observed. While H' and 1-D were
153 directly calculated through the function, J was calculated by dividing H' by the logarithm of the
154 number of taxa observed. These estimates were performed via R software (R Core Team, 2016,
155 version 3.0.2). The Mann-Whitney U-test (U) was used to compare MP1/MP2 chemical and
156 physical parameters and MP1/MP2 number of microarthropods for different taxa, and the
157 comparison of MP1/MP2 richness, biodiversity and soil biological indices.

158

159 **3. Results and discussion**

160

161 *3.1 Soil characteristics*

162 The soil of site MP2 was at early stages of differentiation, and was formed essentially by the
163 material resulting from the physical degradation of serpentine rocks. Skeleton was dominant and the
164 fine earth fraction was only 22% and 9% in horizon A and A/C respectively. The soil organic matter
165 (SOM) in the horizon A was 2.95%. Stones cover more than 90% of soil surface as a result of
166 selective erosion (Fig. SI1 and Table SI1). The soil from site MP1 showed evidence of evolution in
167 terms of physical and chemical weathering and SOM accumulation; the fine earth fraction was 40%
168 and 15% respectively for horizon A and A/C and WW, while the SOM in horizon A was 4.84%
169 (Fig. SI1 and Table SI1).

170 Soil chemical-physical analyses highlighted the differences between the two sites reported in
171 Table 1. The amount of total Ni in the two sites was very high, but MP2 showed a significant higher
172 content in total Ni ($P < 0.05$). The amount of Fe and K was also significantly higher in MP2 than in
173 MP1 ($P < 0.05$), while Co was significantly higher in MP1 ($P < 0.05$). The lowest Ni, Zn and Co
174 DTPA extractable fractions were observed in MP1. Ca, Mg and Zn did not reveal any significant
175 variation between the two sites. SOM and water content were significantly higher in MP1, while pH
176 was similar in the two sites (around 7.00).

177

178 *3.2 Plant diversity*

179 The two sites were different in terms of aboveground biomass, with about 158 ± 32 g m⁻² d.w.
180 in MP1 and about 114 ± 6 g m⁻² d.w. in MP2. Site MP2 is characterized by the typical vegetation of
181 scree slopes, with species highly adapted to extreme conditions (extremely coarse and stony,
182 weakly developed soil, low in organic matter and nutrient content, higher Ni concentration) (Tables
183 1, SI1); the vegetation cover is discontinuous, and the aboveground biomass low. Site MP1 is a
184 weakly developed soil too (Leptosol according to WRB), but the percentage of fine earth and SOM
185 is higher compared to MP2, and this is reflected in plant species composition and aboveground
186 biomass (Tables 1, SI1); the vegetation cover is continuous and the aboveground biomass

187 significantly higher than in MP2. A survey of vascular plant diversity present in sites MP1 and MP2
188 is reported in Table S2. Poaceae, Carofillaceae and Fabaceae were present in MP1 only, while
189 Asteraceae and Brassicaceae were predominant in MP2. Interestingly, most of the genera and
190 species present in MP2 (e.g. *Echinops*, *Thymus*, *Globularia bisnagarica*, *Helichrysum italicum*,
191 *Artemisia annua*, and *Sedum montanum*) are known to be adapted to dry rocky soils and to produce
192 secondary metabolites with anti-inflammatory and antimicrobial proprieties (Pignatti, 1982). The
193 species identified in site MP1 are typical of forest margins and humid basic soils (i.e. *Cornus*
194 *sanguinea*, *Clematis vitalba*, *Hypericum perforatum*) (Pignatti, 1982). Moreover, only three plant
195 species/genera were present in both sites, *Euphorbia cyparissias*, belonging to Euphorbiaceae,
196 *Galium*, belonging to the Rubiaceae, and *N. caerulescens*, belonging to Brassicaceae (Table SI2).
197 The former two are annual and perennial herbaceous plants, quite commonly found in a wide region
198 of Europe, while *Noccaea caerulescens* is a well-known Ni hyperaccumulator, endemic to Mt.
199 Prinzer and present with different populations which colonise various micro-habitats and display
200 distinguishing phenotypic traits (Visioli et al., 2012). *Alyssum bertolonii*, the other Ni
201 hyperaccumulator which colonises serpentinite in the Italian Apennines (Galardi et al., 2007;
202 Lombini et al., 1998), is present in site MP2 only. These analyses confirmed previous studies which
203 indicated that specialised flora is found on serpentine rocks and screes only (Ferrari et al., 1992) and
204 is of great interest for its adaptation to such stress-inducing habitats.

205

206 3.3 Rhizosphere bacterial biodiversity

207 Previous studies have shown that variations in the structure of bacterial communities may be
208 due to the presence of different plants in non-serpentine and serpentine sites (Pessoa-Filho et al.,
209 2015). By comparing 16 soils belonging to both serpentine and non-serpentine geology, Venter et
210 al. (2018) demonstrated that the local heterogeneity in variables such as soil characteristics or
211 microclimate gradients may impact cyanobacteria and algal biodiversity. In addition, when

212 considering the same serpentine site, the structure and functions of the rhizosphere bacterial
213 community varied between hyperaccumulator and tolerant plants (Matarozzi et al., 2017).
214 Considering all these aspects, despite the substantial differences in plant biodiversity between sites
215 MP1 and MP2, the presence of *N. caerulescens* in both sites enabled us to investigate possible
216 changes in bacterial diversity associated with the rhizosphere of the same plants in the two micro-
217 habitats. *N. caerulescens*, endemic to Mt. Prinzera, is a Ni hyperaccumulator accession and different
218 populations were previously analysed for morphological and molecular traits highlighting the high
219 phenotypic plasticity of this accession (Visioli et al., 2012). In particular, *N. caerulescens*
220 individuals in MP1 and MP2 were diversified for both growth parameters and Ni accumulation
221 capacity (Visioli et al., 2012). The 16S rDNA based community profiling highlighted the bacterial
222 diversity in the rhizosphere of *N. caerulescens* sampled in MP1 and MP2. A total 153,482 raw
223 pyrosequencing reads was obtained from the two rhizospheres. After quality trimming 138,756
224 sequences were obtained: 44,307 reads for MP1 and 94,449 reads for MP2 (Table SI3). The
225 phylogenetic analysis revealed high bacterial complexity phylum, family and genus levels, with 11
226 phyla representing >1% in both samples (Fig. 1; supplementary file SI4).

227 Bacterial diversity in sites MP1 and MP2 measured on the basis of OTUs, and calculated
228 bacterial diversity indices, i.e. Shannon diversity index and Chao1 estimator of richness, showed
229 that there are no significant differences between samples. The rarefaction curves using the Chao and
230 Shannon indices reached a plateau, thus indicating that further sequencing would not have resulted
231 in the retrieval of more OTUs (Fig. SI2).

232 Actinobacteria (MP1 37%, MP2 48%), Proteobacteria (MP1 32%, MP2 31%) and
233 Acidobacteria (MP1 7.94%, MP2 6.51%) were the predominant phyla in the rhizosphere. The
234 significant differences observed in the abundance of Actinobacteria sequences between the two sites
235 were not surprising as the soil in site MP2 was expected to be more selective because of the lower
236 SOM (2.95% versus 4.84% in MP1) and the higher content of Ni (2032 $\mu\text{g g}^{-1}$ of soil dry weight
237 versus 1700 $\mu\text{g g}^{-1}$ in MP1) (Table 1), and Actinobacteria exhibit *K-strategist* attributes, i.e. are

238 more successful in resource-limited environments. The higher richness in Actinobacteria of site
239 MP2 can be ascribed to the higher percentage of Mycobacteriaceae, Geodermatophilaceae,
240 Micromonosporaceae, Pseudonocardiaceae, Rubrobacteriaceae families and U. m. of Frankiales
241 order (Fig. 2). Other phyla with > 2% relative abundance were Bacterioidetes, more present in MP1
242 (6.3%) with Cytophagaceae, Chitinophagaceae and Saprospiraceae families, than in MP2 (1.94%).
243 This can probably be attributed to the more abundant aboveground biomass (either plant or animal)
244 in MP1, giving a greater concentration of carbon sources suitable for these bacteria specialised in
245 degradation of cellulose and chitin (Beier and Bertison, 2013).

246 Despite the selective conditions, the bacterial diversity was high in both sites, with forty-
247 four families representing >1% (26% of them belonging to unculturable bacteria) (Fig. SI3). The
248 high bacterial biodiversity found on serpentine soils is in accordance with previous studies on ECM
249 fungal communities, demonstrating that serpentine soils are not depauperate and not
250 characteristically composed of specialised species, compared to non-serpentine ones (Branco,
251 2010).

252 The greater richness in Acidobacteria in site MP1 was mainly due to U. m. of subgroup 6 order
253 (Fig. SI3). Because of the difficulty in cultivating most of its members, this phylum changed from
254 being virtually unknown to being recognized as one of the most abundant and diverse on the planet
255 only after cultivation-independent analysis methods were fully developed (Kielak et al., 2016). In
256 particular, subgroup 6 has been reported as a subgroup of Acidobacteria with a marked preference
257 for soil environments with increased availability of nutrients and higher organic carbon content
258 (Navarrete et al., 2015).

259 By analysing these data, we can conclude that there is no difference in the degree of
260 biodiversity between the rhizosphere of *N. caerulescens* in the two environments, probably because
261 rhizosphere is a “protected” environment for bacteria. Authors who analysed ECM fungal
262 biodiversity on serpentine soils previously speculated that the high diversity may be maintained by
263 a combination of environmental effects and competitive relationships among species (Branco,

264 2010). However, in our situation, environmental factors such as differences in SOM, humidity and
265 Ni concentration shape the bacterial communities in the rhizosphere of the same plant species. This
266 is not surprising as several studies concerning the diversity and community composition of
267 microorganisms (either bacteria or fungi) in the rhizosphere of plants have identified the
268 environment as the most important factor shaping the community and influencing richness and
269 diversity, rather than the root itself (Turrini et al., 2017; Zhang et al., 2018).

270

271 *3.4 Microarthropods biodiversity*

272 We found an average abundance of 13,248 ind/m² belonging to 16 taxa and 6,065 ind/m²
273 belonging to 9 taxa in MP1, in May and October respectively. In MP2 we observed an average
274 abundance of 4,628 ind/m² belonging to 12 taxa in May, and a mean of 524 ind/m² belonging to 8
275 taxa in October. In both periods the number of taxa was higher in MP1 compared to MP2, but the
276 difference was greater in May (Fig.2). This datum is similar to the trend reported in Visioli et al.
277 (2013) and confirms the different arthropod community supported by the two soils. Acari and
278 Collembola showed the highest abundance in both sites, except for MP2 in October (Fig. 2).
279 Comparing MP1 and MP2, the former showed the highest abundance for both taxa and for both
280 periods. This trend is followed by many other taxa. All taxa extracted in May were more abundant
281 in MP1 than in MP2, except for Psocoptera. Statistical analysis showed significant differences for
282 Symphyla and Coleoptera only (both larvae and adult) (Fig. 2a). Moreover, Pseudoscorpiones,
283 Diplopoda and Pauropoda were absent in MP2.

284 Pseudoscorpiones, Diplopoda, Pauropoda, Chilopoda, Diplura and Tysanoptera were found
285 in neither sites in October. The other taxa showed a trend similar to the one observed in May,
286 except for Araneae (Fig. 2b). Symphyla and Protura were found in MP1 only. Statistical
287 computation highlighted significant differences for Symphyla and Coleoptera (adult and larvae) as
288 in May (Fig. 2a, b).

289 As previously established by other studies (Menta et al., 2018; 2013), the more adapted
290 microarthropod taxa were associated with stable soil conditions with a constant supply of organic
291 matter. Moreover, vegetal cover enhanced soil protection, mitigating thermal and hydric stresses
292 (Bird et al., 2004; Eaton et al., 2004). We can hypothesise that the richer and diversified cover
293 vegetation, and the higher organic matter content in MP1 improve soil conditions for the soil fauna
294 community. Indeed, the trend observed for the taxa was confirmed by the index computation (Table
295 2): Margalef richness index, Shannon diversity index, Simpson index and QBS-ar revealed higher
296 values in MP1 than MP2 in both periods. Only Pielou's Evenness was similar in the two soils
297 (Table 2), thereby highlighting the relevant presence of Acari and Collembola as major
298 communities, and thus not an equal distribution between the groups.

299

300 **Conclusion**

301 In conclusion, huge differences were observed in plant genera in the two serpentine micro-
302 habitats considered in this study, located in Mt Prinzera Natural Reserve in the Northern Apennines,
303 with the serpentinite scree rich in peculiar flora such as rare hyperaccumulator plant species. SOM
304 and Ni seem to be the main factors influencing the differences in abundance between bacterial taxa.
305 The results obtained for soil microarthropod communities confirmed this trend, showing that the
306 habitat with lower SOM and higher Ni concentration is less suitable for soil fauna. This may be the
307 result of concerted effects driven by both the higher Ni concentration and the lower SOM content,
308 as a consequence of scanty cover vegetation. Indeed, cover plants play a decisive role in
309 determining soil biodiversity, and due to the scarce vegetation in this area the soil is more affected
310 by chemical-physical variations in terms of temperature, moisture, SOM content, erosion, etc.
311 Investigating which taxa are more sensitive to high Ni concentrations can be a step towards
312 clarifying how serpentine soil affects both above- and belowground living communities. In
313 conclusion, the data observed with plant, microbial and microarthropod communities identify
314 different micro-habitats, in spite of proximal geographical position and the common metal-rich

315 substrate. This study may have been affected by the limitations imposed by the Natural Reserve of
316 Mt. Prinzera. However, the same approach could be repeated in different contexts (e.g. other
317 serpentine outcrops in Italy) to ensure general reliability of these results.

318

319 **Acknowledgements**

320 We thank Corrado Zanni for the plant genera and species classification by floristic relevés and Dr.
321 Michele Zanelli (Director of the Mt. Prinzera Natural Reserve) for giving the permission of
322 sampling the plants of *N. caerulea* Mt. Prinzera populations

323

324 **Funding**

325 This research was supported by funding to Prof. Anna Maria Sanangelantoni and to Dr. Giovanna
326 Visioli from FIL of the University of Parma Local Funding for Research.

327

328 **References**

329

330 Beier, S., Bertilsson, S., 2013. Bacterial chitin degradation-mechanisms and ecophysiological
331 strategies. *Front. Microbiol.* 4:149. doi: 10.3389/fmicb.2013.00149.

332 Bird, S.B., Coulson, R.N., Fischer, R.F., 2004. Change in soil and litter arthropods abundance
333 following tree harvesting and site preparation in a loblolly pine (*Pinus taeda* L.) plantation.
334 *Forest Ecol. Manag.* 202, 195–208.

335 Boyd, R.S., 2009. High-nickel insects and nickel hyperaccumulator plants: a review. *Insect Sci.* 16,
336 19-31.

337 Branco, S., 2010. Serpentine soils promote ectomycorrhizal fungal diversity. *Mol. Ecol.* 19, 5566-
338 5576.

339 Chiarucci, A., Bovini, I., Fattorini, L., 2003. Community dynamics of serpentine vegetation in
340 relation to nutrient addition and climatic variability. *J. Mediterr. Ecol.* 4, 23–30.

341 Coleman, R.G., Jove, C., 1992. Geological origin of serpentinites in: Baker, A.J.M., Proctor, J.,
342 Reeves, R.D. (Eds.), *The Vegetation of Ultramafic (Serpentine) Soils*. Intercept, Andover, pp.
343 1–17.

344 Eaton, R. J., Barbercheck, M., Buford, M, Smith, W., 2004. Effects of organic matter removal, soil
345 compaction, and vegetation control on Collembolan populations. *Pedobiologia* 48, 121-128

346 Edgar, RC., 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*
347 26, 2460-2461.

348 Ferrari, C., Lombini, A., Carpenè, B., 1992. Serpentine flora of the northern Apennines, Italy. In:
349 Baker, A.J.M., Proctor, J., Reeves, R.D. (Eds.), *The Vegetation of Ultramafic (Serpentine)*
350 *Soils*. Intercept, Andover, pp. 159–173.

351 Frizzi F., Masoni A., Çelikkol M., Palchetti E., Ciofi C., Chelazzi G., Santini G., 2017. Serpentine
352 soils affect heavy metal tolerance but not genetic diversity in a common Mediterranean ant.
353 *Chemosphere* 180, 326-334.

354 Galardi, F., Corrales, I., Mengoni, A., Pucci, S., Barletti, L., Barzanti, R., Arnetoli, M., Gabbrielli,
355 R, Gonnelli, C., 2007. Intra-specific differences in nickel tolerance and accumulation in the
356 Ni-hyperaccumulator *Alyssum bertolonii*. *Environ. Exp. Bot.* 60, 377-384

357 IUSS Working Group., 2015. *World Reference Base for Soil Resources 2014, update 2015*
358 *International soil classification system for naming soils and creating legends for soil maps.*
359 *World Soil Resources Reports*, 106.

360 Harrison, S., Rajakaruna, N., 2011. *Serpentine: The Evolution and Ecology of a Model System.*
361 University of California Press

362 Kielak, A. M., Barreto, C. C., Kowalchuk, G. A., van Veen, J. A., & Kuramae, E. E., 2016. The
363 ecology of Acidobacteria: moving beyond genes and genomes. *Front. Microbiol.* 7, 744.
364 <http://doi.org/10.3389/fmicb.2016.00744>

365 Lombini, A., Dinelli, E., Ferrari, C., Simoni, A., 1998. Plant-soil relationships in the serpentinite
366 screes of Mt. Prinzera (Northern Appennines, Italy). *J. Geochem. Explor.* 64, 19–23.

367 Lozupone, C., Knight, R., 2005. UniFrac: a new phylogenetic method for comparing microbial
368 communities. *Appl Environ Microbiol* 71, 8228-8235.

369 Mancabelli, L., Ferrario, C., Milani, C., Mangifesta, M., Turrioni, F., Duranti, S., Lugli, GA.,
370 Viappiani, A., Ossiprandi, M.C., van Sinderen, D, Ventura, M., 2016. Insides into the
371 biodiversity of the gut microbiota of broiler chickens. *Environ. Microbiol.* 18, 4727-4738

372 Margalef, R., 1958. Information theory in ecology. *Int. J. Gen. Syst.* 3, 36-71.

373 Mattarozzi, M., Manfredi, M., Montanini, B., Gosetti, F., Sanangelantoni, A.M., Marengo, E.,
374 Careri, M., Visioli, G., 2017. A metaproteomic approach dissecting major bacterial functions
375 in the rhizosphere of plants living in serpentine soil. *Anal. Bioanal. Chem.* 409, 2327-2339.

376 Mengoni, A., Barzanti, R., Gonnelli, C., Gabbrielli, R., Bazzicalupo, M., 2001. Characterization of
377 nickel-resistant bacteria isolated from serpentine soil. *Environ. Microbiol.* 3, 691-698.

378 Mengoni, A, Grassi, E., Barzanti, R., Biondi, E.G., Gonnelli, C., Kim, C.K., Bazzicalupo, M. 2004.
379 Genetic diversity of bacterial communities of serpentine soil and of rhizosphere of the
380 nickel-hyperaccumulator plant *Alyssum bertolonii*. *Microb. Ecol.* 48, 209-217.

381 Menta, C., Conti, F.D., Pinto, S., Bodini, A., 2018. Soil Biological Quality index (QBS-ar): 15
382 years of application at global scale. *Ecol. Indic.* 85, 773-780.

383 Menta, C., García- Montero, L.G., Conti, F.D., Pinto, S., 2013. The Role and Diversity of Soil
384 Fauna in: Different Woodlands, in *Woodlands. Structure, Species Diversity and Sustainable*
385 *Management.* Manzanares, L., (Ed), Nova Science Publisher, New York

386 Navarrete, A. A., Venturini, A. M., Meyer, K. M., Klein, A. M., Tiedje, J. M., Bohannon, B. J. M.,
387 Nüsslein, K., Tsai, S. M., Rodrigues, J. L. M., 2015. Differential response of Acidobacteria

388 subgroups to forest-to-pasture conversion and their biogeographic patterns in the western
389 Brazilian Amazon. *Front. Microbiol.* 6, 1443. <http://doi.org/10.3389/fmicb.2015.01443>

390 Oksanen, J., Blanchet, G.F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R.,
391 O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2017.
392 Community Ecology Package. Vers. 2.4-3. <https://cran.r-project.org>,
393 <https://github.com/vegandevs/vegan>

394 Pessoa-Filho, M., Barreto, C.C., DosReis Junior, F.B., Fragoso, R.R., Costa, F.S., Mendes De
395 Carvalho, De Andrade, L.R.M., 2015. Microbiological functioning, diversity and structure of
396 bacterial communities in ultramafic soils from a tropical savanna. *Antonie Van Leeuwenhoek*
397 107, 935-949.

398 Pielou, E.C., 1975. *Ecological Diversity*. John Wiley and Sons, New York

399 Pignatti, S., 1982. *Flora d'Italia*. Edagricole, Bologna, Vol.3

400 Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.,
401 2103. The SILVA ribosomal RNA gene database project: improved data processing and web-
402 based tools. *Nucl. Acids Res. (Database issue)*:D590-6. doi: 10.1093/nar/gks1219.

403 R Core Team, 2016. *R: A Language and environment for statistical computing*. The R foundation
404 for statistical computing, Vienna, Austria. <http://www.R-project.org/>

405 Rajkumar, M., Freitas, H., 2008. Influence of metal resistant-plant growth promoting bacteria on the
406 growth of *Ricinus communis* in soil contaminated with heavy metals. *Chemosphere* 71, 834-
407 842.

408 Seneviratne, M., Gunaratne S., Bandara, T, Weerasundara, L., Rajakaruna, N., Seneviratne, G.,
409 Vithanage, M., 2016. Plant growth promotion by *Bradyrhizobium japonicum* under heavy
410 metal stress. 105, 19-24

411 Shannon, C.E., Weaver, W., 1963. The mathematical theory of communication. Bell Syst. Tech. J.
412 27, 379-423 and 623-656.

413 Simpson, E.H., 1949. Measurement of diversity. Nature, 163, 688

414 Turrini, A., Agnolucci, M., Palla, M., Tomé, E., Tagliavini, M., Scandellari, F., Giovannetti, M.,
415 2017. Species diversity and community composition of native arbuscular mycorrhizal fungi in apple
416 roots are affected by site and orchard management. Appl. Soil. Ecol. 116, 42-54

417 Venter, A., Levanets, A., Sieber, S., Rajakaruna, N., 2015. A preliminary survey of the diversity of
418 soil algae and cyanoprokariotes on mafic and ultramafic substrates in South Africa. Aust. J.
419 Bot. 63,341-352.

420 Venter, A., Siebert, S., Rajakaruna, S.B., Levanets, A., Ismail, A., Allam, M., Peterson, B., Sanko,
421 T., 2018. Biological crusts of serpentine and non-serpentine soils from the Barberton
422 Greenstone belt of South Africa. Ecol. Res. 33629-640

423 Visioli, G., Vincenzi, S., Marmioli, M., Marmioli, N., 2012. Correlation between phenotype and
424 proteome in the Ni hyperaccumulator *Noccaea caerulescens* subsp. *caerulescens*. Environ.
425 Exp. Bot. 77, 156–164.

426 Visioli, G., Menta, C., Gardi, C., Conti, F.D., 2013. Metal toxicity and biodiversity in serpentine
427 soils: application of bioassay tests and microarthropod index. Chemosphere 90, 1267-1273.

428 Visioli, G., D'Egidio, S., Vamerali, T., Mattarozzi, M., and Sanangelantoni, A.M., 2014. Culturable
429 endophytic bacteria enhance Ni translocation in the hyperaccumulator *Noccaea caerulescens*.
430 Chemosphere 117, 538-544.

431 Zhang, B., Zhang, J., Liu, Y., Guo, Y., Shi, P., Gehong Wei, G., 2018. Biogeography and ecological
432 processes affecting root-associated bacterial communities in soybean fields across China. Sci.
433 Tot. Environ. 627, 20-27.

434

435 **Figure captions**

436 **Fig. 1:** Taxonomic distribution at phylum levels based on 16S rDNA reads across the two soil
437 samples (MP1 and MP2) according to the percentage of sequences considered >1% in abundance
438 respect to the total number of sequences at least in one sample (x-axis).

439 **Fig. 2:** Mean (\pm Std Dev) of microarthropod density (ind/m²) observed in MP1 and MP2 in May (a)
440 and October (b) (A: adult; L: larvae). * p<0.05

441

442

445

446 **Table 1:** Soil parameters in MP1 and MP2 sites: total Ca, K, Mg, Ni, Fe, Zn, Co contents ($\mu\text{g g}^{-1}$ of soil d.w), DTPA extractable fractions of metals
 447 indicated in brackets, pH values, water content (WC) % (Soil organic matter (SOM)%). Five replicates were taken from the 2 different sites MP1
 448 and MP2 in around an area of 10 m² each. Data are means of measures of five samples (\pm Std Dev). Significant different values between samples
 449 are indicated with an Asterisk * $p < 0.05$

Sites	Ca	K	Mg	Ni	Fe	Zn	Co	pH	WC (%)	SOM (%)
MP1	84 \pm 1.2	146 \pm 3.5	395 \pm 5.1	1700 \pm 24	45500 \pm 341	75 \pm 0.2	112 \pm 1.3*	7.10	25.38 \pm 0.2*	4.84 \pm 0.21*
				(34.8 \pm 3.4)		(1.4 \pm 0.05)	(7.4 \pm 0.04)			
MP2	82 \pm 2.1	160 \pm 2.7*	322 \pm 3.6	2032 \pm 34*	62120 \pm 236*	71 \pm 0.5	97 \pm 1.7	7.20	19.01 \pm 0.4	2.95 \pm 0.32
				(50 \pm 2.5*)		(1.7 \pm 0.06)	(8.5 \pm 0.1)			

450

451

453 **Table 2** – Mean (\pm Std Dev) of Margalef's Index (D), Shannon diversity index (H'), Evenness (J), Simpson dominance index (1-D), Soil Biological Quality
 454 index (QBS-ar) calculated for the microarthropod communities. * $p < 0.05$

Indices	May			October		
	MP1	MP2	p value	MP1	MP2	p value
D	1.86 \pm 0.52	0.92 \pm 0.46	-	1.27 \pm 0.12	0.93 \pm 0.41	*
H'	1.32 \pm 0.35	0.93 \pm 0.28	*	1.37 \pm 0.20	0.79 \pm 0.21	-
J	0.52 \pm 0.12	0.55 \pm 0.05	-	0.66 \pm 0.10	0.66 \pm 0.18	-
1-D	2.89 \pm 0.66	2.16 \pm 0.43	*	3.11 \pm 0.99	1.83 \pm 0.33	-
QBS-ar	182 \pm 31	89 \pm 42	*	102 \pm 8	43 \pm 11	*

455

Figure 1




Figure 2




Figure SI1: Pictures of MP1 and MP2 sites with GPS coordinates, and the corresponding *Nocca caerulea* plants growing at each site.



Figure SI2: Rarefaction curves generated for 16S rRNA gene sequences obtained from MP1 and MP2 samples. Panel a) displays the rarefaction curves using the Chao1 index. Panel b) displays rarefaction curves using the Shannon index




Figure S13: Taxonomic distribution at family levels based on 16S rDNA reads across the two soil samples (MP1 and MP2) according to the percentage of sequences considered $>1\%$ at least in one sample (x-axis)

Taxonomy at Family level (>1%)




Table SI1: Soil mini-pit description

MP1 –Skeletal Mollic Leptosol

Horizon	Depth (cm)	Description
A	0-30	10 YR 3/2, sandy-loam, moderate fine granular structure, common rock fragments, abundant fine roots.
A/C	>30	10 YR 3/3, sandy, massive, abundant rock fragments

MP2 – Orthoskeletal Leptosol

Horizon	Depth (cm)	Description
A	0-25	10 YR 4/3, sandy, moderate fine granular structure, abundant rock fragments, common fine and medium roots.
A/C	>25	10 YR 4/4, sandy, massive, very abundant rock fragments

Table SI2: Plants (Genus and species and the corresponding Families) found in MP1 and MP2 site in Spring 2015 (x represents presence)

Family	Genus/species	MP1 site	MP2 site
Asteraceae	<i>Sonchus</i>	x	
	<i>Artemisia annua</i>		x
	<i>Echinops</i>		x
	<i>Helichrysum italicum</i>		x
	<i>Hieracium pilosella</i>		x
Boraginaceae	<i>Myosotis</i>	x	
Brassicaceae	<i>Biscutella coronopilolia</i>		x
	<i>Noccaea caerulescens</i>	x	x
	<i>Alyssum bertolonii</i>		x
Caryophyllaceae	<i>Silene</i>	x	
Cyperaceae	<i>Carex</i>	x	
Cornaceae	<i>Cornus sanguinea</i>	x	
Crassulaceae	<i>Sedum montanum</i>		x
Euphorbiaceae	<i>Euphorbia cyparissias</i>	x	x
Fabaceae	<i>Lotus cornicatus</i>	x	
	<i>Robinia</i>	x	
Hypericaceae	<i>Hypericum perforatum</i>	x	
Lamiaceae	<i>Thymus</i>		x
Orobanchaceae			x
Plantaginaceae	<i>Globularia bisnagarica</i>		x
Poaceae	<i>Sesleria pichiana</i>		x
	<i>Bromus</i>	x	
	<i>Poa</i>	x	
Ranunculaceae	<i>Clematis vitalba</i>	x	
Rosaceae	<i>Rubus fruticosus</i>	x	
Rubiaceae	<i>Galium</i>	x	x
Umbelliferae			x

Table S13: Data filtering report of single sample runs. MP1=rhizosphere of *N. caerulea* plants at MP1 site MP2= rhizosphere of *N. caerulea* plants at MP2 site

Sample	Number of sequenced pe reads	Number of pe reads with mean quality > 20	Number of merged pe reads	Number of reads removed because of:						Final Read Number
				Human sequences	Outside bounds (100-400)	Ambiguous bases	Homopolymers > 7	Mismatch in primers >1	Reverse primer not found	
MP1	48739	48168	46668	65	0	0	23	2254	19	44307
MP2	104743	103865	100055	36	0	0	8	5532	29	94449