

Review

# Recent Advances in the Chemical Fixation of Carbon Dioxide: A Green Route to Carbonylated Heterocycle Synthesis

Renato Dalpozzo <sup>1,\*</sup>, Nicola Della Ca' <sup>2</sup>, Bartolo Gabriele <sup>1</sup> and Raffaella Mancuso <sup>1</sup>

<sup>1</sup> Dipartimento di Chimica e Tecnologie Chimiche, Università della Calabria, 87030 Arcavacata di Rende (Cs), Italy; bartolo.gabriele@unical.it (B.G.); raffaella.mancuso@unical.it (R.M.)

<sup>2</sup> Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale Parco Area delle Scienze, Università di Parma, 17/a, 43124 Parma, Italy; nicola.dellaca@unipr.it

\* Correspondence: renato.dalpozzo@unical.it; Tel.: +39-0984-49-2055

Received: 13 May 2019; Accepted: 2 June 2019; Published: 6 June 2019



**Abstract:** Carbon dioxide produced by human activities is one of the main contributions responsible for the greenhouse effect, which is modifying the Earth's climate. Therefore, post-combustion CO<sub>2</sub> capture and its conversion into high value-added chemicals are integral parts of today's green industry. On the other hand, carbon dioxide is a ubiquitous, cheap, abundant, non-toxic, non-flammable and renewable C<sub>1</sub> source. Among CO<sub>2</sub> usages, this review aims to summarize and discuss the advances in the reaction of CO<sub>2</sub>, in the synthesis of cyclic carbonates, carbamates, and ureas appeared in the literature since 2017.

**Keywords:** carbon dioxide; carbonylated heterocycles; carbonylation with carbon dioxide; carboxylation; cyclic carbamates; cyclic carbonates; cyclic ureas; oxazolidinones

## 1. Introduction

The combustion of fossil fuels produces large amounts of waste gases. Among them, carbon dioxide is one of the main contributions responsible for the greenhouse effect, which is modifying the Earth's climate [1]. Thus, post-combustion CO<sub>2</sub> capture and its conversion into high value-added chemicals are integral parts of today's green energy industry. In fact, the usage of carbon dioxide as a ubiquitous, cheap, abundant, non-toxic, non-flammable and renewable C<sub>1</sub> source has great importance from the viewpoint of both environmental protection and resource utilization.

Carbon dioxide has two polar carbonyl bonds, but its linear shape makes it non-polar, thus it is a thermodynamically and kinetically relatively stable molecule. This feature represents the major obstacle for CO<sub>2</sub> utilization, which still remains rather limited. The CO<sub>2</sub> conversion mainly depends on efficient activation by appropriate catalysts, to enhance electrophilicity of the central carbon atom.

Over the past decade, the usage of CO<sub>2</sub> as a building block to prepare valuable organic molecules has attracted increasing attention [2–9]. Even a journal completely devoted to CO<sub>2</sub> utilization started to be published by Elsevier in 2013. Moreover, Poliakov and Leitner proposed twelve principles of CO<sub>2</sub> chemistry that are a set of criteria for assessing the viability of different reactions in which CO<sub>2</sub> is the feedstock for making organic chemicals [10].



In addition to the carbon dioxide fixation [11,12], the classical carboxylation reaction [13–17], the reduction to carbon monoxide or formic acid [18–21], and polymerization [22–24], the preparation of useful heterocyclic carbonyl compounds such as cyclic carbonates, carbamates, and ureas attracted the interest of many research groups. Recently, an account has been published on the reaction of CO<sub>2</sub> with alcohols and amines into carbonates, ureas, and carbamates in the presence of ceric oxide as the

catalyst [25]. Another recent report described some porous catalysts for CO<sub>2</sub> capture and conversions and aimed to be a guide to prepare new porous catalysts for the synthesis of cyclic carbonates [26]. Finally, a very recent review covered the synthetic strategies to cyclic urethanes from amines and CO<sub>2</sub> and *N*-methylation, *N*-formylation from amines, CO<sub>2</sub> and H<sub>2</sub> [27].

Heterocyclic systems are widespread structures in natural products and artificial compounds (over 90% of new drugs contain at least one heterocyclic moiety). Heterocycles are also important building blocks for multistep synthesis. Therefore, the development of new methods for their synthesis is always welcome in the chemical community, in particular if these methods benefit from cheap, simple, and readily available starting materials. Notwithstanding the strong effort of Vessally [28–33] and others [34–37] to review large part of the synthesis of heterocycles by CO<sub>2</sub> fixation, many papers appeared in the literature in the last years, making an update necessary. Therefore, this review aims to summarize and discuss the recent advances in the reaction of CO<sub>2</sub>, for the synthesis of cyclic carbonates, carbamates, and ureas.

The synthesis of these heterocycles can occur by two main mechanisms on molecules carrying a nucleophilic and an electrophilic site:

- The nucleophilic site (eventually enforced by a base) can add to the carbon atom, to form the corresponding carboxylate, which in turn adds at the electrophilic site to close the cycle (Scheme 1, Equation (1)).
- A catalyst is able to fix the carbon dioxide leading to a zwitterionic species. The carboxylate moiety of this activated species attacks an electrophilic site of the substrate molecule. Then the nucleophilic site closes the cycle, releasing the catalyst (Scheme 1, Equation (2)). *N*-heterocyclic carbenes (NHCs), *N*-heterocyclic olefins (NHOs), phosphorus ylides, polyoxometalates (POMs), ionic liquids (ILs), frustrated Lewis pairs (FLPs), metal-organic frameworks (MOFs) or superbases have been used for this scope.




Scheme 1. General mechanistic survey.

## 2. Cyclic Carbonates

Cyclic carbonates find applications as electrolytes in lithium ion batteries in pharmaceuticals, products for agriculture, and as starting materials for polycarbonates [38–40]. Moreover, propylene

carbonate is considered as a green polar aprotic solvent, because it has high boiling and flash points, low odor level and toxicity and biodegradability. The production of cyclic carbonates from CO<sub>2</sub> is fully atom-economical and uses a safe gas, instead of toxic gases such as phosgene.

The reaction of carbon dioxide with epoxides [41–45] as well as with propargyl alcohols [2] has been widely reviewed in the past years. However, in the last two years a large production of research articles on this topic has appeared in the literature and they are summarized in Table 1. The generally accepted mechanism for the reaction of epoxides and carbon dioxide described in Table 1 is depicted in Scheme 2.



**Scheme 2.** The mechanism of the addition of CO<sub>2</sub> to epoxides.

**Table 1.** Catalysts for the reaction of epoxides and carbon dioxide (R and R<sup>1</sup> see Scheme 2).

Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous metal catalysts</b>					
1	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	1.2	 Bu <sub>4</sub> NBr (0.3 mol% each), 80 °C, 6 h	16–92	[46]
2	R=H, R <sup>1</sup> =Me, Ph R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	2.0	 (0.2 mol%) Bu <sub>4</sub> NBr (0.4 mol%), 120 °C, 5–18 h	87–99	[47]
3	R=H, R <sup>1</sup> =Me, Et, Bu, Ph, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , CH <sub>2</sub> Cl,  Cl, Me <sub>2</sub> , CH <sub>2</sub> OPh, CH <sub>2</sub> OBn, CH <sub>2</sub> O(9 <i>H</i> -carbazol-4-yl), R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> , CH <sub>2</sub> CH(vinyl)(CH <sub>2</sub> ) <sub>2</sub>	0.5	 (0.3 mol%), Bu <sub>4</sub> NBr (3 mol%), 25 °C (80 °C for internal epoxides), 24 h	49–99	[48]

Table 1. Cont.






Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous metal catalysts</b>					
4	R=H, R <sup>1</sup> =Ph, Me, Bu, <i>n</i> -C <sub>8</sub> H <sub>17</sub> , <i>n</i> -C <sub>10</sub> H <sub>23</sub> , CH <sub>2</sub> OH, CH <sub>2</sub> OPh, CH <sub>2</sub> Cl, 4-ClC <sub>6</sub> H <sub>4</sub> , 4-BrC <sub>6</sub> H <sub>4</sub>	1.0	 Bu <sub>4</sub> NBr, 50 °C, 24 h aliphatic epoxides (0.1:0.2 mol%) aromatic epoxides (0.25:0.5 mol%)	50–94	[49]
5	R=H, R <sup>1</sup> =Me <sub>2</sub> , R=Ph, R <sup>1</sup> =Me, Ph (only <i>trans</i> ) R=R <sup>1</sup> =Me, (CH <sub>2</sub> ) <sub>4</sub> , (CH <sub>2</sub> ) <sub>3</sub>		 Bu <sub>4</sub> NBr, (1:2 mol%), 50 °C, 24 h	17–60	
6	R=H, R <sup>1</sup> =Me, Et, Ph, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , <i>n</i> -C <sub>8</sub> H <sub>17</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> Br, 4-ClC <sub>6</sub> H <sub>4</sub> , CH <sub>2</sub> Ometacrylate, CH <sub>2</sub> Oallyl,	1.0	 ISA6 (1 mol%) 50 °C, 2–12 h	90–99	[50]
7	R=H, R <sup>1</sup> =Ph, Me, Et, Bu, Bn, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-ClC <sub>6</sub> H <sub>4</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> O- <i>i</i> -Pr, CH <sub>2</sub> OBn, CH <sub>2</sub> Oallyl, CH <sub>2</sub> Opropargyl	0.1	 SalenCoI (1 mol%), Ph <sub>3</sub> P=CHCHO (1 mol%), 25 °C, 24–36 h	79–95	[51]
8	R=H, R <sup>1</sup> =Ph, 4-ClC <sub>6</sub> H <sub>4</sub> , 4-FC <sub>6</sub> H <sub>4</sub> , <i>n</i> -C <sub>6</sub> H <sub>13</sub> , Pr, (CH <sub>2</sub> ) <sub>2</sub> CH=CH <sub>2</sub> , CH <sub>2</sub> Oallyl, CH <sub>2</sub> OBn, CH <sub>2</sub> Cl, CH <sub>2</sub> NMe <sub>2</sub>	0.1	 (10 mol%), 40 °C, 24 h	86–99	[52]

Table 1. Cont.






Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous metal catalysts</b>					
9	R=H, R <sup>1</sup> =Et, Me <sub>2</sub> , (R)-Me, (R)-Ph, 4-FC <sub>6</sub> H <sub>4</sub> , <i>n</i> -C <sub>6</sub> H <sub>13</sub> , (CH <sub>2</sub> ) <sub>2</sub> CH=CH <sub>2</sub> , vinyl, CH <sub>2</sub> Cl, CH <sub>2</sub> OH, CH <sub>2</sub> O <i>t</i> -Bu, CH <sub>2</sub> OPropargyl, CH <sub>2</sub> OCH <sub>2</sub> furyl, CH <sub>2</sub> Ometacrylate 	1.0	CaI <sub>2</sub> /PEG-500 (5 mol%), 25 °C, 24 h	84–99	
10	R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>3</sub> , (CH <sub>2</sub> ) <sub>4</sub> , (CH <sub>2</sub> ) <sub>5</sub> , (CH <sub>2</sub> ) <sub>6</sub> , Me, CH <sub>2</sub> OCH <sub>2</sub> , CH <sub>2</sub> CH=CHCH <sub>2</sub> , CH <sub>2</sub> CH(Me)(CH <sub>2</sub> ) <sub>2</sub> , CH <sub>2</sub> CH(vinyl)(CH <sub>2</sub> ) <sub>2</sub> ,  R=C <sub>8</sub> H <sub>17</sub> , R <sup>1</sup> =(CH <sub>2</sub> ) <sub>7</sub> CO <sub>2</sub> Me	A: 2.0 B: 5.0	CaI <sub>2</sub> /PEG-500 (5 mol%) A: 70 °C, 24 h B: 90 °C, 48 h	29–98 <i>cis:trans</i> 53:47 to >99:1 <i>cis:trans</i> 11:89 (R=R <sup>1</sup> = <i>t</i> -Me)	[53]
11	R=H, R <sup>1</sup> =CH <sub>2</sub> O- <i>t</i> -Bu, CH <sub>2</sub> OH, CH <sub>2</sub> OMe, CH <sub>2</sub> OPropargyl, CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl, CH <sub>2</sub> OCH <sub>2</sub> furyl, CH <sub>2</sub> Ometacrylate, CH <sub>2</sub> OSiMe <sub>2</sub> - <i>t</i> -Bu, CH <sub>2</sub> OCH <sub>2</sub> CF <sub>2</sub> CHF <sub>2</sub> , CH <sub>2</sub> OCH <sub>2</sub> C <sub>2</sub> F <sub>5</sub> , CH <sub>2</sub> OCH <sub>2</sub> (CF <sub>2</sub> ) <sub>3</sub> CHF <sub>2</sub> , Me, Et, Bu, <i>n</i> -C <sub>6</sub> H <sub>13</sub> ,  (R <sup>2</sup> = Me, CH <sub>2</sub> Cl, Ph, CO <sub>2</sub> Me)	0.1	 2 I <sup>-</sup> (5 mol%), 23 °C, 24 h	23–99	
12	R=R <sup>1</sup> =Me, (CH <sub>2</sub> ) <sub>3</sub> , (CH <sub>2</sub> ) <sub>4</sub> , (CH <sub>2</sub> ) <sub>5</sub> , (CH <sub>2</sub> ) <sub>6</sub> , CH <sub>2</sub> OCH <sub>2</sub> , CH=CH(CH <sub>2</sub> ) <sub>2</sub> , CH <sub>2</sub> CH(Me)(CH <sub>2</sub> ) <sub>2</sub> , CH <sub>2</sub> CH(vinyl)(CH <sub>2</sub> ) <sub>2</sub> , Me, Ph; R=Ph, R <sup>1</sup> =Me, CO <sub>2</sub> Et; R=4-MeOC <sub>6</sub> H <sub>4</sub> , R <sup>1</sup> = CO <sub>2</sub> Me	1.0	 2 I <sup>-</sup> (5 mol%), 45 °C, 48 h	4–98%	[54]

Table 1. Cont.









Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous metal catalysts</b>					
13	<p>R=C<sub>8</sub>H<sub>17</sub>,  R<sup>1</sup>=(CH<sub>2</sub>)<sub>7</sub>CO<sub>2</sub>Me,  (CH<sub>2</sub>)<sub>7</sub>CO<sub>2</sub>Et,  (CH<sub>2</sub>)<sub>7</sub>CO<sub>2</sub>(CH<sub>2</sub>)<sub>5</sub>-i-Pr,  (CH<sub>2</sub>)<sub>11</sub>CO<sub>2</sub>Me,  C<sub>6</sub>H<sub>11</sub> (CH<sub>2</sub>)<sub>7</sub>CO<sub>2</sub>Me</p>  <p>R=H,  R<sup>1</sup>=(CH<sub>2</sub>)<sub>8</sub>CO<sub>2</sub>Me,  epoxidized methyl  <i>O</i>-acetyl ricinoleate,  methyl soyate,  sunflower, soybean,  linseed oils</p>	0.5	 <p>(5 mol%), Ph<sub>3</sub>P (5 mol%), 45 °C, 24 h</p>	55–98	[55]
14	<p>(+)-limonene oxide,  (+)-limonene dioxide,  epoxidized citronellyl  propionate,</p> 	5.0	 <p>(5 mol%), Ph<sub>3</sub>P (5 mol%), 45 °C, 48 h</p>	19–81	[55]
15	<p>R=H, R<sup>1</sup>=Me, Et, Bu,  Ph, <i>n</i>-C<sub>6</sub>H<sub>13</sub>, CH<sub>2</sub>Cl,  CH<sub>2</sub>OH, CH<sub>2</sub>OPh,  CH<sub>2</sub>OBn, CH<sub>2</sub>OAllyl,</p>  <p>R=R<sup>1</sup>=(CH<sub>2</sub>)<sub>4</sub></p>	0.1	 <p>(10 mol%), 50 °C, 6 h</p>	25–99	[56]
16	<p>R=H, R<sup>1</sup>=Ph,  CH<sub>2</sub>O-<i>i</i>-Pr, CH<sub>2</sub>OAllyl,  CH<sub>2</sub>OPh, CH<sub>2</sub>Cl</p>	0.1	Nanocrystalline MgO (12.5 mg per mmole epoxide), Bu <sub>4</sub> NBr (50 mol%), r. t., 4–8 h	42–99	[57]
17	<p>R=H, R<sup>1</sup>=Me, Et, Ph,  CH<sub>2</sub>Cl</p>	1.6	 <p>(0.1 mol%), Dimethylaminopyridine (0.2 mol%), 100 °C, 2 h</p>	54–97	[58]
18	<p>R=H, R<sup>1</sup>=Me, Bu, Ph,  <i>n</i>-C<sub>10</sub>H<sub>23</sub>, CH<sub>2</sub>OAllyl,  CH<sub>2</sub>OMe, CH<sub>2</sub>Cl,  CH<sub>2</sub>OCH<sub>2</sub>furyl  R=R<sup>1</sup>=(CH<sub>2</sub>)<sub>3</sub>, <i>c</i>-Me</p>	0.1	 <p>(0.2 mol%), 35 °C, 6 h</p>	26–65	[59]

Table 1. Cont.






Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous metal catalysts</b>					
19	R=H, R <sup>1</sup> =Ph, 4-ClC <sub>6</sub> H <sub>4</sub> , Bu, CH <sub>2</sub> Cl	5.0	 (2 mol%), 50 °C, 24 h	98	[60]
20	R=H, R <sup>1</sup> =Me, Ph, Et, CH <sub>2</sub> OAllyl, CH <sub>2</sub> OH, CH <sub>2</sub> OPh, CH <sub>2</sub> Cl, 	3.0	 (0.04 mol%), Bu <sub>4</sub> NBr (0.4 mol%), 100 °C, 6 h	27–100	[61]
21	R=H, R <sup>1</sup> =Me, Et, Bu, Ph, <i>n</i> -C <sub>8</sub> H <sub>17</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl	2.0	 IL-Zn-TPP (0.1 mol%), 60 °C, 8–30 h	72–98	[62]
22	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> ,	1.6	 (0.1 mol%), Dimethylaminopyridine (0.2 mol%), 100 °C, 2 h	6–89	[63]
23	R=H, R <sup>1</sup> =Me, Et, Ph,	0.1	Fe(O <sub>2</sub> CNEt <sub>3</sub> ) <sub>3</sub> (1 mol%), Bu <sub>4</sub> NBr (2 mol%), 25 °C, 24 h	87–90	[64]
<b>Homogenous organocatalysts</b>					
24	R=H, R <sup>1</sup> = Me, Et, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> =CH	0.1–0.5	Ascorbic acid (2–4 mol%), Bu <sub>4</sub> NI (4–8mol%), r.t.–60 °C, 23 h	82–97	[65]

Table 1. Cont.







Entry	R, R <sup>1</sup>	P <sub>CO<sub>2</sub></sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous organocatalysts</b>					
25	R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> , (CH <sub>2</sub> ) <sub>3</sub> , CH <sub>2</sub> OCH <sub>2</sub> , CH <sub>2</sub> N(Bn)CH <sub>2</sub> , Me  (c:t=94:6), R=Ph, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> OMe, CO <sub>2</sub> Et (c:t=12:88 to 1:99)	3.0	 (3 mol%), Et <sub>4</sub> NBr (6 mol%), 80 °C, 18 h	53–90	[66]
26	R=H, R <sup>1</sup> =Ph, Pr, 4-ClC <sub>6</sub> H <sub>4</sub> , 4-FC <sub>6</sub> H <sub>4</sub> , (CH <sub>2</sub> ) <sub>2</sub> CH=CH <sub>2</sub> , CH <sub>2</sub> OAllyl, CH <sub>2</sub> OBn, CH <sub>2</sub> Cl	0.1	NEt <sub>3</sub> •HI (10 mol%), 40 °C, 24 h	82–99	[67]
27	R=H, R <sup>1</sup> =(CH <sub>2</sub> ) <sub>8</sub> CO <sub>2</sub> Me, R=C <sub>8</sub> H <sub>17</sub> , R <sup>1</sup> =(CH <sub>2</sub> ) <sub>7</sub> CO <sub>2</sub> Me, (CH <sub>2</sub> ) <sub>7</sub> CO <sub>2</sub> Et, (CH <sub>2</sub> ) <sub>7</sub> CO <sub>2</sub> (CH <sub>2</sub> ) <sub>5</sub> -i-Pr, (CH <sub>2</sub> ) <sub>9</sub> CO <sub>2</sub> Me, R=CH <sub>2</sub> CHOH(CH <sub>2</sub> ) <sub>5</sub> Me, CH <sub>2</sub> CHOAc(CH <sub>2</sub> ) <sub>5</sub> Me R <sup>1</sup> =(CH <sub>2</sub> ) <sub>7</sub> CO <sub>2</sub> Me,  epoxidized sunflower, soybean, linseed oils, methyl soyate	2.5	 (5 mol%), 80 °C, 24 h	62–99	[68]
28	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl, 4-FC <sub>6</sub> H <sub>4</sub> , 4-ClC <sub>6</sub> H <sub>4</sub> , 4-BrC <sub>6</sub> H <sub>4</sub> R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> , (CH <sub>2</sub> ) <sub>3</sub>	0.1	 (10 mol%), Bu <sub>4</sub> N <sup>+</sup> I <sup>-</sup> (10 mol%), 80 °C, 30 h	78–92	[69]
29	R=H, R <sup>1</sup> =Me, Me <sub>2</sub> , Bu, CH <sub>2</sub> OPh, CH <sub>2</sub> Cl R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	0.1	1,5-diazabicyclo(5.4.0)undec-7-ene (DBU) (5 mol%), N-iodosuccinimide (NIS) (5 mol%) 60 °C, 12 h	71–99	[70]
30	R=H, R <sup>1</sup> =Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> Br, CH <sub>2</sub> OAllyl, CH <sub>2</sub> O- <i>t</i> -Bu, CH <sub>2</sub> OPh	0.1	4-(Dimethylamino)pyridinium bromide (DMAPHBr) (1 mol%) 120 °C, 4 h	94–98	[71]
31	R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>3</sub> , (CH <sub>2</sub> ) <sub>4</sub> , CH <sub>2</sub> CH(vinyl)(CH <sub>2</sub> ) <sub>2</sub> , CH <sub>2</sub> OCH <sub>2</sub> , <i>c</i> -Ph, <i>t</i> -Ph, <i>c</i> -Me, <i>t</i> -Me	0.1–0.4	DBU (5 mol%), <i>n</i> Bu <sub>4</sub> NCl (10 mol%), 120 °C, 24 h.	41–96	[72]
32	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl R=R <sup>1</sup> =H, (CH <sub>2</sub> ) <sub>4</sub>	2.0	 (0.25 mol%), 130 °C, 4 h	52–96	[73]

Table 1. Cont.

Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous organocatalysts</b>					
33	R=H, R <sup>1</sup> =Me, Ph, Bu, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , CH <sub>2</sub> Cl	0.5	 (0.3 mol%), 130 °C, 12 h	96–99	[74]
34	R=H, R <sup>1</sup> =Me, Bu, Ph, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> OH, CH <sub>2</sub> OPh, CH <sub>2</sub> OMe, CH <sub>2</sub> Oallyl R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> , (CH <sub>2</sub> ) <sub>3</sub> , CH <sub>2</sub> OCH <sub>2</sub>	0.5	 (1 mol%), 80 °C, 18 h	41–97	[75]
35	R=H, R <sup>1</sup> =Me, Ph, Bu, CH <sub>2</sub> Cl, CH <sub>2</sub> Oph R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> ,	3.0	 β-cyclodextrin (1.5 mol%), 130 °C, 10 h	63–98	[76]
36	R=H, R <sup>1</sup> =Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> Br, CH <sub>2</sub> Oph	0.1	 (25 mol%), 30–60 °C, 12*30 h	94–95	[77]
37	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> O- <i>t</i> -Bu, CH <sub>2</sub> OBu, CH <sub>2</sub> Oph, CH <sub>2</sub> Oallyl,  R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	2.0	 (0.25 mol%), 120 °C, 3 h	67–99	[78]
38	R=H, R <sup>1</sup> =CH <sub>2</sub> Cl, (CH <sub>2</sub> ) <sub>2</sub> CH=CH <sub>2</sub> , CH <sub>2</sub> Oallyl, CH <sub>2</sub> Oph, Me, Bu, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , Ph R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	0.1	 (5 mol%), Bu <sub>4</sub> Ni (5 mol%), 25–60 °C, 24 h	14–95	[79]

Table 1. Cont.







Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous organocatalysts</b>					
39	R=H, R <sup>1</sup> =Bu, Me, Et, <i>n</i> -C <sub>8</sub> H <sub>17</sub> , Ph, 4-ClC <sub>6</sub> H <sub>4</sub> , 4-BrC <sub>6</sub> H <sub>4</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> OH, CH <sub>2</sub> OPh		 (1 mol%) Bu <sub>4</sub> Ni (1 mol%), 90 °C, 2 h	R <sup>2</sup> =Ph 52–96 R <sup>2</sup> =Bu 97–99	
40	R=H, R <sup>1</sup> =Bu, Me, Et, <i>n</i> -C <sub>8</sub> H <sub>17</sub> , Ph, 4-ClC <sub>6</sub> H <sub>4</sub> , 4-BrC <sub>6</sub> H <sub>4</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> OH, CH <sub>2</sub> OPh, CH <sub>2</sub> OCH <sub>2</sub> (2-furyl), CH <sub>2</sub> OCO(2-furyl),  R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>3</sub> , (CH <sub>2</sub> ) <sub>4</sub>	1.0	 (0.75 mol%), 80 °C, 1 h	49–99	[80]
41	R=H, R <sup>1</sup> =Bu, Me, Ph, CH <sub>2</sub> OAllyl, CH <sub>2</sub> OPh, CH <sub>2</sub> OBn, CH <sub>2</sub> Cl, CH <sub>2</sub> Br, CH <sub>2</sub> Morpholin-4-yl	2.0	 CDC-CO <sub>2</sub> (5 mol%) 100 °C, 12 h	77–96	[81]
42	R=H, R <sup>1</sup> =Bu, Ph, C <sub>8</sub> H <sub>17</sub> , C <sub>12</sub> H <sub>25</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> OEt, CH <sub>2</sub> OPh, CH <sub>2</sub> OCOC(Me)=CH <sub>2</sub> R=R <sup>1</sup> = <i>t</i> -Me, (CH <sub>2</sub> ) <sub>3</sub> , (CH <sub>2</sub> ) <sub>4</sub>	1.7	 (1 mol%) Bu <sub>4</sub> Ni (1 mol%), 100 °C, 15 h	48–98	[82]
43	R=H, R <sup>1</sup> =Me, CH <sub>2</sub> Cl, CH <sub>2</sub> OH, CH <sub>2</sub> OMe, CH <sub>2</sub> O- <i>t</i> -Bu, CH <sub>2</sub> OPh	1.0	 (2 mol%) r. t., 6–12 h	90–99	[83]

Table 1. Cont.





Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Homogenous organocatalysts</b>					
44	R=H, R <sup>1</sup> =Me, Et, Bu, (CH <sub>2</sub> ) <sub>2</sub> CH=CH <sub>2</sub> , <i>n</i> -C <sub>6</sub> H <sub>13</sub> , Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> OBu, CH <sub>2</sub> OPh, CH <sub>2</sub> OAllyl, Me <sub>2</sub> , Me(CH <sub>2</sub> Cl), CH <sub>2</sub> O(9 <i>H</i> -carbazol-4-yl)	0.1	 (4 mol%), r.t., 24 h	95–73	[84]
45	R=R <sup>1</sup> =Me, (CH <sub>2</sub> ) <sub>3</sub> , (CH <sub>2</sub> ) <sub>4</sub> , CH <sub>2</sub> OCH <sub>2</sub> , R=C <sub>8</sub> H <sub>17</sub> , R <sup>1</sup> =(CH <sub>2</sub> ) <sub>7</sub> CO <sub>2</sub> Me, (CH <sub>2</sub> ) <sub>7</sub> CO <sub>2</sub> Et, (CH <sub>2</sub> ) <sub>11</sub> CO <sub>2</sub> Me, 	0.1	 (4 mol%), 80 °C, 24 h	43–95	
46	R=H, R <sup>1</sup> =Ph, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , Bu, CH <sub>2</sub> OAllyl, CH <sub>2</sub> OBn, CH <sub>2</sub> Cl, CH <sub>2</sub> OH,	0.1	 (10 mol%), 70 °C, 4 h	65–87	[85]
<b>Heterogeneous catalysts</b>					
47	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> Br,	0.1	HUST-1-Co (0.8 mg/mmol), Bu <sub>4</sub> NBr (7 mol%), rt, 30–48 h	93–97	[86]
48	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , <i>n</i> -C <sub>10</sub> H <sub>21</sub> , CH <sub>2</sub> OAllyl	1.0	Al-HPC (0.25 mmol%), Bu <sub>4</sub> NBr (2 mol%), 40 °C, 1–24 h	85–99	[87]
49	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , <i>n</i> -C <sub>10</sub> H <sub>21</sub> , CH <sub>2</sub> OAllyl R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	1.0	Al-iPOP-1 or Al-iPOP-2 (0.1 mol%) 40 °C, 3–36 h	8–99 (1) 14–99 (2)	[88]
50	R=H, R <sup>1</sup> =CH <sub>2</sub> Cl	0.1	Zn-Co/ZIF, 80 °C, 24 h	57	[89]
51	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> OPh, CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl, R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> , (CH <sub>2</sub> ) <sub>6</sub>	0.7	Zn-Co/ZIF-67 (Zn:Co 1:9) (50 mg), 100 °C, 2–18 h	8–99	[90]
52	R=H, R <sup>1</sup> =Me, Ph, Bn, CH <sub>2</sub> Cl	1.0	cCTF-500 (4% wt), 90 °C, 12 h	36–99	[91]
53	R=H, R <sup>1</sup> =Ph, Bu, CH <sub>2</sub> OPh, CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	0.4	PGDBr-5-2OH (1.9 mol %), 70 °C, 4–48 h	90–98	[92]
		0.1	PGDBr-5-2OH (1.9 mol %), 70 °C, 24–96 h	90–97	
		0.1	PGDBr-5-2OH (1.9 mol %), <i>n</i> -Bu <sub>4</sub> NI (8 mol %), r. t., 18–120 h	80–93	
54	R=H, R <sup>1</sup> =Me, Et, CH <sub>2</sub> Cl, CH <sub>2</sub> Br	0.1	Cu <sub>2</sub> [(C <sub>20</sub> H <sub>12</sub> N <sub>2</sub> O <sub>2</sub> )(COO) <sub>4</sub> ] (0.2 mol%), Bu <sub>4</sub> NBr (8 mol%), r.t., 48 h	88–96	[93]

Table 1. Cont.






Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Heterogeneous catalysts</b>					
55	R=H, R <sup>1</sup> =Ph, Bu, CH <sub>2</sub> OPh, CH <sub>2</sub> OBu, CH <sub>2</sub> OBn, CH <sub>2</sub> Cl, 	0.1	[Co <sub>2</sub> (resorcin-4-arene <sub>0.5</sub> )V <sub>4</sub> O <sub>12</sub> ] $\cdot$ 3DMF 5H <sub>2</sub> O (0.2 mol%, based on V), Bu <sub>4</sub> NBr (5 mol%), 80 °C, 12 h	87–99	[94]
56	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> OPh	0.8	Zn <sub>2</sub> [1,4-(CO <sub>2</sub> ) <sub>2</sub> C <sub>6</sub> H <sub>4</sub> ](DABCO) (17 mg per mmol substrate), 100 °C, 12–30 h	90–99	[95]
57	R=H, R <sup>1</sup> =Me, Pr, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> Br, CH <sub>2</sub> OPh	0.1	ZnO@NPC-Ox-700 (50 mg per mmol substrate), Bu <sub>4</sub> NBr (20 mol%), 25–60 °C, 1–3 h	85–99	[96]
58	R=H, R <sup>1</sup> =CH <sub>2</sub> Cl	1.0	KCO <sub>3</sub> (C <sub>6</sub> H <sub>4</sub> O <sub>7</sub> ) (C <sub>6</sub> H <sub>5</sub> O <sub>7</sub> ) (H <sub>2</sub> O) <sub>2</sub> (UTSA-16) (0.15 mmol), 120 °C, 6 h	98	[97]
59	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl	1.5	UDIL-I-60%U (5% wt) 120 °C, 3 h	83–99	[98]
60	R=H, R <sup>1</sup> =Bu	0.5	 (0.89 mol%), 80 °C, 18 h	>99	[65]
61	R=H, R <sup>1</sup> =CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl, CH <sub>2</sub> OPh, <i>n</i> -C <sub>6</sub> H <sub>13</sub>	0.1	 (5 mol %) 100 °C, 24 h	78–99	[99]
62	R=H, R <sup>1</sup> =Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl, <i>n</i> -C <sub>6</sub> H <sub>13</sub> R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	2.5	 (0.5 mol%) 130 °C, 15 h (140 °C, 86 h for cyclohexene oxide)	94–99	[100]
63	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> OPh R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	2.5	 (1.4 mol%), 140 °C, 3 h (24 h for cyclohexene oxide)	66–98	[101]
64	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl,	1.0	CBAP-1(EDA) (2 mol% of N sites), 130 °C, 4 h	77–98	[102]
65	(CH <sub>2</sub> ) <sub>2</sub> CH=CH <sub>2</sub>		CBAP-1(EDA) (2 mol% of N sites), Bu <sub>4</sub> NBr (1.8 mol%), 25 °C, 36 h	81–95	

Table 1. Cont.










Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Heterogeneous catalysts</b>					
66	R=H, R <sup>1</sup> =Me, Et, Bu, Ph, <i>n</i> -C <sub>6</sub> H <sub>13</sub> , <i>t</i> -BuO, (CH <sub>2</sub> ) <sub>8</sub> CO <sub>2</sub> Me, CH <sub>2</sub> Cl, (CH <sub>2</sub> ) <sub>2</sub> CH=CH <sub>2</sub> , <i>i</i> -PrO, CH <sub>2</sub> OAllyl, Me(CH <sub>2</sub> Cl) R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> , R=C <sub>8</sub> H <sub>17</sub> , R <sup>1</sup> =(CH <sub>2</sub> ) <sub>7</sub> CO <sub>2</sub> Me	1.0	 (2 mol%), 90 °C, 6 h	23–98	[103]
67	R=H, R <sup>1</sup> =Et, Bu, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> Br, CH <sub>2</sub> OPh	0.1	COF-JLU7 (0.5 mol%), Bu <sub>4</sub> NBr (0.5 mol%), 40 °C, 48 h	61–99	[104]
68	R=H, R <sup>1</sup> =Ph, CH <sub>2</sub> Cl	0.7	 (0.1 mol%), 100 °C, 4–10 h	90–100	[105]
69	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> OAllyl	2.0	 (2% wt), 120 °C, 2–15 h	80–95	[106]
70	R=H, R <sup>1</sup> =Me, Ph, 4-FC <sub>6</sub> H <sub>4</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> Br	0.1	 (6 mg per mmol of substrate), 100 °C, 24 h	95–98	[107]
71	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl	2.0	Zn-C <sub>3</sub> N <sub>4</sub> (25) (3.5 mg per mmol of substrate), KI (1.5 mol%), 130–150 °C, 5 h	92–99	[108]
72	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl, Me <sub>2</sub>	0.8	 (5 mg per mmol of substrate), 80 °C, 4 h	33–93	[109]
73	R=H, R <sup>1</sup> =Me, Ph, Bu, CH <sub>2</sub> Cl, CH <sub>2</sub> OPh R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	1.0	 (1 mol%), 100 °C, 2.5 H	81–99	[110]


Table 1. Cont.

Entry	R, R <sup>1</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
<b>Heterogeneous catalysts</b>					
74	R=H, R <sup>1</sup> =Me, Ph, Bu, CH <sub>2</sub> Cl, CH <sub>2</sub> OBu R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	1.5	SBA-Zn-TPy <sup>+</sup> PBr <sup>-</sup> (0.1 mol%), 120 °C, 4.5–5 h	23–99	[111]
75	R=H, R <sup>1</sup> =Me, Ph, Bu, CH <sub>2</sub> Cl, CH <sub>2</sub> OBu R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	1.5	ZnTPy-BIM4/CNTs-3, (POSS-Imi, 0.07 mol%) 120 °C, 2.5–24 h	51–98	[112]
76	R=H, R <sup>1</sup> =Ph, Bn, 4-ClC <sub>6</sub> H <sub>4</sub> , 4-FC <sub>6</sub> H <sub>4</sub> , CH <sub>2</sub> Cl, CH <sub>2</sub> OPh, CH <sub>2</sub> OBn	0.1	 (0.5 mol%), 120 °C, 10 h	94–96	[113]
77	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub> , (CH <sub>2</sub> ) <sub>3</sub>	3.0	RH or RD Au/Zn-MOF nanocages (3.2 mg/mmol), 70 °C, 6 h	95–99	[114]
78	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> Br, CH <sub>2</sub> OBu	0.1	[Ni(4,6-bis(triazol-1-yl)isophthalate)(4,6- bis(triazol-1-yl)isophthalic acid)] 2DMF·2H <sub>2</sub> O (1 mol%), Bu <sub>4</sub> NBr (10 mol%), 25 °C, 48 h	40–99	[115]
79	R=H, R <sup>1</sup> =Me, Et, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> Br, CH <sub>2</sub> OPh	3.0	CNT-NHC-Ag (8 mg/mmol) 4-dimethylaminopyridine (0.5 mol%) 120 °C, 8 h	30–92	[116]
80	R=H, R <sup>1</sup> =Me, Ph, CH <sub>2</sub> Cl, CH <sub>2</sub> OH,  R=R <sup>1</sup> =(CH <sub>2</sub> ) <sub>4</sub>	4.0	POSS-Imi (0.013–0.133 mol%), 150 °C, 3–16 h	30–99	[117]

Entry 1: the activity and selectivity of Zn(3,5-dimethyl-1,2,4-triazole)F was maintained after five recycles, while yield decreased in each cycle with Zn(3-amino-1,2,4-triazole)<sub>2</sub>. Entry 3: enantiopure (*S*)-benzyloxymethylepoxide was converted into the corresponding cyclic carbonates with retention of the configuration, while both (*R*) and (*S*)-styrene oxides gave partial racemization. An 82% yield of propylene oxide carbonate was obtained at 100 °C, 1 h without a co-catalyst. For internal epoxides, the *cis/trans* ratio ranged from 98:2 to 96:4. The catalyst could be precipitated from the reaction mixture with ethyl acetate, filtered and reused six times with no decrease in carbonate yield. Entry 4: the *trans*-isomers gave exclusively the *trans*-cyclic carbonates; the *cis*-isomers gave a 90:10 *cis:trans* mixture of cyclic carbonates. The use of monometallic aluminum catalysts resulted in much lower conversions (27% vs 85–94%), thus supporting a cooperative intervention of both aluminum ions in the reaction mechanism, one activating the epoxide and the other the carbon dioxide. Entry 6: catalyst can be separated by precipitation in Et<sub>2</sub>O and recycled eight times without any significant loss in activity. Cyclohexene oxide did not react. Entry 7: enantiopure (*S*)-styrene oxide gave cyclic carbonate in 93% ee. The reaction was scaled up to 50 mmol with 79% yield after 15 days. Entry 8: enantiopure epoxides gave carbonates with no loss of enantiomeric purity. *Trans*-stilbene oxide reacted in very low yield. Product was distilled from reaction mixture and to the resultant residual catalyst in the reaction flask was added a fresh substrate and CO<sub>2</sub>. This procedure was repeated 10 times without any significant yield decrease. Entry 9: the ee was >99% for R<sup>1</sup>=(*R*)-Me and 90% for R<sup>1</sup>=(*R*)-Ph. Reaction scale-up allowed the isolation of 98% yield of product starting from 42 g of substrate. Bis-epoxides gave bis-carbonates. Entry 11: (*R*)-propylene oxide was converted to carbonate with full retention of the configuration, but (*R*)-styrene oxide underwent partial racemization (42% ee). Entry 12:

internal epoxides retained configuration (>99:1) except *cis*-stilbene oxide which gave *c:t* = 43:57.  Gave also a rearranged product (81% overall yield, 12:88 ratio). Entry 13: mixtures of *cis:trans* cyclic carbonates or of all possible diastereomers were always obtained. Entry 14: cyclic carbonate from limonene oxides was recovered as a *c:t* = 14:86 mixture. Entry 15: epichlorohydrin and cyclohexene oxide gave the worse yields, the former because of

substitution of the chlorine with *N*-methyldiethanolamine, the latter for steric hindrance. The reaction was scaled up to 100 mmol. (*R*)-styrene oxide was converted into cyclic (*R*)-carbonate in 97% yield with 99% ee. Catalyst could be separated by precipitation in Et<sub>2</sub>O and recycled six times with very low decrease in activity. Entry 17: cyclohexene oxide gave only 4% yield. Other cobaloximes were tested with lower results. Entry 18: conversions, not yields (selectivity >99%), *cis*-butene oxide gave cyclic carbonate in *c:t* = 98:2 ratio. Epichlorohydrin and disubstituted substrates required a catalyst loading of 0.4 mol%. Entry 19: it should be noted that the catalyst recovered by CCl<sub>4</sub> precipitation and reused produced only 33–40% yield of styrene carbonate, but increased the yield of hexane oxide from 40% to 98%. Very likely some water molecules remained coordinated to the catalyst. The catalyst then remained active in, at least, five consecutive runs without loss of its activity. Entry 20: conversion not yields, with >99% selectivity. Cyclohexene oxide did not react. Entry 21: cyclohexene oxide gave only 21% yield after 48 h. Catalyst could be recovered by solvent precipitation and reused in ten consecutive runs without loss of its activity. Entry 22: complexes with Fe and Cu or the presence of a second *ortho*-methyl group on the *S*-phenyl group were less efficient. Interestingly when this second ligand was allowed to react with copper, it is reduced from Cu(II) to Cu(I). Entry 23: epichlorohydrin and cyclohexene oxide were also tested but the isolated yields were not determined. Styrene oxide required 48 h. Entry 24: the best reaction conditions depended on the substrate. Cyclohexene epoxide was also tested and required 2.0 MPa at 100 °C. A DFT calculation of the reaction pathway was performed. Entry 25: all reactions occurred with retention of configuration, according to a double-inversion mechanism as depicted in Scheme 2. Even styrene carbonate from (*S*)-styrene oxide was obtained with 99% ee, (45 °C, 1.0 MPa). Entry 26: substituted styrene oxides required 60 °C. The reactions with 1,2-disubstituted epoxides led to very low yields (5–8%), with retention of configuration. Enantiopure epoxides gave cyclic carbonates with no loss of enantiomeric purity. The reaction is scalable up to 50 mmol. The catalyst was reused five times, by distilling off the product, with slow decrease of activity, because of partial sublimation property under the distillation conditions. Entry 27: *c:t* ratios ranged from 65:35 to 50:50. For epoxidized oils: 5 mol% with respect to oxirane number. Entry 29: DBU/NIS adduct was insoluble in ether, thus, after the extraction of products with ether the catalyst could be dried and reused for five times with no significant loss in its catalytic activity. To achieve good yields 2,2-dimethyloxirane and cyclohexene oxide required also Bu<sub>4</sub>Ni (5 mol%). Entry 30: cyclohexene oxide required 12 h and was recovered in 33% yield. A mixture of 15% CO<sub>2</sub> and 85% N<sub>2</sub> afforded styrene carbonate in 91% yield after 14 h. Catalyst could be separated from the aqueous phase after pouring the reaction mixture in water. After drying, it was reused five times without decrease of activity. Entry 31: low-boiling epoxides gave higher yields in a 20 mmol scale with respect to a 5 mmol scale with a refluxing condenser. Higher CO<sub>2</sub> pressure was requested by more sterically hindered epoxides. Stereochemistry was generally maintained but at harsher conditions it decreased, very likely by partial S<sub>N</sub>1 reaction. Cyclooctene oxide gave only a 6% yield. Entry 32: the catalyst could be separated from the reaction mixture by distillation under reduced pressure and reused five times with no significant loss in its catalytic activity. Entry 33: the catalyst could be separated by vacuum distillation, dried and reused five times with no significant loss in its catalytic activity. Entry 34: disubstituted epoxides required 64 h, 100 °C, and 3 mol% of catalyst, to avoid diol formation. Entry 35: the catalyst was precipitated with Et<sub>2</sub>O, dried and reused four times with low decrease in its catalytic activity, due to some leaching of the salt from dextrin cavity. Entry 36: only epoxides with strong electron donating groups reacted and the less the electron donating power were, the harsher the reaction conditions must be. Propene oxide was already unreactive. Entry 37: isobutylene oxide gave only 10% yield. The authors affirmed in the text that also cyclohexene oxide gave low yield but 91% was reported in the table. Most epoxides exhibited good conversion also under atmospheric pressure. Entry 38: higher temperatures were requested by less reactive epoxides such as styrene and cyclohexene oxides. Enantiopure epoxides gave carbonate in 64–92% ee and enantiomeric excess decreased with increasing temperature. Entry 39: higher yields with R<sup>2</sup>=Bu were due to the high solubility of the catalyst. Entry 40: other imidazolium salt gave lower yields. Products could be purified by distillation and the residue containing the catalyst could be used five times with no significant loss in its catalytic activity. Entry 41: the actual catalytic species



was Entry 42: disubstituted epoxides needed a higher catalyst loading (3%) and longer reaction times (48 h). Entry 43: the catalyst could be precipitated and used five times with no significant loss in its catalytic activity. Entry 44: styrene, isobutene oxides, and carbazolylglycididol

required 40 °C. (*R*)-styrene oxide gave (*R*)-carbonate with 70% ee. Entry 45: fatty acid oxides required 0.5 MPa of CO<sub>2</sub> and 100 °C and trans-carbonate was the most abundant isomer (83:17 to 99:1). Entry 46: when [HDBU]I was insoluble in epoxide dimethylformamide was added as the solvent. Cyclohexene oxide reacted at 140 °C, with 3 MPa after 48 h in dimethylformamide leading to the product in 76% yield. The catalyst could be recovered after reduced pressure distillation of the products and reused five times with no significant loss in its catalytic activity. DFT calculations were performed to elucidate the mechanism. Entry 47: HUST-1-Co was a crosslinked cobalt porphyrin obtained from 5,10,15,20-tetraphenylporphyrin, dichloromethane as cross-linker and cobalt acetate. The catalyst could be used fifteen times with no significant loss in its catalytic activity. Entry 48: Al-HPC was a crosslinked aluminum porphyrin obtained from 5,10,15,20-tetraphenylporphyrin, formaldehyde dimethyl acetal as cross-linker and diethylaluminum chloride. The catalyst could be used ten times with no significant loss in its catalytic activity. Cyclohexene oxide gave only 28% yield. Entry 49: Al-iPOP was a crosslinked aluminum porphyrin obtained from 5,10,15,20-tetra(4-bromophenyl)porphyrin linked by Yamamoto–Ullmann coupling, coupled with a polymeric ionic liquid from (4-bromophenyl)- (1) and (4-bromobenzyl)imidazolium bromide (2). Al-iPOP-2 was generally more efficient. The catalysts could be used six times with no significant loss in its catalytic activity. Entry 50: the catalyst was obtained from 2-methylimidazole (8 mmol), Co(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O (0.5 mmol) and Zn(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O (0.5 mmol) in methanol. The catalyst could be used four times with no significant loss in its catalytic activity. Entry 51: the catalyst was obtained from 2-methylimidazole, Co(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O and Zn(NO<sub>3</sub>)<sub>2</sub>·4H<sub>2</sub>O in water. The amount refers to a 9 mmol scale. It was also recycled four times with little deactivation (from >99 to 91% yield). The substrates were lacking in the paper table and were furnished by private communication. Entry 52: porous charged covalent triazine framework obtained at 500 °C by reaction of 1,1'-bis(4-cyanophenyl)-[4,4'-bipyridine]-1,1'-dium dichloride in melted anhydrous ZnCl<sub>2</sub>. It was also recycled 4 times without deactivation. Entry 53: a mesoporous poly(ionic liquid) obtained from 1-glycidyl-3-vinylimidazolium bromide and divinylbenzene then opened in hot water. The catalyst was recycled 10 times without deactivation. Entry 54: larger R<sup>1</sup> groups gave very poor yields (6–10%). The catalyst was recycled 5 times without deactivation. Entry 55: samples of Cu-DABCO, Ni-DABCO, and Co-DABCO were also tested but with worse results. The catalyst was used three times without deactivation. Entry 57: ZnO@NPC-Ox was obtained from ZIF-8 (see entries 42–43) which was pyrolyzed and subsequently oxidized by NaOCl to produce ZnO nanoparticles (NPs) encapsulated into N-doped porous carbon. The catalyst was used 10 times without deactivation. Entry 58: the catalyst was used five times without deactivation, but it required a tedious purification procedure from the reaction mixture. Other epoxides gave very low conversions (24–26%), albeit with high selectivity (99%) and styrene oxide did not react. Entry 59: UDIL= urea-derivative-based ionic liquids, I=iodide, 60%U is the relative mass of urea added to UDIL. Cyclohexene oxide gave 45% yield, at 130 °C, after 9 h and with 3.0 MPa of CO<sub>2</sub>. The catalyst was used five times without deactivation and was tolerant of the presence of water. Entry 60: The catalyst was used for 12 runs. Quantitative yield was recovered in the first four runs, then a slight decrease (up to 85% in the twelfth run) together a loss of catalyst (0.69 mol% recovered from the last run) was observed. Entry 61 other imidazolium-based ionic polymers gave worse results. The catalyst was used 10 times without deactivation. Entry 62: the catalyst was used four times without deactivation. Entry 63: the catalyst was used 10 times without deactivation and was also used in a continuous flow reactor for 120 h without deactivation. Entry 64: CBAP-1(EDA) was prepared from 1,3,5-triphenylbenzene and terephthaloyl chloride. The obtained polymer was then reductively aminated with ethylene diamine. Cyclohexene oxide gave only 14% yield after 6 h at 140 °C. The catalyst was used 5 times without deactivation. Entry 65: CBAP-1(EDA-Zn) was obtained by treatment of CBAP-1(EDA) with Zn(OAc)<sub>2</sub>. Cyclohexene oxide gave 45% yield after 48 h at 60 °C. The catalyst was used five times without deactivation. Entry 66: most of reactions were carried out with the same catalyst sample recycled from the previous reaction. Its efficiency was verified repeating the reaction with butane oxide every seven times and comparing the yield. Catalyst efficiency significantly decreased after 14 runs. Oleate oxide gave 43:57 *c:t* mixture. Entry 67: COF-JLU7 was prepared from 2,4,6-tris(4-aminophenoxy)triazine and 2,5-dihydroxy-1,4-benzenedicarboxaldehyde under acidic catalysis. The reaction with (*R*)- or (*S*)-styrene oxide exhibited excellent enantioselectivity (97% and 93% ee, respectively). The catalyst was used five times without deactivation. Entry 68: cyclohexene oxide gave only 30% of conversion. The catalyst could be reused but poisoning of the active sites was observed. Thus, to maintain high conversion and selectivity, the catalyst had to be regenerated, by treatment with diluted NaOH. Entry 69: cyclohexene oxide gave only 9% yield. The yield of carbonate decreased


sharply from 97% to 64% in the third cycle and then slowly (until 59% after other two runs). After the five cycles, less than the original catalyst amount was recovered, thus authors explained the lesser efficiency with this loss of catalyst amount. [Entry 70](#): the catalyst was used 10 times without deactivation. [Entry 71](#): Zn-C<sub>3</sub>N<sub>4</sub>(25) was prepared by thermal polymerization of Zn(OAc)<sub>2</sub>·2H<sub>2</sub>O/dicyandiamine in mass ratio of 25%. The catalyst was used five times without deactivation. [Entry 72](#): cyclohexene oxide gave only 6% yield together many byproducts. The catalyst was used five times without deactivation. [Entry 73](#): the addition of *N*-(3-aminopropyl)-imidazole increased the fiber weight of 20%. The fiber was intertwined on the stirring paddle of the reactor and could be used 21 times in gram-scale reactions. [Entry 74](#): obtained by refluxing 5,10,15,20-tetrakis(4-pyridyl)porphyrin zinc(II), mesoporous silica SBA-15, and 3-(trimethoxysilyl)propyl bromide in dimethylformamide. Other solvents gave a catalyst with worse catalytic activity. The catalyst was used five times without deactivation. [Entry 75](#): obtained by the reaction of 5,10,15,20-tetrakis(4-pyridyl)porphyrin zinc(II), di(1*H*-imidazol-1-yl)methane, and 1,4-bis(bromomethyl)benzene in the presence of carbon nanotubes (CNTs). The catalyst was used seven times without deactivation. [Entry 76](#): the catalyst was used five times without deactivation. 4-nitrostyrene oxide and (but-3-en-1-yl)oxirane did not react or decomposed. [Entry 77](#): regular hexahedral (RH) ZIF-8 and rhombic dodecahedral (RD) ZIF-8 homometallic nanoparticles (see entries 50-51) underwent cation exchange with Au<sup>3+</sup> ions from NaAuCl<sub>4</sub>. The catalyst was used 6 times without deactivation. [Entry 78](#): there is a discrepancy between text (chloro- and bromo-methyloxirane reported) and table (bromo- and chloro-oxirane reported). We think that the text is right. Catalyst was prepared from nickel wires and 4,6-bis(triazol-1-yl)isophthalic acid in acidic solution. [Entry 79](#): the catalyst was obtained from reaction of carbon nanotubes (CNT) with *N*-vinyl-*N'*-allylimidazole silver complex (NHC-Ag). Cyclohexene and isobutene oxides gave poor yields (10 and 11% respectively). The reusability was tested with propargyl alcohols (see footnote Table 2, entry 5). [Entry 80](#): obtained by reaction between triethoxy-3-(2-imidazol-1-yl)propylsilane and octakis(3-bromopropyl)-octasilsesquioxane then grafted onto SiO<sub>2</sub> and finally by reaction with 1-methylimidazole. The catalyst was used five times without significant deactivation.

In this mechanism, the rate determining step is the epoxide ring opening, and the nucleophilicity of the generated alkoxide species becomes significant for the attack at CO<sub>2</sub>. This pathway involves two consecutive S<sub>N</sub>2 steps, thus the configuration is generally maintained with enantiopure epoxides. However, styrene oxide, as well as other epoxides giving stable carbocations, often gave partial racemization, which is accounted for a partial S<sub>N</sub>1 mechanism. It should be noted that the scorpionate catalyst (entry 43) was rationally designed by authors on the basis of the mechanism and combining experimental and computational efforts [84].

A comparative study on the activity of a series of fifteen aluminum-based complexes has been recently reported, providing a useful comparison of activity metrics and explaining what are the most important features of the catalyst for cyclic organic carbonate formation [118].

Another challenging task was the research of organocatalysts as efficient as metal catalysts. In these cases, the intervention of hydrogen bond donors in the catalyst is often necessary to activate the ring opening step (see entries 24, 25, 27, 29, 34, 35, 37, 38, 39, 40, 44, 45, 46, 62, 66, 69, 70). Very recently, the activity of a diverse selection of hydrogen bond donors has been correlated to their pK<sub>a</sub> [119]. It was found that hydroxyl protons with Brønsted acidity in the range 9 < pK<sub>a</sub> < 11 gave the best catalytic performance, therefore phenol and ascorbic acid derivatives are ideal for cycloaddition reaction of epoxides and CO<sub>2</sub>. Density functional theory (DFT) calculations supported this hypothesis, low energy barriers have been calculated for the reaction catalyzed by phenols and the occurrence of aggregation between molecules of ascorbic acid further lowers the energy barriers increasing the catalytic activity. The halide source can be external (e.g., an ammonium halide salt) or internal (e.g., the counterion of the metal ion or of salt moieties in the catalyst). When no halide source is present as co-catalyst or catalyst counterion, a labile nucleophilic moiety has to be anyway present, for instance a tertiary amine (entries 17, 22, 26, 29, 35, 41, 44, 45, 56, 68, 69, 70, 75) or the carboxylate ions in histidine (entry 28) or aspartate (entry 33) catalysts.

A quite different mechanism was proposed by Liu, Wei, Dai and co-workers, whose paper is mentioned in entries 44–45 [85]. On the basis of their results, of some experiments with modified catalysts and DFT calculations they proposed the mechanism depicted in Scheme 3.




**Scheme 3.** Mechanism proposed for cooperative multifunctional organocatalysts.

Conditions set up by Werner and co-workers (entry 11) [55] allowed the selective reaction of the monosubstituted oxirane in the presence of a di-substituted one. In fact, 3-(oxiran-2-yl)-7-oxabicyclo[4.1.0]heptane gave only the monocarbonate at the external epoxide moiety in 92% yield at 23 °C, and the bis-carbonate in 67% yield at 45 °C.

Moreover, the re-use of the catalyst is very important. Thus some simple catalysts were recovered by distillation or precipitation from the reaction mixture (entries 3, 6, 8, 15, 19, 26, 29, 32, 33, 35, 46), and many examples of heterogeneous catalysts have been reported (Table 1, third part). It should be particularly outlined the apparatus setup by Shi, Hu and co-workers (entry 73), who interweaved their catalyst in the stirring paddle, thus providing a very simple method to recover the catalyst [111]. In addition, the interweaved paddle could be reused more than twenty times, just increasing reaction times after the 20th time, in a gram scale reaction.

The coupling of internal epoxides and CO<sub>2</sub> is generally more difficult for the steric hindrance of the two substituents. Thus reaction described in entries 10, 12, 13, 14, 25, 27, 31, 45 are particularly worthy of note for their good results.

In this regard, it should be mentioned the work of Kleij and co-workers, who were able to obtain trisubstituted cyclic organic carbonates with well-defined stereochemical configurations, starting from cyclic (5–8 membered) hydroxy epoxides and CO<sub>2</sub> (Scheme 4) [120]. *Syn/cis* carbonates were always obtained only from addition of 25 mol% of NBu<sub>4</sub>Br. The *anti/cis* bicyclic carbonates instead required individual reaction conditions (different aluminum catalyst, halide source, solvent) and only five-, six-, and seven-membered cyclic epoxides reacted. Larger seven- and eight-membered cyclic epoxides could be converted to *anti/trans* carbonate in the presence of aluminum catalyst and dimethylaminopyridine (DMAP).




Scheme 4. Preparation of highly substituted organic carbonates.

These different behaviors were attributed to different mechanisms of formation: *syn/cis* from carbonate polymer depolymerization, *anti/cis* from OH-assisted depolymerization (larger ring did not allow a conformation in which OH is near enough) and *anti/trans* from substrate-assisted activation of  $\text{CO}_2$ . Then some different hydroxycyclohexene oxides were successfully transformed into *anti/cis* bicyclic carbonates. The reaction suffers from steric hindrance (*ortho*-substituted phenyl groups needed longer reaction times as 66 h, and afforded only 41% yield). Some products were then submitted for further modifications.

The style used by authors to present the results reported in Table 1 was quite different. Some were more oriented to the synthesis, other to the catalyst performance. Therefore it is not simple to compare results and efficiency at a glance. For example, turn over number (TON), a key to evaluating a catalyst, if a green, sustainable atom-economy is pursued, is often not reported. So we suggest that readers point their attention to yields, catalyst amount and  $\text{CO}_2$  pressure to evaluate the best catalyst among the eighty entries of Table 1. This difficulty in the choice of the best catalyst was also shown by D'Elia in his fine review, which showed also the harsh increase of documents in the last few years [45].

Kim and co-workers set up a divergent synthesis to cyclic carbonate and oxazolidinones in the coupling reactions between  $\text{CO}_2$  and epoxy alcohols or amines, respectively (Scheme 5) [121]. It is also worth noting that the starting materials are enantiopure and the reaction is completely enantiospecific. The divergent pathway is induced by the cocatalyst: tetrabutylammonium iodide triggers a mechanism superimposable with Scheme 2, while dimethylaminopyridine acts as a base on the amino group, which in turn attacks carbon dioxide. On the basis of this mechanism, it is clear that *N*-aryl groups highly influence the reactivity (electron-donating groups worked better than electron-withdrawing groups). Finally, the inversion of the stereochemistry, when oxazolidinones are prepared, allowed the synthesis of both enantiomers of 3-(arylamino)-propane-1,2-diols from a single enantiopure epoxyamine.



**Scheme 5.** Stereodivergent synthesis starting from epoxyamines and CO<sub>2</sub>.

Reactions reported in Table 1 are general, although more or less efficient. There are also some papers that reported only one example of conversion of epoxides into carbonates. Two cases reported in Table 1 are the work of Verpoort on Zn-Co/ZIF (entry 50) [88], because it was then extended to other substrates (entry 51) [89]; and the work of Kleij and co-workers on bifunctional resorcinarenes [74], because they reported the only example for their catalyst on polymeric support (entry 60). Other catalysts have been proposed for CO<sub>2</sub> addition to epoxides in which only one substrate was studied. For instance, Rownaghi and co-workers were able to permanently immobilize bromide on microstructured zirconium doped polyamide-imide hollow fibers crosslinked with 3-aminopropyltrimethoxysilane, followed by alkylation with 1,2-dibromoethane. This fiber acts as bifunctional catalysts for the cycloaddition of CO<sub>2</sub> to styrene oxide (100% conversion of oxide and 98% selectivity of carbonate). This catalyst did not significantly leach ZrO<sub>2</sub>, amine, or bromine under the employed reaction conditions (120 °C, 8 h, 2.0 MPa) [122]. This work should be further developed to better compare these results with the catalytic and efficient fibers described in entry 73 [109].

The nitrogen doped ordered mesoporous polymers prepared by the Huang and Dai group showed excellent catalytic activity (conversion >95%) in the reaction of CO<sub>2</sub> with propylene oxide (100 °C, 1.5 h, 1.2 MPa), when charged with zinc or cobalt ions [123].

The porous cationic polymers (5% wt.), obtained from condensation of tris(1,10-phenanthroline-5,6-dione)Ru(II) dichloride with *ortho*-aromatic amines in AlCl<sub>3</sub> at 400 °C, allowed the conversion of propylene oxide to carbonate (almost quantitative conversion of oxide and exclusive selectivity toward carbonate, at 100 °C, 24 h, 0.1 MPa) [124].


Dufaud and co-workers presented a study on the binding strength of cavitand receptors of ammonium salts. Thus, tetraphosphonate showed a higher binding strength toward the ammonium cation. However, the presence of acidic phenol groups, which cooperatively activated the hexene oxide in a triphosphonate cavitand host, gave quantitative yields of cyclic carbonate at atmospheric pressure of CO<sub>2</sub> and 100 °C. Moreover, Bu<sub>4</sub>NI turned out to be the most efficient halide donor [125].

Propylene carbonate was also obtained from propylene oxide at atmospheric CO<sub>2</sub> pressure and at ambient temperature, by using various metal carbamates, such as Ti(O<sub>2</sub>CNEt<sub>2</sub>)<sub>4</sub>, Al(O<sub>2</sub>CNR<sub>2</sub>)<sub>3</sub> (R=Et, *i*-Pr), Cu(O<sub>2</sub>CNEt<sub>2</sub>)<sub>2</sub> and Sn(O<sub>2</sub>CNEt<sub>2</sub>)<sub>4</sub>, in combination with NBu<sub>4</sub>X (X=Br or Cl) as a cocatalyst [126]. With different catalyst and cocatalyst amounts, conversion up to 71% and selectivity up to >99% were

reached. Solid catalysts were prepared from titanium and zirconium carbamates with amorphous silica, but these heterogeneous catalysts were less efficient than the homogeneous ones.

Graphitic carbon nitride (prepared with 1:1 urea–thiourea mixture) treated with aqueous  $H_2SO_4$  produced a catalyst bifunctional acidic ( $-SO_3H$ ) and basic ( $-NH_2$ ) sites. The concentration of  $H_2SO_4$  influenced the number of sites and the highest catalytic activity was reached with 60 wt. % acid. At 100 °C and 1.0 MPa  $CO_2$  pressure for 1 h, the synthesis of cyclic carbonates from epichlorohydrin (92.8% conversion, 99.2% selectivity) and propylene oxide (61.5% conversion, 99.3% selectivity), was obtained [127]. The catalyst was easily recovered and recycled six times with negligible loss of activity.

Seven alkanol amine catalysts derived from reactants containing two or three epoxy moieties and secondary amines were synthesized by Chung and co-workers [128]. Among them, bis(methylpiperazinyl) triol (1.4 mol%, at 1 MPa, 120 °C, 3 h) revealed the best catalyst in the formation of propylene carbonate (98% yield). Under milder conditions (5.6 mol% catalyst 0.5 MPa, 100 °C, 8 h) yield decreased to 90%. Comparing all the set of catalysts, the catalyst performance was affected by the number of hydroxyl and amine groups, and by the synergistic effects of these groups as demonstrated by DFT calculations of the stable conformational state. The proposed reaction mechanism is depicted in Scheme 6.




**Scheme 6.** Plausible mechanism of the cycloaddition of  $CO_2$  to propylene oxide catalyzed by bis(methylpiperazinyl)triol.

The synthesis of 1,2-butylene carbonate from 1,2-butylene oxide was obtained in a selectivity of 76% and 64% yield at 135 °C and 7.5 MPa pressure of  $CO_2$  in 20 h in the absence of organic solvent with a ceria-lanthana-zirconia/graphene nanocomposite catalyst [129].

The merit of the synthesis of carbonates from internal epoxides has already been outlined above. Recently, polyethylene glycol (PEG-400) and KI (4 mol%) were found able to catalyze the reaction of methyl soyate epoxide to the corresponding carbonated fatty acid methyl esters in 99% yield at 120 °C with 3.0 MPa pressure of  $CO_2$  in 20 h. Longer reaction times (30 h) are requested to obtain the methyl

linoleate tricarbonatate [130]. However, for these catalysts further studies would be required to better understand their importance in this reaction.


The kinetic resolution of epoxides was employed to prepare enantioenriched epoxides [131]. Recently, Ema and co-workers reported the kinetic resolution of epoxides with CO<sub>2</sub> by a chiral macrocyclic organocatalyst (catalyst-1, Scheme 7) [132]. The *s* factor  $\{(\ln[1-c(1+ee)])/(\ln[1-c(1-ee)])\}$ , where *c* is the conversion, is satisfactory for disubstituted epoxides (9–13), but it is low for terminal epoxides (2.5–4.1). These values make this reaction very interesting for internal epoxides, taking also into account the trouble in performing the synthesis of 4,5-disubstituted 1,3-dioxolan-2-ones. The X-ray analysis showed multiple hydrogen-bonding sites for the enantioselective activation of epoxides into the chiral cavity of the catalyst. DFT calculations confirmed a classical mechanism as depicted in Scheme 2.



Scheme 7. Kinetic resolution of epoxides with organocatalyst.

Later they reported another macrocyclic organocatalyst (catalyst-2, Scheme 7) but its efficiency was comparable for terminal epoxides (*s* = 1.7–5.0). However, with respect to catalyst-1, catalyst-2 showed some advantages, that are the low catalyst loading (only 0.2–0.5 mol%) and the opposite resolution of the epoxide and some drawbacks, that are higher CO<sub>2</sub> pressure and longer reaction times [133].

On the other hand, Meggers and co-workers set up reaction conditions for the kinetic resolution of chiral terminal epoxides with *s* factors up to 16.6 [134]. They employed an iridium(III) complex (1 mol%) of iridium catalyst and 1.5 mol% of NEt<sub>4</sub>Br as the co-catalyst (Scheme 8). It should be noted that no polymerization side reaction is observed. Authors provided a mechanism in agreement with Scheme 2. While the decrease of enantiomeric excess with the progress in conversion was expected, the decrease of the *s* factor with increasing conversion was surprising, but authors did not give a satisfactory explanation for this phenomenon. Moreover, they did not also explain the different enantiomers obtained with simple alkyl epoxides.




**Scheme 8.** Kinetic resolution of epoxides with iridium catalyst.

The synthesis of six membered cyclic carbonates from the addition of CO<sub>2</sub> to oxetanes is more difficult, because the four-membered ring is less reactive than epoxide and for the higher thermodynamic stability of the polymers with respect to the cyclic carbonates. Therefore, examples of this reaction are less common in the literature. In the last two years isolate examples are reported:

- under the conditions described in Table 1 entry 4, 1,3-dioxan-2-one was recovered in 67% yield at 80 °C, and 1.0 MPa CO<sub>2</sub> pressure [47];
- with the POSS-Imi catalyst (0.067 mol%, Table 1, entry 70) but the conversion is very low (16%) albeit selectivity was almost quantitative [116].

Tassaing, Jérôme et al. studied the influence of the main reaction parameters (organocatalytic mixture, pressure, and temperature) on the yield and the selectivity in the product distribution of the reaction of CO<sub>2</sub> with oxetanes with an organocatalytic combination of ammonium salts and aromatic alcohols [135]. Cyclic carbonate was favored at lower temperatures, pressures, and conversions, as well as in the absence of alcohol. It should also be noted that substituted oxetanes were unreactive. They also investigated the mechanism of the synthesis of the six membered cyclic carbonate using DFT calculations performed at the M06-2X/6-311G(d,p) level. The mechanism is not far from that accepted for epoxides (Scheme 9).





**Scheme 9.** Mechanism of the reaction of oxetane with CO<sub>2</sub>.


Very recently, Dove and Coulembier found trimethylene carbonate as byproduct in the co-polymerization of oxetane and CO<sub>2</sub> catalyzed by 1,5,7-triazabicyclo[4.4.0]dec-5-ene and I<sub>2</sub> [136]. However, trimethylene carbonate was the sole product, albeit in low yields (30% after 5 days) when


1-tert-butyl-4,4,4-tris(dimethylamino)-2,2-bis-[tris(dimethylamino)-phosphoranylideneamino]-2 $\lambda^5,4\lambda^5$ -catenadi(phosphazene) was used as a cocatalyst together with iodine. Authors affirmed that the strong complexation between the two catalysts was very likely the reason of this selectivity.


The cycloaddition of propargylic alcohols with CO<sub>2</sub> is another manner to prepare five-membered cyclic carbonates. As for the addition to epoxides, this is also a 100% atom economical reaction. Moreover, the resulting  $\alpha$ -alkylidene cyclic carbonates are important compounds with many applications in organic synthesis. The reaction has been widely studied in the past years and reviewed in the literatures cited in the introduction, thus here we restrict the description to the last two years again (Table 2). The reaction is generally catalyzed by a base which deprotonates the alcohol to start the catalytic cycle, as described in Scheme 10. Sometimes the alcoholate can open the cyclic carbonate leading to an oxoalkyl acyclic carbonate.

**Table 2.** Catalysts for the reaction of propargyl alcohols and carbon dioxide (R, R<sup>1</sup> and R<sup>2</sup> see Scheme 10).

Entry	R	R <sup>1</sup> , R <sup>2</sup>	P <sub>CO<sub>2</sub></sub> (MPa)	Conditions	Yield (%)	Ref
1	Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-MeOC <sub>6</sub> H <sub>4</sub> , 4-ClC <sub>6</sub> H <sub>4</sub> , 4-MeCOC <sub>6</sub> H <sub>4</sub> , 4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub> , 3-piridyl	R <sup>1</sup> =R <sup>2</sup> =Me, (CH <sub>2</sub> ) <sub>4</sub> R <sup>1</sup> =Me, R <sup>2</sup> =Et, <i>i</i> -Pr	2.0	 (5 mol%), 60 °C, 3 h	55–96	[137]
2	Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-FC <sub>6</sub> H <sub>4</sub> , 4-C <sub>6</sub> H <sub>13</sub> C <sub>6</sub> H <sub>4</sub> , 4- <i>t</i> -BuC <sub>6</sub> H <sub>4</sub>	R <sup>1</sup> =R <sup>2</sup> =Me, Et <i>i</i> -Pr	2.5	 (200 mol%) 60 °C, 24 h	68–99	[138]
3	H, Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-ClC <sub>6</sub> H <sub>4</sub> , 4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub> , 4-MeOC <sub>6</sub> H <sub>4</sub> , 4-pyridyl	R <sup>1</sup> =R <sup>2</sup> =Me, (CH <sub>2</sub> ) <sub>3</sub> , (CH <sub>2</sub> ) <sub>4</sub> R <sup>1</sup> =Me, R <sup>2</sup> =Ph	2.0	CDC-CO <sub>2</sub> (5 mol%), (entry 39, Table 1), 80 °C, 12 h	51–94	[80]
4	H	R <sup>1</sup> =R <sup>2</sup> =Me, Et, (CH <sub>2</sub> ) <sub>4</sub> R <sup>1</sup> =Me, R <sup>2</sup> =Et	1.0	Zn/Fe <sub>3</sub> O <sub>4</sub> /ECS (133 mg/mmol), NEt <sub>3</sub> (Equation (1)) 30 °C, 12–20 h	90–93	[139]
5	H	R <sup>1</sup> =R <sup>2</sup> =(CH <sub>2</sub> ) <sub>4</sub> R <sup>1</sup> =Me, R <sup>2</sup> =Et, <i>i</i> -Bu	3.0	CNT-NHC-Ag (151 mg/mmol) (entry 71, Table 1), 80 °C, 24 h	97–99	[115]
6	H	R <sup>1</sup> =R <sup>2</sup> =Me, Ph R <sup>1</sup> =Me, R <sup>2</sup> =Et, Ph, <i>i</i> -Bu, Allyl	5.0	( <i>n</i> -Bu <sub>4</sub> N) <sub>2</sub> (CO <sub>2</sub> ) <sub>2</sub> , (2.5–5 mol%) 80 °C, 6 h	60–99	[140]

Entry 1: the active catalyst was . Entry 2: 1-(2-phenylethynyl)cyclohexanol was unreactive; 2-methylbut-3-yn-2-ol gave quantitatively 1,1-dimethyl-2-oxopropyl-1',1'-dimethyl-2'-propynyl carbonate. The catalyst could be separated with water, then after drying reused four times without significant deactivation.

Entry 3: the active catalyst  acted as a base on the propargyl alcohol. Entry 4: Zn/Fe<sub>3</sub>O<sub>4</sub>/ECS was a magnetically separable catalyst prepared from corn starch, magnetite and ZnI<sub>2</sub>. 4-Phenyl-2-methylbut-3-yn-2-ol was unreactive. The catalyst was recovered with an external magnet, and used 4 times without significant deactivation. Entry 5: the NHC-Ag complex was also supported on graphene (GN) and the reaction gave similar results. CNT-NHC-Cu and GN-NHC-Cu complexes gave lesser yields. However, CNT-NHC-Cu was used in the recycle tests and reused eight times without significant deactivation. Entry 6: Propargyl alcohol and but-3-yn-2-ol were unreactive, owing to the well-known Thorpe-Ingold effect [141], 1-(2-Phenylethynyl)cyclohexanol and 3,6-dimethylocta-1,7-diyne-3,6-diol gave low conversion (12% and 35%, respectively), but selectivity near 100%. A higher catalyst loading was necessary for sterically hindered alcohols.




**Scheme 10.** Mechanism of the cycloaddition of propargylic alcohols with CO<sub>2</sub>.


A detailed mechanistic investigation was recently performed by Tassaing's research group for the conversion of 2-methyl-3-butyn-2-ol [142]. Authors found an influence of the temperature, pressure, solvent, and catalyst nature and loading on the rates and yields of the reaction. Moreover, in contrast to our previous results [143], they did not detect other secondary products. Finally they confirmed the mechanism depicted in Scheme 10 by DFT calculation at the M06-2X/6-311G(d, p) level. Then, the same research group tested a series of *n*-tetrabutylammonium organic salts as catalysts for the same reaction [144]. In particular, tetrabutylammonium acetate and azide allowed yields up to 98% in less than 10 h at 80 °C and 3.0 MPa under solvent-free conditions. Once more, they supported their data by kinetic studies and mechanism by DFT calculations.

The mechanism described in Scheme 10 involved, after cyclization a vinyl anion intermediate, which sequentially underwent protonation. However, this anion could be trapped by every electrophile. In fact, under palladium catalysis it was trapped by aryl halides [145]. The reaction suffered from steric factors. In fact, 2-chloro-1-iodobenzene and 1,2,4-trichloro-5-iodobenzene gave 45% and 6% yields, respectively. Iodomethane gave the corresponding product, albeit in only 31% yield. Among the propargyl alcohol tested, only 2-methyl-4-(pyridin-2-yl)but-3-yn-2-ol gave no reaction. The reaction exclusively gave the stereoisomer in which the Ar group arising from the aryl halide is located trans to the oxygen attached to the double bond. Authors after some control experiments proposed the mechanism depicted in Scheme 11.

The reaction of diols and CO<sub>2</sub> under basic conditions is another method to obtain cyclic carbonates of different ring size. Ongoing their study on this reaction [146], Buchard and co-workers reported the synthesis of 5-, 6-, 7- and 8-membered cyclic carbonates using 2,2,6,6-tetramethylpiperidine as the base catalyst (Scheme 12, Equation (1)) [147]. As expected, larger cycles were recovered with increasing amounts of polymers.



**Scheme 11.** Arylcarboxylation of propargylic alcohols with  $\text{CO}_2$  and aryl halides.



**Scheme 12.** Synthesis of 5-, 6-, 7- and 8-membered cyclic carbonates.

From DFT calculations, authors suggested a mechanism in which  $\text{CO}_2$  is attacked by the alcoholate, then tosylation of the carbonate occurred and, finally, the second alcoholate moiety closed the cycle by addition/elimination process. It should be noted that sugar diols also reacted, which gave poor yields with the classical reaction with phosgene.

Another reaction of sugar diols and CO<sub>2</sub> was introduced by Feng, Gnanou and co-workers. They already found that unprotected  $\alpha$ -methyl D-glucopyranoside reacted with CO<sub>2</sub>, leading to water soluble oligoglycocarbonates [148]. Further studies from the same group showed that the reaction of CO<sub>2</sub> with *cis* vicinal hydroxy groups of sugars afforded instead 5-membered bicyclic glycocarbonates, in particular in the *cis*-2,3 or *cis*-3,4 positions of methyl  $\alpha$ -D-mannopyranoside and methyl  $\alpha$ -D-galactopyranoside, respectively (Scheme 12, Equations (2)–(4)) [149]. Also, a six-membered bicyclic glycocarbonate was prepared from methyl 6-bromo-6-deoxy-2,3-di-O-methyl  $\alpha$ -D-galactopyranoside. The cyclic glycocarbonates were further allowed to react in good yield with amines or polymerized.

These two interesting procedures summarize two environmental benign features: the use of CO<sub>2</sub> as starting material and the avoidance of phosgene or its derivatives


### 3. Cyclic Carbamates

Carbamates are widely employed in agricultural chemistry, medicinal chemistry, and polyurethane preparation. In particular, cyclic carbamates (2-oxazolidinones, 2-oxazinanones) are the core structures of many valuable drugs. For a long time phosgene was the classical C1 synthon for their synthesis; but its high toxicity has led researchers to explore greener C1 sources. Thus the use of CO<sub>2</sub> has been gaining increasing importance.

Many strategies for CO<sub>2</sub>-based preparation of cyclic carbamates parallel those already seen above: for instance the cycloaddition to aziridines, oxetanes, or amino epoxides or the addition to alkenes, alkynes, and propargylic amines or alcohols. Most of these syntheses are already reviewed in the literature cited in the introduction, but in the last two years other papers appeared in the literature and they are collected in this section.


The synthesis of substituted 2-oxazolidinones by coupling of CO<sub>2</sub> with aziridines is another reaction which allows the chemical fixation of CO<sub>2</sub> with 100% atom efficiency. A first example has been reported in the previous section (Scheme 5) [106]. Other syntheses of substituted 2-oxazolidinones by coupling of CO<sub>2</sub> with aziridines published in the last two years are collected in Table 3.

While regioselectivity is not a problem with epoxides, because the oxygen atoms are indistinguishable, the synthesis of oxazolidinones often gives mixture of regioisomers (Scheme 13). The most striking difference with the mechanism of epoxides is the higher nucleophilicity of the nitrogen, which give raise a zwitterionic intermediate.



**Scheme 13.** A simplified mechanism of the addition of CO<sub>2</sub> to aziridines.

**Table 3.** Catalysts for the reaction of aziridines and carbon dioxide (R, R<sup>1</sup>, and R<sup>2</sup> see Scheme 13).

Entry	R	R <sup>1</sup> , R <sup>2</sup>	P <sub>CO<sub>2</sub></sub> (MPa)	Conditions	Yield (%)	Regio. (R <sup>2</sup> 5:R <sup>2</sup> 4)	Ref
1	R=Pr, Bu, <i>n</i> -C <sub>5</sub> H <sub>11</sub> , (CH <sub>2</sub> ) <sub>2</sub> - <i>i</i> -Pr	R <sup>1</sup> =H, R <sup>2</sup> =Ph, 4-MeC <sub>6</sub> H <sub>4</sub>	1.0	ISA 6 (1 mol%), (entry 6, Table 1), 50 °C, 2–8 h	80–99	95.5 to 98:2	[49]
2	R=H, Me, Et, Bn	R <sup>1</sup> =H, R <sup>2</sup> =Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-ClC <sub>6</sub> H <sub>4</sub> , 4-(MeOCH <sub>2</sub> )C <sub>6</sub> H <sub>4</sub> , 2-naphthyl	0.1	SalenCoI (2.5 mol%), Ph <sub>3</sub> P=CHCOPh (2.5 mol%), (entry 7, Table 1), 25 °C, 48 h IL-Zn-TPP (0.1 mol%), (entry 21, Table 1) 90 °C, 2–10 h	38–90	100:0	[50]
3	R=Pr, Bu, <i>i</i> -Bu, <i>n</i> -C <sub>5</sub> H <sub>11</sub>	R <sup>1</sup> =H, R <sup>2</sup> =Ph, 4-MeC <sub>6</sub> H <sub>4</sub>	2.0	CDC-CO <sub>2</sub> (5 mol%), (entry 39, Table 1), 80 °C, 12 h	82–96	97.3 to 98:2	[61]
4	Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-ClC <sub>6</sub> H <sub>4</sub> , 4-BrC <sub>6</sub> H <sub>4</sub> , CH <sub>2</sub> OPh, 2-thienyl	R <sup>1</sup> =H, R <sup>2</sup> =Et, Pr, Bu, <i>i</i> -Pr, <i>c</i> -C <sub>6</sub> H <sub>11</sub> , Bn	2.0	(1 mol%), DBU (2 mol%), 50 °C, 48 h	52–99	90:10 to 97:3	[115]
5	Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-ClC <sub>6</sub> H <sub>4</sub> , 4-BrC <sub>6</sub> H <sub>4</sub> , 4-MeOC <sub>6</sub> H <sub>4</sub> , 4- <i>t</i> -BuC <sub>6</sub> H <sub>4</sub>	R <sup>1</sup> =H, R <sup>2</sup> =Et, Pr, Bu, <i>i</i> -Bu	0.1		43–92	93:7 to 99:1	[150]






Entry 2: the reaction with (*R*)-*N*-methyl-2-phenylaziridine completely retained the ee. Entry 4: the four-substituted oxazolidinone (86:14) was preferentially produced when R = CH<sub>2</sub>OPh very likely for its electron-donating nature. Halo-substituted phenyl rings needed high catalyst loading (10 mol%).

The positive charge on the nitrogen atom generally favors the attack of the halide source on the carbon which better carries the positive charge, leading to the five-substituted product from *N,C*-disubstituted aziridines. Recently, Pinhas and co-workers reported DFT calculations at the B3LYP/6-31+G(d,p) level of theory, kinetic studies and experiments in order to elucidate the mechanism [151]. Their calculations demonstrated that the reaction could not proceed at room temperature without a catalyst. Moreover, their data were unable to distinguish between two possible pathways involving either initial capture of CO<sub>2</sub> or ring opening by X<sup>-</sup> from the catalyst. In fact, both mechanisms are influenced by polar solvents. The reaction is zero order with respect to aziridine, thus isolate aziridine is not involved in the rate determining step, which was the addition of CO<sub>2</sub>.


The synthesis of oxazolidin-2-ones from unsaturated amines by using CO<sub>2</sub> has attracted much attention in the last years [152]. A study of the reaction mechanism with the ωB97XD functional theory was conducted by Yuan and co-workers under Ag(I) catalysis [153]. Authors discovered that the substrate is incorporated into the AgNO<sub>3</sub> salt, followed by isomerization (Scheme 14). Then this intermediate attacks CO<sub>2</sub> leading to a carbamate intermediate. The five-membered versus the six-membered ring closure has transition states the energy of which is influenced by the diazabicycloundecene (DBU) amount (an increasing of DBU amount increases the energy gap between the two transition states, favoring the oxazolidinone). Moreover, solvents with larger proton affinities also favor the formation of oxazolidinones.

The papers regarding propargylamines and published in the last two years are collected in Table 4.

**Table 4.** Catalysts for the reaction of propargylamines and carbon dioxide (R, R<sup>1</sup>, R<sup>2</sup> see Scheme 14, NR<sup>3</sup> indicates the nitrogen substituent).

Entry	R	R <sup>1</sup> ,R <sup>2</sup>	NR <sup>3</sup>	P <sub>CO2</sub> (MPa)	Conditions	Yield (%)	Ref
1	H, Ph, Me	R <sup>1</sup> =R <sup>2</sup> =H, Me, (CH <sub>2</sub> ) <sub>4</sub>	H, Me, <i>i</i> -Pr, <i>t</i> -Bu, Bn, 4-MeOC <sub>6</sub> H <sub>4</sub>	0.1	 (1 mol%), 40–80 °C, 0.16–90 h	74–98	[154]
2	H, Me, Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-MeOC <sub>6</sub> H <sub>4</sub> , 4-CNC <sub>6</sub> H <sub>4</sub>	R <sup>1</sup> =R <sup>2</sup> =H, Me, Et, (CH <sub>2</sub> ) <sub>4</sub>	H, Me	0.5	MOF (0.4 mol%), 60 °C, 24 h	31–99	[155]
3	Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-BrC <sub>6</sub> H <sub>4</sub>	R <sup>1</sup> =R <sup>2</sup> =Me, (CH <sub>2</sub> ) <sub>4</sub> R <sup>1</sup> =H, R <sup>2</sup> = <i>c</i> -C <sub>6</sub> H <sub>11</sub> R <sup>1</sup> =Me, R <sup>2</sup> =Et	Bu, <i>c</i> -C <sub>5</sub> H <sub>9</sub> , Bn	0.1	 (11–22 mol%), 100 °C, 24 h	89–93	[156]
4	H, Ph	R <sup>1</sup> =R <sup>2</sup> =H R <sup>1</sup> =H, R <sup>2</sup> =Et	Bn, Bu, Ph, <i>c</i> -C <sub>6</sub> H <sub>11</sub> , PhCH(Me), Ph(CH <sub>2</sub> ) <sub>2</sub> , 4-ClC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub> , 4-MeOC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>	0.1	 (5 mol%), 60 °C, 12 h	8–96	[157]
5	H, Ph, 4-MeC <sub>6</sub> H <sub>4</sub>	R <sup>1</sup> =R <sup>2</sup> =H, Me R <sup>1</sup> =H, R <sup>2</sup> =Me, Et, Pr, <i>i</i> -Pr	Bu, Bn	0.1	 (10 mol%), 90 °C, 10 h	71–99	[158]
6	H	R <sup>1</sup> =R <sup>2</sup> =H, Me, (CH <sub>2</sub> ) <sub>4</sub> R <sup>1</sup> =H, R <sup>2</sup> =Me	Me, Bn	0.5	Bu <sub>4</sub> NF (1 mol%), 110 °C, 12 h	94–99	[159]
7	Ph, 4-MeC <sub>6</sub> H <sub>4</sub> , 4-MeOC <sub>6</sub> H <sub>4</sub> , 4-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub> , 4-CNC <sub>6</sub> H <sub>4</sub>	R <sup>1</sup> =R <sup>2</sup> =H	Me, Bn	0.5	Bu <sub>4</sub> NF (1–2 mol%), 110 °C, 24–48 h	77–91	[160]
8	H, Me, Et, Ph	R <sup>1</sup> =R <sup>2</sup> =H, Me	Bn, 4-MeOC <sub>6</sub> H <sub>4</sub> CH <sub>2</sub>	Air	 (0.5 mol%) DBU (10 mol%) r. t., 48 h	35–90	[161]
9	H, Me, Ph, 4-MeC <sub>6</sub> H <sub>4</sub>	R <sup>1</sup> =R <sup>2</sup> =H, R <sup>1</sup> =H, R <sup>2</sup> =Me, Ph, 4-MeC <sub>6</sub> H <sub>4</sub>	Me, <i>i</i> -Pr, Bu, Bn,	1.0	KCC-1/IL/Ni@Pd NPs (0.1 mg/mmol), 15 W compact fluorescent lamp r. t., 3 h	83–96	[162]


Entry 1: Longer reaction times were requested by more sterically hindered nitrogen atoms. Internal alkynes reacted slower than terminal alkynes. When R=Me an 8.25:1, 5-*exo*:6-*endo* ratio was observed. Entry 2: MOF



was a supramolecular structure obtained from


, 1,4-benzenedicarboxylic acid and

$\text{Cd}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ . Increasing the substrate size the yields decreased and with  $\text{R} = 1\text{-naphthyl}$  no reaction occurred. Diprop-2-ynylamine gave 5-methylene-3-(prop-2-ynyl)oxazolidin-2-one in 62% isolated yield. The catalyst could be reused up to four times without deactivation. **Entry 5:** both amino-activated and  $\text{CO}_2$ -activated mechanisms were investigated by DFT calculations and  $\text{CO}_2$ -activated resulted in a low energy barrier in the first step. **Entry 6:** primary amines were unreactive. **Entry 7:** electron withdrawing groups in  $\text{R}$  favored tautomerization to 2-oxazolones. (oxazolidinone/oxazolone ratio 12:57 and 2:64 for  $\text{R} = 4\text{-CF}_3\text{C}_6\text{H}_4$ ,  $4\text{-CNC}_6\text{H}_4$ , respectively. In these cases oxazolidinones were obtained at  $70^\circ\text{C}$  in 24 and 3 h for  $\text{R} = 4\text{-CF}_3\text{C}_6\text{H}_4$ ,  $4\text{-CNC}_6\text{H}_4$ , respectively. **Entry 8:** internal bulky alkynes gave the lowest yields. *N*-Phenyl amines were unreactive and primary amines gave only 4% yield. **Entry 9:** KCC-1/IL/Ni@Pd NPs was prepared in three steps: KCC-1 NPs from tetraethyl orthosilicate, cetylpyridinium bromide and urea; KCC-1/IL NPs from KCC-1 NPs, bis(trimethoxysilylpropyl)imidazolium iodide and NaH by sonication; finally KCC-1/IL/Ni@Pd NPs from KCC-1/IL NPs,  $\text{Ni}(\text{OAc})_2 \cdot 4\text{H}_2\text{O}$  and  $\text{PdBr}_2$ . The catalyst could be reused up to ten times without deactivation.



**Scheme 14.** Mechanism of diazabicycloundecene (DBU)- $\text{AgNO}_3$ -catalyzed  $\text{CO}_2$  incorporation into propargylic amine.

With the catalyst reported in entry 1, some homopropargylic amines were tested (Scheme 15) [134]. However, at  $40^\circ\text{C}$  and with 5 mol% of catalyst loading only 36–37% yields were recovered, albeit the starting amine was completely consumed. Authors also observed that, in the absence of  $\text{CO}_2$ , internal alkynes were unreactive, while terminal alkynes gave some side reactions. Thus, in order to improve yield, they tested a homopropargylamine with an internal (phenyl-substituted) alkyne moiety. However,  $\text{CO}_2$  was not incorporated and dehydropyrrole was favored over the carboxylative cyclization. The same reaction was observed with 2-alkynylaniline, which led to indole.




**Scheme 15.** Other reactions carried out with the catalyst described in Table 4 entry 1.

The reactions described in entries 3 [156] and 8 [161] are very interesting, because they were carried out under air. All reactions described in entry 8 were carried out under these conditions, while only three different oxazolidinones were prepared in 66–95% yields with cyanuric acid. Actually, the 2-oxazolidinone syntheses in low-concentration CO<sub>2</sub> in air are rare (See also: [163,164]).

The observation by Fujita and co-workers [140] that some amounts of tautomer 2-oxazolinone was found under their reaction conditions, prompted them to set up the best conditions for this tautomerization (5 mol% of Bu<sub>4</sub>NOH, at 80 °C for 6 h).


As shown in Table 4, *N*-alkyl propargylamines are generally employed in these reactions, because *N*-aryl derivatives undergo alkyne hydroarylation more easily than CO<sub>2</sub> incorporation owing to the lower nucleophilicity of the nitrogen atom. Only in Table 4, entry 1, *N*-*p*-methoxyphenylpropargylamine was reported to react in 74% yield, but simple *N*-phenylpropargylamine did not give any oxazolidinone [134]. In the recent literature, however, an interesting work was performed by Zhou and his research group, who were able to carry out a four component one-pot sequence reaction to give enantioenriched oxazolidinones (Scheme 16) [165]. Authors separately tested the preparation of the enantioenriched propargyl amines and their cyclization, and, when the best conditions were found, they successfully performed the one-pot reaction. The absolute configuration of the chiral 2-oxazolidinone was determined to be *R* by X-ray analysis. The reaction was scaled up to 4.0 mmol scale with only 5 mol % of copper catalyst, and product was achieved in 84% yield and 94% ee.

Finally, the CO<sub>2</sub>-fixation to propargyl amines-pyridine co-polymers promoted by a two-fold amount of DBU should be mentioned [166,167]. The polyoxazolidinone was obtained in 100% conversion (86% yields) in 48 h, even under air. In the same paper also three examples of propargyl amines are reported and it should be noted that *N*-benzyl-2-methyl-4-(pyridin-2-yl)but-3-yn-2-amine afforded a *E/Z* mixture of oxazolidinone that is the only example in the two years covered by this review.



**Scheme 16.** Tandem asymmetric aldehyde-alkyne-amine coupling-carboxylative cyclization one-pot sequence.


Another three-component reaction involving propargylic amines, CO<sub>2</sub> and *N*-bromosuccinimide (NBS) allowed the stereoselective synthesis of (*E*)-bromovinylloxazolidinones [168]. The reaction was catalyzed by silver acetate in the presence of a base. Among the bases, a guanidine derivative was found the most efficient (Scheme 17). The reaction mechanism is similar to that reported in Scheme 14, but the vinylsilver intermediate is trapped by bromine atom and not by hydrogen from the amine. This is another example of vinyl anion trapping by an electrophile different from hydrogen as already reported in Section 2 (Scheme 11) [145]. It should also be noted that in this reaction only the (*E*)-vinylsilver isomer is formed.



**Scheme 17.** Synthesis of (*E*)-bromovinylloxazolidinones.

Also allylamines have been employed for carbon dioxide fixation into cyclic carbamates.


The reactions had three components and also involved alkyl halides and the presence of light, because generally the alkyl halide is homolytically broken by an irradiated metal catalyst. For instance tris(bipyridine)ruthenium(II) chloride allowed the synthesis of many oxazolidinones (Scheme 18, Equation (1)), as well as 3-benzyl-6-phenyl-6-(2,2-difluoro-3-ethoxy-3-oxopropyl)-1,3-oxazinan-2-one and 3-benzyl-6-phenyl-5-(2,2-difluoro-3-ethoxy-3-oxopropyl)-1,3-oxazinan-2-one (54 and 38% yields, respectively) [169]. The reaction of primary allylamines, allylamines with electron withdrawing nitrogen protecting groups or aliphatic R<sup>2</sup> substituents failed as well as the synthesis of cyclic carbonates from allyl alcohols and of oxazinone from 2-(prop-1-en-2-yl)aniline. The reaction was scaled up to 1.31 g of recovered product. Some manipulations of the products were also successfully carried out. Then the same research group carried out a similar reaction under palladium catalysis and found the same features, except for primary allylamines, which smoothly reacted. (Scheme 18, Equation (2)) [170]. Scheme 18 also reported the mechanism surmised by authors for the reaction. Both reactions were scaled up to gram-scale and oxazolidinone was recovered in 84% yield under ruthenium catalysis and 77% yield under palladium catalysis. Products of both reactions were also manipulated to give open-chain compounds.



**Scheme 18.** Visible-light photoredox induced reaction of allylamines with CO<sub>2</sub>.


He and co-workers demonstrated that, in the presence of 1,5,7-triazabicyclo[4.4.0]dec-5-ene TBD as the base, the metal catalyst is unnecessary with perfluoroalkyl halides [171]. They carried out some

experiments to elucidate the mechanism and from the evidence they proposed the mechanism depicted in Scheme 19.



**Scheme 19.** Visible-light photoredox induced reaction of allylamines with CO<sub>2</sub> and perfluoro compounds.

The *N*-heterocyclic carbene copper and silver complexes supported on graphene or nanotubes, introduced by Chen and co-workers (see also Table 1, entry 72 and Table 2, entry 5), allowed the synthesis of oxazolidinones in a three-component reaction (Scheme 20) [115]. Copper complexes gave generally higher yields.



**Scheme 20.** Three-component coupling of CO<sub>2</sub>, primary amine, and various propargyl alcohols.

An alternative and attractive strategy to produce oxazolidinones from less-toxic and easily available starting materials is a three-component reaction of CO<sub>2</sub>, epoxides, and amines. However, this reaction suffers from large amounts of epoxides because at least a molecule is transformed into a diol by-product (Scheme 21). Two alternative pathways have been suggested:

- the first one involved the formation of cyclic carbonate by the classical mechanism (see Scheme 2), then amine opened the carbonate by releasing of diol and formation of a isocyanate, which in turn added another molecule of epoxide (path a);
- the second one involved the interaction between the cyclic carbonate and the amino-alcohol from nucleophilic opening of epoxide by amine (path b).




( $R = C_6H_{13}, (CH_2)_6CH=CH_2$ ) afforded a single regioisomer albeit in 45% and 38% yields, respectively. Also *p*-nitroaniline, benzyl- and octyl- amines gave a single regioisomer. In all cases, the five-substituted oxazolidinone is recovered. Both isocyanates and aminoalcohols gave the products when submitted to the reaction conditions.

Diazabicycloundecene, instead, allowed the reaction of aliphatic epoxides. However, increasing the bulkiness of the epoxides yields lowered and cyclohexene oxide led to unidentified mixture of products (Scheme 21, Equation (2)) [173]. The analysis of the reaction course by GC-MS did not show the presence of isocyanate intermediates. Also, a Fe(II) *N*-heterocyclic carbene (NHC) complex was found able to perform this reaction (Scheme 21, Equation (3)) [174]. Authors did not attempt reaction with internal epoxides, and also found that styrene oxides gave lower yields than aliphatic epoxides. Some experiments, carried out to elucidate the mechanism, suggested path b for the reaction.

KCC-1 nanoparticles have been already encountered in this review in Table 4, entry 9. On this fibrous material was supported Cu (II)- $\beta$ -cyclodextrin and the obtained catalyst afforded oxazolidinones from reaction of anilines, CO<sub>2</sub> and ethylene oxide (Scheme 21, Equation (4)) [175]. Albeit restricted only to ethylene oxide and *p*-substituted anilines, this reaction is interesting because it did not use epoxide excess. In fact, in the proposed mechanism, authors affirmed that the diol by-product could be captured by the catalyst. Then, a molecule of aniline could nucleophilically remove it to give the ethanolamine intermediate for a further catalytic cycle.

Conversely from CO, which can favorably add the ethanolamine [176], the direct reaction between amino alcohols and CO<sub>2</sub> is thermodynamically disfavored. To shift the equilibrium to 2-oxazolidinones dehydrating agents or auxiliaries are often needed, with generation of waste. Among dehydrating agents in the last two years, He and co-workers proposed propargyl alcohols and published some papers with different catalysts to carry out this reaction [177–179].

The catalyst or the catalyst mixture should activate the OH groups both of propargyl alcohol and of amino alcohol by hydrogen bonds and the CO<sub>2</sub>, as well as coordinate the triple bond and act as a Lewis acid on the carbonyl group. In all the papers, yields of oxazolidinones and ketones by-product are comparable. Authors performed control experiments, DFT calculation, kinetic and NMR studies and from these data suggested the reaction pathway depicted in Scheme 22 [177], which was confirmed in the other papers. The first proposed catalyst was the couple Ag<sub>2</sub>O and tetramethylguanidine (Scheme 22) [177]. The reaction was also applied to diols; 1-phenylethan-1,2-diol and 3-phenoxypropan-1,2-diol gave the corresponding cyclic carbonates in 95% and 58%, respectively.




**Scheme 22.** Cascade reaction of propargyl alcohols, carbon dioxide, and 2-aminoethanols.

Then, the same research group proposed another catalytic mixture to perform the same reaction, that are: CuI (5 mol%) 1,10-phenantroline (5 mol%) *t*-BuOK (10 mol%), CO<sub>2</sub> (0.5 MPa), 80 °C, 12 h [178].

Yields ranged 18–95% for hydroxyketones and 24–97% for the same oxazolidinones reported in Scheme 22, as well as the diols. Only 1,2-propanediol was added extra. The third catalyst was 1,5,7-triazabicyclo[4.4.0]dec-5-ene trifluoroethanol (15 mol%), CO<sub>2</sub> (0.1 MPa), 80 °C, 12 h [179]. For the same set of products, yields ranged 23–99% for hydroxyketones and 24–99% for the oxazolidinones and cyclic carbonates. The easy recovering of this last catalyst allowed its recycle and the target product was still obtained in 75% yield, after five reuses.


Repo and co-workers used tosyl chloride as the auxiliary [180]. The most important feature of this reaction is the formation and the stabilization of the carbamate species before the competing *N*-tosylation reaction, which is then irreversible (Scheme 23). Lower reaction temperatures and higher CO<sub>2</sub> pressure favored the oxazolidinone formation. The presence of a good leaving group such as in 3-chloropropan-2-ol-1-amine favors the 6-membered ring notwithstanding the higher stability of the 5-membered ring. In fact, propan-1,2-diol-1-amine gave mainly the five-membered ring. Starting from chiral compounds the S<sub>N</sub>2 nature of the reaction allowed the preparation of enantioenriched compounds in good purity. The mechanism was then studied by DFT calculation confirming author's hypothesis.



**Scheme 23.** Synthesis of cyclic carbamates from amino alcohols.


Finally, oxazolidinones could be obtained in high yields from a three-component reaction among CO<sub>2</sub>, 1,2-dichloroethane and aromatic amines catalyzed by *N*-heterocyclic carbene obtained in situ from an ionic liquid (Scheme 24) [181]. However, the reaction with 2,3-dichlorobutane afforded the corresponding 4,5-dimethyl-3-aryloxazolidin-2-ones in very low yields (28–40%). On the other hand, 1,2-dichloropropane gave good yields (about 70%) at higher temperatures, but without regioselectivity and a mixture of 5-methyl-3-aryloxazolidin-2-one and 4-methyl-3-aryloxazolidin-2-one was always recovered. The reaction occurred also with CO<sub>2</sub> diluted in water, air and nitrogen. The catalyst was used five times without significant deactivation. Interestingly the reaction was extended to other dichloroalkanes. Many six-membered 3-aryl-1,3-oxazinan-2-ones (19 examples) were obtained in 70–93% yields at 90 °C for 6 h. The seven-membered 3-aryl-1,3-oxazepan-2-one was also formed but in lower yield (22–35%, 3 examples). Larger cycles such as those from 1,5-dichloropentane, 1,6-dichlorohexane and 1,10-dichlorodecane were not obtained, but ω-chloroalkylcarbamates were recovered in 63, 55, 37%, respectively. Some experiments were carried out to elucidate the mechanism

depicted in Scheme 24. Authors also demonstrated by means of control experiments that carbonate ions were not the C1 source of the reaction.




**Scheme 24.** Synthesis of oxazolidinones from 1,2-dichloroethanes.

These last two reactions have been employed also for larger cyclic carbamates than oxazolidinones. Very recently, an enantioselective cyclization was developed to prepare six-membered cyclic carbamates from homoallylic amines (Scheme 25) [182]. The catalyst was studied in order to have Brønsted basicity enough to avoid amine to form an achiral carbamate salt intermediate. Some amounts of water favored the catalyst-carbamic acid complex, but an excess allowed the formation of an unreactive crystalline ligand. The optimum amount was 90 ppm. Allylic amines gave low enantiomeric excess under these conditions. Unsubstituted homoallylamines ( $\text{R}^1 = \text{H}$ ) gave low yield (43%) and ee (13%). The enantiomeric catalyst gave the enantiomeric product in comparable yield and selectivity. The absolute configuration was determined by X-ray analysis.



**Scheme 25.** Synthesis of enantioenriched 1,3-oxazinan-2-ones.

Other six-membered rings, recently obtained by CO<sub>2</sub> fixation, are some azo-linked 4*H*-benzo[*d*][1,3]oxazine-2,4-diones from azo-linked aminobenzoic acids [183]. The catalyst was called by authors CuO@RHA/MCM-41 nanocomposite and was prepared from CuO nanoparticles and a MCM-41 matrix obtained from rice husk ash (RHA). The best yields were obtained with electron-withdrawing substituted substrates. This heterogeneous catalyst was recycled up to six times without significant loss of activity. In the proposed mechanism (Scheme 26), Cu ions act as a Lewis acid to activate CO<sub>2</sub>. Then a carbamate ion is formed, which in turn undergoes intramolecular reaction and dehydration to give the product.




**Scheme 26.** Synthesis of 4*H*-benzo[*d*][1,3]oxazine-2,4-diones.

Another cyclic carbamate, which shows biological and pharmaceutical activities, is 1,3,4-oxadiazole-2(3*H*)-one. Its synthesis could be performed by 1,3-dipolar cycloaddition of nitrile imine with CO<sub>2</sub>. However, the low reactivity of carbon dioxide toward 1,3-dipoles and the fast dimerization of nitrile imines make this reaction rare. In the last two years three papers appeared in the literature anyway. In 2017, Zhang found that CsF/18-crown-6 are able to enhance both the reactivity of CO<sub>2</sub> as a 1,3-dipolarophile and the in situ formation of nitrile imines from hydrazone chloride (Scheme 27, Equation (1)) [184]. Among the prepared products, a potential drug for Parkinson's disease therapy and the commercial herbicide oxadiazon were obtained in 89% and 88% isolated yield, respectively. The same procedure can be successfully applied to the synthesis of 3,4-thiadiazol-2(3*H*)-one from COS.

The reaction mechanism was studied by Fernández-Herrera, Merino and co-workers at the SMD/M06-2X/def2-TZVP level. They found that the reaction proceeds by a three-step mechanism and not by 1,3-dipolar cycloaddition (Scheme 27, Equation (2)), thus explaining how the low reactivity of carbon dioxide toward 1,3-dipoles is overcome. The rate-determining step is the final five-membered ring closure. However, authors affirmed that increasing the concentration and the temperature the concerted pathway becomes more likely [185].

Very recently, a new synthesis of substituted 1,3,4-oxadiazol-2(3*H*)-ones from aryl hydrazines and  $\alpha$ -oxocarboxylic acids, with KI as the catalyst, has been developed (Scheme 27, Equation (3)) [186]. Authors made some control experiments and they found that the extra pressurized CO<sub>2</sub> is necessary for obtaining good yield (with the CO<sub>2</sub> only released by the  $\alpha$ -oxocarboxylic acid, the product was obtained in 38% yield); the hypoiodite ions is a key active species to give the hydrazone iodide; and that methyl acrylate instead of CO<sub>2</sub> led to a cycloadduct in 74% yield, thus suggesting a cycloaddition reaction also with CO<sub>2</sub>. However, it is surprising that after this evidence, authors did not start from more available aldehydes instead of  $\alpha$ -oxocarboxylic acids.



Scheme 27. Synthesis of 1,3,4-oxadiazole-2(3H)-one.

Finally, the synthesis of 2-benzoxazolone (60% yield) from 2-aminophenol in the presence of tributylamine should be mentioned [187]. Reaction details will be discussed in the next section (Scheme 31).


#### 4. Cyclic Ureas

Ureas framework is present in natural products, agricultural pesticides, herbicides, and pharmaceuticals. The synthesis of urea itself was the founder of modern organic chemistry. However, the selective and efficient synthesis of substituted ureas and in particular of cyclic ureas from CO<sub>2</sub> and amines is still challenging [188,189].

Quinazoline-2,4(1H,3H)-diones, important intermediates in the synthesis of pharmaceuticals, can be obtained from reaction of 2-aminobenzonitriles and CO<sub>2</sub>. This synthesis is particularly important because of the direct utilization of carbon dioxide as well as the high atom economy. In the last two years many different methods have been reported in the literature. For instance, the bi-functional graphitic carbon nitride prepared by Samanta and Srivastava [127] was efficient only in dimethylformamide (90.2% yield) or dimethylsulfoxide (93.2% yield), with 50 mg/mmol of catalyst, at 130 °C and 2.5 MPa CO<sub>2</sub> pressure, after 12 h in the reaction of 2-aminobenzonitrile. The aprotic dipolar solvent should enhance the activation of the CO<sub>2</sub> molecules as well as favor the abstraction of an acidic proton from the NH<sub>2</sub> group of 2-aminobenzonitrile. Examples of reaction with larger applicability are collected in Table 5.

The accepted mechanism for this reaction is depicted in Scheme 28. Many reactions showed a relationship between the pK<sub>a</sub> of the basic catalyst and the reaction rate. For example, in ionic

liquid the  $pK_a = 14.7$  (the acidity of the quinazoline-2,4-dione product) is the borderline between the base-catalyzed reaction and the quinazoline-2,4-dione ion catalysis [190]. A further computational study was conducted at M06-2X level and 6-31G (d) basis set utilizing organic bases as the catalyst, which confirmed the mechanism [191]. Other studies on the mechanism were carried out with the M06 functional owing to its recognized ability in the description of organometallic chemistry with noncovalent interactions [192].







**Scheme 28.** Mechanism of the reaction of aminobenzonitriles and  $\text{CO}_2$  catalyzed by bases.

Cesium carbonate was chosen as the catalyst and the LanL2DZ basis set was used for Cs atoms and the 6-311 + G(d,p) basis set was used to describe other atoms. The energy involved in different reaction pathways was calculated and the most favorable pathway is very similar to the depicted in Scheme 28. A “naked nitrogen ion” is necessary for the attack to  $\text{CO}_2$  and the rearrangement 6-imino-1,3-oxazine-2-one to pyrimidine-2,4(1*H*,3*H*)-dione occurs intramolecularly. In this study, the isocyanate intermediate is not found along the reaction pathway, but the complex A was invoked as the key intermediate for the rearrangement.

Fujita reported just an example of pyrido[2,3-*d*]pyrimidine-2,4(1*H*,3*H*)-dione (footnote to entry 1, Table 5) [160], but almost at the same time Zhang and co-workers published a detailed study on the synthesis of many dihydropyrimidine-2,4(1*H*,3*H*)-diones fused with heterocycles (Scheme 29) [193].

An ionic liquid was used both as the catalyst and the solvent (6 mmol/mmol substrate). The reaction temperatures (60–90 °C) and times (3–96 h) clearly depended from the substrate. Among the various amino-carbonitrile heterocycles, only 2-amino-1*H*-indole-3-carbonitrile and 3-amino-1*H*-indole-2-carbonitrile did not react. However, *N*-acetyl-3-amino-1*H*-indole-2-carbonitrile gave the expected product, suggesting that an N-H group on the heterocycle prevented the reaction. In fact, no other free N-H heterocycle was tested.

**Table 5.** Catalysts for the reaction of 2-aminobenzonitriles and CO<sub>2</sub>.


Entry	X	P <sub>CO<sub>2</sub></sub> (MPa)	Conditions	Yield (%)	Ref
1	H, 4-Cl, 4,5-(MeO) <sub>2</sub>	0.1	 (22 mol%), 100 °C, 24 h	78–86	[156]
2	H, 4-Me, 4-NO <sub>2</sub> , 5-Cl	2.0	 (0.5 g/mmol), 100 °C, 20 h	56–83	[68]
3	H, 5-Me, 5-F, 5-Cl, 5-Br, 4,5-(MeO) <sub>2</sub>	2.0	Bu <sub>4</sub> NF (1 mol%), 110 °C, 24 h	96->99	[160]
4	H	1.0	Bu <sub>3</sub> N (20 mol%), 100 °C, 36 h	75	[187]
4	H, 4-Cl, 4-Me, 5-F, 5-Me, 4,5-(MeO) <sub>2</sub>	0.1	 (100% mol%), 80 °C, 24 h	87–98	[194]
5	H, 4-Cl, 5-Cl, 5-Br, 4,5-(MeO) <sub>2</sub>	1.0	Zhabuye basic salt-lake brine (1.5 mL/mmol), 140 °C, 8–30 h	96–98	[195]

Entry 3: pyrido[2,3-*d*]pyrimidine-2,4-(1*H*,3*H*)-dione was also obtained in 59% yield. Entry 4: product was obtained after bubbling CO<sub>2</sub> at 15 KPa and 50 °C for 36 h in 93% yield, even at a gram scale and in 87% under simulated flue gas system containing 15% of CO<sub>2</sub> and 3% of H<sub>2</sub>O. Entry 5: other salt lake waters were tested with worse results. In fact, the large amounts of basic B<sub>4</sub>O<sub>7</sub><sup>2-</sup> and CO<sub>3</sub><sup>2-</sup> ions in Zhabuye salt-lake brine are responsible of the best efficiency. Electron-withdrawing groups required longer reaction times. A carbonate solution was unable to give the reaction, thus CO<sub>3</sub><sup>2-</sup> was not the C1 source. The Zhabuye basic salt-lake brine can be separated by centrifugation from the reaction mixture and reused directly for five times without significant loss of activity.


Both CO<sub>2</sub> and RNC are classical C1 sources and their use in a multicomponent reaction is an attractive way to prepare heterocycles introducing carbonyl and imine moiety at the same time. However, the difference in kinetic and thermodynamic stability of both C1-reactants makes their contemporary use rare, because are scarce the catalytic system which sufficiently activates CO<sub>2</sub> and tunes the RNC insertion. In particular, the reaction of 2-haloanilines, RNC, and CO<sub>2</sub> could give rise to two reaction products: 2-amino-4*H*-benzo[*d*][1,3]oxazin-4-one and 4-imino-1,4-dihydro-2*H*-benzo[*d*][1,3]oxazin-2-one. Tuning both regio- and chemoselectivity to give the second product is particularly interesting, because it spontaneously rearranges to quinazoline-2,4-(1*H*,3*H*)-diones (see Scheme 28). Palladium catalysis was able to perform this cascade reaction affording *N*3-substituted quinazoline-2,4-(1*H*,3*H*)-diones (Scheme 30). Three papers appeared in the literature starting from *o*-bromo- or *o*-iodoanilines [196–198]. All research groups carried out some control experiments to elucidate the reaction mechanism and all were in agreement to propose the mechanism depicted in Scheme 30. After the best reaction conditions and the scope of the reaction (Scheme 30, Equation (1)), the first paper reported also some post-functionalization, giving rise to *N*1-substituted quinazolinodiones, and 2,4-dichloro-6,7-dimethoxyquinazoline a key intermediate of some drugs. Moreover, by using the cheap <sup>13</sup>CO<sub>2</sub>, labelled quinazolinodiones were also synthesized.

The reaction was scaled up to 5 mmol [196]. Almost simultaneously, another synthesis was developed starting from *o*-iodoanilines (Scheme 30, Equation (2)). It is worth to note that the reaction worked at atmospheric pressure of CO<sub>2</sub>, while the other worked at overpressure. The reaction was also carried out at a half gram scale and afforded product in 77% yield. Under these reaction conditions, *o*-bromo- or *o*-chloro-aniline, strong electron-withdrawing substituted *o*-iodoanilines, 4-nitrophenylisonitrile, and 2,6-dimethylphenylisonitrile did not react [197].


Later, Zhang and coworkers used 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl, (SPhos) as the palladium ligand. Under these conditions, *o*-bromoanilines with electron-donating substituents gave satisfactory yield, but electron-withdrawing substituents afforded low yields (Scheme 30, Equation (3)). It should be noted that electron-deficient *o*-bromo- or *o*-iodoanilines gave good yields with PPh<sub>3</sub> as the ligand, but with CsF as the base, conversely from Equation 2 (Scheme 30, Equation (4)) [198].

The direct carboxylation of diamines with carbon dioxide is another attractive manufacture of cyclic ureas. In the past years some catalysts have been introduced for this reaction and have been discussed in the reviews cited at the top of this section [188,189]. However, in the time range covered by us other interesting papers appeared in the literature. For instance, Lee, Kim and co-workers found that, in the presence of carbonate or bicarbonate as the bases, the synthesis of cyclic ureas is greatly enhanced by the presence of some amounts of imidazolidin-2-one. In particular, the reaction of ethylenediamine showed a classical autocatalytic rate, while 1,2- and 1,3-propanediamine yields increased from about 50% to 75%, if 10 mol% of imidazolidin-2-one was added at 200 °C, after 2 h and 5.0 MPa of CO<sub>2</sub>. The reactions went to completion in 4 h [199].

Since the carboxylation of amines by CO<sub>2</sub> is known to proceed with almost no activation barrier, the role of imidazolidin-2-one cannot favor an increase of the nucleophilicity of the diamine, but rather to assist the catalyst in the protonation and deprotonation steps of the catalytic cycle, very likely via its enolic form.



Scheme 29. Synthesis of heterocycle-fused pyrimidine-2,4-(1H,3H)-diones.




**Scheme 30.** Three-component synthesis of quinazoline-2,4-(1H,3H)-diones.

Also the solvent play an important role, because it should be polar enough to stabilize anionic species, but it must not interact strongly with the base catalyst via hydrogen bonding. Authors

performed a theoretical calculation to support these findings. On these bases, authors introduced 2-pyrrolidone as the best solvent for this reaction [200]. In fact, it can give rise to a keto-enol tautomerism and it is polar enough to stabilize the anionic species. The cyclic ureas from ethylenediamine, 1,2- and 1,3-propanediamine were recovered in 83–95% yields at 200 °C, after 2 h and 5.0 MPa of CO<sub>2</sub> even in the absence of a base.

Benzimidazolones have been obtained from the reaction of *o*-phenylenediamines with CO<sub>2</sub> in the presence of tributylamine as the base catalyst (Scheme 31) [187]. The catalyst can be recovered from the reaction mixture at 210–214 °C for 1 h and then reused without loss of activity, but slight reduction in yield was observed owing to the incomplete recovery. Electron-withdrawing substituents decrease nucleophilicity of the *o*-phenylenediamine, thus leading to less efficient reactions. Reaction with *N*-phenyl and *N*-methylphenylenediamine gave the expected product in 80 and 84% yields, respectively.




**Scheme 31.** Synthesis of 2-benzimidazolones.


We already reported that [<sup>13</sup>C]-labeled quinazolinones have been synthesized with the method described in Scheme 30, Equation (1) [196], but in the time range covered by this review, *o*-azidoamines and radiolabeled [<sup>11</sup>C and <sup>14</sup>C] as well as labelled [<sup>13</sup>C] CO<sub>2</sub> were used to prepare pharmaceutically important radiolabeled cyclic ureas [201]. The reaction proceeded in the presence of a phosphine, which reacted with the azide to give an iminophosphorane, which in turn underwent an aza-Wittig reaction to give an isocyanate. Finally the isocyanate is intramolecularly attacked by the amino group to cyclize (Scheme 32). During the experiments for the synthesis of *N*-alkylated heterocycles in a continuous flow reactor on a bed of  $\gamma$ -Al<sub>2</sub>O<sub>3</sub> in supercritical CO<sub>2</sub>, authors found that under particular reaction conditions diethanolamine (1 M, at 250 °C, 15.0 MPa, and 0.2 mL/min) afforded 3-(2-hydroxyethyl)oxazolidin-2-one with a 56% of conversion and 73% selectivity [202]. At lower pressure and flow rate the oxazolidinone is converted in the expected 2,2'-(piperazine-1,4-diyl)diethanol. Owing to the higher nucleophilicity of amino groups the reaction was then carried out with *N*-(2-aminoethyl)ethanolamine. Actually at 250 °C, 1-(2-hydroxyethyl)imidazolidin-2-one was recovered with 85% selectivity and 70% yield. In a saturated CO<sub>2</sub> atmosphere, but without supercritical CO<sub>2</sub> as the solvent, the imidazolidinone was formed in 62% selectivity, 15% yield from 24% conversion. Conversely from oxazolidinone, imidazolidinone was stable in all the tested reaction conditions. Further studies are necessary to better understand this reaction.

Finally, our research group was able to setup an easy access to imidazolidin-2-ones from the three-component reaction of propargylamines, primary amine and CO<sub>2</sub> with 1,5,7-triazabicyclo[4.4.0]dec-5-ene (TBD) as the catalyst under solvent-free conditions (Scheme 33) [203]. The most interesting features of this reaction were: (i) secondary alkyl amines as well as allylamine gave worse yields even increasing temperature to 120 °C. (ii) The higher stability of carbamate arising from benzylamine led to very low yields (<5%). However, changing the base to 7-methyl-1,5,7-triazabicyclo[4.4.0]dec-5-ene and increasing temperature to 120 °C yield increased to 80%. (iii) Anilines reacted despite their low nucleophilicity, but yields greatly depended from electronic properties of the substituents, ranging from

81% for *p*-methoxyaniline to 22% for *p*-bromoaniline. (iv) Propargylamine led to 1*H*-imidazol-2(3*H*)-one, that is the most stable endocyclic double bond. (v) Also internal triple bond gave the product but in a 1:1 mixture of *E/Z* diastereoisomers. (vi) Only *N*-methylprop-2-yn-1-amine afforded 57% yield of cyclic urea, while other secondary propargylamines afforded only oxazolidinone. The reaction could be carried out in a gram scale (87% yield).




**Scheme 32.** Synthesis of labelled pharmaceutically relevant cyclic ureas.



Scheme 33. Urea derivatives from carbon dioxide and amines.


## 5. Other Heterocycles

In this last section, we report the synthesis of other heterocycles, which can be obtained by capture of carbon dioxide. For instance, a polythiazolium-based polymer was able to catalyze the cyclization of 2-aminobenzenethiols to benzothiazole in the presence of DBU and  $\text{CO}_2$  and phenylsilane as the reductant [204]. Authors found that primary amides used as the solvent reacted with 2-aminobenzenethiol in the presence of silane, so they used a cyclic amide, *N*-methyl-2-pyrrolidone (NMP). Moreover, temperatures  $>70^\circ\text{C}$  also reduced the yield. Four-substituted benzothiazoles were recovered in low yields, very likely for steric effects, but authors claimed that this was the first example of the use of three-substituted-2-aminobenzenethiols in the synthesis of four-substituted benzothiazoles. The polymer precatalyst was recovered adding excess HI and precipitation from methanol and reused for 7 times without losing its activity at 12 mol% of catalyst loading. Regarding the mechanism, authors proposed that DBU generated *in situ* a free carbene from the polymer precatalyst, which bound  $\text{CO}_2$ . The following steps are reported in Scheme 34.




Scheme 34. Cyclization of 2-aminobenzenethiols to benzothiazole.

In addition, developing the reaction already cited in Section 2 (Scheme 11) [127], 2-alkynylanilines, aryl iodides, and  $\text{CO}_2$  provided a series of 3,3-diaryl 2,4-quinolinediones (Scheme 35) [205]. It should be noted that this palladium catalyst allowed the incorporation of  $\text{CO}_2$ , conversely from the palladium catalyst described in Section 3 (Scheme 15) [134]. The reaction was scaled up to a 1 mmol scale and product was recovered in 75% yield. In addition, bromobenzene afforded 3,3-diphenyl-2,4-quinolinedione in 72% yield. Some control experiments allowed authors the formulation of the mechanism depicted in Scheme 35.



**Scheme 35.** Multicomponent synthesis of 3,3-diaryl 2,4-quinolinediones.

The reaction of *o*-allylanilines with CO<sub>2</sub> in the presence of a strong base led to 2-quinolinones. The detailed mechanism has been recently studied by density functional theory calculations. The calculated minimum energy reaction pathway is depicted in Scheme 36 [206]. Moreover, the base was found to play a significant role in reducing the energy barriers. The presence of urea and Boc-protected aniline among the by-products was explained by the competitive addition of *tert*-butanol or aniline to the isocyanate intermediate. Calculation also predicted that weaker bases such as Na<sub>2</sub>CO<sub>3</sub> or NaHCO<sub>3</sub> might promote the reaction as well.



**Scheme 36.** Lactamization of sp<sup>2</sup> C–H bonds with CO<sub>2</sub>.

## 6. Conclusions

The importance of CO<sub>2</sub> as C1 synthon in chemical reactions is greatly increasing in recent years. In fact, human activity in its industrial processes produces as by products about 3.3 10<sup>10</sup> metric tons of CO<sub>2</sub> increasing the deleterious greenhouse effect. Thus, since sequestration of some of the circulating carbon dioxide is possible, its use as feedstock in chemical process creates added value to a waste material and positive effect on the environment. Among the many transformations of carbon dioxide into valuable chemicals, the synthesis of heterocycles undoubtedly plays an indisputable role, for the widespread presence of these moieties in new drugs and as building blocks for multistep synthesis [207,208]. The next future should address this research field towards sustainable methods such as the use of recoverable organocatalysts instead of precious metal ones, and towards asymmetric reaction [209], since most of the organic active pharmaceuticals are chiral molecules.

**Author Contributions:** Conceptualization, B.G. and R.D.; Resources, and bibliographic research R.M.; Data Curation, R.M.; Writing-Original Draft Preparation, R.D.; Writing-Review & Editing, R.D.; Visualization, B.G., N.D.C.; Supervision, N.D.C.

**Funding:** This research received no external funding.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Solomon, S.; Plattner, G.-K.; Knutti, R.; Friedlingstein, P. Irreversible climate change due to carbon dioxide emissions. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 1704–1709. [CrossRef]
2. Zhou, Z.; Xia, S.; He, L. Green Catalysis for Three-Component Reaction of Carbon Dioxide, Propargylic Alcohols and Nucleophiles. *Acta Phys. Chim. Sin.* **2018**, *34*, 838–844. [CrossRef]
3. Cantat, T.; He, L.-N. (Eds.) CO<sub>2</sub> Capture and Chemistry 2017. *Curr. Opin. Green Sustain. Chem.* **2017**, *3*, 1–66. Available online: <https://www.sciencedirect.com/journal/current-opinion-in-green-and-sustainable-chemistry/vol/3/suppl/C> (accessed on 5 June 2019).
4. Song, Q.; Zhou, Z.; He, L. Efficient: Selective and sustainable catalysis of carbon dioxide. *Green Chem.* **2017**, *19*, 3707–3728. [CrossRef]
5. ChemSusChem: Hot Topic: Carbon Dioxide First published: 1 April 2017, Last updated: 23 April 2018. Available online: [https://onlinelibrary.wiley.com/doi/toc/10.1002/\(ISSN\)1864-564X.hottopic-co2](https://onlinelibrary.wiley.com/doi/toc/10.1002/(ISSN)1864-564X.hottopic-co2) (accessed on 5 June 2019).
6. Lu, X.B. Carbon Dioxide and Organometallics. In *Topics in Organometallic Chemistry*; Springer: Cham, Switzerland, 2015; Volume 53.
7. Fiorani, G.; Guo, W.; Kleij, A.W. Sustainable conversion of carbon dioxide: The advent of organocatalysis. *Green Chem.* **2015**, *17*, 1375–1389. [CrossRef]
8. Zhou, H.; Lu, X. Lewis base-CO<sub>2</sub> adducts as organocatalysts for CO<sub>2</sub> transformation. *Sci. China Chem.* **2017**, *60*, 904–911. [CrossRef]
9. Tortajada, A.; Juliá-Hernández, F.; Börjesson, M.; Moragas, T.; Martín, R. Transition-Metal-Catalyzed Carboxylation Reactions with Carbon Dioxide. *Angew. Chem. Int. Ed.* **2018**, *57*, 15948–15982. [CrossRef]
10. Poliakov, M.; Leitner, W.; Streng, E.S. The Twelve Principles of CO<sub>2</sub> Chemistry. *Faraday Discuss.* **2015**, *183*, 9–17. [CrossRef]
11. Kumar, M.; Sundaram, S.; Gnansounou, E.; Larroche, C.; Thakura, I.S. Carbon dioxide capture, storage and production of biofuel and biomaterials by bacteria: A review. *Bioresour. Technol.* **2018**, *247*, 1059–1068. [CrossRef]
12. Muthuramalingam, S.; Velusamy, M.; Mayilmurugan, R. Fixation and sequestration of carbon dioxide by copper(II) complexes. *J. Chem. Sci.* **2018**, *130*, 78. [CrossRef]
13. Speight, J.G. Carbonylation. In *Environmental Organic Chemistry for Engineers*; Elsevier: Oxford, UK, 2017.
14. Qiao, C.; Cao, Y.; He, L.-N. Transition Metal-Catalyzed Carboxylation of Terminal Alkynes with CO<sub>2</sub>. *Mini-Rev. Org. Chem.* **2018**, *15*, 283–290. [CrossRef]
15. Zou, B.; Hu, C. Halogen-free processes for organic carbonate synthesis from CO<sub>2</sub>. *Curr. Opin. Green Sustain. Chem.* **2017**, *3*, 11–16. [CrossRef]

16. Vessally, E.; Hosseinian, A.; Babazadeh, M.; Edjlali, L.; Hosseinzadeh-Khanmiri, R. Metal Catalyzed Carboxylative Coupling of Terminal Alkynes, Organohalides and Carbon Dioxide: A Novel and Promising Synthetic Strategy Toward 2-Alkynoates (A Review). *Curr. Org. Chem.* **2018**, *22*, 315–322. [[CrossRef](#)]
17. Buttner, H.; Longwitz, L.; Steinbauer, J.; Wulf, C.; Werner, T. Recent Developments in the Synthesis of Cyclic Carbonates from Epoxides and CO<sub>2</sub>. *Top. Curr. Chem.* **2017**, *375*, 50. [[CrossRef](#)]
18. Francke, R.; Schille, B.; Roemelt, M. Homogeneously Catalyzed Electroreduction of Carbon Dioxide—Methods, Mechanisms, and Catalysts. *Chem. Rev.* **2018**, *118*, 4631–4701. [[CrossRef](#)]
19. Cotton, C.A.R.; Edlich-Muth, C.; Bar-Even, A. Reinforcing carbon fixation: CO<sub>2</sub> reduction replacing and supporting carboxylation. *Curr. Opin. Biotechnol.* **2018**, *49*, 49–56. [[CrossRef](#)]
20. Pander, J.E., III; Ren, D.; Huang, Y.; Loo, N.W.X.; Hong, S.H.L.; Yeo, B.S. Understanding the Heterogeneous Electrocatalytic Reduction of Carbon Dioxide on Oxide-Derived Catalysts. *ChemElectroChem* **2018**, *5*, 219–237. [[CrossRef](#)]
21. Voiry, D.; Shin, H.S.; Loh, K.P.; Chhowalla, M. Low-dimensional catalysts for hydrogen evolution and CO<sub>2</sub> reduction. *Nat. Rev. Chem.* **2018**, *2*, 0105. [[CrossRef](#)]
22. Pagliaro, M.; Fidalgo, A.; Palmisano, L.; Ilharco, L.M.; Parrino, F.; Ciriminna, R. Polymers of limonene oxide and carbon dioxide: Polycarbonates of the solar economy. *Preprints* **2018**, *2018*, 040067. [[CrossRef](#)]
23. Muthuraj, R.; Mekonnen, T. Recent progress in carbon dioxide (CO<sub>2</sub>) as feedstock for sustainable materials development: Co-polymers and polymer blends. *Polymer* **2018**, *145*, 348–373. [[CrossRef](#)]
24. Kamphuis, A.J.; Piccioni, F.; Pescarmona, P.P. CO<sub>2</sub>-fixation into cyclic and polymeric carbonates: Principles and applications. *Green Chem.* **2019**, *21*, 406–448. [[CrossRef](#)]
25. Tomishige, K.; Tamura, M.; Nakagawa, Y. CO<sub>2</sub> Conversion with Alcohols and Amines into Carbonates, Ureas, and Carbamates over CeO<sub>2</sub> Catalyst in the Presence and Absence of 2-Cyanopyridine. *Chem. Rec.* **2018**, *18*, 1–26. [[CrossRef](#)]
26. Liang, J.; Huang, J.-B.; Cao, R. Metal-organic frameworks and porous organic polymers for sustainable fixation of carbon dioxide into cyclic carbonates. *Coord. Chem. Rev.* **2019**, *378*, 32–65. [[CrossRef](#)]
27. Li, J.-Y.; Song, Q.-W.; Zhang, K.; Liu, P. Catalytic Conversion of Carbon Dioxide through C-N Bond Formation. *Molecules* **2019**, *24*, 182. [[CrossRef](#)]
28. Farshbaf, S.; Fekri, L.Z.; Nikpassand, M.; Mohammadi, R.; Vessally, E. Dehydrative condensation of β-aminoalcohols with CO<sub>2</sub>: An environmentally benign access to 2-oxazolidinone derivatives. *J. CO<sub>2</sub> Util.* **2018**, *25*, 194–204. [[CrossRef](#)]
29. Didehbana, K.; Vessally, E.; Salary, M.; Edjlali, L.; Babazadeh, M. Synthesis of a variety of key medicinal heterocyclic compounds via chemical fixation of CO<sub>2</sub> onto *o*-alkynylaniline derivatives. *J. CO<sub>2</sub> Util.* **2018**, *23*, 42–50. [[CrossRef](#)]
30. Vessally, E.; Babazadeh, M.; Hosseinian, A.; Arshadi, S.; Edjlali, L. Nanocatalysts for chemical transformation of carbon dioxide. *J. CO<sub>2</sub> Util.* **2017**, *21*, 491–502. [[CrossRef](#)]
31. Vessally, E.; Didehban, K.; Babazadeh, M.; Hosseinian, A.; Edjlali, L. Chemical fixation of CO<sub>2</sub> with aniline derivatives: A new avenue to the synthesis of functionalized azole compounds (A review). *J. CO<sub>2</sub> Util.* **2017**, *21*, 480–490. [[CrossRef](#)]
32. Vessally, E.; Soleimani-Amiri, S.; Hosseinian, A.; Edjlali, L.; Babazadeh, M. Chemical fixation of CO<sub>2</sub> to 2-aminobenzonitriles: A straightforward route to quinazoline-2,4(1*H*,3*H*)-diones with green and sustainable chemistry perspectives. *J. CO<sub>2</sub> Util.* **2017**, *21*, 342–352. [[CrossRef](#)]
33. Arshadi, S.; Vessally, E.; Sobati, M.; Hosseinian, A.; Bekhradnia, A. Chemical fixation of CO<sub>2</sub> to *N*-propargylamines: A straightforward route to 2-oxazolidinones. *J. CO<sub>2</sub> Util.* **2017**, *19*, 120–129. [[CrossRef](#)]
34. Hosseinian, A.; Ahmadi, S.; Mohammadi, R.; Monfared, A.; Rahmani, Z. Three-component reaction of amines, epoxides, and carbon dioxide: A straightforward route to organic carbamates. *J. CO<sub>2</sub> Util.* **2018**, *27*, 381–389. [[CrossRef](#)]
35. Liu, X.-F.; Wang, M.-Y.; He, L.-N. Heterogeneous Catalysis for Oxazolidinone Synthesis from Aziridines and CO<sub>2</sub>. *Curr. Org. Chem.* **2017**, *21*, 698–707. [[CrossRef](#)]
36. Arshadi, S.; Vessally, E.; Hosseinian, A.; Soleimani-Amiri, S.; Edjlali, L. Three component coupling of CO<sub>2</sub>, propargyl alcohols, and amines: An environmentally benign access to cyclic and acyclic carbamates (A Review). *J. CO<sub>2</sub> Util.* **2017**, *21*, 108–118. [[CrossRef](#)]
37. Yu, B.; He, L.-N. Upgrading Carbon Dioxide by Incorporation into Heterocycles. *ChemSusChem* **2015**, *8*, 52–62. [[CrossRef](#)]

38. Darensbourg, D.J. Making Plastics from Carbon Dioxide: Salen Metal Complexes as Catalysts for the Production of Polycarbonates from Epoxides and CO<sub>2</sub>. *Chem. Rev.* **2007**, *107*, 2388–2410. [[CrossRef](#)]
39. Poland, S.J.; Darensbourg, D.J. A quest for polycarbonates provided via sustainable epoxide/CO<sub>2</sub> copolymerization processes. *Green Chem.* **2017**, *19*, 4990–5011. [[CrossRef](#)]
40. Zhang, S.S. A review on electrolyte additives for lithium-ion batteries. *J. Power Sources* **2006**, *162*, 1379–1394. [[CrossRef](#)]
41. He, Q.; O'Brien, J.W.; Kitselman, K.A.; Tompkins, L.E.; Curtis, G.C.T.; Kerton, F.M. Synthesis of cyclic carbonates from CO<sub>2</sub> and epoxides using ionic liquids and related catalysts including choline chloride–metal halide mixtures. *Catal. Sci. Technol.* **2014**, *4*, 1513–1528. [[CrossRef](#)]
42. Lan, D.-H.; Fan, N.; Wang, Y.; Gao, X.; Zhang, P.; Chen, L.; Au, C.-T.; Yin, S.-F. Recent advances in metal-free catalysts for the synthesis of cyclic carbonates from CO<sub>2</sub> and epoxides. *Chin. J. Catal.* **2016**, *37*, 826–845. [[CrossRef](#)]
43. Bobbink, F.D.; Dyson, P.J. Synthesis of carbonates and related compounds incorporating CO<sub>2</sub> using ionic liquid-type catalysts: State-of-the-art and beyond. *J. Catal.* **2016**, *343*, 52–61. [[CrossRef](#)]
44. Alves, M.; Grignard, B.; Mereau, R.; Jerome, C.; Tassaing, T.; Detrembleur, C. Organocatalyzed coupling of carbon dioxide with epoxides for the synthesis of cyclic carbonates: Catalyst design and mechanistic studies. *Catal. Sci. Technol.* **2017**, *7*, 2651–2684. [[CrossRef](#)]
45. Shaikh, R.R.; Pornpraprom, S.; D'Elia, V. Catalytic Strategies for the Cycloaddition of Pure, Diluted, and Waste CO<sub>2</sub> to Epoxides under Ambient Conditions. *ACS Catal.* **2018**, *8*, 419–450. [[CrossRef](#)]
46. Kim, H.-U.; Babu, R.; Roshan, R.; Park, D.-W. Catalytic performance of metal azolate frameworks in the solventless synthesis of cyclic carbonates from CO<sub>2</sub> and epoxides. *Appl. Catal. A Gen.* **2017**, *538*, 59–65. [[CrossRef](#)]
47. Elkurtehi, A.I.; Kerton, F.M. Coupling Reactions of Carbon Dioxide with Epoxides Catalyzed by Vanadium Aminophenolate Complexes. *ChemSusChem* **2017**, *10*, 1249–1254. [[CrossRef](#)]
48. Chen, F.; Liu, N.; Dai, B. Iron(II) Bis-CNN Pincer Complex-Catalyzed Cyclic Carbonate Synthesis at Room Temperature. *ACS Sustain. Chem. Eng.* **2017**, *5*, 9065–9075. [[CrossRef](#)]
49. Wu, X.; North, M. A Bimetallic Aluminium(Salphen) Complex for the Synthesis of Cyclic Carbonates from Epoxides and Carbon Dioxide. *ChemSusChem* **2017**, *10*, 74–78. [[CrossRef](#)]
50. Luo, R.; Yang, Z.; Zhang, W.; Zhou, X.; Ji, H. Recyclable bifunctional aluminum salen catalyst for CO<sub>2</sub> fixation: The efficient formation of five-membered heterocyclic compounds. *Sci. China Chem.* **2017**, *60*, 979–989. [[CrossRef](#)]
51. Zhou, F.; Xie, S.-L.; Gao, X.-T.; Zhang, R.; Wang, C.-H.; Yin, G.-Q.; Zhou, J. Activation of (salen)CoI complex by phosphorane for carbon dioxide transformation at ambient temperature and pressure. *Green Chem.* **2017**, *19*, 3908–3915. [[CrossRef](#)]
52. Kaneko, S.; Shirakawa, S. Potassium Iodide–Tetraethylene Glycol Complex as a Practical Catalyst for CO<sub>2</sub> Fixation Reactions with Epoxides under Mild Conditions. *ACS Sustain. Chem. Eng.* **2017**, *5*, 2836–2840. [[CrossRef](#)]
53. Steinbauer, J.; Werner, T. Poly(ethylene glycol)s as Ligands in Calcium-Catalyzed Cyclic Carbonate Synthesis. *ChemSusChem* **2017**, *10*, 3025–3029. [[CrossRef](#)]
54. Steinbauer, J.; Spannenberg, A.; Werner, T. An in situ formed Ca<sup>2+</sup>–crown ether complex and its use in CO<sub>2</sub>-fixation reactions with terminal and internal epoxides. *Green Chem.* **2017**, *19*, 3769–3779. [[CrossRef](#)]
55. Longwitz, L.; Steinbauer, J.; Spannenberg, A.; Werner, T. Calcium-Based Catalytic System for the Synthesis of Bio-Derived Cyclic Carbonates under Mild Conditions. *ACS Catal.* **2018**, *8*, 665–672. [[CrossRef](#)]
56. Zhao, T.-X.; Zhang, Y.-Y.; Liang, J.; Li, P.; Hu, X.-B.; Wu, Y.-T. Multisite activation of epoxides by recyclable CaI<sub>2</sub>/N-methyldiethanolamine catalyst for CO<sub>2</sub> fixation: A facile access to cyclic carbonates under mild conditions. *Mol. Catal.* **2018**, *450*, 87–94. [[CrossRef](#)]
57. Chowdhury, A.H.; Bhanja, P.; Salam, N.; Bhaumik, A.; Islam, S.M. Magnesium oxide as an efficient catalyst for CO<sub>2</sub> fixation and N-formylation reactions under ambient conditions. *Mol. Catal.* **2018**, *450*, 46–54. [[CrossRef](#)]
58. Kilic, A.; Durgun, M.; Aytar, E.; Yavuz, R. Synthesis and characterization of novel positively charged organocobaloximes as catalysts for the fixation of CO<sub>2</sub> to cyclic carbonates. *J. Organomet. Chem.* **2018**, *858*, 78–88. [[CrossRef](#)]

59. Della Monica, F.; Buonerba, A.; Paradiso, V.; Milione, S.; Grassi, A.; Capacchione, C. [OSSO]-Type Fe(III) Metallate as Single-Component Catalyst for the CO<sub>2</sub> Cycloaddition to Epoxides. *Adv. Synth. Catal.* **2018**, *361*, 283–288. [[CrossRef](#)]
60. Kuznetsova, S.A.; Rulev, Y.A.; Larionov, V.A.; Smol'yakov, A.F.; Zubavichus, Y.V.; Maleev, V.I.; Li, H.; North, M.; Saghyan, A.S.; Belokon, Y.N. Self-Assembled Ionic Composites of Negatively Charged Zn(salen) Complexes and Triphenylmethane Derived Polycations as Recyclable Catalysts for the Addition of Carbon Dioxide to Epoxides. *ChemCatChem* **2019**, *11*, 511–519. [[CrossRef](#)]
61. Milani, J.L.S.; Oliveira, I.S.; Dos Santos, P.A.; Valdo, A.K.S.M.; Martins, F.T.; Cangussu, D.; Das Chagas, R.P. Chemical fixation of carbon dioxide to cyclic carbonates catalyzed by zinc(II) complex bearing 1,2-disubstituted benzimidazole ligand. *Chin. J. Catal.* **2018**, *39*, 245–249. [[CrossRef](#)]
62. Chen, Y.; Luo, R.; Yang, Z.; Zhou, X.; Ji, H. Imidazolium-based ionic liquid decorated zinc porphyrin catalyst for converting CO<sub>2</sub> into five-membered heterocyclic molecules. *Sustain. Energy Fuels* **2018**, *2*, 125–132. [[CrossRef](#)]
63. Sogukomerogullari, H.G.; Aytar, E.; Ulusoy, M.; Demir, S.; Dege, N.; Richeson, D.S.; Sönmez, M. Synthesis of complexes Fe, Co and Cu supported by “SNS” pincer ligands and their ability to catalytically form cyclic carbonates. *Inorg. Chim. Acta* **2018**, *471*, 290–296. [[CrossRef](#)]
64. Bresciani, G.; Marco Bortoluzzi, M.; Marchetti, F.; Pampaloni, G. Iron(III) *N,N*-Dialkylcarbamate-Catalyzed Formation of Cyclic Carbonates from CO<sub>2</sub> and Epoxides under Ambient Conditions by Dynamic CO<sub>2</sub> Trapping as Carbamate Ligands. *ChemSusChem* **2018**, *11*, 2737–2743. [[CrossRef](#)]
65. Arayachukiat, S.; Kongtes, C.; Barthel, A.; Vummaleti, S.V.C.; Poater, A.; Wannakao, S.; Cavallo, L.; D'Elia, V. Ascorbic Acid as a Bifunctional Hydrogen Bond Donor for the Synthesis of Cyclic Carbonates from CO<sub>2</sub> under Ambient Conditions. *ACS Sustain. Chem. Eng.* **2017**, *5*, 6392–6397. [[CrossRef](#)]
66. Sopeña, S.; Martin, E.; Escudero-Adán, E.C.; Kleij, A.W. Pushing the Limits with Squaramide-Based Organocatalysts in Cyclic Carbonate Synthesis. *ACS Catal.* **2017**, *7*, 3532–3539. [[CrossRef](#)]
67. Kumatabara, Y.; Okada, M.; Shirakawa, S. Triethylamine Hydroiodide as a Simple Yet Effective Bifunctional Catalyst for CO<sub>2</sub> Fixation Reactions with Epoxides under Mild Conditions. *ACS Sustain. Chem. Eng.* **2017**, *5*, 7295–7301. [[CrossRef](#)]
68. Bettner, H.; Steinbauer, J.; Wulf, C.; Dindaroglu, M.; Schmalz, H.-G.; Werner, T. Organocatalyzed Synthesis of Oleochemical Carbonates from CO<sub>2</sub> and Renewables. *ChemSusChem* **2017**, *10*, 1076–1079. [[CrossRef](#)]
69. Saptal, V.B.; Bhanage, B.M. Bifunctional Ionic Liquids Derived from Biorenewable Sources as Sustainable Catalysts for Fixation of Carbon Dioxide. *ChemSusChem* **2017**, *10*, 1145–1151. [[CrossRef](#)]
70. Ma, R.; He, L.-N.; Liu, X.-F.; Liu, X.; Wang, M.-I. DBU as activator for the N-iodosuccinimide promoted chemical fixation of carbon dioxide with epoxides. *J. CO<sub>2</sub> Util.* **2017**, *19*, 28–32. [[CrossRef](#)]
71. Zhang, Z.; Fan, F.; Xing, H.; Yang, Q.; Bao, Z.; Ren, Q. Efficient Synthesis of Cyclic Carbonates from Atmospheric CO<sub>2</sub> Using a Positive Charge Delocalized Ionic Liquid Catalyst. *ACS Sustain. Chem. Eng.* **2017**, *5*, 2841–2846. [[CrossRef](#)]
72. Hirose, T.; Qu, S.; Kodama, K.; Wang, X. Organocatalyst system for disubstituted carbonates from cycloaddition between CO<sub>2</sub> and internal epoxides. *J. CO<sub>2</sub> Util.* **2018**, *24*, 261–265. [[CrossRef](#)]
73. Wang, T.; Zheng, D.; Ma, Y.; Guo, J.; He, Z.; Ma, B.; Liu, L.; Ren, T.; Wang, L.; Zhang, J. Benzyl substituted imidazolium ionic liquids as efficient solvent-free catalysts for the cycloaddition of CO<sub>2</sub> with epoxides: Experimental and Theoretic study. *J. CO<sub>2</sub> Util.* **2017**, *22*, 44–52. [[CrossRef](#)]
74. Yue, S.; Hao, X.J.; Wang, P.-P.; Li, J. Amino acid-based ionic liquids for CO<sub>2</sub> conversion to form cyclic carbonate under solvent-free conditions. *Mol. Catal.* **2017**, *433*, 420–429. [[CrossRef](#)]
75. Jose, T.; Cañellas, S.; Pericàs, M.A.; Kleij, A.W. Polystyrene-supported bifunctional resorcinarenes as cheap, metal-free and recyclable catalysts for epoxide/CO<sub>2</sub> coupling reactions. *Green Chem.* **2017**, *19*, 5488–5493. [[CrossRef](#)]
76. Li, K.; Wu, X.; Gu, Q.; Zhao, X.; Yuan, M.; Ma, W.; Ni, W.; Hou, Z. Inclusion complexes of organic salts with β-cyclodextrin as organocatalysts for CO<sub>2</sub> cycloaddition with epoxides. *RSC Adv.* **2017**, *7*, 14721–14732. [[CrossRef](#)]
77. Hu, J.; Ma, J.; Liu, H.; Qian, Q.; Xiea, C.; Han, B. Dual-ionic liquid system: An efficient catalyst for chemical fixation of CO<sub>2</sub> to cyclic carbonates under mild conditions. *Green Chem.* **2018**, *20*, 2990–2994. [[CrossRef](#)]

78. Peng, J.; Wang, S.; Yang, H.-J.; Ban, B.; Wei, Z.; Wang, L.; Lei, B. Highly efficient fixation of carbon dioxide to cyclic carbonates with new multi-hydroxyl bis-(quaternary ammonium) ionic liquids as metal-free catalysts under mild conditions. *Fuel* **2018**, *224*, 481–488. [[CrossRef](#)]
79. Wang, X.; Wang, L.; Zhao, Y.; Kodama, K.; Hirose, T. Efficient and practical organocatalytic system for the synthesis of cyclic carbonates from carbon dioxide and epoxides: 3-hydroxypyridine/tetra-*n*-butylammonium iodide. *Tetrahedron* **2017**, *73*, 1190–1195. [[CrossRef](#)]
80. Castro-Osma, J.A.; Martínez, J.; de la Cruz-Martínez, F.; Caballero, M.P.; Fernández-Baeza, J.; Rodríguez-López, J.; Otero, A.; Lara-Sánchez, A.; Tejada, J. Development of hydroxy-containing imidazole organocatalysts for CO<sub>2</sub> fixation into cyclic carbonates. *Catal. Sci. Technol.* **2018**, *8*, 1981–1987. [[CrossRef](#)]
81. Liu, A.-H.; Dang, Y.-L.; Zhou, H.; Zhang, J.-J.; Lu, X.-B. CO<sub>2</sub> Adducts of Carbodicarbenes: Robust and Versatile Organocatalysts for Chemical Transformation of Carbon Dioxide into Heterocyclic Compounds. *ChemCatChem* **2018**, *10*, 2686–2692. [[CrossRef](#)]
82. Maeda, C.; Sasaki, S.; Takaishi, K.; Ema, T. Calix[4]pyrroles as macrocyclic organocatalysts for the synthesis of cyclic carbonates from epoxides and carbon dioxide. *Catal. Sci. Technol.* **2018**, *8*, 4193–4198. [[CrossRef](#)]
83. Hong, M.; Kim, Y.; Kim, H.; Cho, H.J.; Baik, M.-H.; Kim, Y. Scorpionate Catalysts for Coupling CO<sub>2</sub> and Epoxides to Cyclic Carbonates: A Rational Design Approach for Organocatalysts. *J. Org. Chem.* **2018**, *83*, 9370–9380. [[CrossRef](#)]
84. Liu, N.; Xie, Y.-F.; Wang, C.; Li, S.-J.; Wei, D.; Li, M.; Dai, B. Cooperative Multifunctional Organocatalysts for Ambient Conversion of Carbon Dioxide into Cyclic Carbonates. *ACS Catal.* **2018**, *8*, 9945–9957. [[CrossRef](#)]
85. Mosteirín, N.F.; Jehanno, C.; Ruipérez, F.; Sardon, H.; Dove, A.P. Rational study of DBU salts for the CO<sub>2</sub> insertion into epoxides for the synthesis of cyclic carbonates. *ACS Sustain. Chem. Eng.* **2019**. [[CrossRef](#)]
86. Wang, S.; Song, K.; Zhang, C.; Shu, Y.; Li, T.; Tan, B. A novel metalporphyrin-based microporous organic polymer with high CO<sub>2</sub> uptake and efficient chemical conversion of CO<sub>2</sub> under ambient conditions. *J. Mater. Chem. A* **2017**, *5*, 1509–1515. [[CrossRef](#)]
87. Chen, Y.; Luo, R.; Xu, Q.; Zhang, W.; Zhou, X.; Ji, H. State-of-the-Art Aluminum Porphyrin-based Heterogeneous Catalysts for the Chemical Fixation of CO<sub>2</sub> into Cyclic Carbonates at Ambient Conditions. *ChemCatChem* **2017**, *9*, 767–773. [[CrossRef](#)]
88. Chen, Y.; Luo, R.; Xu, Q.; Jiang, J.; Zhou, X.; Ji, H. Charged Metalloporphyrin Polymers for Cooperative Synthesis of Cyclic Carbonates from CO<sub>2</sub> under Ambient Conditions. *ChemSusChem* **2017**, *10*, 2534–2541. [[CrossRef](#)]
89. Zhou, K.; Mousavi, B.; Luo, Z.; Phatanasri, S.; Chaemchuen, S.; Verpoort, F. Characterization and properties of Zn/Co zeolitic imidazolate frameworks vs. ZIF-8 and ZIF-67. *J. Mater. Chem. A* **2017**, *5*, 952–957. [[CrossRef](#)]
90. Zanon, A.; Chaemchuen, S.; Mousavi, B.; Verpoort, F. 1 Zn-doped ZIF-67 as catalyst for the CO<sub>2</sub> fixation into cyclic carbonates. *J. CO<sub>2</sub> Util.* **2017**, *20*, 282–291. [[CrossRef](#)]
91. Buyukcakir, O.; Je, S.H.; Talapaneni, S.N.; Kim, D.; Coskun, A. Charged Covalent Triazine Frameworks for CO<sub>2</sub> Capture and Conversion. *ACS Appl. Mater. Interfaces* **2017**, *9*, 7209–7216. [[CrossRef](#)]
92. Guo, Z.; Jiang, Q.; Shi, Y.; Li, J.; Yang, X.; Hou, W.; Zhou, Y.; Wang, J. Tethering Dual Hydroxyls into Mesoporous Poly(ionic liquid)s for Chemical Fixation of CO<sub>2</sub> at Ambient Conditions: A Combined Experimental and Theoretical Study. *ACS Catal.* **2017**, *7*, 6770–6780. [[CrossRef](#)]
93. Li, P.-Z.; Wang, X.-J.; Liu, J.; Phang, H.S.; Li, Y.; Zhao, Y. Highly Effective Carbon Fixation via Catalytic Conversion of CO<sub>2</sub> by an Acylamide-Containing Metal–Organic Framework. *Chem. Mater.* **2017**, *29*, 9256–9261. [[CrossRef](#)]
94. Lu, B.-B.; Yang, J.; Liu, Y.-Y.; Ma, J.F. A Polyoxovanadate–Resorcin[4]arene-Based Porous Metal–Organic Framework as an Efficient Multifunctional Catalyst for the Cycloaddition of CO<sub>2</sub> with Epoxides and the Selective Oxidation of Sulfides. *Inorg. Chem.* **2017**, *56*, 11710–11720. [[CrossRef](#)]
95. Mousavi, B.; Chaemchuen, S.; Moosavi, B.; Zhou, K.; Yusubov, M.; Verpoort, F. CO<sub>2</sub> Cycloaddition to Epoxides by using M-DABCO Metal–Organic Frameworks and the Influence of the Synthetic Method on Catalytic Reactivity. *ChemistryOpen* **2017**, *6*, 674–680. [[CrossRef](#)]
96. Ding, M.; Chen, S.; Liu, X.-Q.; Sun, L.-B.; Lu, J.; Jiang, H.-L. Metal–Organic Framework-Templated Catalyst: Synergy in Multiple Sites for Catalytic CO<sub>2</sub> Fixation. *ChemSusChem* **2017**, *10*, 1898–1903. [[CrossRef](#)]
97. Song, L.; Zhang, X.; Chen, C.; Liu, X.; Zhang, N. UTSA-16 as an efficient microporous catalyst for CO<sub>2</sub> conversion to cyclic carbonates. *Microporous Mesoporous Mater.* **2017**, *241*, 36–42. [[CrossRef](#)]

98. Liu, M.; Lan, J.; Liang, L.; Sun, J.; Arai, M. Heterogeneous catalytic conversion of CO<sub>2</sub> and epoxides to cyclic carbonates over multifunctional tri-s-triazine terminal-linked ionic liquids. *J. Catal.* **2017**, *347*, 138–147. [[CrossRef](#)]
99. Zhong, W.; Bobbink, F.D.; Fei, Z.; Dyson, P.J. Polyimidazolium Salts: Robust Catalysts for the Cycloaddition of Carbon Dioxide into Carbonates in Solvent-Free Conditions. *ChemSusChem* **2017**, *10*, 2728–2735. [[CrossRef](#)]
100. Bobbink, F.D.; Van Muyden, A.P.; Gopakumar, A.; Fei, Z.; Dyson, P.J. Synthesis of Cross-linked Ionic Poly(styrenes) and their Application as Catalysts for the Synthesis of Carbonates from CO<sub>2</sub> and Epoxides. *ChemPlusChem* **2017**, *82*, 144–151. [[CrossRef](#)]
101. Zhang, X.; Su, D.; Xiao, L.; Wu, W. Immobilized protic ionic liquids: Efficient catalysts for CO<sub>2</sub> fixation with epoxides. *J. CO<sub>2</sub> Util.* **2017**, *17*, 37–42. [[CrossRef](#)]
102. Ravi, S.; Puthiaraj, P.; Ahn, W.-S. Cyclic carbonate synthesis from CO<sub>2</sub> and epoxides over diamine functionalized porous organic frameworks. *J. CO<sub>2</sub> Util.* **2017**, *21*, 450–458. [[CrossRef](#)]
103. Steinbauer, J.; Longwitz, L.; Frank, M.; Epping, J.; Kragld, U.; Werner, T. Immobilized bifunctional phosphonium salts as recyclable organocatalysts in the cycloaddition of CO<sub>2</sub> and epoxides. *Green. Chem.* **2017**, *19*, 4435–4445. [[CrossRef](#)]
104. Zhi, Y.; Shao, P.; Feng, X.; Xia, H.; Zhang, Y.; Shi, Z.; Mua, Y.; Liu, X. Covalent organic frameworks: Efficient, metal-free, heterogeneous organocatalysts for chemical fixation of CO<sub>2</sub> under mild conditions. *J. Mater. Chem. A* **2018**, *6*, 374–382. [[CrossRef](#)]
105. Maya, M.M.; Rangel-Rangel, E.; Díaz, U.; Iglesias, M. Efficient cycloaddition of CO<sub>2</sub> to epoxides using novel heterogeneous organocatalysts based on tetramethylguanidine-functionalized porous polyphenylenes. *J. CO<sub>2</sub> Util.* **2018**, *25*, 170–179. [[CrossRef](#)]
106. Song, X.; Wu, Y.; Pan, D.; Wei, R.; Gao, L.; Zhang, J.; Xiao, G. Melem based multifunctional catalyst for chemical fixation of carbon dioxide into cyclic carbonate. *J. CO<sub>2</sub> Util.* **2018**, *24*, 287–297. [[CrossRef](#)]
107. Subramanian, S.; Park, J.; Byun, J.; Jung, Y.; Yavuz, C.T. Highly Efficient Catalytic Cyclic Carbonate Formation by Pyridyl Salicylimines. *ACS Appl. Mater. Interfaces* **2018**, *10*, 9478–9484. [[CrossRef](#)]
108. Wang, X.; Liu, M.S.; Yang, L.; Lan, J.W.; Chen, Y.L.; Sun, J.M. Synthesis of Zn Modified Carbon Nitrides Heterogeneous Catalyst for the Cycloaddition of CO<sub>2</sub> to Epoxides. *ChemistrySelect* **2018**, *3*, 4101–4109. [[CrossRef](#)]
109. Liu, D.; Li, G.; Liu, H. Functionalized MIL-101 with imidazolium-based ionic liquids for the cycloaddition of CO<sub>2</sub> and epoxides under mild condition. *Appl. Surf. Sci.* **2018**, *428*, 218–225. [[CrossRef](#)]
110. Shi, X.-L.; Chen, Y.; Duan, P.; Zhang, W.; Hu, Q. Conversion of CO<sub>2</sub> into Organic Carbonates over a Fiber-Supported Ionic Liquid Catalyst in Impellers of the Agitation System. *ACS Sustain. Chem. Eng.* **2018**, *6*, 7119–7127. [[CrossRef](#)]
111. Jayakumar, S.; Li, H.; Tao, L.; Li, C.; Liu, L.; Chen, J.; Yang, Q. Cationic Zn-Porphyrin Immobilized in Mesoporous Silicas as Bifunctional Catalyst for CO<sub>2</sub> Cycloaddition Reaction under Cocatalyst Free Conditions. *ACS Sustain. Chem. Eng.* **2018**, *6*, 9237–9245. [[CrossRef](#)]
112. Jayakumar, S.; Li, H.; Chen, J.; Yang, Q. Cationic Zn-Porphyrin Polymer Coated onto CNTs as a Cooperative Catalyst for the Synthesis of Cyclic Carbonates. *ACS Appl. Mater. Interfaces* **2018**, *10*, 2546–2555. [[CrossRef](#)]
113. Tiffner, M.; Häring, M.; Díaz Díaz, D.; Waser, M. Cationic Polymers Bearing Quaternary Ammonium Groups-Catalyzed CO<sub>2</sub> Fixation with Epoxides. *Top. Catal.* **2018**, *61*, 1545–1550. [[CrossRef](#)]
114. Tang, L.; Zhang, S.; Wu, Q.; Wang, X.; Wu, H.; Jiang, Z. Heterobimetallic Metal-Organic Framework Nanocages as Highly Efficient Catalysts for CO<sub>2</sub> Conversion under Mild Condition. *J. Mater. Chem. A* **2018**, *6*, 2964–2973. [[CrossRef](#)]
115. Li, Y.-Z.; Wang, H.-H.; Yang, H.-Y.; Hou, L.; Wang, Y.-Y.; Zhu, Z. An Uncommon Carboxyl-Decorated Metal-Organic Framework with Selective Gas Adsorption and Catalytic Conversion of CO<sub>2</sub>. *Chem. Eur. J.* **2018**, *24*, 865–871. [[CrossRef](#)]
116. Zhang, Z.; Gao, H.; Wu, H.; Qian, Y.; Chen, L.; Chen, J. Chemical Fixation of CO<sub>2</sub> by Using Carbon Material-Grafted N-Heterocyclic Carbene Silver and Copper Complexes. *ACS Appl. Nano Mater.* **2018**, *1*, 6463–6476. [[CrossRef](#)]
117. Calabrese, C.; Liotta, L.F.; Giacalone, F.; Gruttadauria, M.; Aprile, C. Supported Polyhedral Oligomeric Silsesquioxane-Based (POSS) Materials as Highly Active Organocatalysts for the Conversion of CO<sub>2</sub>. *ChemCatChem* **2019**, *11*, 560–567. [[CrossRef](#)]

118. Rintjema, J.; Kleij, A.W. Aluminum-mediated formation of cyclic carbonates: Benchmarking catalytic performance metrics. *ChemSusChem* **2017**, *10*, 1274–1282. [[CrossRef](#)]
119. Yingcharoen, P.; Kongtes, C.; Arayachukiat, S.; Suvarnapunya, K.; Vummaleti, S.V.C.; Wannakao, S.; Cavallo, L.; Poater, A.; D'Elia, V. Assessing the pK<sub>a</sub>-Dependent Activity of Hydroxyl Hydrogen Bond Donors in the Organocatalyzed Cycloaddition of Carbon Dioxide to Epoxides: Experimental and Theoretical Study. *Adv. Synth. Catal.* **2019**, *361*, 366–373. [[CrossRef](#)]
120. Laserna, V.; Martin, E.; Escudero-Adán, E.C.; Kleij, A.W. Substrate-Triggered Stereoselective Preparation of Highly Substituted Organic Carbonates. *ACS Catal.* **2017**, *7*, 5478–5482. [[CrossRef](#)]
121. Lee, Y.; Choi, J.; Kim, H. Stereocontrolled, Divergent, Al(III)-Catalyzed Coupling of Chiral *N*-Aryl Epoxy Amines and CO<sub>2</sub>. *Org. Lett.* **2018**, *20*, 5036–5039. [[CrossRef](#)]
122. Jawad, A.; Rezaei, F.; Rowanghi, A.A. Porous polymeric hollow fibers as bifunctional catalysts for CO<sub>2</sub> conversion to cyclic carbonates. *J. CO<sub>2</sub> Util.* **2017**, *21*, 589–596. [[CrossRef](#)]
123. Liu, F.; Huang, K.; Wu, Q.; Dai, S. Solvent-Free Self-Assembly to the Synthesis of Nitrogen-Doped Ordered Mesoporous Polymers for Highly Selective Capture and Conversion of CO<sub>2</sub>. *Adv. Mater.* **2017**, *29*, 1700445. [[CrossRef](#)]
124. Kim, K.; Kim, S.; Talapaneni, S.N.; Buyukcakir, O.; Almutawa, A.M.I.; Polychronopoulou, K.; Coskun, A. Transition metal complex directed synthesis of porous cationic polymers for efficient CO<sub>2</sub> capture and conversion. *Polymer* **2017**, *126*, 296–302. [[CrossRef](#)]
125. Mirabaud, A.; Mulatier, J.-C.; Martinez, A.; Dutasta, J.-P.; Dufaud, V. Merging host-guest chemistry and organocatalysis for the chemical valorization of CO<sub>2</sub>. *Catal. Today* **2017**, *281*, 387–391. [[CrossRef](#)]
126. Bresciani, G.; Marchetti, F.; Rizzi, G.; Gabbani, A.; Pineider, F.; Pampaloni, G. Metal *N,N*-dialkylcarbamates as easily available catalytic precursors for the carbon dioxide/propylene oxide coupling under ambient conditions. *J. CO<sub>2</sub> Util.* **2018**, *28*, 168–173. [[CrossRef](#)]
127. Samanta, S.; Srivastava, R. A novel method to introduce acidic and basic bifunctional sites in graphitic carbon nitride for sustainable catalysis: Cycloaddition, esterification, and transesterification reactions. *Sustain. Energy Fuels* **2017**, *1*, 1390–1404. [[CrossRef](#)]
128. Kim, H.G.; Limb, C.S.; Kim, D.-W.; Cho, D.H.; Lee, D.K.; Chung, J.S. Multifunctional alkanolamine as a catalyst for CO<sub>2</sub> and propylene oxide cycloaddition. *Mol. Catal.* **2017**, *438*, 121–129. [[CrossRef](#)]
129. Onyenkeadi, V.; Kellici, S.; Saha, B. Greener synthesis of 1,2-butylene carbonate from CO<sub>2</sub> using graphene-inorganic nanocomposite catalyst. *Energy* **2018**, *165A*, 867–876. [[CrossRef](#)]
130. Liu, W.; Lu, G.; Xiao, B.; Xie, C. Potassium iodide–polyethylene glycol catalyzed cycloaddition reaction of epoxidized soybean oil fatty acid methyl esters with CO<sub>2</sub>. *RSC Adv.* **2018**, *8*, 30860–30867. [[CrossRef](#)]
131. Liu, S.; Suematsu, N.; Maruoka, K.; Shirakawa, S. Design of bifunctional quaternary phosphonium salt catalysts for CO<sub>2</sub> fixation reaction with epoxides under mild conditions. *Green Chem.* **2016**, *18*, 4611–4615. [[CrossRef](#)]
132. Ema, T.; Yokoyama, M.; Watanabe, S.; Sasaki, S.; Ota, H.; Takaishi, K. Chiral Macrocyclic Organocatalysts for Kinetic Resolution of Disubstituted Epoxides with Carbon Dioxide. *Org. Lett.* **2017**, *19*, 4070–4073. [[CrossRef](#)]
133. Maeda, C.; Mitsuzane, M.; Ema, T. Chiral Bifunctional Metalloporphyrin Catalysts for Kinetic Resolution of Epoxides with Carbon Dioxide. *Org. Lett.* **2019**, *21*, 1397–1401. [[CrossRef](#)]
134. Qin, J.; Larionov, V.A.; Harms, K.; Meggers, E. Kinetic Resolution of Epoxides with CO<sub>2</sub> Catalyzed by a Chiral-at-Iridium Complex. *ChemSusChem* **2019**, *12*, 320–325. [[CrossRef](#)]
135. Alves, M.; Grignard, B.; Boyaval, A.; Méreau, R.; De Winter, J.; Gerbaux, P.; Detrembleur, C.; Tassaing, T.; Jérôme, C. Organocatalytic Coupling of CO<sub>2</sub> with Oxetane. *ChemSusChem* **2017**, *10*, 1128–1138. [[CrossRef](#)]
136. Huang, J.; de Winter, J.; Dove, A.P.; Coulembier, O. Metal-free synthesis of poly(trimethylene carbonate) by efficient valorization of carbon dioxide. *Green Chem.* **2019**, *21*, 472–477. [[CrossRef](#)]
137. Zhou, H.; Wang, G.-X.; Lu, X.-B. CO<sub>2</sub> Adducts of  $\alpha$ -Carbon Alkylated *N*-Heterocyclic Olefins: Highly Active Organocatalysts for CO<sub>2</sub> Chemical Transformation. *Asian. J. Org. Chem.* **2017**, *6*, 1264–1269. [[CrossRef](#)]
138. Qiu, J.; Zhao, Y.; Li, Z.; Wang, H.; Fan, M.; Wang, J. Efficient Ionic-Liquid-Promoted Chemical Fixation of CO<sub>2</sub> into  $\alpha$ -Alkylidene Cyclic Carbonates. *ChemSusChem* **2017**, *10*, 1120–1127. [[CrossRef](#)]
139. Panwar, V.; Jain, S.L. Zinc grafted to magnetic nanostarch for cyclic carbonate synthesis from propargylic alcohols and CO<sub>2</sub> at room temperature. *J. CO<sub>2</sub> Util.* **2018**, *24*, 306–314. [[CrossRef](#)]

140. Grignard, B.; Ngassamtounzoua, C.; Gennen, S.; Gilbert, B.; Méreau, R.; Jerome, C.; Tassaing, T.; Detrembleur, C. Boosting the Catalytic Performance of Organic Salts for the Fast and Selective Synthesis of  $\alpha$ -Alkylidene Cyclic Carbonates from Carbon Dioxide and Propargylic Alcohols. *ChemCatChem* **2018**, *10*, 2584–2592. [[CrossRef](#)]
141. Jung, M.E.; Piizzi, G. *gem*-Disubstituent Effect: Theoretical Basis and Synthetic Applications. *Chem. Rev.* **2005**, *105*, 1735–1766. [[CrossRef](#)]
142. Boyaval, A.; Méreau, R.; Grignard, B.; Detrembleur, C.; Jerome, C.; Tassaing, T. Organocatalytic Coupling of CO<sub>2</sub> with a Propargylic Alcohol: A Comprehensive Mechanistic Study. *ChemSusChem* **2017**, *10*, 1241–1248. [[CrossRef](#)]
143. Della Ca', N.; Gabriele, B.; Ruffolo, G.; Veltri, L.; Zanetta, T.; Costa, M. Effective Guanidine-Catalyzed Synthesis of Carbonate and Carbamate Derivatives from Propargyl Alcohols in Supercritical Carbon Dioxide. *Adv. Synth. Catal.* **2011**, *353*, 133–146. [[CrossRef](#)]
144. Méreau, R.; Grignard, B.; Boyaval, A.; Detrembleur, C.; Jerome, C.; Tassaing, T. Tetrabutylammonium Salts: Cheap Catalysts for the Facile and Selective Synthesis of  $\alpha$ -Alkylidene Cyclic Carbonates from Carbon Dioxide and Alkynols. *ChemCatChem* **2018**, *10*, 956–960. [[CrossRef](#)]
145. Sun, S.; Wang, B.; Gu, N.; Yu, J.-T.; Cheng, J. Palladium-Catalyzed Arylcarboxylation of Propargylic Alcohols with CO<sub>2</sub> and Aryl Halides: Access to Functionalized  $\alpha$ -Alkylidene Cyclic Carbonates. *Org. Lett.* **2017**, *19*, 1088–1091. [[CrossRef](#)] [[PubMed](#)]
146. Gregory, G.L.; Ulmann, M.; Buchard, A. Synthesis of 6-membered cyclic carbonates from 1,3-diols and low CO<sub>2</sub> pressure: A novel mild strategy to replace phosgene reagents. *RSC Adv.* **2015**, *5*, 39404–39408. [[CrossRef](#)]
147. McGuire, T.M.; López-Vidal, E.M.; Gregory, G.L.; Buchard, A. Synthesis of 5- to 8-membered cyclic carbonates from diols and CO<sub>2</sub>: A onestep, atmospheric pressure and ambient temperature procedure. *J. CO<sub>2</sub> Util.* **2018**, *27*, 283–288. [[CrossRef](#)]
148. Pati, D.; Chen, Z.; Feng, X.; Hadjichristidis, N.; Gnanou, Y. Synthesis of polyglycocarbonates through polycondensation of glucopyranosides with CO<sub>2</sub>. *Polym. Chem.* **2017**, *8*, 2640–2646. [[CrossRef](#)]
149. Pati, D.; Feng, X.; Hadjichristidis, N.; Gnanou, Y. CO<sub>2</sub> as versatile carbonation agent of glycosides: Synthesis of 5- and 6-membered cyclic glycocarbonates and investigation of their ring-opening. *J. CO<sub>2</sub> Util.* **2018**, *24*, 564–571. [[CrossRef](#)]
150. Xie, Y.; Lu, C.; Zhao, B.; Wang, Q.; Yao, Y. Cycloaddition of Aziridine with CO<sub>2</sub>/CS<sub>2</sub> Catalyzed by Amidato Divalent Lanthanide Complexes. *J. Org. Chem.* **2019**, *84*, 1951–1958. [[CrossRef](#)]
151. Phung, C.; Tantillo, D.J.; Hein, J.E.; Pinhas, A.R. The mechanism of the reaction between an aziridine and carbon dioxide with no added catalyst. *J. Phys. Org. Chem.* **2018**, *31*, e3735. [[CrossRef](#)]
152. Zhang, Z.; Ye, J.-H.; Wu, D.-S.; Zhou, Y.-Q.; Yu, D.-G. Synthesis of Oxazolidin-2-ones from Unsaturated Amines with CO<sub>2</sub> by Using Homogeneous Catalysis. *Chem. Asian. J.* **2018**, *13*, 2292–2306. [[CrossRef](#)]
153. Yuan, R.; Wei, B.; Fu, G. How the Coordinated Structures of Ag(I) Catalysts Affect the Outcomes of Carbon Dioxide Incorporation into Propargylic Amine: A DFT Study. *J. Org. Chem.* **2017**, *82*, 3639–3647. [[CrossRef](#)]
154. Brunel, P.; Monot, J.; Kefalidis, C.E.; Maron, L.; Martin-Vaca, B.; Bourissou, D. Valorization of CO<sub>2</sub>: Preparation of 2-Oxazolidinones by Metal-Ligand Cooperative Catalysis with SCS Indenediide Pd Complexes. *ACS Catal.* **2017**, *7*, 2652–2660. [[CrossRef](#)]
155. Zhao, D.; Liu, X.-H.; Zhu, C.; Kang, Y.-S.; Wang, P.; Shi, Z.; Lu, Y.; Sun, W.-Y. Efficient and Reusable Metal–Organic Framework Catalysts for Carboxylative Cyclization of Propargylamines with Carbon Dioxide. *ChemCatChem* **2017**, *9*, 4598–4606. [[CrossRef](#)]
156. Yu, B.; Kim, D.; Kim, S.; Hong, S.H. Cyanuric Acid-Based Organocatalyst for Utilization of Carbon Dioxide at Atmospheric Pressure. *ChemSusChem* **2017**, *10*, 1080–1084. [[CrossRef](#)] [[PubMed](#)]
157. Liu, X.; Wang, M.-Y.; Wang, S.-Y.; Wang, Q.; He, L.-N. In Situ Generated Zinc(II) Catalyst for Incorporation of CO<sub>2</sub> into 2-Oxazolidinones with Propargylic Amines at Atmospheric Pressure. *ChemSusChem* **2017**, *10*, 1210–1216. [[CrossRef](#)] [[PubMed](#)]
158. Zhao, Y.; Qiu, J.; Li, Z.; Wang, H.; Fan, M.; Wang, J. An Experimental and Theoretical Study on the Unexpected Catalytic Activity of Triethanolamine for the Carboxylative Cyclization of Propargylic Amines with CO<sub>2</sub>. *ChemSusChem* **2017**, *10*, 2001–2007. [[CrossRef](#)] [[PubMed](#)]
159. Fujii, A.; Choi, J.-C.; Fujita, K.-i. Quaternary ammonium salt-catalyzed carboxylative cyclization of propargylic amines with CO<sub>2</sub>. *Tetrahedron Lett.* **2017**, *58*, 4483–4486. [[CrossRef](#)]

160. Fujii, A.; Matsuo, H.; Choi, J.-C.; Fujitani, T.; Fujita, K.-i. Efficient synthesis of 2-oxazolidinones and quinazoline-2,4(1*H*,3*H*)-diones from CO<sub>2</sub> catalyzed by tetrabutylammonium fluoride. *Tetrahedron* **2018**, *74*, 2914–2920. [[CrossRef](#)]
161. Inagaki, F.; Maeda, K.; Nakazawa, K.; Mukai, C. Construction of the Oxazolidinone Framework from Propargylamine and CO<sub>2</sub> in Air at Ambient Temperature: Catalytic Effect of a Gold Complex Featuring an L<sub>2</sub>/Z-Type Ligand. *Eur. J. Org. Chem.* **2018**, 2972–2976. [[CrossRef](#)]
162. Sadeghzadeh, S.M.; Zhiani, R.; Emrani, S. Ni@Pd nanoparticles supported on ionic liquid-functionalized KCC-1 as robust and recyclable nanocatalysts for cycloaddition of propargylic amines and CO<sub>2</sub>. *Appl. Organometal. Chem.* **2018**, *32*, e3941. [[CrossRef](#)]
163. Yoshida, M.; Mizuguchi, T.; Shishido, K. Synthesis of Oxazolidinones by Efficient Fixation of Atmospheric CO<sub>2</sub> with Propargylic Amines by using a Silver/1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU) Dual-Catalyst System. *Chem. Eur. J.* **2012**, *18*, 15578–15581. [[CrossRef](#)]
164. Inagaki, F.; Okada, Y.; Matsumoto, C.; Yamada, M.; Nakazawa, K.; Mukai, C. Energyless CO<sub>2</sub> Absorption, Generation, and Fixation Using Atmospheric CO<sub>2</sub>. *Chem. Pharm. Bull.* **2016**, *64*, 8–13. [[CrossRef](#)] [[PubMed](#)]
165. Gao, X.-T.; Gan, C.-C.; Liu, S.-I.; Zhou, F.; Wu, H.-H.; Zhou, J. Utilization of CO<sub>2</sub> as a C1 Building Block in a Tandem Asymmetric A<sup>3</sup> Coupling-Carboxylative Cyclization Sequence to 2-Oxazolidinones. *ACS Catal.* **2017**, *7*, 8588–8593. [[CrossRef](#)]
166. Kim, N.-K.; Sogawa, H.; Yamamoto, K.; Hayashi, Y.; Kawauchi, S.; Takata, T. DBU-mediated Highly Efficient CO<sub>2</sub>-fixation to Propargylamines and Polypropargylamine. *Chem. Lett.* **2018**, *47*, 1063–1066. [[CrossRef](#)]
167. Kim, N.-K.; Sogawa, H.; Felicia, M.D.; Takata, T. DBU-catalyzed CO<sub>2</sub> fixation in polypropargylamines under solvent-free conditions. *Polym. J.* **2019**, *51*, 351–357. [[CrossRef](#)]
168. Sugiyama, N.; Ohseki, M.; Kobayashi, R.; Sekine, K.; Saito, K.; Yamada, T. Silver-catalyzed Three-component Reaction of Propargylic Amines, Carbon Dioxide, and *N*-Bromosuccinimide for Stereoselective Preparation of (*E*)-Bromovinylloxazolidinones. *Chem. Lett.* **2017**, *46*, 1323–1326. [[CrossRef](#)]
169. Yin, Z.-B.; Ye, J.-H.; Zhou, W.-J.; Zhang, Y.-H.; Ding, L.; Gui, Y.-Y.; Yan, S.-S.; Li, J.; Yu, D.-G. Oxy-Difluoroalkylation of Allylamines with CO<sub>2</sub> via Visible-Light Photoredox Catalysis. *Org. Lett.* **2018**, *20*, 190–193. [[CrossRef](#)]
170. Sun, L.; Ye, J.-H.; Zhou, W.-J.; Zeng, X.; Yu, D.-G. Oxy-Alkylation of Allylamines with Unactivated Alkyl Bromides and CO<sub>2</sub> via Visible-Light-Driven Palladium Catalysis. *Org. Lett.* **2018**, *20*, 3049–3052. [[CrossRef](#)]
171. Wang, M.-Y.; Cao, Y.; Liu, X.; Wang, N.; He, L.-N.; Li, S.-H. Photoinduced radical-initiated carboxylative cyclization of allyl amines with carbon dioxide. *Green Chem.* **2017**, *19*, 1240–1244. [[CrossRef](#)]
172. Seo, U.R.; Chung, Y.K. Potassium phosphate-catalyzed one-pot synthesis of 3-aryl-2-oxazolidinones from epoxides, amines, and atmospheric carbon dioxide. *Green Chem.* **2017**, *19*, 803–808. [[CrossRef](#)]
173. Lv, M.; Wang, P.; Yuan, D.; Yao, Y. Conversion of Carbon Dioxide into Oxazolidinones Mediated by Quaternary Ammonium Salts and DBU. *ChemCatChem* **2017**, *9*, 4451–4455. [[CrossRef](#)]
174. Chen, F.; Li, M.; Wang, J.; Dai, B.; Liu, N. Fe(II) complexes: Reservoirs for Lewis acids and carbenes and their utility in the conversion of CO<sub>2</sub> to oxazolidinones. *J. CO<sub>2</sub> Util.* **2018**, *28*, 181–188. [[CrossRef](#)]
175. Sadeghzadeh, S.M.; Zhiani, R.; Moradi, M. KCC-1 Supported Cu(II)-β-Cyclodextrin Complex as a Reusable Catalyst for the Synthesis of 3-Aryl-2-oxazolidinones from Carbon Dioxide, Epoxide, Anilines. *ChemistrySelect* **2018**, *3*, 3516–3522. [[CrossRef](#)]
176. Mancuso, R.; Raut, D.S.; Della Ca', N.; Fini, F.; Carfagna, C.; Gabriele, B. Catalytic oxidative carbonylation of amino moieties to ureas, oxamides, 2-oxazolidinones, and benzoxazolones. *ChemSusChem* **2015**, *8*, 2204–2211. [[CrossRef](#)] [[PubMed](#)]
177. Li, X.-D.; Song, Q.-W.; Lang, X.-D.; Chang, Y.; He, L.-N. AgI/TMG-Promoted Cascade Reaction of Propargyl Alcohols, Carbon Dioxide, and 2-Aminoethanols to 2-Oxazolidinones. *ChemPhysChem* **2017**, *18*, 3182–3188. [[CrossRef](#)] [[PubMed](#)]
178. Li, X.-D.; Cao, Y.; Ma, R.; He, L.-N. Thermodynamically favorable protocol for the synthesis of 2-oxazolidinones via Cu(I)-catalyzed three-component reaction of propargylic alcohols, CO<sub>2</sub> and 2-aminoethanols. *J. CO<sub>2</sub> Util.* **2018**, *25*, 338–345. [[CrossRef](#)]
179. Xia, S.; Song, Y.; Li, X.; Li, H.; He, L.-N. Ionic Liquid-Promoted Three-Component Domino Reaction of Propargyl Alcohols, Carbon Dioxide and 2-Aminoethanols: A Thermodynamically Favorable Synthesis of 2-Oxazolidinones. *Molecules* **2018**, *23*, 3033. [[CrossRef](#)] [[PubMed](#)]

180. Niemi, T.; Fernández, I.; Steadman, B.; Mannisto, J.K.; Repo, T. Carbon dioxide-based facile synthesis of cyclic carbamates from amino alcohols. *Chem. Commun.* **2018**, *54*, 3166–3169. [[CrossRef](#)] [[PubMed](#)]
181. Mei, C.; Zhao, Y.; Chen, Q.; Cao, C.; Pang, G.; Shi, Y. Synthesis of Oxazolidinones and Derivatives through Three-Component Fixation of Carbon Dioxide. *ChemCatChem* **2018**, *10*, 3057–3068. [[CrossRef](#)]
182. Yousefi, R.; Struble, T.J.; Payne, J.L.; Vishe, M.; Schley, N.D.; Johnston, J.N. Catalytic, Enantioselective Synthesis of Cyclic Carbamates from Dialkyl Amines by CO<sub>2</sub>-Capture: Discovery, Development, and Mechanism. *J. Am. Chem. Soc.* **2019**, *141*, 618–625. [[CrossRef](#)]
183. Nikpassand, M.; Fekri, L.Z.; Pourahmad, A. One-pot Synthesis of new azo-linked 4*H*-benzo[*d*][1,3]oxazine-2,4-diones from carbon dioxide using CuO@RHA/MCM-41 nanocomposite in green media. *J. CO<sub>2</sub> Util.* **2018**, *27*, 320–325. [[CrossRef](#)]
184. Guo, C.-X.; Zhang, W.-Z.; Zhang, N.; Lu, X.-B. 1,3-Dipolar Cycloaddition of Nitrile Imine with Carbon Dioxide: Access to 1,3,4-Oxadiazole-2(3*H*)-ones. *J. Org. Chem.* **2017**, *82*, 7637–7642. [[CrossRef](#)]
185. Murillo, F.; Barroso, J.; de los Santos, M.G.; Ávila, G.; Pan, S.; Fernández-Herrera, M.A.; Merino, G. Revisiting the Formation Mechanism of 1,3,4-Oxadiazole-2(3*H*)-ones from Hydrazonyl Chloride and Carbon Dioxide. *J. Org. Chem.* **2018**, *83*, 13045–13050. [[CrossRef](#)] [[PubMed](#)]
186. Yang, N.; Zhang, H.; Yuan, G. KI-catalyzed reactions of aryl hydrazines with  $\alpha$ -oxocarboxylic acids in the presence of CO<sub>2</sub>: Access to 1,3,4-oxadiazol-2(3*H*)-ones. *Org. Chem. Front.* **2019**, *6*, 532–536. [[CrossRef](#)]
187. Brahmaya, M.; Dai, S.A.; Suen, S.-Y. Facile synthesis of 2-benzimidazolones via carbonylation of *o*-phenylenediamines with CO<sub>2</sub>. *J. CO<sub>2</sub> Util.* **2017**, *22*, 135–142. [[CrossRef](#)]
188. Wang, H.; Xin, Z.; Li, Y. Synthesis of Ureas from CO<sub>2</sub>. *Top. Curr. Chem.* **2017**, *375*, 49. [[CrossRef](#)]
189. He, X.; Li, X.-Y.; Song, Y.; Xia, S.-M.; Lang, X.-D.; He, L.-N. Synthesis of Urea Derivatives using Carbon Dioxide as Carbonylation Reagent in Ionic Liquids. *Curr. Organocat.* **2017**, *4*, 112–121. [[CrossRef](#)]
190. Hulla, M.; Chamam, S.M.A.; Laurenczy, G.; Das, S.; Dyson, P.J. Delineating the Mechanism of Ionic Liquids in the Synthesis of Quinazoline-2,4(1*H*,3*H*)-dione from 2-Aminobenzonitrile and CO<sub>2</sub>. *Angew. Chem. Int. Ed.* **2017**, *56*, 10559–10563. [[CrossRef](#)] [[PubMed](#)]
191. Sabet-Sarvestani, H.; Eshghi, H.; Izadyar, M. A theoretical study on the efficiency and role of guanidines-based organic superbases on carbon dioxide utilization in quinazoline-2,4(1*H*, 3*H*)-diones synthesis. *Struct. Chem.* **2017**, *28*, 675–686. [[CrossRef](#)]
192. Yan, C.; Ren, Y.; Jia, J.-F.; Wu, H.-S. Mechanism of the chemical fixation of carbon dioxide with 2-aminobenzonitrile catalyzed by cesium carbonate: A computational study. *Mol. Catal.* **2017**, *432*, 172–186. [[CrossRef](#)]
193. Li, C.; Lu, X.; Yang, Y.; Yang, S.; Zhang, L. Ionic liquid promoted synthesis of heterocycle-fused pyrimidine-2,4-(1*H*,3*H*)-diones utilising CO<sub>2</sub>. *Tetrahedron Lett.* **2018**, *59*, 2463–2466. [[CrossRef](#)]
194. Shi, G.; Chen, K.; Wang, Y.; Li, H.; Wang, C. Highly Efficient Synthesis of Quinazoline-2,4(1*H*,3*H*)-diones from CO<sub>2</sub> by Hydroxyl Functionalized Aprotic Ionic Liquids. *ACS Sustain. Chem. Eng.* **2018**, *6*, 5760–5765. [[CrossRef](#)]
195. Hu, J.; Chen, S.; Guo, Y.; Li, L.; Deng, T. Basic Salt-Lake Brine: An Efficient Catalyst for the Transformation of CO<sub>2</sub> into Quinazoline-2,4(1*H*,3*H*)-diones. *ChemSusChem* **2018**, *11*, 4219–4225. [[CrossRef](#)] [[PubMed](#)]
196. Mampuy, P.; Neumann, H.; Sergeev, S.; Orru, R.V.A.; Jiao, H.; Spannenberg, A.; Maes, B.U.W.; Beller, M. Combining Isocyanides with Carbon Dioxide in Palladium-Catalyzed Heterocycle Synthesis: *N*3-Substituted Quinazoline-2,4-(1*H*,3*H*)-diones via a Three-Component Reaction. *ACS Catal.* **2017**, *7*, 5549–5556. [[CrossRef](#)]
197. Xu, P.; Wang, F.; Wei, T.-Q.; Yin, L.; Wang, S.-Y.; Ji, S.J. Palladium-Catalyzed Incorporation of Two C1 Building Blocks: The Reaction of Atmospheric CO<sub>2</sub> and Isocyanides with 2-Iodoanilines Leading to the Synthesis of Quinazoline-2,4(1*H*,3*H*)-diones. *Org. Lett.* **2017**, *19*, 4484–4487. [[CrossRef](#)] [[PubMed](#)]
198. Zhang, W.-Z.; Li, H.; Zeng, Y.; Tao, X.; Lu, X. Palladium-Catalyzed Cyclization Reaction of *o*-Haloanilines, CO<sub>2</sub> and Isocyanides: Access to Quinazoline-2,4(1*H*,3*H*)-diones. *Chin. J. Chem.* **2018**, *36*, 112–118. [[CrossRef](#)]
199. Jin, S.-J.; Khan, Y.; Maeng, J.-H.; Kim, Y.J.; Hwang, J.; Cheong, M.; Lee, J.S.; Kim, H.S. Efficient catalytic systems for the carboxylation of diamines to cyclic ureas using ethylene urea as a promoter. *Appl. Catal. B Environ.* **2017**, *209*, 139–145. [[CrossRef](#)]
200. Hwang, J.; Han, D.; Oh, J.J.; Cheong, M.; Koo, H.-J.; Lee, J.S.; Kim, H.S. Efficient Non-Catalytic Carboxylation of Diamines to Cyclic Ureas Using 2-Pyrrolidone as a Solvent and a Promoter. *Adv. Synth. Catal.* **2019**, *361*, 297–306. [[CrossRef](#)]

201. Del Vecchio, A.; Caillé, F.; Chevalier, A.; Loreau, O.; Horkka, K.; Halldin, C.; Schou, M.; Camus, N.; Kessler, P.; Kuhnast, B.; et al. Late-Stage Isotopic Carbon Labeling of Pharmaceutically Relevant Cyclic Ureas Directly from CO<sub>2</sub>. *Angew. Chem. Int. Ed.* **2018**, *57*, 9744–9748. [[CrossRef](#)] [[PubMed](#)]
202. Streng, E.S.; Lee, D.S.; George, M.W.; Poliakoff, M. Continuous N-alkylation reactions of amino alcohols using  $\gamma$ -Al<sub>2</sub>O<sub>3</sub> and supercritical CO<sub>2</sub>: Unexpected formation of cyclic ureas and urethanes by reaction with CO<sub>2</sub>. *Beilstein J. Org. Chem.* **2017**, *13*, 329–337. [[CrossRef](#)]
203. Marchegiani, M.; Nodari, M.; Tansini, F.; Massera, C.; Mancuso, R.; Gabriele, B.; Costa, M.; Della Cà, N. Urea derivatives from carbon dioxide and amines by guanidine catalysis: Easy access to imidazolidin-2-ones under solvent-free conditions. *J. CO<sub>2</sub> Util.* **2017**, *21*, 553–561. [[CrossRef](#)]
204. Chun, S.; Yang, S.; Chung, Y.K. Synthesis of benzothiazoles from 2-aminobenzenethiols in the presence of a reusable polythiazolium precatalyst under atmospheric pressure of carbon dioxide. *Tetrahedron* **2017**, *73*, 3438–3442. [[CrossRef](#)]
205. Wang, B.; Sun, S.; Yu, J.-T.; Jiang, Y.; Cheng, J. Palladium-Catalyzed Multicomponent Reactions of o-Alkynylanilines, Aryl Iodides, and CO<sub>2</sub> toward 3,3-Diaryl 2,4-Quinolinediones. *Org. Lett.* **2017**, *19*, 4319–4322. [[CrossRef](#)] [[PubMed](#)]
206. Li, W.; Li, C.; Lyu, Y. Lactamization of sp<sup>2</sup> C–H bonds with CO<sub>2</sub> under transition-metal-free and redox-neutral conditions: A computational mechanistic study. *Org. Chem. Front.* **2018**, *5*, 2189–2201. [[CrossRef](#)]
207. Wang, S.; Xi, C. Recent advances in nucleophile-triggered CO<sub>2</sub>-incorporated cyclization leading to heterocycles. *Chem. Soc. Rev.* **2019**, *48*, 382–404. [[CrossRef](#)] [[PubMed](#)]
208. Niemi, T.; Repo, T. Antibiotics from Carbon Dioxide: Sustainable Pathways to Pharmaceutically Relevant Cyclic Carbamates. *Eur. J. Org. Chem.* **2019**, 1180–1188. [[CrossRef](#)]
209. Vaitla, J.; Guttormsen, Y.; Mannisto, J.K.; Nova, A.; Repo, T.; Bayer, A.; Hopmann, K.H. Enantioselective Incorporation of CO<sub>2</sub>: Status and Potential. *ACS Catal.* **2017**, *7*, 7231–7244. [[CrossRef](#)]



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).