



# UNIVERSITÀ DI PARMA

## ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Novel insights on pink discoloration in cheese: The case of Pecorino Toscano

This is the peer reviewed version of the following article:

*Original*

Novel insights on pink discoloration in cheese: The case of Pecorino Toscano / Martelli, F.; Bancalari, E.; Neviani, E.; Bottari, B.. - In: INTERNATIONAL DAIRY JOURNAL. - ISSN 0958-6946. - 111:(2020), p. 104829. [10.1016/j.idairyj.2020.104829]

*Availability:*

This version is available at: 11381/2879781 since: 2022-01-21T14:28:31Z

*Publisher:*

Elsevier Ltd

*Published*

DOI:10.1016/j.idairyj.2020.104829

*Terms of use:*

Anyone can freely access the full text of works made available as "Open Access". Works made available

*Publisher copyright*

note finali coverpage

(Article begins on next page)

02 May 2026

# Novel insights on pink discoloration in cheese: the case of Pecorino Toscano

Francesco Martelli<sup>1</sup>, Elena Bancalari<sup>1\*</sup>, Erasmo Neviani<sup>1</sup>, Benedetta Bottari<sup>1</sup>

<sup>1</sup> Department of Food and Drug, University of Parma, Parco Area delle Scienze 47-49/A, 43124, Parma, Italy

\*Corresponding: [elena.bancalari@unipr.it](mailto:elena.bancalari@unipr.it)

## Abstract

Pink discoloration in cheese has been the subject of wide research, but the basis for this phenomenon still remain elusive. This defect impacts a wide range of ripened cheeses, resulting in the rejection of cheese and a consequent economic loss for dairy industries. As multiple causes for pink discoloration have been reported for different cheeses in the literature, the aim of this research was to investigate the cause of a pink discoloration found on Pecorino Toscano cheese rind. The results of microbiological analysis revealed the presence of high microbial counts associated to the rind and the nearest inner part. Strains isolated from the colored part were mainly identified as *Serratia liquefaciens*, suggesting that an environmental contamination of the cheese rind by this species could be the cause of the observed defect and expanding the knowledge related to pink discoloration in cheeses.

**Keywords:** Cheese defects, *Serratia liquefaciens*, environmental contaminations, Pecorino Toscano cheese

## 1. Introduction

The occurrence of defect in cheeses is of great concern for the dairy industry as it could lead to products recall and thus representing a significant economic loss for dairy companies (Carminati et al., 2019). Defects may have different origin and may occur in diverse step of cheese making process. They can be related to variations in milk quality, milk treatment, hygiene practices and cheese processing parameters (manufacture technology and/or ripening). Defects can also be linked to differences in the starter cultures activity and acidity profiles, as well as to the nature and number of the non-starter microorganisms which come from raw milk or other environmental sources (O'Sullivan et al., 2013).

To date, several studies have been focused on the understanding of cheese discoloration' causes. In particular, the reason for the formation of pink defect in cheeses without colorant addition, in some cases still remains unknown (Quigley et al., 2016). To date, it has been mainly attributed to the metabolism of certain strains of thermophilic lactobacilli, propionic acid bacteria, *Thermus* spp., or *Glutamicibacter arilaitensis* as well as the activity of residual microbial enzymes or to Maillard reactions, particularly related to the presence of unfermented galactose (Cleary et al., 2018; Daly et al., 2012).

During summer 2018, a Tuscan dairy producing Pecorino Toscano, a PDO pasteurized sheep's milk medium ripened Italian cheese, observed an unusual pink discoloration on the rind of several cheese wheels (Personal communication; Figure 1). With the aim of determining the cause of discoloration, microbiological analysis followed by molecular identification of the species recovered in the pink area, were carried out.

## 2. Materials and Methods

### 2.1 Sampling

In order to identify the bacteria species potentially responsible for the alteration, the crust and the closer inner part of six representative cheeses were sampled and homogenized 1:10 with sterile Ringer's solution,

45 by means of a blender (STOMACHER, 400 circulator, UK) (Oxoid, Basingstoke, UK) for 2.30 min at 230  
46 rpm. Each cheese was analyzed separately. The samples dilutions were then plated in triplicated on different  
47 growth media. Pseudomonas CFC agar (PS) (Oxoid, UK) was used in order to count *Pseudomonadaceae*,  
48 incubating plates at 25°C for 48 hours. Plate Count Agar (PCA) (Oxoid, UK) was used for the determination  
49 of total microbial count by incubating the plates at 30°C for 24hours. Yeast and molds were counted on  
50 Yeast extract dextrose Chloramphenicol Agar (YEDC). For the determination of aerobic spore-forming  
51 bacteria, the homogenized samples were heated to 80°C for 10 min and then cooled before analysis and then  
52 plated on Tryptone Soy Agar (TSA, Merck KGaA, Darmstadt, Germany), followed by incubation for 24 h at  
53 30 °C under aerobic condition (Abdelmassih et al., 2011).

## 54 **2.2 Microbial isolation from pink areas**

55 Fourteen colonies with pinkish or reddish color present only on PS plates were selected and isolated through  
56 three purification steps on the same medium. The purity of the strains was checked by verifying the  
57 morphology by means of optical microscope Olympus BX51 (Olympus, Waltham, United States) then  
58 stored at – 80° C in Tryptic Soy Broth (TSB) (Oxoid, Basingstoke, UK) added with 20% (v/v) glycerol.

## 59 **2.3. Genotypic identification of isolated strains**

60 The isolated strains were cultured twice at 30 °C for 18 h in TSB (Oxoid) in aerobic conditions and then the  
61 DNA was extracted and purified by using DNeasy and Blood Tissue (Qiagen, Germany) and checked on  
62 agarose gel with TAE 1X running buffer (1% w/v). Then, the 16S rRNA gene was amplified by PCR  
63 (Takahashi et al., 2014).

64 Primers based on conserved regions of the 16S rRNA gene, 16S 46Fw: CAG GCC TAA CAC ATG CAA  
65 GTC and 16S 536Rv: GGG CGG WGT CAA GGC (Marchesi et al., 1998), were used for direct PCR  
66 amplification of a 1300-bp portion of the 16S rRNA gene. In brief, samples underwent an initial denaturation  
67 of 2 min at 94°C and 30sec at 94°C, then 30 cycles of 30-sec at 55°C, 30 sec at 72°C, followed by 10 min at  
68 72°C. Reaction products were separated by electrophoresis in 1.0% (w/v) agarose gels and the amplification  
69 products were sequenced (Macrogen Europe Inc.). Sequences comparison was performed against NCBI  
70 database using BLASTN -(Basic Local Alignment Tool, BLAST, p://www. ncbi.nlm.nih.gov/blast.cgi).  
71 Criteria for identification at the species level were defined as a 16S rDNA sequence similarity of  $\geq 98\%$   
72 (Drancourt et al., 2000) with that of the type strains sequences in GenBank.

## 73 **3 Results and discussion**

### 74 **3.1 Microbial counts**

75 In the present study, the total microbial counts was of 7.82 Log CFU/g (mean value of the samples) (fig.2),  
76 which corresponds to the results found by Proroga and colleagues (2009) on Pecorino. Yeast and moulds  
77 count was 3 Log CFU/g, which is typical for ripened cheeses such as Pecorino Toscano PDO (Todaro et al.,  
78 2011). The presence of spore forming microorganisms was lower than 3 Log CFU/g (Fig. 2), confirming  
79 what has been observed in the literature for similar cheeses (Palmas et al., 1999). Microbial counts on  
80 Pseudomonas agar were 1 log unit higher than the average data available in literature, reporting counts of no  
81 more than 5 Log CFU/g in the same medium (Todaro et al., 2011). In all the analyzed cheese samples, pink  
82 colonies were present on PS plates.

### 84 **3.2 Genotypic identification of isolated strains**

85 Fourteen isolated strains were identified at species level as follows: 10 strains belonging to *Serratia*  
86 *liquefaciens*, 2 *Enterobacter cloacae*, 2 *Enterobacter hormaechei* (Table 1).

87 The presence of *Enterobacteriaceae* is not surprising as, despite they are considered as indicative of a poor  
88 microbiological quality of cheese, they are also recognized as part of the natural microbiota of many dairy  
89 products (Chaves-Lopez et al., 2006).  
90 *S. liquefaciens* is a psychrotrophic and highly motile organism commonly found in water, soil, plants and  
91 more specifically in dairies environment and raw milk (Ikumapayi et al., 2015; Machado et al., 2015) *S.*  
92 *liquefaciens* is a pathogenic microorganism able to cause infections in humans, especially in  
93 immunocompromised hosts (Momose et al.,2018; Caneschi et al., 2019). A possible role of this species in  
94 discoloration defects had been previously observed, but mainly related to the presence of its hydrolytic heat  
95 resistant enzymes, rather than the viable cells (Decimo et al., 2014). In the present study, viable *S.*  
96 *liquefaciens* cells were isolated from the rind and the nearest inner part, where the pink discoloration was  
97 present. Furthermore, as no discoloration was found in the deep inner part of the cheeses, despite the species  
98 is able to grow both under aerobic and anaerobic conditions (Bergey et al.,2000), an environmental  
99 contamination of the rind was hypothesized. Pecorino Toscano is ripened between 5 and 15°C with a relative  
100 humidity ranging between 75 and 95%. Such conditions could encourage the growth of *S.liquefaciens* on  
101 cheese rind and the production of a red pigment in the later stages of bacterial growth. The production of red  
102 pigments such as prodigiosin has in fact been reported to be regulated by a quorum sensing mechanism, by  
103 which, the increasing number of cells is the signal to start the production of the pigment (Thomson et al.,  
104 2000).

105  
106

#### 107 **4. Conclusion**

108

109 The microbial analysis of each sample from the six Pecorino Toscano cheese revealed the presence of *S.*  
110 *liquefaciens*, suggesting that an environmental contamination of the cheese rind could be the cause of the  
111 observed defect. In particular, it seems of central importance, when studying this kind of defect, the  
112 evaluation of brine and ripening chambers hygiene as well as the overall hygiene of the working areas.  
113 Another important aspect is the control of rind humidity, which may promote the development of alternative  
114 microorganisms coming from the environment. The result of our study could expand the knowledge related  
115 to pink discoloration in cheese, which is still ambiguous in the literature.

116

117

118

#### 119 **References**

120

Abdelmassih, M., Planchon, V., Anceau, C., & Mahillon, J. (2011). Development and validation of stable

121

reference materials for food microbiology using *Bacillus cereus* and *Clostridium perfringens* spores:

122

Spores as stable reference material. *Journal of Applied Microbiology*, 110(6), 1524–1530.

123

<https://doi.org/10.1111/j.1365-2672.2011.05007.x>

124

Bergey, D. H.Holt, John G., eds. *Bergey's Manual Of Determinative Bacteriology*. Philadelphia : Lippincott

125

Williams & Wilkins, 2000. Print. (9th Edition,p.217,Table 5.2)

126

127 Carminati, D., Bonvini, B., Rossetti, L., Zago, M., Tidona, F., & Giraffa, G. (2019). Investigation on the  
128 presence of blue pigment-producing *Pseudomonas* strains along a production line of fresh  
129 mozzarella cheese. *Food Control*, *100*, 321–328. <https://doi.org/10.1016/j.foodcont.2019.02.009>

130 Chaves-Lopez, C., De Angelis, M., Martuscelli, M., Serio, A., Paparella, A., & Suzzi, G. (2006).  
131 Characterization of the Enterobacteriaceae isolated from an artisanal Italian ewe's cheese  
132 (Pecorino Abruzzese). *Journal of Applied Microbiology*, *101*(2), 353–360.  
133 <https://doi.org/10.1111/j.1365-2672.2006.02941.x>

134 Cleary, J. L., Kolachina, S., Wolfe, B. E., & Sanchez, L. M. (2018). Coproporphyrin III Produced by the  
135 Bacterium *Glutamicibacter arilaitensis* Binds Zinc and Is Upregulated by Fungi in Cheese Rinds.  
136 *MSystems*, *3*(4), e00036-18, /msystems/3/4/msys.00036-18.atom.  
137 <https://doi.org/10.1128/mSystems.00036-18>

138 Daly, D. F. M., McSweeney, P. L. H., & Sheehan, J. J. (2012). Pink discolouration defect in commercial  
139 cheese: a review. *Dairy Science & Technology*, *92*(5), 439–453. [https://doi.org/10.1007/s13594-](https://doi.org/10.1007/s13594-012-0079-0)  
140 [012-0079-0](https://doi.org/10.1007/s13594-012-0079-0)

141 Decimo, M., Morandi, S., Silvetti, T., & Brasca, M. (2014). Characterization of Gram-Negative Psychrotrophic  
142 Bacteria isolated from Italian Bulk Tank Milk: Gram-negative psychrotrophs in milk.... *Journal of*  
143 *Food Science*, *79*(10), M2081–M2090. <https://doi.org/10.1111/1750-3841.12645>

144 Drancourt, M., Bollet, C., Carlioz, A., Martelin, R., Gayral, J.-P., & Raoult, D. (2000). 16S Ribosomal DNA  
145 Sequence Analysis of a Large Collection of Environmental and Clinical Unidentifiable Bacterial  
146 Isolates. *Journal of Clinical Microbiology*, *38*(10), 3623–3630.  
147 <https://doi.org/10.1128/JCM.38.10.3623-3630.2000>

148 Ikumapayi, U. N., Kanteh, A., Manneh, J., Lamin, M., & Mackenzie, G. A. (2016). An outbreak of *Serratia*  
149 *liquefaciens* at a rural health center in The Gambia. *The Journal of Infection in Developing Countries*,  
150 *10*(08), 791–798. <https://doi.org/10.3855/jidc.7184>

151 Machado, S. G., da Silva, F. L., Bazzolli, D. M. S., Heyndrickx, M., Costa, P. M. de A., & Vanetti, M. C. D.  
152 (2015). *Pseudomonas* spp. and *Serratia liquefaciens* as Predominant Spoilers in Cold Raw Milk:

153 Predominant spoilers in cold raw milk. *Journal of Food Science*, 80(8), M1842–M1849.  
154 <https://doi.org/10.1111/1750-3841.12957>

155 Marchesi, J. R., Sato, T., Weightman, A. J., Martin, T. A., Fry, J. C., Hiom, S. J., & Wade, W. G. (1998). *Design*  
156 *and Evaluation of Useful Bacterium-Specific PCR Primers That Amplify Genes Coding for Bacterial*  
157 *16S rRNA*. 64, 7.

158 Momose, T., Masutani, S., Oshima, A., Kawasaki, H., Tanaka, R., Iwamoto, Y., Ishido, H., & Senzaki, H.  
159 (2018). First Pediatric Case of Infective Endocarditis Caused by *Serratia Liquefaciens*. *International*  
160 *Heart Journal*, 59(6), 1485–1487. <https://doi.org/10.1536/ihj.17-595>

161 O’Sullivan, J.D., Giblin, L., Paul L.H. Mc Sweeney P.L.H., Sheehan, J.J., & Cotter, P.D. (2013). Nucleic acid-  
162 based approaches to investigate microbial-related cheese quality defects. Review article doi:  
163 10.3389/fmicb.2013.00001

164 Palmas, F., Cosentino, S., Fadda, M. E., Deplano, M., & Mascia, V. (1999). Microbial characteristics of  
165 Pecorino processed cheese spreads. *Le Lait*, 79(6), 607–613. <https://doi.org/10.1051/lait:1999650>

166 Proroga, Y. T. R., Cerrone, A., Valvini, O., Castellano, S., Carullo, M. R., Bove, D., Guarino, A., & Iovane, G.  
167 (2009). MICROBIOLOGICAL CHARACTERIZATION OF A TYPICAL ITALIAN CHEESE: PECORINO DI  
168 LATICAUDA (PART I). *Italian Journal of Food Safety*, 1(5), 55. <https://doi.org/10.4081/ijfs.2009.5.55>

169 Quigley, L., O’Sullivan, D. J., Daly, D., O’Sullivan, O., Burdikova, Z., Vana, R., Beresford, T. P., Ross, R. P.,  
170 Fitzgerald, G. F., McSweeney, P. L. H., Giblin, L., Sheehan, J. J., & Cotter, P. D. (2016). Thermus and  
171 the Pink Discoloration Defect in Cheese. *mSystems*, 1(3), mSystems.00023-16, e00023-16.  
172 <https://doi.org/10.1128/mSystems.00023-16>

173 Takahashi, S., Tomita, J., Nishioka, K., Hisada, T., & Nishijima, M. (2014). Development of a Prokaryotic  
174 Universal Primer for Simultaneous Analysis of Bacteria and Archaea Using Next-Generation  
175 Sequencing. *PLoS ONE*, 9(8), 9. <https://doi.org/10.1371/journal.pone.0105592>

176 Thomson, N. R., Crow, M. A., McGowan, S. J., Cox, A., & Salmond, G. P. C. (2000). Biosynthesis of  
177 carbapenem antibiotic and prodigiosin pigment in *Serratia* is under quorum sensing control:  
178 Production of antibiotic and pigment by *Serratia*. *Molecular Microbiology*, 36(3), 539–556.  
179 <https://doi.org/10.1046/j.1365-2958.2000.01872.x>

180 Todaro, M., Francesca, N., Reale, S., Moschetti, G., Vitale, F., & Settanni, L. (2011). Effect of different salting  
181 technologies on the chemical and microbiological characteristics of PDO Pecorino Siciliano cheese.  
182 *European Food Research and Technology*, 233(6), 931–940. [https://doi.org/10.1007/s00217-011-](https://doi.org/10.1007/s00217-011-1593-7)  
183 1593-7

184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198

199 **Figure captions**

200 Figure 1. Example of pink discoloration on Pecorino Toscano cheese

201 Figure 2. Box plot of microbial counts, reported as mean value for each media

202 **Table 1.** Identification of isolated strains.

203

Strain number	Bacterial Genus and Species	Score	Query cover	Accession number
5006	<i>Serratia liquefaciens</i>	730	98%	MH669247.1
5007	<i>Serratia liquefaciens</i>	819	98%	MT279350.1

---

5008	<i>Serratia liquefaciens</i>	1099	99%	MH669144.1
5009	<i>Serratia liquefaciens</i>	926	98%	MH669139.1
5010	<i>Serratia liquefaciens</i>	1003	98%	MH668086.1
5011	<i>Serratia liquefaciens</i>	1120	99%	MT279350.1
5012	<i>Enterobacter hormaechei</i>	1051	98%	KC154051.1
5013	<i>Enterobacter cloacae</i>	1280	99%	MT145960.1
5014	<i>Enterobacter hormaechei</i>	1253	98%	MT258990.1
5015	<i>Serratia liquefaciens</i>	1375	99%	MH668086.1
5016	<i>Serratia liquefaciens</i>	1375	99%	MT279350.1
5017	<i>Enterobacter cloacae</i>	1373	99%	MT145960.1
5018	<i>Serratia liquefaciens</i>	1282	99%	MT279350.1
5019	<i>Serratia liquefaciens</i>	1554	99%	MH669139.1

---