



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Adventitious shoot organogenesis and encapsulation technology in hop (*Humulus lupulus* L.)

This is the peer reviewed version of the following article:

Original

Adventitious shoot organogenesis and encapsulation technology in hop (*Humulus lupulus* L.) / Liberatore, CLAUDIA MARIA; Rodolfi, Margherita; Beghè, Deborah; Fabbri, Andrea; Ganino, Tommaso; Chiancone, Benedetta. - In: SCIENTIA HORTICULTURAE. - ISSN 0304-4238. - 270:(2020).
[10.1016/j.scienta.2020.109416]

Availability:

This version is available at: 11381/2879921 since: 2021-12-29T07:21:19Z

Publisher:

Elsevier B.V.

Published

DOI:10.1016/j.scienta.2020.109416

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

1 **Adventitious shoot organogenesis and encapsulation technology in hop (*Humulus lupulus* L.)**

2 Claudia Maria Liberatore¹, Margherita Rodolfi¹, Deborah Beghè¹, Andrea Fabbri¹, Tommaso
3 Ganino^{1,2}, Benedetta Chiancone¹

4 ¹Department of Food and Drug, University of Parma, Italy.

5 ²Institute of BioEconomy (IBE), National Research Council (CNR), Sesto Fiorentino, Italy

6

7 ***Corresponding author:** benedetta.chiancone@unipr.it tel. 0039 0521 905661

8 Orcid Benedetta Chiancone: 0000-0003-2731-2349

9

10

11 **Abstract**

12 Due to the increasing interest of growers in hop cultivation, there is a rising demand for hop plants
13 that is not fully satisfied by nursery activity. Biotechnological methods and, specifically, *in vitro*
14 propagation, could offer new opportunities to overcome these limitations, allowing the production
15 of a great number of plantlets, in reduced space and independently of the season. In this research,
16 organogenesis from two types of explant, petioles and internodes, and, for the first time to our
17 knowledge, unipolar explants encapsulation have been studied to evaluate alternative methods for
18 hop propagation and to try to satisfy hop nursery need. Our results confirmed the possibility of
19 resorting to organogenesis as a propagation method, only if a precise and continuous check of
20 somaclonal variation is carried out, due to the occurrence of a small percentage of tetraploids.
21 Moreover, hop microcuttings were demonstrated to be a suitable starting material for encapsulation;
22 indeed, in less than one month, it was possible to obtain a very high regrowth (up to 100%) and
23 conversion (up to 82%). These results represent a first step towards the application of this
24 technology to hop, also for other purposes, such as short and long term preservation.

25 **Keywords:** flow cytometry, hop, internodes, *in vitro* regeneration, petioles, synthetic seed.

26 **1. Introduction**

27 All over the world, the commercial value of hop (*Humulus lupulus* L.) is in its essential oils and
28 resins that confer flavour, bitterness and aroma to beer (Zanoli and Zavatti, 2008).

29 In line with consumer growing interest for craft beers and for hops endowed with particular
30 phytochemical profiles (Barth-Hass, 2016), recently, also in Italy, the culture of hop has attracted
31 the interest of growers in increasing areas dedicated to hop. Unfortunately, the specialized nursery
32 activity on the Italian territory is scarce, there is a lack of certified plants and, for this reason, hop
33 growers resort to self-produced rhizome or purchase propagation material on parallel markets,
34 regardless of genetic correspondence and phytosanitary quality (Carbone and Cherubini, 2016).
35 Moreover, the problem of hop plant lack is, further, exacerbated by the plant growing slowness and

36 by the season-dependency of traditional propagation material, such as dormant rhizomes and soft
37 wood cuttings (Neve, 1991). To support the nursery sector, micropropagation can represent a
38 valuable alternative to hop traditional multiplication, allowing the obtainment of a high number of
39 true-to-type plants out of the natural season, in a relatively short time and in limited space (Barlass
40 and Skene, 1982). *In vitro* plant tissue culture can be performed starting from different types of
41 explants: meristems, characterized by the presence of undifferentiated cells, and differentiated
42 tissues, such as leaf portions, stems, petioles and buds (Smith, 2000; Thakur and Kanwar, 2018;
43 Hesami et al., 2019).

44 Numerous are the published studies carried out by exploiting all the potentialities of hop *in vitro*
45 tissue culture (Gurriarán et al., 1999; Faragò et al., 2009; Fortes et al., 2010; Roy et al., 2001); some
46 of them report the use of micropropagation to eradicate viruses, resorting to meristem culture and
47 cloning (Vine and Jones, 1969; Adams, 1975; Samyn and Welvaert 1983), others evaluated the
48 organogenic ability of petioles and stems (Batista et al., 1996) or leaf portions (Liberatore, 2020)
49 testing different basal media compositions.

50 Generally, micropropagation led to the obtainment of plants true to the starting material, but,
51 sometimes, due to several conditions of *in vitro* tissue culture and, most of all, to the induction of
52 de- and redifferentiation processes (Standardi and Piccioni, 1998), various type of changes, known
53 as somaclonal variation, may occur (Larkin and Scowcroft, 1981). In hop, Šuštar-Vozlič et al.
54 (1999) and Liberatore (2020) reported the obtainment of tetraploids, respectively from petiole and
55 leaf portion culture.

56 To overcome this problem, it would be possible to resort, as starting material, to vegetative
57 propagules (bulbs, tubers, corms) or shoot cuttings and buds, exploiting the natural ability of the
58 plant species to produce already organized meristematic tissues (Standardi et al., 1999). Actually, in
59 this way, the de- re-differentiation phase will be avoided, highly reducing the risks of somaclonal
60 variation (Standardi and Piccioni, 1998). Encapsulation technology, when, as starting material,
61 unipolar plant propagules are used, represents one of the solutions to overcome somaclonal

62 variation and, moreover, presents numerous advantages. Specifically, artificial seed or synthetic
63 seed defined as “artificially encapsulated somatic embryos, shoots or other tissues which can be
64 used for sowing under *in vitro* or *ex vitro* conditions” (Aitken-Christie et al., 1995), combines the
65 advantages of clonal propagation (high efficiency production, genetic uniformity of plant material,
66 sanitary plant conditions, perfect reduced spaces requirements) with those of zygotic seeds (easy
67 handling and transportability, storability, reduced dimensions, mechanization potentiality)
68 (Standardi and Micheli, 2013). Moreover, encapsulation technology represents a valuable tool to
69 facilitate the exchange of sterile plant material among different laboratories, thanks to the reduced
70 dimensions of propagules and to the easiness of transportability (Redenbaugh, 1993; Gray et al.,
71 1995). Several are the studies reporting the use of encapsulation technology for different plant
72 species, resorting to *in vitro* derived unipolar explants, including ornamental plants (Lambardi et al.,
73 2006; Benelli et al., 2017; Micheli and Benelli, 2019), *Morus indica* L. (Micheli et al., 2017), *Olea*
74 *europaea* L. (Micheli et al., 1998; Micheli et al., 2007; Micheli et al., 2019), Carrizo citrange
75 (*Citrus sinensis* Osb. × *Poncirus trifoliata* L. Raf.) (Germanà et al., 2011) and *Vitis vinifera* L.
76 (Benelli, 2016). In hop, this technique has been used to evaluate the response of several genotypes
77 to cryopreservation, for germplasm storage (Reed et al., 2003; Reed and Hummer, 2013). However,
78 to the best of our knowledge, nothing has been published about the use of encapsulation technology
79 applied to hop.

80 The main aims of this ~~study-research was-were the hop *in vitro* propagation improvement to~~
81 ~~evaluate the possibility of propagating hop~~, resorting to *in vitro* petiole and internode regeneration
82 and, ~~for the first time, to study the study, for the first time, of~~ the encapsulation technology in this
83 species. With ~~these aims objectives~~, different types of growth regulators, at different concentrations,
84 were applied, studying their effect on vegetative performance of the different explant types and
85 methodologies tested. To verify the possible somaclonal variation occurrence among regenerants, a
86 ploidy analysis was carried out within shoots obtained from petiole and internode *in vitro* culture.

87

88 2. Material and methods

89 2.1 Plant material

90 Petioles and internodes, for *in vitro* regeneration, and microcuttings, for encapsulation, were
91 isolated from two-month-old *in vitro* cultured plantlets of hop, genotype “Gianni” (Mongelli et al.,
92 2015). Plantlets were cultured in 500-ml glass jars containing 100 ml of **MS-HF** ([Hormone Free](#))
93 culture medium: Murashige and Skoog (MS) salt and vitamin mixture (1x) (Murashige and Skoog,
94 1962), 30 g L⁻¹ of sucrose, 8 g L⁻¹ of agar (pH 5.8); *in vitro* cultures were maintained in a growth
95 chamber, at 25±1 °C and light intensity of 20 µmol m⁻² s⁻¹, under 16 h photoperiod.

96

97 2.2 In vitro petiole and internode regeneration

98 Isolated petioles and internodes, deprived of the leaves, were cut in sections (0.5 mm) and put in
99 culture, in sterile Petri dishes (ten explants per each Petri dish, ten Petri dishes per treatment). In
100 order to evaluate the influence of type and concentration of growth regulators, added in the culture
101 medium on hop petiole and internode regeneration, five culture media, with the following
102 composition, were tested: **MS-BAP0**: MS-HF culture medium supplemented with 0.1 µM 1-
103 Naphthaleneacetic acid (NAA); **MS-BAP2**: MS-BAP0 culture medium supplemented with 8.88 µM
104 6-Benzylaminopurine (BAP); **MS-BAP4**: MS-BAP0 culture medium supplemented with 17.77 µM
105 BAP; **MS-BAP6**: MS-BAP0 culture medium supplemented with 26.66 µM BAP; **MS-BAP8**: MS-
106 BAP0 culture medium supplemented with 35.55 µM BAP. All culture media, after adjusting the pH
107 to 5.8, were sterilized in autoclave for 20 min at 121 °C. Petri dishes were then sealed and placed in
108 growth chamber, at 25±1 °C and light intensity of 20 µmol m⁻² s⁻¹, under 16 h photoperiod.

109

110 2.2.1 Flow cytometry analysis of petiole and internode regenerants

111 The ploidy level of regenerants obtained from petiole and internode culture was evaluated using a
112 NovoCyte (Acea Biociences) and compared with that of the mother plant. Each regenerant was
113 analysed three times. About 0.5 cm² of sample (a young leaf from a developed shoot or from the

114 mother plant, genotype Gianni) was chopped using a razor blade in a Petri dish containing 0.5 mL
115 of extraction buffer (Partec CyStain PI Absolute P Nuclei Extraction Buffer; Partec GMBH,
116 Münster, Germany). The suspension was filtered through a 30- μ m filter into a 3.5 mL plastic tube,
117 to which was then added 2.0 mL of Partec CyStain PI Absolute P Staining Buffer, containing 12 μ l
118 of Propidium Iodide Solution and 6 μ l RNase. Samples were maintained in the dark for 30 min and
119 then analyzed by flow cytometry. At least 5000 nuclei were analysed in each sample (NovoCyte
120 Flow Cytometer Operator's Guide), Acea NovoExpress v.1.25 software was used.

121

122 2.3 Microcutting encapsulation

123 Uninodal microcuttings (3-4 mm long) from *in vitro* proliferated shoots, without leaves and with
124 two axillary buds, were excised and, subsequently, subjected to encapsulation, using artificial
125 endosperm (AE) with the following composition: half-strength MS medium, 50 g l⁻¹ sucrose and 0.1
126 μ M NAA (pH 5.8). Single microcuttings were immersed, firstly, in the encapsulating solution (AE
127 enriched with sodium alginate - 2.5%, w/v - alginic acid sodium salt, medium viscosity; code
128 366551 Carlo Erba)); secondly, the alginate coated propagules were immersed in the complexing
129 solution (AE enriched with CaCl₂ - 1.1% w/v) for 35 min. After complexation, the capsules were
130 rinsed three times (15 min each time) with sterile AE (Micheli and Standardi, 2005; Standardi and
131 Micheli, 2013).

132 In order to test the effect of type and concentration of growth regulators on hop encapsulated
133 microcuttings vegetative parameters, the following thesis were assessed: **EMC-0**: microcuttings
134 encapsulated in AE; **EMC-0.5**: microcuttings encapsulated in AE enriched with 2.22 μ M BAP;
135 **EMC-1.0**: microcuttings encapsulated in AE enriched with 4.44 μ M BAP; **EMC-1.5**: microcuttings
136 encapsulated in AE enriched with 6.66 μ M with BAP; **EMC-2.0**: microcuttings encapsulated in AE
137 enriched with 8.88 μ M. Ten capsules were placed in each sterile Petri dish, containing **MS-HF**
138 culture medium; ten Petri dishes were prepared per each treatment. As control, 100 not encapsulated

139 microcuttings (naked, **MC**) were put in culture on the same culture medium (ten microcuttings per
140 Petri dish).

141

142 2.4 Statistical analysis of data

143 2.4.1 *In vitro* petiole and internode regeneration

144 The following parameters were measured every week for 14 weeks, monitoring petioles and
145 internodes in culture: viability (percentage of explants with a green or light brown appearance,
146 without necrosis or yellowing), callogenesis (percentage of explants producing callus), rooting
147 (percentage of explants developing roots), shooting (percentage of explants developing shoots),
148 number and length of roots per each explant and number and length of shoots per each explant.

149 At the end of the experiment, Mean Regeneration Time (MRT) and Regeneration Energy (RE)
150 were calculated (formulae below reported were adapted to organogenesis, respectively from Kader
151 (2005) and from Paul (1972) who developed them for seed germination); the formulae used were
152 the following: $MRT = \sum f \cdot x / f_{tot}$ (f=number of explants with shoots on a given day, x= days needed
153 for explant to regenerate; f_{tot} = number of all explants with shoots); $RE = \sum f \cdot x$ before the
154 peak*100)/ f_{tot} (f=number of explants with shoots on a given day; x= days needed for explant to
155 regenerate; peak= time at which regeneration is the highest; f_{tot} = number of all explants with
156 shoots).

157 Two-way ANOVA (influence of “Type of explant” (TE) and of “Culture Medium Composition”
158 (CMC) was used to calculate the differences between treatments per each parameter considered;
159 Tukey’s test ($p \leq 0.05$) was used for mean separation (SYSTAT 13.1, Systat Software, Inc; Pint
160 Richmond, CA).

161

162 2.4.2 Microcutting encapsulation

163 The following parameters were measured every week for 4 weeks: viability (percentage of
164 encapsulated explants with a green appearance, without necrosis or yellowing), regrowth

165 (percentage of encapsulated explants producing shoots >4 mm), conversion (percentage of explants
166 with extrusion of shoots and 4 mm long roots), number and length of shoots per explant, and
167 number and length of roots per explant.

168 At the end of the experiment, Mean Regrowth Time (MRET), Regrowth Energy (REE), Mean
169 Conversion Time (MCT) and Conversion Energy (CE) were calculated (formulae below reported
170 were adapted to organogenesis respectively from Kader (2005) (MRET and MCT) and from Paul
171 (1972) (MRET and CE) who developed them for seed germination); the formulae used were the
172 following: $MRET = \sum f \cdot x / f_{tot}$ (f=number of microcuttings with shoots on a given day, x= days
173 needed for explant to regenerate; f_{tot} = number of all microcuttings with shoots); $REE = \sum f \cdot x$
174 before the peak*100)/ f_{tot} (f=number of microcuttings with shoots on a given day, x= days needed
175 for microcuttings to sprout; peak= time at which regeneration is the highest; f_{tot} = number of all
176 microcuttings with shoots); $MCT = \sum f \cdot x / f_{tot}$ x= days needed for explant to regenerate; f_{tot} =
177 number of all microcuttings with shoots); $CE = \sum f \cdot x$ before the peak*100)/ f_{tot} (f=number of
178 microcuttings with roots on a given day; x= days needed for microcuttings to root; peak= time at
179 which regeneration is the highest; f_{tot} = number of all microcuttings with roots). Data were used to
180 calculate means.

181 One-way ANOVA was used to calculate the differences for the factor “Culture Medium
182 Composition”, per each parameter considered; Tukey’s test ($p \leq 0.05$) was used for mean separation;
183 when data were not normally distributed, non-parametric tests (Kruskal-Wallis’ test, $p \leq 0.05$) were
184 carried out and Dwass-Steel-Critchlow-Fligner’s test was used for mean separation (SYSTAT 13.1,
185 Systat Software, Inc; Pint Richmond, CA).

186

187 **3. Results**

188 *3.1 In vitro petiole and internode regeneration*

189 In the course of the experiment, almost all explants remained viable, preserving their green colour
190 for the first week, then turning light brown. Statistical analysis evidenced significant differences for

191 the factor “TE” (Table 1), with internodes statistically more viable than petioles (100% and 91.3%,
192 respectively; data not shown).

193 Callus formation was the first response observed in the cultured explants, after about one week of
194 culture; indeed, a yellowish/greenish spongy callus started to develop at the cut ends of explants,
195 from which it increased rapidly, up to cover the entire surface of the explants (Fig. 1a). Statistical
196 analysis carried out, at the end of the experiment, on the percentage of callus producing explants did
197 not evidence a significant difference for both factors considered, TE and CMC (Table 1).

198 Rhizogenesis was observed, after one week of culture, from internodes (Fig. 1b) and, after two
199 weeks of culture, from petioles. Statistical analysis evidenced a significant interaction between the
200 two factors, TE and CMC (Table 1): ~~considering the factor TE, t~~the significant highest percentage
201 of explants with roots was recorded in the MS-BAP0 for petioles and in MS-BAP0 and MS-BAP2
202 for internodes; ~~whilst, considering the factor CMC~~Moreover, the percentage of explants with roots
203 was statistically higher in internodes than in petioles, for both culture media MS-BAP0 (70.8% vs
204 40.0%) and MS-BAP2 (52.9% vs. 10.9%); (Table 1).

205 The number of roots produced per single explant was very variable, depending, mainly, on BAP
206 concentration. Statistical analysis confirmed that CMC was the factor that significantly influenced
207 the explant response (Table 1): indeed, both for petioles and internodes, in explants cultured on MS-
208 BAP0 (3.6) a number of roots statistically higher than in those cultured on MS-BAP4 (1.6) and MS-
209 BAP8 (1.3) was recorded (~~data not shown~~Fig. 2a).

210 Regarding the root length, statistical analysis evidenced a significant interaction between the two
211 parameters; for both types of explant, petioles and internodes, culture media MS-BAP2 and MS-
212 BAP4 induced roots to grow statistically more than MS-BAP0 (Table 1). Comparing the two types
213 of explant, for every culture medium tested, it seems that internodes regenerated statistically longer
214 roots than petioles (Table 1).

215 Together with the emergence of the first roots, in both types of explants, calli formed organogenic
216 centers, some of which, after 2 weeks, started to turn into small shoots; for petioles, organogenesis

217 was observed in explants cultured in all culture media, while, for internodes, explants cultured in
218 MS-BAP0 did not show any shoot regeneration. Mostly indirect organogenesis was observed (Fig.
219 1c); but, even though at very low rate (4.5%, ~~data not shown~~), and only from petioles cultured in
220 MS-BAP0, also direct shoot regeneration was observed (Fig 1d).

221 Shoot regeneration continued for all the period in which the explants were kept in culture and it
222 was observed from the explants cultured in all culture media; a total of 90 shoots (67 regenerants
223 obtained from petioles and 23 from internodes, data not shown) were obtained at the end of
224 experiment; shoots carried on growing in length and forming well developed leaves (Fig. 1e).

225 Statistical analysis carried out on the percentage of shoot producing explants evidenced a
226 significant difference only for the factor TE (Table 1). Indeed, within petioles, a statistically higher
227 percentage of shoot regenerating explants was recorded (8.1% vs. 3.9%, ~~data not shown~~) (Fig. 2b).

228 Regarding the parameter “number of shoots”, no significant differences were detected; on the
229 average, shoot number was 1.8 for petioles and 1.6 for internodes (data not shown).

230 Analyzing the shoot length, statistical analysis evidenced a significant interaction between the two
231 factors: in petioles, significant differences were recorded between shoots obtained in MS-BAP8 (9.4
232 mm) and those in MS-BAP2 and MS-BAP4 (respectively 3.9 mm and 4.3 mm); whilst, in
233 internodes, the statistically longest shoots were obtained from explants cultured in MS-BAP2 (13.5
234 mm). Moreover, significant differences were observed in the length of shoots obtained from
235 explants cultured on MS-BAP2; indeed, from internodes, shoots statistically longer than those
236 obtained from petioles were obtained (13.5 mm vs. 3.9 mm) (Table 1).

237 A statistically significant interaction was detected for the parameter MTR, calculated considering
238 the explant response during the 14 weeks of culture. The analysis evidenced that petioles cultured
239 on medium MS-BAP0 needed, to produce new shoots, statistically less time (9.5 dd) than those
240 cultured on media with BAP; within explants in culture on BAP-containing culture media, the
241 highest concentration of BAP (MS-BAP8) statistically slowed down the regeneration process (61.6
242 dd) (Table 1), mainly if compared to MS-BAP2 and MS-BAP6 (25.3 dd and 32.2 dd respectively);

243 regarding internodes, explants cultured on MS-BAP4 needed a statistically higher number of days
244 (63.0 dd) to start organogenesis respect to MS-BAP6 (17.5 dd) and MS-BAP8 (14.0 dd). Moreover,
245 internodes began to regenerate shoots statistically earlier than petioles, when cultured on media MS-
246 BAP6 (17.5 dd vs 32.2 dd) and MS-BAP8 (14.0 dd vs. 61.6 dd). Considering RE, in petioles, MS-
247 BAP0 induced explants to regenerate much more synchronously and faster than the other media
248 considered; culturing petioles on a medium containing a high amount of BAP (MS-BAP8)
249 determined the statistically lower RE (3.3%), an opposite trend was observed in internodes that
250 gave better performances in terms of RE, when cultured MS-BAP8 (Table 1).

251

252 3.1.1 Flow cytometry analysis of petiole regenerants

253 Cytofluorimetric analysis, carried out on 90 regenerants, revealed that 89 of them were diploid,
254 like the mother plant (Fig. [2a3a](#)) and only one (regenerated from a petiole explant cultured on MS-
255 BAP8) was tetraploid (Fig. [2b3b](#)); whereas, other kind of ploidy variation, such as mixoploid or
256 octoploid, were not recorded. Because of the limited number of obtained tetraploid, it was not
257 possible to carry out a statistical analysis.

258

259 3.2 Microcutting encapsulation

260 In the thirty days of the experiment (at the end of this period, no more changes were observed in
261 the cultured explants), all microcuttings, both naked and encapsulated, maintained their viability,
262 showing a bright green color over the entire period in which they were kept in culture (Fig. [4a](#)).
263 Regrowth (Fig. [4b](#)) varied from 74% to 100% and it was statistically lower for EMC-0 and EMC-
264 2 encapsulated microcuttings (Table 2). Together with the MC, also the encapsulated ones produced
265 shoots, demonstrating that hop
266 microcuttings well respond to the encapsulation process (Table 2). Statistical analysis revealed that
267 the parameter “number of shoots” was influenced by BAP concentration; indeed, by increasing the
268 concentration of this growth regulator, the number of produced shoots decreased (Table 2). AE

269 containing 6.66 μ M BAP (EMC-1.5) induced shoots to elongate statistically more than those
270 encapsulated in others AEs (Table 2).

271 The analysis of MRET and REE, carried out considering the explant response during 30 days in
272 culture, showed that 6.66 μ M BAP containing AE (EMC-1.5) determined the statistically shortest
273 MRET (10.8 dd) and the highest REE (9.6%) respect to EMC-0. It means that the best combination
274 of MRET and REE is obtained with EMC-1.5, meaning that this artificial endosperm composition
275 lead microcuttings to regrow faster and in a more synchronous way (Table 2).

276 Overall the conversion was high for all explants (Fig. 3e4c), except for EMC-0 explants which
277 showed a conversion value significantly lower than the others (8.0%) (Table 2). Statistical analysis
278 revealed that BAP concentration influenced the number of roots produced per single explant;
279 indeed, the statistically highest number of roots was recorded in microcuttings encapsulated with
280 AE containing 6.66 μ M BAP (1.8), while the absence of BAP or its highest concentration in the AE
281 seem to inhibit root formation. Moreover, it seems that BAP induces root elongation; as a matter of
282 fact, the significantly shorter roots were observed in naked microcuttings (MC) and in those
283 encapsulated in artificial endosperm without BAP (EMC-0) (Table 2).

284 Considering the combination of both parameters, MCT and CE, EMC-1.5 seems to reduce
285 significantly the time needed for encapsulated microcuttings to produce roots, making the
286 conversion a much more synchronous process.

287 Overall, EMC-1.0 and EMC-1.5 had the best performance for most of the parameters considered,
288 although by adding only 2.22 μ M BAP (EMC-0.5) it was possible to obtain results almost
289 comparable with those achieved with a much higher concentration of BAP, reducing significantly
290 the production costs.

291

292 **4. Discussion**

293 As a rule, plant regeneration is influenced by several factors including culture medium
294 composition, genotype and explant type (Ganeshan et al., 2002; Liu et al., 2010). In hop, numerous

295 are the studies reporting somatic regeneration, starting from different types of explants, such as leaf
296 portions, internodes and petioles (Batista et al., 1996; Gurriarán et al., 1999; Motegi, 1979; Roy et
297 al., 2001; Smýkalová et al., 2001; Peredo et al., 2006). In literature, many authors reported
298 outstanding results in hop regeneration using internodal explants (Batista et al., 1996; Faragò et al.,
299 2009; Gurriarán et al., 1999; Motegi, 1979). However, it is well known that in hop, as in other
300 species, one of the main factors influencing regeneration response is the genotype (Batista et al.,
301 1996; Gurriarán et al., 1999; Faragò et al., 2009).

302 Therefore, the purpose of this research was to study the regeneration ability from petioles and
303 internodes of hop, genotype Gianni, evaluating the effect of the type and the concentration of
304 growth regulators, added in the basal medium. In this study, callus initiation represented the first
305 manifestation of the organogenic process. As reported by Batista et al. (1996), for stems and
306 petioles of the cultivars Bragança and Brewer's Gold, also for the genotype Gianni, callus started to
307 develop at the extremities of explants and increased notably up to cover the entire petiole and
308 internode surface. The percentage of callus producing explants was very high, although there were
309 no significant differences among considered media; this is in contrast with results obtained by
310 Gurriarán et al. (1999), who reported that, when IAA was added in media containing BAP,
311 callogenic response decreased for cv. Nugget explants and increased for cv. Brewers Gold explants,
312 pointing out that, in hop, different genotypes have different reaction to regeneration inductive
313 treatment.

314 Together with callus, explants produced roots, either directly or through callus formation. In hop,
315 there is a lack in literature about root induction from petioles; however, even as petioles would
316 seem to be less responsive than other explants sources (Škof et al., 2007), in this investigation no
317 significant differences between the two types of explant have been recorded; indeed, a high rooting
318 potential of petioles has been observed. Similar result was reported in a previous study, on
319 regeneration from leaves of the same hop genotype (Liberatore, 2020).

320 Type and concentration of cytokinin represent a key factor affecting the organogenic ability in hop
321 (Batista et al., 1996; Gurriarán et al., 1999; Šuštar-Vozlič et al., 1999); indeed, shoot regeneration
322 has not been observed in internodal explants cultured in MS-BAP0. Moreover, in this study, only
323 from petioles cultured in MS-BAP0, direct organogenesis has been observed; result comparable
324 with those reported by previous studies, in which they describe how the presence of BAP in the
325 culture medium promotes indirect organogenesis (Kazeroonian et al., 2018; Fujii and Shimizu 1990;
326 Zayova et al., 2012).

327 However, it is also known that organogenic ability depends on the type of explants (Šuštar-Vozlič
328 et al., 1999); actually, in this investigation shoot regeneration was detected in both types of explants
329 tested, petiole and internodes, but with different response.

330 Since indirect organogenesis is well known to induce genetic variability (Larkin and Scowcroft,
331 1981), regenerated shoots were subjected to cytofluorimetric analysis to check their ploidy level. In
332 this study only one regenerant, of 90 analyzed (1.1%), was tetraploid; a notably lower rate in
333 comparison to that recorded in a previous study, in which 7.2% of regenerants obtained from
334 genotype Gianni hop leaves was tetraploid (Liberatore, 2020); moreover, in this study, the only
335 tetraploid regenerant was obtained from a petiole cultured on the culture medium containing the
336 highest concentration of BAP (MS-BAP8); also in the above cited study on hop leaf regeneration,
337 the highest percentage of tetraploids (66.7%) was obtained from leaf portions cultured on the same
338 culture medium (Liberatore, 2020). Nevertheless, due to the low mutant rate, probably the induction
339 of regeneration from petioles and internodes could represent a valid method to produce plants true
340 to original starting material; but, anyway, before using petiole or internode regeneration as true-to-
341 type propagation method, it will always be necessary to check the genetic correspondence of
342 regenerants.

343 In order to avoid the occurrence of mutations in hop micropropagation process, resorting to
344 unipolar propagules can be a valuable solution; indeed, skipping the de- and re-differentiation phase
345 reduces significantly mutation occurrence (Standardi and Piccioni, 1998). A simple and efficient

346 method for *in vitro* propagation is the synthetic seed technology, resorting to unipolar explants; as a
347 matter of fact, several are the scientific studies, also in a specie strictly related to hop, such as
348 *Cannabis sativa* L., reporting the obtainment of true-to type plantlets (Nyende et al., 2003; Lata et
349 al., 2011; Chandrasekhara Reddy et al., 2012). Other than, encapsulation technology has other
350 numerous advantages potential applications, among which the most interesting is its use as a
351 valuable method for mass propagation of plant species; indeed, to date, numerous are the studies
352 concerning the application of encapsulation technology to several plant species, including fruit tree
353 crops, ornamentals, cereals and vegetables (Lambardi et al., 2006; Rai et al., 2009), but until now,
354 this technology was never applied to hop propagation. Several are the factors that markedly
355 influence *in vitro* encapsulated propagule behaviour, among which the most important are the initial
356 choice of plant materials, both in terms of genotype and of type of explant, artificial endosperm and
357 culture medium composition and growth conditions (Rai et al., 2009).

358 In this study, in which hop microcuttings were encapsulated, different artificial endosperm
359 compositions were tested, in order to evaluate, first of all, the suitability of hop, Gianni,
360 microcuttings to be encapsulated, then to study how their vegetative performances are influenced by
361 the artificial endosperm composition, in terms of BAP concentration. All the encapsulated explants,
362 independently on artificial endosperm composition, showed a high viability (100%), throughout the
363 experiment, demonstrating that hop microcuttings could absorb the water and nutrients they need
364 from the capsule; results reported in this study are in line with those obtained in other species, such
365 as *Actinidia deliciosa*, *Malus domestica*, *Olea europaea*, Carrizo citrange (Gardi et al., 1999;
366 Micheli and Standardi, 2005; Germanà et al., 2011; Micheli et al., 2019).

367 Researchers working on synthetic seed technology (Adriani et al., 2000; Micheli et al., 2019)
368 agree on the importance of regrowth, but most of all, of conversion in making this technology really
369 valuable. Our results showed that hop microcutting regrowth and conversion were strongly
370 influenced by the presence of growth regulators. Being this the first study on hop synthetic seed
371 technology, there is a complete lack of literature to compare our results with, the only study that

372 could be useful is the one about the nodal segment encapsulation of *Cannabis sativa*, a plant species
373 genetically close to hop, both species belonging to *Cannabaceae* family (Lata et al., 2009). In the
374 study on encapsulation of *Cannabis sativa*, in which the cytokinin (Thidiazuron) was added to
375 artificial endosperm, at a concentration 10 fold higher than that used in this study (BAP), up to 77%
376 of conversion was obtained; moreover, in *Cannabis*, around 21 days were needed to capsules to
377 convert (Lata et al., 2009); both results are similar to those reported in this research; indeed, in hop,
378 up to 82% of conversion was obtained in 20 dd, on the average, but, adding in the artificial
379 endosperm a concentration 10 fold lower of cytokinin. Other than in *Cannabis sativa*, the
380 importance of type, concentration and ratio of auxins and cytokinins is ~~highlighted~~highlighted for
381 several species, as reported by Lambardi et al. (2006).

382 In this study, BAP was added at different concentrations to induce regrowth and conversion in
383 encapsulated hop, cv. Gianni, microcuttings. After one month of culture, the highest BAP
384 concentration used (8.88 μmol) appears to inhibit the regrowth process in hop, as reported by Badr-
385 Elden (2013) in strawberry capsules; on the contrary, in *Mimosa pudica* L., Banu et al. (2014)
386 observed the best results, in terms of regrowth, from microcuttings encapsulated with artificial
387 endosperm containing the same BAP concentration (8.88 μmol). Conversion of encapsulated hop,
388 ~~ex-genotype~~ Gianni, was highly stimulated by the presence of BAP in the artificial endosperm,
389 independently on its concentration; results in contrast with those reported in this study are those
390 reported in *Celastrus paniculatus* (Fonseka et al., 2019) and in Prata-anã' banana's microshoots,
391 clone Gorutuba, (Pereira et al., 2017) which the best performance, in terms of conversion, was
392 obtained from explants encapsulated in 8.88 μmol BAP enriched artificial endosperm.

393 As above reported in some examples, consulting the literature about the influence of artificial
394 endosperm composition on vegetative parameters of several plant species encapsulated
395 microcuttings, results are extremely different, demonstrating, once more, the strong influence of the
396 genotype on their *in vitro* response.

397

398 **5. Conclusion**

399 A biotechnological approach to propagate *Humulus lupulus* L. represents a valid instrument to
400 obtain a relevant number of plants in a relatively short time. In this work, two methods such as
401 organogenesis from petioles and internodes and, for the first time to our knowledge, encapsulation
402 technology were described. ~~With the first one, it was possible to confirm the possibility of~~
403 ~~using~~ petioles and internodes as starting material for hop, genotype Gianni, *in vitro* propagation;
404 however, since ploidy analysis detected the presence of a certain, even though low, percentage of
405 tetraploids, there is the need of a continuous check of genetic correspondence, due to the possibility
406 of the occurrence of genetic mutations.

407 Encapsulation technology, by resorting to non-embryogenic (unipolar) plant propagules that allow
408 to by-pass the de-re-differentiation phase, represents an alternative to traditional methods,
409 exploiting the regeneration process from meristematic centres.

410 In less than one month, a very high regrowth (up to 100%) and conversion (up to 82%) was
411 obtained, proving that hop microcuttings are suitable for encapsulation. This result represents a first
412 step towards the application of the encapsulation technology for other purposes, such as hop
413 germplasm conservation. However, further investigations and insights are required to assess the
414 practical applicability of these techniques to different hop genotypes.

415

416 **Funding**

417 This research did not receive any specific grant from funding agencies in the public, commercial or
418 not-for-profit sectors.

419

420 **Acknowledgments**

421 Authors would like to thank Dr. Mirca Lazzaretti (Department of Chemistry, Life Sciences and
422 Environmental Sustainability, University of Parma) for the support in flow cytometric analysis.

423

424 **References**

- 425 Adams, A.N., 1975. Elimination of viruses from the hop (*Humulus lupulus*) by heat therapy and
426 meristem culture. J. Hort. Sci. 50, 151-160. <https://doi.org/10.1080/00221589.1975.11514616>
- 427 Adriani, M.; Piccioni, E.; Standard, A. (2000) Effect of different treatments on the conversion of
428 'Hayward' kiwifruit synthetic seeds to whole plants following encapsulation of in vitro-derived
429 buds. 28(1), 59-67. <https://doi.org/10.1080/01140671.2000.9514123>.
- 430 Aitken-Christie, J., Kozai, T., Smith, M.A.L., 1995. Glossary. In: Aitken-Christie J., Kozai, T.,
431 Smith, M.A.L., (Eds.) Automation and environmental control in plant tissue culture. Kluwer,
432 Dordrecht, pp. 9-12.
- 433 Badr-Elden, A.M., 2013. An effective protocol for *in vitro* storage and *ex vitro* re-growth of
434 strawberry capsules. Atlas J. Chem. Biochem. 1 (2), 30–38.
435 <http://dx.doi.org/10.5147/ajcb.2013.0081>
- 436 Banu, L.A., Rashid, M.H.O., Miah, M.A.B. 2014. Development of artificial seed and preservation
437 in *Mimosa pudica* L., an important medicinal plant in Bangladesh. J. Bio-Sci. 22:89-99.
438 <https://doi.org/10.3329/jbs.v22i0.30013>
- 439 Barlass, M., Skene, K.G.M., 1982. *In vitro* plantlet formation from *Citrus* species and hybrids. Sci
440 Hortic. 17, 333-341. [https://doi.org/10.1016/0304-4238\(82\)90114-5](https://doi.org/10.1016/0304-4238(82)90114-5)
- 441 Barth-Hass, G., 2016. The Barth Report, Nuremberg, Germany.
- 442 Batista, D., Sousa, M.J., Pais, M.S., 1996. Plant regeneration from stem and petiole-derived callus
443 of *Humulus lupulus* (hop) clone Bragança and var. Brewers Gold. In Vitro Cell Dev. Biol. Plant.
444 32, 37-41. <https://doi.org/10.1007/BF02823011>
- 445 Benelli, C., 2016. Encapsulation of shoot tips and nodal segments for in vitro storage of “Kober
446 5BB” grapevine rootstock. Horticulturae 2, 10. doi:10.3390/horticulturae2030010
- 447 Benelli, C., Micheli, M., De Carlo, A. 2017. An improved encapsulation protocol for regrowth and
448 conservation of four ornamental species. Acta Soc. Bot. Pol. 86(3), 3559.
449 <https://doi.org/10.5586/asbp.3559>

450 Carbone, K., Cherubini, D., 2016. Il luppolo fresco “made in Italy”: qualità e competitività per
451 l’industria brassicola. In *Conference, Criticità e opportunità per lo sviluppo sostenibile della*
452 *filiera brassicola*, Roma, 26 Ottobre 2016.

453 Chandrasekhara Reddy, M., Sri Rama Murthy, K., Pullaiah, T., 2012. Synthetic seeds: A review in
454 agriculture and forestry. *Afr. J. Biotechnol.* 11, 14254-14275. <https://doi.org/10.5897/AJB12.770>

455 ~~Czabator, F.J., 1962: Germination Value: an index combining speed and completeness of pine~~
456 ~~germination. *Forest Science* 8, 386-396.~~

457 Fonseka, D.L.C.K, Wickramaarachchi, W.W.U.I., Madushani R.P.S., 2019. Synthetic seed
458 production as a tool for the conservation and domestication of *Celastrus paniculatus*: A rare
459 medicinal plant. *Annu Res Rev Biol* 32(4), 1-8. <https://doi.org/10.9734/arrb/2019/v32i430092>

460 Fortes, A.M., Santos, F., Pais, M.S., 2010. Organogenic nodule formation in hop: a tool to study
461 morphogenesis in plants with biotechnological and medicinal applications. *Biomed. Res. Int.*
462 2010, 1-16. <https://doi.org/10.1155/2010/583691>

463 Fujii, Y., Shimizu, K., 1990. Regeneration of plants from achenes and petals of *Chrysanthemum*
464 *coccineum*. *Plant Cell Rep.* 8(10), 625-627. doi: 10.1007/BF00270069

465 Ganeshan, S., Caswell, K.L., Kartha, K.K., Chibbar, R.N., 2002. Shoot regeneration and
466 proliferation, in: Khachatourians, G.G., McHughen, A., Scorza, R., Nip, W.K., Hui, Y.H. (Eds.),
467 *Transgenic plants and crops*. Marcel Dekker, Inc, New York, pp. 69–84.

468 Gardi, T., Piccioni, E., Standardi, A., 1999. Effect of bead nutrient composition on regrowth ability
469 of stored vitro-derived encapsulated microcuttings of different woody species. *J*
470 *Microencapsulation* 16(1), 13-25.

471 Gercheva, P., Andinova, T., Ivanova, K., 2000. Plant regeneration from leaf tissues of apple
472 cultivars Granny Smith, Morspur Golden and Starkrimson. *Bulg. J. Agric. Sci.* 6, 637-643.

473 Germanà, M.A., Micheli, M., Chiancone, B., Macaluso, L., Standardi, A., 2011. Organogenesis and
474 encapsulation of *in vitro*-derived propagules of *Carrizo citrange* (*Citrus sinensis* (L.) Osb. ×

475 *Poncirus trifoliata* (L.) Raf.). Plant Cell Tiss. Org. Cult. 106, 299-307.
476 <https://doi.org/10.1007/s11240-011-9921-y>

477 Gray, D.J., Compton, M.E., Harrell, R.C., Cantliffe, D.J., 1995. Somatic embryogenesis and the
478 technology of synthetic seed, in: Bajaj, Y.P.S. (Ed.), Biotechnology in agriculture and forestry.
479 Vol 30. Somatic embryogenesis and synthetic seed I. Springer, Berlin, pp. 127-151.

480 Gurriarán, M.J., Revilla, M.A., Tamés, R.S., 1999. Adventitious shoot regeneration in cultures of
481 *Humulus lupulus* L. (hop) cvs. Brewers Gold and Nugget. Plant Cell Rep. 18, 1007-11.
482 <https://doi.org/10.1007/s002990050699>.

483 Kader, M., 2005. Varying temperature regimes affect osmotically primed sorghum seeds and
484 seedlings. Int. Sorghum Millets Newslett., 42, 39.

485 Kazeroonian, R., Mousavi, A., Jari, S.K., Tohidfar, M., 2018. Factors Influencing *in Vitro*
486 Organogenesis of *Chrysanthemum morifolium* cv. 'Resomee Splendid'. Iran J Biotechnol.
487 16(2):e1454.. doi:10.21859/ijb.1454

488 Korban, S.S., O'Connor, P.A., Elobeidy, A., 1992. Effects of thidiazuron, naphthaleneacetic acid,
489 dark incubation and genotype on shoot organogenesis from *Malus* leaves. J. Hortic. Sci. 67, 341-
490 349. <https://doi.org/10.1080/00221589.1992.11516257>

491 Lambardi, M., Benelli, C., Ozudogru, E.A., Ozden-Tokatli, Y., 2006. Synthetic seed technology in
492 ornamental plants. Floric. Ornam. Plant Biotechnol. Adv. Top. Iss. 2, 347-354.
493 <https://doi.org/10.20546/ijcmas.2017.611.079>

494 Larkin, P.J., Scowcroft, W.R., 1981. Somaclonal variation- a novel source of variability from cell
495 cultures for plant improvement. Theor Appl Genet 60, 197-214.
496 <https://doi.org/10.1007/BF02342540>.

497 Lata, H., Chandra, S., Natascha, T., Khan, I.A., ElSohly, M.A., 2011. Molecular analysis of genetic
498 fidelity in *Cannabis sativa* L. plants grown from synthetic (encapsulated) seeds following *in vitro*
499 storage. Biotechnol. Lett. 33, 2503-2508. <https://doi.org/10.1007/s10529-011-0712-7>

500 Lata, H., Chandra, S., Khan, I.A., ElSohly, M.A., 2009. Propagation through alginate encapsulation
501 of axillary buds of *Cannabis sativa* L. -an important medicinal plant. *Physiol. Mol. Boil. Plant* 15,
502 79-86. <https://doi.org/10.1007/s12298-009-0008-8>.

503 Liberatore, CM., 2020. Biotechnological methods to increase and preserve agrobiodiversity in hop.
504 PhD Thesis. University of Parma.

505 Liu, C., Callow, P., Rowland, L.J., Hancock, J.F., Song, G-q., 2010. Adventitious shoot
506 regeneration from leaf explants of southern highbush blueberry cultivars. *Plant Cell Tiss. Org.*
507 *Cult.* 103, 137-144. <https://doi.org/10.1007/s11240-010-9755-z>

508 Micheli, M., Benelli, C., 2019. Synthetic seeds of two aquatic plants, in: *Synthetic Seeds: recent*
509 *Trends and Future Perspectives. Methods in Molecular Biology.* in: Faisal, M., Alatar, A., (Eds.)
510 *Synthetic Seeds.* Springer, Cham. The Netherlands, pp. 233-239. [https://doi.org/10.1007/978-3-](https://doi.org/10.1007/978-3-030-24631-0_9)
511 [030-24631-0_9](https://doi.org/10.1007/978-3-030-24631-0_9)

512 Micheli M., Standardi A., 2005. Encapsulation of *in vitro*-derived explants of olive (cv. Moraiolo).
513 In: *Effect of pretreatments, their size and the coating.* *Curr. Top Biotechnol.* 2, 81-8.

514 Micheli, M., Standardi, A., Fernandes da Silva, D., 2019. Encapsulation and Synthetic Seeds of
515 Olive (*Olea europaea* L.): Experiences and Overview, in: Faisal, M., Alatar, A., (Eds.) *Synthetic*
516 *Seeds.* Springer, Cham. The Netherlands, pp. 347-361. [https://doi.org/10.1007/978-3-030-24631-](https://doi.org/10.1007/978-3-030-24631-0_9)
517 [0_9](https://doi.org/10.1007/978-3-030-24631-0_9)

518 Micheli, M., Bececco, V., Gardi, T., Martorana, L., Chiancone, B., Germanà, M.A., 2017.
519 Encapsulation of black mulberry microcuttings: studies on capsules and synthetic seeds. *Acta*
520 *Hortic.* 1155, 65-70. <https://doi.org/10.17660/ActaHortic.2017.1155.8>

521 Micheli, M., Hafiz, I.A., Standardi, A., 2007. Encapsulation of *in vitro* derived explants of olive
522 (*Olea europaea* L. cv. Moraiolo) II. Effects of storage on capsule and derived shoots performance.
523 *Sci. Hortic.* 113, 286-292. <https://doi.org/10.1016/j.scienta.2007.04.001>

524 Micheli, M., Mencuccini, M., Standardi, A., 1998. Encapsulation of “*in vitro*” proliferated buds of
525 olive. *Adv. Hortic. Sc.* 12(4), 163-168.

526 Mongelli, A., Rodolfi, M., Ganino, T., Marieschi, M., Dall'Asta, C., Bruni, R., 2015. Italian hop
527 germplasm: characterization of wild *Humulus lupulus* L. genotypes from Northern Italy by means
528 of phytochemical, morphological traits and multivariate data analysis. *Industrial Crops and*
529 *Products* 70, 16-27. <http://dx.doi.org/10.1016/j.indcrop.2015.02.036>

530 Motegi, T., 1979. Differentiation of shoots from hop stem callus culture. *Kyoyubokenkyu Neupo*
531 *Iwate Ika Daigaku* 14, 15-17.

532 Murashige, T., Skoog, F., 1962. A revised medium for rapid growth and bioassays with tobacco
533 tissue cultures. *Physiol Plant.* 15, 473-479. <http://dx.doi.org/10.1111/j.1399-3054.1962.tb08052.x>

534 Neve, R.A., 1991. *Hops*. Chapman and Hall, New York. <https://doi.org/10.1007/978-94-011-3106-3>

535 Nyende, A.B., Schittenhelm, S., Mix-Wagner, G., Greef, J., 2003. Production, storability, and
536 regeneration of shoot tips of potato (*Solanum tuberosum* L.) encapsulated in calcium alginate
537 hollow beads. *In vitro Cell. Dev. Biol. Plant* 39, 540-544. <https://doi.org/10.1079/IVP2003442>

538 [Paul, A. 1972. A guide to forest seed handling \(with special reference to the tropics\). *FAO Forestry,*](#)
539 [20:387.](#)

540 Peredo, E.L., Ángeles-Revilla, M., Arroyo-García, R., 2006. Assessment of genetic and epigenetic
541 variation in hop plants regenerated from sequential subcultures of organogenic calli. *Journal of*
542 *Plant Physiology* 163, 1071–1079

543 Pereira, J.C.G., Rocha, S.S., Cardoso Nogueira Londe, L., Batista Da Mota, M.C., Santos Alves,
544 P.F., Silva Viana, W., 2017. Growth of 'prata-ana' banana's microshoots clone gorutuba from
545 synthetic seeds: substrates and bap concentration. *Rev. Bras. Frutic.* 39(5), e-892.
546 <http://dx.doi.org/10.1590/0100-29452017892>.

547 Rai, M.K., Asthana, P., Singh, S.K., Jaiswal, V.S., Jaiswal, U., 2009. The encapsulation technology
548 in fruit plants—a review. *Biotech. Adv.* 27, 671-679.
549 <https://doi.org/10.1016/j.biotechadv.2009.04.025>

550 Redenbaugh, K., 1993. Introduction, in: Redenbaugh, K., (Ed.), *Synseeds: applications of synthetic*
551 *seeds to crop improvement*. CRC, Boca Raton, Fla. pp. 3-7.

552 Reed, B.M., Hummer, K.E., 2013. Cryopreservation and maintenance of hop material in the USDA
553 germplasm collection. Acta Hort. 1010, 191-197.
554 <https://doi.org/10.17660/ActaHortic.2013.1010.22>

555 Reed, B.M., Okut, N., D'Achino, J., Narver, L., DeNoma, J., 2003. Cold storage and
556 cryopreservation of hops (*Humulus L.*) shoot cultures through application of standard protocols.
557 CryoLetters 24, 389-396.

558 Roy, A.T., Leggett, G., Koutoulis, A., 2001. Development of a shoot multiplication system for hop
559 (*Humulus lupulus L.*). *In Vitro Cell.Dev.Biol.-Plant* 37, 79-83. [https://doi.org/10.1007/s11627-](https://doi.org/10.1007/s11627-001-0015-0)
560 [001-0015-0](https://doi.org/10.1007/s11627-001-0015-0)

561 Samyn, G, Welvaert, W., 1983. Producing a "nuclear stock" of virus free hop plants. Med. Fac.
562 Landbouw. Rijksuniv. Gent. 48, 877-881.

563 Škof, S., Bohanec, B., Kastelec, D., Luthar, Z. 2007. Spontaneous induction of tetraploidy in hop
564 using adventitious shoot regeneration method. Plant Breed. 126, 416-421.
565 <https://doi.org/10.1111/j.1439-0523.2007.01378.x>

566 Smith, R.H., 2000. Plant Tissue Culture: Techniques and Experiments, Ed 2. Academic Press, San
567 Diego.

568 Smýkalová, I., Ortová, M., Lipavská, H., Patzak, J., 2001. Efficient in vitro micropropagation and
569 regeneration of *Humulus lupulus* on low sugar, starch-gelrite media. Biol. Plant. 44, 7-12.

570 Standardi, A., Micheli, M., 2013. Encapsulation of *in vitro*-derived explants: an innovative tool for
571 nurseries, in: Lambardi, M., Ozudogru, E.A., Jain, S.M. (Eds.), Protocols for micropropagation of
572 selected economically-important horticultural plants. Springer, New York, 397-418.
573 https://doi.org/10.1007/978-1-62703-074-8_31.

574 Standardi, A., Piccioni, E., 1998. Recent perspectives on synthetic seed technology using non
575 embryogenic in vitro-derived explants. Int. J. Plant Sci. 159, 968-978.
576 <https://doi.org/10.1086/314087>

577 Šuštar-Vozlič, J., Javornik, B., Bohanec, B., 1999. Studies of somaclonal variation in hop (*Humulus*
578 *lupulus* L.). *Phyton. – Ann. Rei. Bot.* 39, 283-287.

579 Thakur, K., Kanwar, K., 2018. *In Vitro* Plant Regeneration by Organogenesis from Leaf Callus of
580 Carnation, *Dianthus caryophyllus* L. cv. 'Master'. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci.*
581 88, 1147–1155. <https://doi.org/10.1007/s40011-017-0851-2>

582 Vine, S.J., Jones, O.P., 1969. The culture of shoot tips of hop (*Humulus lupulus* L) to eliminate
583 viruses. *J. Hort. Sci.* 44, 181-28.

584 Zanolli, P., Zavatti, M., 2008. Pharmacognostic and pharmacological profile of *Humulus lupulus* L.
585 *J. Ethnopharmacol.*, 116, 383-396. <https://doi.org/10.1016/j.jep.2008.01.011>

586 Zayova, E., Vassilevska-Ivanova, R., Kraptchev, B., Stoeva, D., 2012. Indirect shoot organogenesis
587 of eggplant (*Solanum melongena* L.). *J. Cent. Eur. Agr.*, 13(3), 446-457.
588 <https://doi.org/10.5513/JCEA01/13.3.1070>
589

Table 1. Influence of “Type of explant” and “Culture medium composition” on several vegetative parameters of *in vitro* cultured hop, cv. Gianni, petioles and internodes, after 14 weeks of culture

Type of explant	Culture medium composition	Viability (%)	Explants with callus (%)	Explants with roots (%)	Explants with shoots (%)	Roots (n°)	Root length (mm)	Shoots (n°)	Shoots length (mm)	MRT (dd)	RE (%)
Petioles	MS-BAP0	84.1	90.7	40.0	3.0	3.2	5.3	2	5.0	9.5	11.3
	MS-BAP2	94.9	94.9	10.9	7.1	2.5	11.7	2.8	3.9	25.3	7.1
	MS-BAP4	95.7	95.7	7.3	7.1	1.4	10.8	2.6	4.3	57.3	6.0
	MS-BAP6	87.3	96.4	5.8	8.1	3.0	7.3	2.4	5.1	32.2	5.8
	MS-BAP8	94.5	91.5	2.7	15.0	1.7	8.4	2.4	9.4	61.6	3.3
Internodes	MS-BAP0	100.0	89.6	70.8	0.0	4.1	7.4	-	-	-	-
	MS-BAP2	100.0	98.3	52.9	10.4	2.7	17.2	2.2	13.5	40.8	5.5
	MS-BAP4	100.0	100.0	20.9	2.1	1.7	18.2	1.0	1.0	63.0	1.6
	MS-BAP6	100.0	93.2	6.7	3.5	1.3	11.4	2.0	6.6	17.5	6.0
	MS-BAP8	100.0	100.0	3.3	3.5	1.0	11.3	2.5	7.3	14.0	7.1
<i>Statistical analysis of factors</i>											
TE		0.016	0.424	0.000	0.027	0.632	0.000	0.363	0.393	0.279	0.510
CMC		0.777	0.526	0.000	0.103	0.000	0.000	0.892	0.062	0.076	0.174
TE*CMC		0.777	0.741	0.002	0.249	0.390	0.036	0.959	0.005	0.047	0.040
Two-way analysis of variance (ANOVA), followed by Tukey's test; $p \leq 0.05$.											
TE: Type of explant; CMC: Culture Medium; MS-BAP0: MS-HF culture medium supplemented with 0.1 μM of NAA; MS-BAP2: MS-BAP0 culture medium supplemented with 8.88 μM of BAP; MS-BAP4: MS-BAP0 culture medium supplemented with 17.77 μM of BAP; MS-BAP6: MS-BAP0 culture medium supplemented with 26.66 μM of BAP. MS-BAP8: MS-BAP0 culture medium supplemented with 35.55 μM of BAP; MRT: Mean Regeneration Time. RE: Regeneration Energy.											

Table 2. Influence of encapsulation and artificial endosperm composition on several vegetative parameters of hop, cv. Gianni, microcuttings, after 4 weeks of culture

Thesis	Viability (%)	Regrowth (%)	Conversion (%)	N° of shoots (n°)	L. of shoots (mm)	N° of roots (n°)	L of roots (mm)	MRET (dd)	REE (%)	MCT (dd)	CE (%)
MC-HF	100	94.0 a	72.0 a	1.3 a	5.5 c	1.0 d	5.8 b	15.0 ab	6.4 ab	17.1 b	5.9 ab
EMC-0	100	74.0 b	8.0 b	1.2 a	2.5 d	1.2 c	2.3 b	18.1 a	4.5 b	28.0 a	3.6 b
EMC-0.5	100	98.0 a	78.0 a	1.1 ab	13.4 b	1.3 b	9.5 a	6.4 b	7.6 ab	24.0 a	4.8 b
EMC-1.0	100	100.0 a	68.0 a	1.1 ab	11.2 b	1.3 b	9.9 a	13.4 ab	8.9 ab	18.3 b	6.6 ab
EMC-1.5	100	96.0 a	82.0 a	1.0 b	19.3 a	1.8 a	13.2 a	10.8 b	9.6 a	15.8 b	7.2 a
EMC-2.0	100	90.0 b	67.5 a	1.0 b	14.9 b	1.0 c	11.1 a	12.1 ab	8.2 ab	16.0 b	6.7 ab

One-way analysis of variance (ANOVA), followed by Tukey's test; $p \leq 0.05$. Per each column, values followed by different letters are statistically different.

MC-HF: naked microcuttings, cultured on Hormon Free (HF) culture medium; **EMC-0**: microcuttings encapsulated with Artificial Endosperm (AE); **EMC-0.5**: microcuttings encapsulated in Artificial Endosperm (AE) with 2.22 μM BAP; **EMC-1.0**: microcuttings encapsulated AE in 4.44 μM BAP; **EMC-1.5**: microcuttings encapsulated in AE with 6.66 μM with BAP; **EMC-2.0**: microcuttings encapsulated in AE with 8.88 μM BAP. **MRET**: Mean Regrowth Time. **REE**: Regrowth Energy; **MCT**: Mean Regrowth Time. **CE**: Conversion Energy.

Figures




Figure 1. Organogenesis from hop, cv. Gianni, petioles and internodes: **a)** callus covering the petiole surface; **b)** first root regeneration from an internode; **c)** indirect organogenesis from a petiole; **d)** direct organogenesis from a petiole; **e)** shoot development from an internode.




Figure 2. a) One-way ANOVA of the parameter “n° of roots” for the factor “Culture Medium Composition”; b) One-way ANOVA of the parameter “Percentage of explants with shoots” for the factor “Type of Explant”.




Figure 23. Cytofluorimetric analysis: histograms of fluorescence intensity of nuclei from diploid leaf tissue of genotype Gianni **a**) mother plant **b**) tetraploid regenerant from a petiole.




Figure 34. Hop, cv. Gianni, encapsulated microcutting development: **a**) viable encapsulated microcutting; **b**) microcutting regrowth; **c**) microcutting conversion.