



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Long time behavior of discrete volume preserving mean curvature flows

This is the peer reviewed version of the following article:

Original

Long time behavior of discrete volume preserving mean curvature flows / Morini, M.; Ponsiglione, M.; Spadaro, E.. - In: JOURNAL FÜR DIE REINE UND ANGEWANDTE MATHEMATIK. - ISSN 0075-4102. - (2022). [10.1515/crelle-2021-0076]

Availability:

This version is available at: 11381/2915788 since: 2024-11-19T15:25:39Z

Publisher:

De Gruyter Open Ltd

Published

DOI:10.1515/crelle-2021-0076

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

02 May 2026

LONG TIME BEHAVIOUR OF DISCRETE VOLUME PRESERVING MEAN CURVATURE FLOWS

MASSIMILIANO MORINI, MARCELLO PONSIGLIONE, AND EMANUELE SPADARO

ABSTRACT. In this paper we analyse the Euler implicit scheme for the volume preserving mean curvature flow. We prove the exponential convergence of the scheme to a finite union of disjoint balls with equal volume for any bounded initial set with finite perimeter.

KEYWORDS: Geometric evolutions; rigidity results; minimizing movements; variational methods.

AMS SUBJECT CLASSIFICATIONS: 53C44; 53C24; 49M25; 49Q20.

CONTENTS

Introduction	1
0.1. Exponential convergence of dissipations	2
0.2. Comments and open questions	3
0.3. Structure of the paper	3
1. A quantitative Alexandrov Theorem for nearly spherical sets	4
2. The incremental problem	7
3. The discrete volume preserving flow	11
3.1. Construction of the discrete flow	11
3.2. Stationary sets for the discrete flow	12
3.3. Long-time behaviour	13
Acknowledgements	21
References	21

INTRODUCTION

We study the asymptotic behaviour of a discrete-in-time approximation of the mean curvature flow with constrained volume. This geometric evolution consists in a family of evolving sets $[0, +\infty) \ni t \mapsto E_t \subset \mathbb{R}^N$, whose normal velocities $V(x, t)$ at any point $x \in \partial E_t$ are proportional to the scalar mean curvature $H_{\partial E_t}(x)$ of ∂E_t at x corrected by a constant forcing term which guarantees that the measure of $|E_t|$ is preserved during the flow:

$$(0.1) \quad V(x, t) = \bar{H}_{\partial E_t} - H_{\partial E_t}(x), \quad \bar{H}_{\partial E_t} := \frac{1}{\mathcal{H}^{N-1}(\partial E_t)} \int_{\partial E_t} H_{\partial E_t}(x) d\mathcal{H}^{N-1}(x).$$

Under suitable assumptions, this geometric flow is a model for coarsening phenomena in physical systems. For example, one can consider mixtures that, after a first relaxation time, can be described by two subdomains of nearly pure phases far from equilibrium, evolving in a way to minimize the total interfacial area between the phases while keeping their volume constant (further details on the physical background can be found in [4, 20, 24, 25]).

It is well-known that a typical evolution of (0.1) develops singularities of different kinds in finite time: components shrinking to points and disappearance, collisions and merging of domains, pinch-offs etc... Compared with the more familiar unconstrained mean curvature flow, the possible changes in the topology of the evolving sets E_t of (0.1) are even wider, because of the nonlocal character of the flow and the subsequent lack of comparison principles. There exist singular solutions also in the two dimensional case (see [18, 19]).

It is then clear that the analysis of the long time behaviour of systems exhibiting coarsening requires the introduction of suitable notions of weak solutions which allow the formation of singularities and extend the flow past the singular times. This is a well-established feature of curvature flows, and many definitions of weak solutions have been introduced in the literature. Here, we follow the method proposed independently by Almgren, Taylor and Wang [2] and by Luckhaus and Sturzenhecker [16], based on De Giorgi's minimizing movement approach. The authors consider an implicit time-discretization of the flow, which is regarded as a gradient flow of the perimeter functional with respect to a metric resembling the L^2 -distance. The limiting time-continuous flow constructed with this method is usually referred to as flat flow. This approach has the advantage to be easily adapted to volume constrained mean curvature flow as shown in [21], producing global-in-time solutions which then permit to analyse the equilibrium configurations reached in the long time asymptotics.

Previous results on the long time behaviour of the volume constrained mean curvature flow are mostly confined to the case of smooth solutions, starting from specific classes of initial regular sets ensuring the existence of global regular solutions; for example uniformly convex and nearly spherical initial sets (see [22, 11]), or nearly strictly stable initial sets in the three and four dimensional flat torus (see [23]). For more general initial data, the long time behaviour in the context of flat flows of convex and star-shaped sets (see [5, 14]) has been characterized only up to (possibly diverging in the case of [5]) translations.

In this paper we characterize the long-time limits of the discrete-in-time approximate flows constructed by the Euler implicit scheme introduced in [2, 16] plus the volume constraint. Our main result in Theorem 3.4 establishes that for every initial bounded set E_0 with finite perimeter, any discrete-in-time mean curvature flow with volume constraint converges exponentially fast to a finite union of disjoint balls with equal radii.

0.1. Exponential convergence of dissipations. The main issue here is to prove that the evolving sets remain uniformly bounded for all times and that the long-time limit is truly unique (and not just up to translations). This is a non-trivial fact, since the discrete scheme (as well as its continuous counterpart) does not satisfy comparison principles. To the best of our knowledge, the boundedness of the flow was not known previously for any weak notion of the nonlocal mean curvature flow. The problem is that summing the dissipations of the discrete schemes formally provides a L^2 (with respect to time) bound on the velocity of the evolving set, rather than the necessary L^1 bound: roughly speaking, one needs to sum the square roots of the dissipations rather than the dissipations themselves.

By a compactness argument and a characterization of the stationary configurations of the discrete flow (based on a recent theorem by Delgadino and Maggi [9] on Caccioppoli sets with constant weak mean curvature), the connected components of the evolving sets converge up to translations to balls.

Therefore, in order to improve the estimate on the dissipations, we compare at each time step the energy of the actual minimizers with that of a properly placed union of balls (one nearby each connected component of the evolving sets). In this way, it turns out to be necessary to estimate the dissipation of a nearly spherical set with respect to a close-by ball (this condition is definitively reached by each connected component of the flow thanks to the compactness argument above) with the actual dissipation of the flow. Indeed, if this happens, one can control the sum of the dissipations for all times larger than any time t from above by the energy dissipated at time t , thus clearly leading to an exponential decay of the dissipations and in turn of the velocity of the evolving sets.

The comparison of the dissipation of the flow and the one with respect to balls is equivalent to prove a control of the L^2 -norm of the function parametrizing a nearly spherical set with the L^2 -norm of the oscillation of its mean curvature. In the case of zero oscillation, this is nothing but Alexandrov's theorem on the characterization of the spheres as the unique embedded hypersurfaces with constant mean curvature and thus the aforementioned estimate may be regarded as a quantitative version of it. Its proof in turn relies on a result proven by Krummel and Maggi [15, Theorem 1.10], which can be seen as a sort of higher order Fuglede-type estimate involving the first variation of the perimeter rather than the perimeter itself. For different quantitative versions of Alexandrov's theorem see also [8, 6, 10].

0.2. Comments and open questions. It remains an open question to extend the results of the present paper to the flat flows, that is to obtain uniform estimates as the time step converges to zero. We conjecture the result to hold true, but the compactness arguments used in this paper in order to gain the closeness to sphere are not quantitative and do not allow to pass to the limit. Moreover, it is worth mentioning that, both nonlocal flat flows and discrete-in-time nonlocal flows are not uniquely defined. There are different ways to impose the volume constraint (by constraining the volume at each time-step, or by penalization, or by tuning suitably a forcing term). The flows so defined can be different (also because of the congenital non-uniqueness of mean curvature flows), but we expect that the asymptotics are always finite unions of equal volume balls.

Finally, we stress that the main Theorem 3.4 is sharp for what concerns the limiting configurations. Indeed, we can show that the union of suitably distant balls with equal radii is a stationary solution for the discrete flow and, hence, a possible asymptotics. However, such solutions are clearly unstable, because whatever small increase in one of the radii (and consequent decrease in the others) will drive the evolution towards a different asymptotic limit. This is easily seen for two disjoint balls with equal radii: any configuration of concentric balls with the same total volume but slightly perturbed radii will have strictly lower perimeter and thus will converge to a single ball (see Corollary 3.5). In this regard, we conjecture that, generically, the evolution of an initial set E_0 will converge to a *unique* limiting ball with the same volume.

0.3. Structure of the paper. In the first section we prove the Fuglede-type inequality for nearly spherical sets with almost constant mean curvature. Section 2 is then devoted to the introduction of the incremental problem at the basis of the discrete-in-time mean curvature flow. Finally, in the last section of the paper we prove the exponential convergence of the discrete flow to a union of well-separated balls with the equal volume.

1. A QUANTITATIVE ALEXANDROV THEOREM FOR NEARLY SPHERICAL SETS

In this section we prove a variant of a stability inequality by Krummel and Maggi [15, Theorem 1.10] related to Alexandrov's theorem, whose statement we recall for the reader's convenience. To this aim, and for later use, we fix the following notation: B denotes the unit ball of \mathbb{R}^N . Given $m > 0$, we denote by $B^{(m)}$ the ball centered at the origin with volume m and we denote by $r(m)$ its radius $\left(\frac{m}{\omega_N}\right)^{\frac{1}{N}}$. For any measurable function $f : \partial B^{(m)} \rightarrow (-1, +\infty)$, we denote by $E_{f,m}$ the set in \mathbb{R}^N whose boundary is described through the radial graph of φ , namely

$$(1.1) \quad E_{f,m} := \{tx \in \mathbb{R}^N : x \in \partial B^{(m)}, 0 \leq t \leq 1 + f(x)\}.$$

In the case $m = \omega_N$ we will write B instead of $B^{(\omega_N)}$ (to denote the unit ball) and E_f instead of E_{f,ω_N} . If the parametrizing function f has small L^∞ -norm, then the set $E_{f,m}$ can be regarded as a graphical radial perturbation of the ball $B^{(m)}$. We finally recall that, given a sufficiently regular set E , $H_{\partial E}$ stands for the sum of the principal curvatures of ∂E (with respect to the orientation given by the outward normal) and, for any bounded measurable set $E \subset \mathbb{R}^N$, we denote by $\text{bar}(E)$ denotes its barycenter, i.e., $\text{bar}(E) = |E|^{-1} \int_E x dx \in \mathbb{R}^N$.

Theorem 1.1 ([15]). *There exist $\delta \in (0, \frac{1}{2})$ and $C > 0$ with the following property: For any $f \in C^1(\partial B) \cap H^2(\partial B)$ such that $\|f\|_{C^1} \leq \delta$ and $\text{bar}(E_f) = 0$, we have*

$$\|f\|_{H^1(\partial B)} \leq C \|H_{\partial E_f} - (N-1)\|_{L^2(\partial B)}.$$

Remark 1.2. *In fact, in [15] the theorem is stated under the assumption $f \in C^{1,1}(\partial B)$. However, it is immediate to see by a standard approximation argument that the statement holds true under the weaker hypothesis $f \in C^1(\partial B) \cap H^2(\partial B)$.*

The stability inequality for nearly spherical sets with almost constant mean curvature that we are going to use in the next sections is the following.

Theorem 1.3. *There exist $\delta \in (0, \frac{1}{2})$ and $C > 0$ with the following property: For any $f \in C^1(\partial B) \cap H^2(\partial B)$ such that $\|f\|_{C^1} \leq \delta$, $|E_f| = \omega_N$ and $\text{bar}(E_f) = 0$, we have*

$$(1.2) \quad \|f\|_{H^1(\partial B)} \leq C \|H_{\partial E_f} - \bar{H}_{\partial E_f}\|_{L^2(\partial B)},$$

where we have set

$$\bar{H}_{\partial E_f} := \int_{\partial B} H_{\partial E_f}(x + f(x)x) d\mathcal{H}^{N-1}.$$

Proof. First of all we notice that, if we take the constant C in (1.2) large enough (e.g., $\sqrt{N\omega_N}$), then it is enough to consider only the case $\|H_{\partial E_f} - \bar{H}_{\partial E_f}\|_2 \leq 1$.

We write explicit formulas for the perimeter and its first variations for the sets E_f : by using the area formula we get

$$(1.3) \quad P(E_f) = \int_{\partial B} (1+f)^{N-1} (1+(1+f)^{-2}|\nabla f|^2)^{\frac{1}{2}} d\mathcal{H}^{N-1}.$$

Recall that $H_{\partial E_f}$ is the first variation of the perimeter at E_f , that is,

$$(1.4) \quad \begin{aligned} \delta P(E_f)[\psi] &:= \frac{d}{dt} \Big|_{t=0} P(E_{f+t\psi}) = \int_{\partial E_f} (H_{\partial E_f} \nu_{\partial E_f}) \cdot \frac{x}{|x|} \psi\left(\frac{x}{|x|}\right) d\mathcal{H}^{N-1}(x) \\ &= \int_{\partial B} (H_{\partial E_f} \nu_{\partial E_f})(p) \cdot x \psi(1+f)^{N-1} (1+(1+f)^{-2}|\nabla f|^2)^{\frac{1}{2}} d\mathcal{H}^{N-1}, \end{aligned}$$

where for simplicity of notation we have set $p = (1 + f(x))x$. On the other hand, by differentiating under the integral sign, from (1.3) we get

$$(1.5) \quad \begin{aligned} \delta P(E_f)[\psi] &= \int_{\partial B} (N-1)(1+f)^{N-2}(1+(1+f)^{-2}|\nabla f|^2)^{\frac{1}{2}}\psi \, d\mathcal{H}^{N-1} \\ &\quad - \int_{\partial B} \frac{(1+f)^{N-4}}{(1+(1+f)^{-2}|\nabla f|^2)^{\frac{1}{2}}} |\nabla f|^2 \psi \, d\mathcal{H}^{N-1} \\ &\quad + \int_{\partial B} \frac{(1+f)^{N-3}}{(1+(1+f)^{-2}|\nabla f|^2)^{\frac{1}{2}}} \nabla f \cdot \nabla \psi \, d\mathcal{H}^{N-1}, \end{aligned}$$

for all $\psi \in C^1(\partial B)$, where ∇f and $\nabla \psi$ are the tangent gradient of f and ψ on ∂B , respectively.

In the following, with a slight abuse of notation, by the symbol $O(g)$ we mean any function h of the form $h(x) = r(x)g(x)$, where $|r(x)| \leq C$ for all $x \in \partial B$, with $C > 0$ being a constant depending only on the a priori C^1 -bound $\|f\|_{C^1} \leq \frac{1}{2}$.

Since the normal to E_f at a point $p = (1 + f(x))x$ with $x \in \partial B$ is given by

$$\nu_{\partial E_f}(p) = \frac{1}{\sqrt{1 + (1+f)^{-2}|\nabla f|^2}} \left(-\frac{\nabla f}{1+f} + x \right),$$

one gets

$$(1.6) \quad \nu_{\partial E_f}(p) \cdot x = \frac{1}{\sqrt{1 + (1+f)^{-2}|\nabla f|^2}}.$$

Therefore, by (1.4), (1.6) and a simple Taylor expansion, we have that

$$(1.7) \quad \delta P(E_f)[\psi] = \int_{\partial B} H_{\partial E_f}(p) \left(1 + (N-1)f + R_1 \right) \psi \, d\mathcal{H}^{N-1},$$

with $|R_1| = O(|f|^2)$. Similarly, using (1.5) and a Taylor expansion, we get

$$(1.8) \quad \begin{aligned} \delta P(E_f)[\psi] &= \int_{\partial B} \left((N-1) + (N-1)(N-2)f + O(|f|^2) + O(|\nabla f|^2) \right) \psi \, d\mathcal{H}^{N-1} \\ &\quad + \int_{\partial B} (\nabla f + h) \cdot \nabla \psi \, d\mathcal{H}^{N-1}, \end{aligned}$$

where h is a vector field satisfying

$$(1.9) \quad |h| = O((|f| + |\nabla f|^2)|\nabla f|).$$

By comparing (1.7) and (1.8), and recalling that $|R_1| = O(|f|^2)$, we infer that

$$(1.10) \quad \begin{aligned} &\int_{\partial B} (-(N-1)f\psi + \nabla f \cdot \nabla \psi) \, d\mathcal{H}^{N-1} \\ &= \int_{\partial B} (H_{\partial E_f}(p) - (N-1))(1 + (N-1)f + R_1)\psi \, d\mathcal{H}^{N-1} \\ &\quad + \int_{\partial B} (-h \cdot \nabla \psi + (O(|f|^2) + O(|\nabla f|^2))\psi) \, d\mathcal{H}^{N-1}. \end{aligned}$$

Observe that testing (1.10) with $\psi = 1$ and setting $R_2 = (N-1)f + R_1$, we get $R_2 = O(|f|)$ and

$$(1.11) \quad \int_{\partial B} (1 + R_2)(H_{\partial E_f}(p) - (N-1)) \, d\mathcal{H}^{N-1} = \int_{\partial B} (O(|f|) + O(|\nabla f|^2)) \, d\mathcal{H}^{N-1}.$$

In particular, for $F := (1 + R_2)H_{\partial E_f}$ and recalling that the overline denotes the average on ∂B , we infer that for every $\eta \in (0, \frac{1}{2})$, if δ is sufficiently small, we have that

$$\left| \overline{F} - (N-1) \int_{\partial B} (R_2 + 1) \right| \leq \eta$$

and, always assuming δ small enough,

$$\begin{aligned}
|\overline{H}_{\partial E_f} - (N-1)| &\leq |\overline{H}_{\partial E_f} - \overline{F}| + \left| \overline{F} - (N-1) \int_{\partial B} (R_2 + 1) \right| + (N-1) \left| \int_{\partial B} R_2 \right| \\
&\leq \left| \int_{\partial B} R_2 H_{\partial E_f} \right| + 2\eta \leq \\
&\leq \omega_N^{-1} \|R_2\|_2 \|H_{\partial E_f} - \overline{H}_{\partial E_f}\|_2 + |\overline{H}_{\partial E_f} - (N-1)| \int_{\partial B} |R_2| d\mathcal{H}^{N-1} + \\
&\quad + (N-1) \int_{\partial B} |R_2| d\mathcal{H}^{N-1} + 2\eta \\
&\leq \eta \|H_{\partial E_f} - \overline{H}_{\partial E_f}\|_2 + \eta |\overline{H}_{\partial E_f} - (N-1)| + 3\eta \\
&\leq \eta |\overline{H}_{\partial E_f} - (N-1)| + 4\eta,
\end{aligned}$$

where in the last inequality we have used $\|H_{\partial E_f} - \overline{H}_{\partial E_f}\|_2 \leq 1$ and that for small δ we have $\int_{\partial B} |R_2| \leq \frac{\eta}{N-1}$. In particular, there exists a constant λ such that

$$(1.12) \quad \overline{H}_{\partial E_f} = N-1 + \lambda, \quad |\lambda| \leq \frac{4\eta}{1-\eta}.$$

Note that λ is arbitrarily small, if η (and hence δ) is chosen accordingly.

Now the proof can be concluded as follows. Consider $\kappa = 1 + \frac{\lambda}{N-1}$. If δ is small enough, then $\kappa \in (\frac{1}{2}, 2)$. Consider also the set $\kappa E_f = E_u$ with $u := \kappa - 1 + \kappa f$. Then,

$$\overline{H}_{\partial E_u} = N-1, \quad \|u\|_{H^1} \leq \kappa \|f\|_{H^1} + \sqrt{N\omega_N} \frac{\lambda}{N-1}, \quad \text{and } \|u\|_{C^1} \leq \kappa \|f\|_{C^1} + \frac{\lambda}{N-1}.$$

In particular, if δ and η are sufficiently small (and, therefore, such is λ), we are in position to apply Theorem 1.1 (because $\|u\|_{C^1}$ becomes arbitrarily small) and infer that

$$\|u\|_{H^1} \leq C \|H_{\partial E_u} - N + 1\|_{L^2} = \kappa^{-1} C \|H_{\partial E_f} - \overline{H}_{\partial E_f}\|_{L^2} \leq 2C \|H_{\partial E_f} - \overline{H}_{\partial E_f}\|_{L^2}.$$

In particular, we can estimate the L^2 norm of the gradient of f :

$$(1.13) \quad \|\nabla f\|_{L^2} \leq \kappa^{-1} \|u\|_{H^1} \leq 4C \|H_{\partial E_f} - \overline{H}_{\partial E_f}\|_{L^2}.$$

Finally, since $|E_f| = |B|$, i.e.,

$$\frac{1}{N} \int_{\partial B} (1+f)^N d\mathcal{H}^{N-1} = \omega_N,$$

we get by Taylor expansion

$$(1.14) \quad \left| \int_{\partial B} f d\mathcal{H}^{N-1} \right| = \int_{\partial B} O(|f|^2) d\mathcal{H}^{N-1}.$$

This implies that

$$(1.15) \quad |\overline{f}| \leq C \|f\|_{L^2}^2 \leq C\delta \|f\|_{L^2}.$$

By Poincaré inequality we have that

$$(1.16) \quad \|f\|_{L^2} \leq 2\|f - \overline{f}\|_{L^2} + 2|\overline{f}|(N\omega_N)^{\frac{1}{2}} \leq C\|\nabla f\|_{L^2} + 2|\overline{f}|(N\omega_N)^{\frac{1}{2}}.$$

Inserting (1.15) in (1.16) and combining with (1.13) we deduce (1.2) if δ is sufficiently small. \square

Remark 1.4. *By a simple rescaling argument it is clear that Theorem 1.3 holds also for the sets $E_{\varphi,m}$ parametrized over a ball $B^{(m)}$ with volume m , with constants δ and C depending on m . Clearly, the dependence of such constants on m can be made uniform when m varies in compact subsets of $(0, +\infty)$.*

Remark 1.5. *The estimate (1.2) is optimal for what concerns the power of the norms. To see this, it is enough to consider sets $E_{\varepsilon f}$ with f a functions in the second eigenspace of the Laplace-Beltrami operator on the sphere (with corresponding eigenvalue $2N$) and $\varepsilon > 0$ sufficiently small: then, computing the L^2 -norm of the average of the mean curvature yields*

$$\|H_{\partial E_{\varepsilon f}} - \overline{H}_{\partial E_{\varepsilon f}}\|_{L^2(\partial B)} \leq C \|\varepsilon f\|_{L^2(\partial B)},$$

for a dimensional constant $C > 0$.

2. THE INCREMENTAL PROBLEM

We start by introducing the incremental minimum problem which defines the discrete-in-time volume preserving mean curvature flow. To this purpose, let $E \neq \emptyset$ be a bounded measurable set. Notice that the topological boundary of E depends on its representative; from now on, we will assume that E coincides with its Lebesgue representative, i.e., with its points of density equal to one. We let

$$(2.1) \quad d_E(x) = \text{dist}(x, E) - \text{dist}(x, \mathbb{R}^N \setminus E)$$

be the signed distance function to ∂E . Let $m > 0$ be fixed, representing the volume of the evolving set.

Fix a time step $h > 0$ and consider the problem

$$(2.2) \quad \min \left\{ P(F) + \frac{1}{h} \int_F d_E(x) dx : |F| = m \right\}.$$

Note that

$$\int_F d_E(x) dx - \int_E d_E(x) dx = \int_{E \Delta F} \text{dist}(x, \partial E) dx$$

so that (2.2) is equivalent to

$$\min \left\{ P(F) + \frac{1}{h} \int_{E \Delta F} \text{dist}(x, \partial E) dx : |F| = m \right\}.$$

Given two sets E, F , we let

$$\mathcal{D}(F, E) := \int_{E \Delta F} \text{dist}(x, \partial E) dx.$$

In order to prove the existence of a solution to (2.2), we need some preliminary results. We start with the following non-vanishing estimate for sets of finite perimeter and finite measure.

Lemma 2.1. *There exists a constant $C = C(N) \in (0, \frac{1}{2})$ such that if $E \subset \mathbb{R}^N$ is a set of finite perimeter and finite measure, then, setting $Q := (0, 1)^N$, we have*

$$\sup_{z \in \mathbb{Z}^N} |E \cap (z + Q)| \geq c(N) \min \left\{ \left(\frac{|E|}{P(E)} \right)^N, 1 \right\}.$$

Proof. Setting

$$\beta := \sup_{z \in \mathbb{Z}^N} |E \cap (z + Q)|,$$

assume that $\beta \leq \frac{1}{2}$. Then, by the Relative Isoperimetric Inequality we have

$$\begin{aligned} P(E) &\geq \sum_{z \in \mathbb{Z}^N} P(E, z + Q) \geq c(N) \sum_{z \in \mathbb{Z}^N} |E \cap (z + Q)|^{\frac{N-1}{N}} \\ &\geq \frac{c(N)}{\beta^{\frac{1}{N}}} \sum_{z \in \mathbb{Z}^N} |E \cap (z + Q)| = \frac{c(N)}{\beta^{\frac{1}{N}}} |E| \end{aligned}$$

and the conclusion easily follows. \square

We are now in a position to prove the following proposition, which in particular establishes the existence of a solution to (2.2). The crucial point in the following statement is that the choice of the penalization parameter Λ depends only on the bounds on the perimeter and on the prescribed measure m , and thus can be made uniform along the minimizing movements scheme.

Proposition 2.2. *Given $m, M > 0$, there exists $\Lambda_0 = \Lambda_0(m, M, h, N) > 0$ such that for any bounded set E of finite perimeter, with $P(E) \leq M$ and $|E| = m$, and for any $\Lambda \geq \Lambda_0$, any solution \bar{F} of*

$$(2.3) \quad \min \left\{ P(F) + \frac{1}{h} \mathcal{D}(F, E) + \Lambda ||F| - m| : F \subset \mathbb{R}^N \text{ measurable} \right\}$$

satisfies the volume constraint $|\bar{F}| = m$. In particular, (2.2) and (2.3) are equivalent.

Proof. First of all, note that the existence of a solution to (2.3) follows by standard arguments, see for instance [21]. We argue by contradiction by assuming for every n the existence of a set E_n , with $|E_n| = m$ and $P(E_n) \leq M$, and a set

$$\bar{F}_n \in \operatorname{argmin} \left\{ P(F) + \frac{1}{h} \mathcal{D}(F, E_n) + n ||F| - m| : F \subset \mathbb{R}^N \text{ measurable} \right\}$$

such that $|\bar{F}_n| \neq m$. In the sequel, we may assume $|\bar{F}_n| < m$, as the other case can be treated analogously. Testing with E_n , by minimality we have

$$(2.4) \quad P(\bar{F}_n) + \frac{1}{h} \mathcal{D}(\bar{F}_n, E_n) + n ||\bar{F}_n| - m| \leq P(E_n) \leq M.$$

In particular,

$$(2.5) \quad |\bar{F}_n| \rightarrow m$$

as $n \rightarrow \infty$. In turn, by Lemma 2.1 there exist a constant $c_0 > 0$ (depending only on m, M and N) and $z_n \in \mathbb{Z}^N$ such that

$$|\bar{F}_n \cap (z_n + Q)| \geq c_0$$

for all n sufficiently large. Thus, by replacing \bar{F}_n, E_n by $\bar{F}_n - z_n, E_n - z_n$, respectively, and appealing to the well-known compactness properties of sets of finite perimeter, we may assume that up to a (not relabelled) subsequence we have $\bar{F}_n \rightarrow F_\infty$ in L^1_{loc} , with $|F_\infty| \geq |F_\infty \cap Q| \geq c_0 > 0$.

The idea now is to modify the sets \bar{F}_n by applying a local dilation in order to impose the volume constraint. In this construction, we follow closely [12, Section 2]. We only outline the main steps.

First of all, arguing as in Step 1 of [12, Section 2], for any fixed $y_0 \in \partial^* F_\infty$ and for any given small $\varepsilon > 0$ we may find a radius $r > 0$ and a point x_0 in a small neighborhood of y_0 such that

$$|\bar{F}_n \cap B_{r/2}(x_0)| < \varepsilon r^N, \quad |\bar{F}_n \cap B_r(x_0)| > \frac{\omega_N r^N}{2^{N+2}}$$

for all n sufficiently large. In the following, to simplify the notation we assume that $x_0 = 0$ and we write B_r instead of $B_r(0)$. For a sequence $0 < \sigma_n < 1/2^N$ to be chosen, we introduce the following bilipschitz maps:

$$\Phi_n(x) := \begin{cases} (1 - \sigma_n(2^N - 1))x & \text{if } |x| \leq \frac{r}{2}, \\ x + \sigma_n \left(1 - \frac{r^N}{|x|^N}\right)x & \frac{r}{2} \leq |x| < r, \\ x & |x| \geq r. \end{cases}$$

Setting $\tilde{F}_n := \Phi_n(\bar{F}_n)$, we have as in Step 3 of [12, Section 2]

$$(2.6) \quad P(\bar{F}_n, B_r) - P(\tilde{F}_n, B_r) \geq -2^N N P(\bar{F}_n, B_r) \sigma_n \geq -2^N N M \sigma_n.$$

Moreover, as in Step 4 of [12, Section 2] we have

$$|\tilde{F}_n| - |\bar{F}_n| \geq \sigma_n r^N \left[c \frac{\omega_N}{2^{N+2}} - \varepsilon(c + (2^N - 1)N) \right]$$

for a suitable constant c depending only on the dimension N . If we fix ε (and thus r) so that the negative term in the square bracket does not exceed half the positive one, then we have

$$(2.7) \quad |\tilde{F}_n| - |\bar{F}_n| \geq \sigma_n r^N C_1,$$

with $C_1 > 0$ depending on N . In particular, from this inequality it is clear that, by a continuity argument, we can choose σ_n so that $|\tilde{F}_n| = m$ for n large; this implies $\sigma_n \rightarrow 0$, thanks to (2.5).

Now, arguing as in [1, Equations (2.12) and (2.13)], we obtain

$$(2.8) \quad |\tilde{F}_n \Delta \bar{F}_n| \leq C_3 \sigma_n P(\bar{F}_n, B_r) \leq C_3 \sigma_n M.$$

Set now

$$i_n := \min_{\bar{B}_r} \text{dist}(\cdot, \partial E_n)$$

and note that if $i_n > 0$, then either $B_r \subset E_n$ or $B_r \subset E_n^c$. In the first case, we have $(E_n \Delta \bar{F}_n) \cap B_r = B_r \setminus \bar{F}_n$ and, in turn,

$$|B_r \setminus \bar{F}_n| \geq |B_{r/2} \setminus \bar{F}_n| = |B_{r/2}| - |\bar{F}_n \cap B_{r/2}| > \left(\frac{\omega_N}{2^N} - \varepsilon \right) r^N \geq \frac{\omega_N}{4^N} r^N,$$

by choosing ε smaller if needed. In turn, recalling (2.4), we may estimate

$$M \geq \frac{1}{h} \mathcal{D}(\bar{F}_n, E_n) \geq \frac{1}{h} \int_{B_r \setminus \bar{F}_n} \text{dist}(x, \partial E_n) dx \geq \frac{i_n}{h} \frac{\omega_N}{4^N} r^N,$$

from which we easily deduce

$$\text{dist}(\cdot, \partial E_n) \leq 4^N r^{-N} M \omega_N^{-1} h + 2r \quad \text{on } B_r.$$

Arguing in a similar way also in the case $B_r \subset E_n^c$, we can finally conclude the existence of a positive constant $C_4 = C_4(r)$ (depending also on N and h) such that, in all cases, and for n large enough we have $\text{dist}(\cdot, \partial E_n) \leq h C_4$ on B_r and thus

$$(2.9) \quad \left| \frac{1}{h} \mathcal{D}(\tilde{F}_n, E_n) - \frac{1}{h} \mathcal{D}(\bar{F}_n, E_n) \right| \leq C_4 |\tilde{F}_n \Delta \bar{F}_n| \leq C_4 C_3 \sigma_n M,$$

where in last inequality we have used (2.8). Combining (2.6), (2.7), and (2.9), we conclude that for n sufficiently large

$$\begin{aligned} P(\tilde{F}_n) + \frac{1}{h} \mathcal{D}(\tilde{F}_n, E_n) &\leq P(\bar{F}_n) + \frac{1}{h} \mathcal{D}(\bar{F}_n, E_n) + n \left| |\bar{F}_n| - m \right| \\ &\quad + \sigma_n \left[(2^N N + C_4 C_3) M - n r^N C_1 \right] \\ &< P(\bar{F}_n) + \frac{1}{h} \mathcal{D}(\bar{F}_n, E_n) + n \left| |\bar{F}_n| - m \right|, \end{aligned}$$

a contradiction to the minimality of \bar{F}_n , since $|\tilde{F}_n| = m$. \square

As a consequence of the previous proposition, together with some standard arguments from the regularity theory of almost minimal sets, we have the following:

Proposition 2.3 (Regularity properties of minimizers). *Let E be a bounded set with $|E| = m$ and $P(E) \leq M$ for some $m, M > 0$. Then, any solution $F \subset \mathbb{R}^N$ to (2.2) satisfies the following regularity properties:*

- i) *There exist $c_0 = c_0(N) > 0$ and a radius $r_0 = r_0(m, M, h, N) > 0$ such that for every $x \in \partial^* F$ and $r \in (0, r_0]$ we have*

$$(2.10) \quad |B_r(x) \cap F| \geq c_0 r^N \quad \text{and} \quad |B_r(x) \setminus F| \geq c_0 r^N.$$

In particular, F admits an open representative whose topological boundary coincides with the closure of its reduced boundary, i.e., $\partial F = \overline{\partial^ F}$. From now on we will always assume that F coincides with its open representative.*

- ii) *There exists $c_1 = c_1(m, M, h, N) > 0$ such that*

$$(2.11) \quad \sup_{E \Delta F} \text{dist}(\cdot, \partial E) \leq c_1.$$

- iii) *There exists $\bar{\Lambda} = \bar{\Lambda}(m, M, h, N) > 0$ such that F is a $\bar{\Lambda}$ -minimizer of the perimeter, that is,*

$$(2.12) \quad P(F) \leq P(F') + \bar{\Lambda} |F \Delta F'|$$

for all measurable set $F' \subset \mathbb{R}^N$ such that $\text{diam}(F \Delta F') \leq 1$.

- iv) *The following Euler-Lagrange condition holds: There exists $\lambda \in \mathbb{R}$ such that for all $X \in C_c^1(\mathbb{R}^N; \mathbb{R}^N)$*

$$(2.13) \quad \int_{\partial^* F} \frac{d_E(x)}{h} X \cdot \nu_F d\mathcal{H}^{N-1} + \int_{\partial^* F} \text{div}_\tau X d\mathcal{H}^{N-1} = \lambda \int_{\partial^* F} X \cdot \nu_F d\mathcal{H}^{N-1}.$$

- v) *There exists a closed set Σ whose Hausdorff dimension is less than or equal to $N - 8$, such that $\partial^* F = \partial F \setminus \Sigma$ is an $(N - 1)$ -submanifold of class $C^{2,\alpha}$ for all $\alpha \in (0, 1)$ with*

$$|H_{\partial F}(x)| \leq \bar{\Lambda} \quad \text{for all } x \in \partial F \setminus \Sigma.$$

- vi) *The set F is bounded and more precisely there exist $k_0 = k_0(m, M, h, N) \in \mathbb{N}$ and $d_0 = d_0(m, M, h, N) > 0$ such that F is made up of at most k_0 connected components, each one having diameter bounded from above by d_0 (a bound on the diameter from below follows from i)).*

Proof. By Proposition 2.2, there exists $\Lambda_0 = \Lambda_0(m, M, h, N)$ such that F is a solution to

$$(2.14) \quad \left\{ P(F) + \frac{1}{h} \mathcal{D}(F, E) + \Lambda_0 | |F| - m | : F \subset \mathbb{R}^N \text{ measurable} \right\}.$$

The density estimates and (2.11) follow arguing as in [21, Lemma 4.4 and Proposition 3.2], respectively. Now, (2.12) easily follows from the fact that F solves (2.14), taking into account (2.11).

Equation (2.13) can be derived by a standard first variation argument (see for instance [17]). In view of (iii) and the classical regularity theory for almost minimizers of the perimeter (see for instance [17] and the references therein), we deduce that there exists a closed set Σ whose Hausdorff dimension is less than or equal to $N - 8$, such that $\partial^* F = \partial F \setminus \Sigma$ is an $(N - 1)$ -submanifold of

class $C^{1,\alpha}$ for all $\alpha \in (0, 1)$. The $C^{2,\alpha}$ regularity stated in (v) follows now from the additional elliptic regularity implied by (2.13), taking into account that d_E is Lipschitz continuous.

Item (vi) follows rather easily from the density estimates. Indeed, let $r_0 = r_0(m, M, h, N) > 0$ be the radius given in i). By an application of Vitali's Covering Lemma, we may find a subset $C \subset F$ such that the closed balls of the family $\{\overline{B_{r_0}(x)}\}_{x \in C}$ are pairwise disjoint and

$$F \subset \bigcup_{x \in C} \overline{B_{5r_0}(x)}.$$

Since by (2.10) we have $|B_{r_0}(x) \cap F| \geq c_0 r_0^N$ for every $x \in C$, it follows that $\#C \leq \frac{|F|}{c_0 r_0^N} = \frac{m}{c_0 r_0^N}$. In turn, since for each connected component \hat{F} of F we have $\hat{F} \subset \cup_{x \in C} \overline{B_{5r_0}(x)}$, we infer that

$$\text{diam } \hat{F} \leq 10r_0 \#C \leq \frac{10m}{c_0 r_0^{N-1}} =: d_0(m, M, h, N).$$

It remains to bound the number of connected components. To this aim, it is enough to show that the measure of any connected component \hat{F} is bounded below by a positive constant depending only on m, M, h , and N . Set $F' := F \setminus \hat{F}$. Then, using (2.12) and the Isoperimetric Inequality, we deduce

$$\bar{\Lambda} |\hat{F}| = \bar{\Lambda} |F \Delta F'| + P(F') - (P(F) - P(\hat{F})) \geq P(\hat{F}) \geq N \omega_N^{\frac{1}{N}} |\hat{F}|^{\frac{N-1}{N}},$$

and thus

$$|\hat{F}| \geq \left(\frac{N}{\bar{\Lambda}}\right)^N \omega_N.$$

This concludes the proof of the proposition. \square

Remark 2.4. Note that (2.13) and the regularity properties given by Proposition 2.3 imply that

$$(2.15) \quad \frac{d_E(x)}{h} + H_{\partial F}(x) = \lambda \quad \text{for every } x \in \partial^* F.$$

3. THE DISCRETE VOLUME PRESERVING FLOW

3.1. Construction of the discrete flow. For any bounded set $E \neq \emptyset$ with finite perimeter we let $T_h E$ denote a solution of (2.2), with $m = |E|$. It is convenient to fix a precise representative for $T_h E$; as done for the set E , we assume that $T_h E$ coincides with its Lebesgue representative (i.e., the set of points of density 1). Now, we construct by induction the discrete-in-time evolution $\{E_h^n\}_{n \in \mathbb{N}}$. Let $E_h^1 := T_h E$ be a solution (arbitrarily chosen) to problem (2.2); assuming that E_h^k is defined for $k \in \{1, \dots, n-1\}$, we let E_h^n be a solution (arbitrarily chosen) to problem (2.2) with E replaced by E_h^{n-1} .

Remark 3.1. The sequence E_h^n defines a piecewise constant in time flow:

$$E_t^{(h)} := E_h^{\lfloor \frac{t}{h} \rfloor} \quad t \in [0, +\infty).$$

Any limit $E_t := \lim_{h_k \rightarrow 0} E_t^{(h_k)}$ of the discrete-in-time approximated flow along subsequences $h_k \rightarrow 0$ is referred to as volume preserving flat flow (see [21] for the convergence of a slightly different scheme). The flat flows can be thought of as weak solutions to the gradient flow of the area functional with a volume constraint.

3.2. Stationary sets for the discrete flow. In this section we characterize the stationary sets E for the volume preserving discrete flow, i.e., such that the constant sequence $E_h^n \equiv E$ is a discrete volume preserving flow.

Proposition 3.2 (Fixed points of the discrete scheme). *Given $m, M, h > 0$, there exists $s_0 = s_0(m, M, h, N) > 0$ such that every stationary bounded set E for the volume preserving discrete flow with time step h , with $|E| = m$ and $P(E) \leq M$ is made up by the union of k disjoint balls with mutual distances larger than s_0 and equal volume $\frac{m}{k}$, for some $k \leq k_0$, with k_0 as in item vi) of Proposition 2.3.*

Viceversa, if E is given by the union of finitely many disjoint balls with positive mutual distances and equal volumes, then there exists $h^ > 0$ such that, for all $h \leq h^*$, the volume preserving flow $\{E_h^n\}$ starting from E is unique and given by the constant sequence $E_h^n = E$ for all $n \in \mathbb{N}$.*

Proof. By (2.13) any stationary set E satisfies

$$\int_{\partial^* E} \operatorname{div}_\tau X \, d\mathcal{H}^{N-1} = \lambda \int_{\partial^* E} X \cdot \nu_E \, d\mathcal{H}^{N-1}$$

for all $X \in C_c^1(\mathbb{R}^N; \mathbb{R}^N)$ and for some $\lambda \in \mathbb{R}$. We may then apply [9, Theorem 1] to conclude that E is given (up to a negligible set) by a finite union of disjoint (open) balls of equal volume. Moreover, by (vi) of Proposition 2.3 the number of such balls is bounded by k_0 , and therefore, in view of (2.10), their radius is bounded from below by a constant R depending only on m, M, h, N . Now, if two balls of radius larger than or equal to R are at a distance s small enough, then E cannot satisfy the second density estimate in (2.10) (for example, (2.10) cannot hold for $x \in \partial E$ of minimal distance between the two balls and $r > 0$ depending on R and the constants r_0, c_0 in Proposition 2.3-i)). Therefore, (2.10) is violated, whenever $s \leq s_0$ for a suitable s_0 depending only on m, M, h, N , thus establishing the first part of the statement.

Assume now that E is union of finitely many disjoint balls with equal radius R and positive mutual distances. We want to show that, for h small enough, E is the unique volume constrained global minimizer of the functional

$$J_h(F) := P(F) + \frac{1}{h} \mathcal{D}(F, E) = P(F) + \frac{1}{h} \int_F d_E(x) \, dx - \frac{1}{h} \int_E d_E(x).$$

We start by showing that for h small enough the second variation of J_h is positive definite with respect to volume preserving variations. To this purpose, let $X \in C_c^1(\mathbb{R}^N; \mathbb{R}^N)$ be a divergence free vector field and let $\Phi(t, \cdot)$ be the associated flow satisfying $\frac{\partial \Phi}{\partial t} = X(\Phi)$ with initial condition $\Phi(0, x) = x$ for all $x \in \mathbb{R}^N$. Then $|\Phi(t, E)| = |E| = m$ for all t , and by standard computations (see for instance [1, 7]) we have

$$\begin{aligned} \frac{\partial^2}{\partial t^2} \Big|_{t=0} J_h(\Phi(t, E)) &= \int_{\partial E} |\nabla(X \cdot \nu_E)|^2 - \frac{N-1}{R^2} (X \cdot \nu_E)^2 \, d\mathcal{H}^{N-1} \\ &\quad + \frac{1}{h} \int_{\partial E} \partial_{\nu_E} d_E (X \cdot \nu_E)^2 \, d\mathcal{H}^{N-1} \\ &= \int_{\partial E} |\nabla(X \cdot \nu_E)|^2 + \left(\frac{1}{h} - \frac{N-1}{R^2} \right) (X \cdot \nu_E)^2 \, d\mathcal{H}^{N-1} \\ &=: \partial^2 J_h(E)[X \cdot \nu_E], \end{aligned}$$

where in the second equality we have used the fact that $\partial_{\nu_E} d_E \equiv 1$ on ∂E . From the above expression it is clear that $\partial^2 J_h(E)$ is positive definite on $H^1(\partial E)$, provided that $h < \frac{R^2}{N-1}$.

Fix $h_0 < \frac{R^2}{N-1}$. Arguing as in [1] there exists $\varepsilon > 0$ such that $J_{h_0}(E) < J_{h_0}(F)$ for all measurable F such that $|F| = |E|$ and $|E\Delta F| \leq \varepsilon$. Now notice that for all $0 < h < h_0$ we have

$$(3.1) \quad J_h(E) = J_{h_0}(E) < J_{h_0}(F) \leq J_h(F) \quad \text{for all } |F| = |E| \text{ and } 0 < |E\Delta F| \leq \varepsilon,$$

i.e., E is an isolated local minimizer for J_h in L^1 , with minimality neighborhood uniform with respect to $h \leq h_0$. Now, given any sequence $\{h_n\}$ going to zero, let F_n be a volume constrained minimizer of J_{h_n} ; it is easy to see that $|E\Delta F_n| \rightarrow 0$ as $n \rightarrow +\infty$, and therefore, for n large enough, $|E\Delta F_n| \leq \varepsilon$, so that by (3.1) $F_n = E$. □

Remark 3.3. *Let us also observe that for $N \leq 7$ we have full regularity of ∂E and thus, in particular, the connected components of E have positive mutual distances and the conclusion of the first part of Proposition 3.2 would also follow by applying the classical Alexandrov's Theorem instead of the more refined results of [9].*

3.3. Long-time behaviour. In the following, we denote by P_∞ the limit of the monotone non-increasing sequence $\{P(E_h^n)\}_{n \in \mathbb{N}}$:

$$(3.2) \quad P_\infty := \lim_{n \rightarrow \infty} P(E_h^n).$$

The following is the main result of the paper on the long time behaviour of the discrete-in-time nonlocal mean curvature flow.

Theorem 3.4. *Let E be a bounded set of finite perimeter with $|E| = m$ and let $h > 0$. Consider any discrete volume constrained mean curvature flow $\{E_h^n\}_{n \in \mathbb{N}}$ starting from E . Then, setting $L := N^{-N} \omega_N^{-1} m^{1-N} P_\infty^N \in \mathbb{N}$, where P_∞ is given in (3.2), there exist L distinct balls B^1, \dots, B^L with the same radius and at positive distance to each other, such that the sets E_h^n converge to $E_\infty := \bigcup_{i=1}^L B^i$ in C^ℓ for every $\ell \in \mathbb{N}$. Moreover, the convergence is exponentially fast.*

We recall that E_h^n converge to E_∞ in C^ℓ if for n large enough the boundaries of the sets E_h^n can be parametrized by normal graphs of C^ℓ functions on ∂E_∞ converging to zero in C^ℓ . We remark that the estimates provided by Theorem 3.4 on the speed of convergence of E_h^n to E_∞ depend on h and are not uniform as $h \rightarrow 0$, thus preventing us to pass to the limit of flat flows. The following is an immediate corollary.

Corollary 3.5. *If $P(E) < 2P(\bar{B})$, where \bar{B} is a ball with volume $\frac{m}{2}$, or if E is union of two tangent balls, then E_∞ is a ball.*

Proof. If $P(E) < 2P(\bar{B})$, then by definition $P_\infty < 2P(\bar{B})$ and $L < 2$. On the other hand, two tangent balls are not stationary for the discrete flow, as shown in Proposition 3.2. This implies that $P(E_h^n) < P(E) = 2P(\bar{B})$, so that one can argue as before. □

The rest of this section is devoted to the proof of Theorem 3.4. We start with the following lemma.

Lemma 3.6. *Let $\{E_h^n\}_{n \in \mathbb{N}}$ be a volume preserving discrete flow starting from E and let $E_h^{k_n}$ be a subsequence such that $\chi_{E_h^{k_n} - \tau_n} \rightarrow \chi_F$ in $L^1(\mathbb{R}^N)$ for some set F and a suitable sequence $(\tau_n)_{n \in \mathbb{N}} \subset \mathbb{R}^N$. Then $d_{E_h^{k_n-1}(\cdot + \tau_n)} \rightarrow d_F$ locally uniformly in \mathbb{R}^N .*

Proof. We start by observing that for every n , by the minimality of E_h^n , using E_h^{n-1} as a competitor, we have

$$(3.3) \quad \frac{1}{h} \mathcal{D}(E_h^n, E_h^{n-1}) \leq P(E_h^{n-1}) - P(E_h^n).$$

Therefore, summing over n we get

$$\sum_{n=1}^{\infty} \frac{1}{h} \mathcal{D}(E_h^n, E_h^{n-1}) \leq P(E).$$

In particular,

$$(3.4) \quad \mathcal{D}(E_h^n, E_h^{n-1}) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Passing to a further (not relabelled) subsequence we may assume that there exists G such that $E_h^{k_n-1} - \tau_n \rightarrow G$ in $L^1_{loc}(\mathbb{R}^N)$. In fact, by the density estimates (2.10) and standard arguments, we get

$$\text{dist}(\cdot, \partial E_h^{k_n-1} - \tau_n) \rightarrow \text{dist}(\cdot, \partial G) \quad \text{locally uniformly in } \mathbb{R}^N,$$

(which defines the *Kuratowski convergence* of the boundaries $\partial E_h^{k_n-1} - \tau_n$ to ∂G). Combining the latter information with (3.4), for every $R > 0$ we easily get

$$0 = \lim_{n \rightarrow \infty} \int_{(E_h^{k_n} \Delta E_h^{k_n-1} - \tau_n) \cap B_R} \text{dist}(x, \partial E_h^{k_n-1} - \tau_n) d\mathcal{H}^{N-1} = \int_{(F \Delta G) \cap B_R} \text{dist}(x, \partial G) d\mathcal{H}^{N-1}.$$

The last equality yields $F \Delta G \subset \partial G$. As G still satisfies the density estimates we have $|\partial G| = 0$ and the conclusion follows. \square

In the next proposition we start by showing that for any discrete volume preserving flow the evolving sets are eventually made up of a constant number L of connected components, each one converging up to translations to the same ball of volume m/L . After establishing such a result, it will remain to show that the magnitude of such translations decays to zero exponentially fast.

Proposition 3.7 (Long-time behavior up to translations). *Let $\{E_h^n\}_{n \in \mathbb{N}}$ be a discrete flat flow with time step $h > 0$ and prescribed volume $m > 0$. Let P_∞ and L be as in the statement of Theorem 3.4. Then, for n sufficiently large, E_h^n has L connected components $E_{h,1}^n, \dots, E_{h,L}^n$, such that $\text{dist}(E_{h,i}^n, E_{h,j}^n) \geq \frac{s_0}{2}$ for $i \neq j$ (with s_0 the constant of Proposition 3.2), and $E_{h,i}^n - \text{bar}(E_{h,i}^n)$ converge to the ball centered at the origin with volume $\frac{m}{L}$ in C^ℓ for every $\ell \in \mathbb{N}$.*

Proof. It suffices to prove that, given any subsequence $\{E_h^{k_n}\}_{n \in \mathbb{N}}$, there exists a sub-subsequence (not relabelled) satisfying the conclusions of the proposition.

To this purpose, let $\{E_h^{k_n}\}_{n \in \mathbb{N}}$ be a given subsequence. By Proposition 2.3-vi), each set $E_h^{k_n}$ is made up of $l_n \leq k_0$ connected components with uniformly bounded diameter less than or equal to d_0 . Therefore, there exist l_n balls $\{B_{d_0}(\xi_i^n)\}_{i=1, \dots, l_n}$ (not necessarily disjoint one from the other), each containing a different component of $E_h^{k_n}$ and such that $E_h^{k_n} \subset \cup_{i=1}^{l_n} B_{d_0}(\xi_i^n)$. Up to passing to a subsequence (not relabelled), we can assume that $l_n = \tilde{l}$, and for all $1 \leq i < j \leq \tilde{l}$ the following limits exist

$$\lim_n |\xi_i^n - \xi_j^n| =: d_{i,j}.$$

Now we define the following equivalence classes: $i \equiv j$ if and only if $d_{i,j} < +\infty$. Denote by l the number of such equivalence classes, let $j(i)$ be a representative for each class $i \in \{1, \dots, l\}$, and set $\sigma_i^n := \xi_{j(i)}^n$ for $i = 1, \dots, l$. We have constructed a subsequence $E_h^{k_n}$ satisfying for n large $E_h^{k_n} \subset \cup_{i=1}^l B_R(\sigma_i^n)$, where $R = d_0 + \max\{d_{i,j} : d_{i,j} < \infty\} + 1$, and for all $i \neq j$ there holds $|\sigma_i^n - \sigma_j^n| \rightarrow +\infty$ as $n \rightarrow +\infty$. We remark that, while $E_h^{k_n-1}$ are in general not elements of the

subsequence $\{E_h^{k_n}\}_{n \in \mathbb{N}}$, they will still play a role in our arguments, since they will be involved in exploiting the minimality properties of $E_h^{k_n}$.

Now, fix $1 \leq i \leq l$, and set

$$F_i^n := (E_h^{k_n} - \sigma_i^n), \quad \tilde{F}_i^n := (E_h^{k_n} - \sigma_i^n) \cap B_R, \quad m_i^n := |\tilde{F}_i^n|.$$

Up to a subsequence, we have $m_i^n \rightarrow m_i$ for some $m_i > 0$. Moreover, by Lemma 3.6 and the compactness of sets of equi-bounded perimeters, there exist measurable sets $\tilde{F}_i \subset\subset B_R$ such that (again up to a subsequence)

$$(3.5) \quad \tilde{F}_i^n \rightarrow \tilde{F}_i \text{ in } L^1, \quad d_{E_h^{k_n-1}}(\cdot + \sigma_i^n) \rightarrow d_{\tilde{F}_i}(\cdot) \text{ locally uniformly.}$$

We shall show that \tilde{F}_i is stationary for the discrete volume preserving flow. To this aim, let \tilde{G}_i be any bounded set with $|\tilde{G}_i| = m_i$. We define the homotetically rescaled sets $\tilde{G}_i^n := \left(\frac{m_i^n}{m_i}\right)^{\frac{1}{N}} \tilde{G}_i$ such that $|\tilde{G}_i^n| = m_i^n$. Note that $\tilde{G}_i^n \rightarrow \tilde{G}_i$ in L^1 and $P(\tilde{G}_i^n) \rightarrow P(\tilde{G}_i)$ as $n \rightarrow +\infty$. We set now $G_i^n := F_i^n \cup \tilde{G}_i^n \setminus \tilde{F}_i^n$: notice that, since \tilde{G}_i^n is bounded and the connected components of $F_i^n \setminus \tilde{F}_i^n$ diverge, it follows that, for sufficiently large n , G_i^n is made up by the same connected components of F_i^n except those of \tilde{F}_i^n which is replaced by \tilde{G}_i^n ; in particular, $|F_i^n| = |G_i^n|$. By the minimality of $E_h^{k_n}$ we have

$$P(F_i^n) + \frac{1}{h} \int_{F_i^n} d_{E_h^{k_n-1}}(x + \sigma_i^n) dx \leq P(G_i^n) + \frac{1}{h} \int_{G_i^n} d_{E_h^{k_n-1}}(x + \sigma_i^n) dx.$$

Since the connected components of $F_i^n \setminus \tilde{F}_i^n$ diverge and the two sets F_i^n and G_i^n differ only by the components \tilde{F}_i^n and \tilde{G}_i^n , by additivity the previous inequality is equivalent to

$$(3.6) \quad P(\tilde{F}_i^n) + \frac{1}{h} \int_{\tilde{F}_i^n} d_{E_h^{k_n-1}}(x + \sigma_i^n) dx \leq P(\tilde{G}_i^n) + \frac{1}{h} \int_{\tilde{G}_i^n} d_{E_h^{k_n-1}}(x + \sigma_i^n) dx.$$

Passing to the limit as $n \rightarrow +\infty$, using (3.5) and the lower semicontinuity of the perimeter, since the sets \tilde{F}_i^n and \tilde{G}_i^n are uniformly bounded, we deduce that

$$(3.7) \quad P(\tilde{F}_i) + \frac{1}{h} \int_{\tilde{F}_i} d_{\tilde{F}_i}(x) dx \leq P(\tilde{G}_i) + \frac{1}{h} \int_{\tilde{G}_i} d_{\tilde{F}_i}(x) dx.$$

This minimality property extends by density to all competitors \tilde{G}_i with finite perimeter and the same volume m_i , so that we deduce that \tilde{F}_i is a fixed point for the discrete scheme with prescribed volume m_i , and whence by Lemma 3.2 \tilde{F}_i is given by the union of disjoint balls with positive mutual distances and equal volume. Moreover, since \tilde{F}_i^n are uniform $\bar{\Lambda}$ -minimal by Proposition 2.3, from the classical regularity theory (see [17]) we also deduce that \tilde{F}_i^n converge to \tilde{F}_i in $C^{1,\alpha}$ for every $\alpha \in (0,1)$. In particular, for n large enough, \tilde{F}_i^n has the same number of connected components of \tilde{F}_i .

Summarizing, we have shown that, for a subsequence (not relabelled) of $E_h^{k_n}$ and for n large enough, $E_h^{k_n}$ is made up by a fixed number K of connected components $E_{h,1}^{k_n}, \dots, E_{h,K}^{k_n}$, each converging to a ball (possibly with different radius, if the components belong to different equivalent classes according to the relation introduced above). Therefore, for $i \leq K$ we have $E_{h,i}^{k_n} - \text{bar}(E_{h,i}^{k_n}) \rightarrow B_{R_i}$, where B_{R_i} is the ball centered at the origin with radius $R_i > 0$.

It remains to show that all the radii R_i are equal to R : from this and the C^1 convergence of the translated components to B_R , it follows that $KP(B_R) = P_\infty$, i.e., $K = L$. To this aim, we

consider the Euler-Lagrange equation (2.13) for $E_h^{k_n}$: for every $X \in C_c^1(\mathbb{R}^n; \mathbb{R}^n)$,

$$\int_{\partial E_h^{k_n}} \frac{d_{E_h^{k_n-1}}(x)}{h} X \cdot \nu_{E_h^{k_n}} d\mathcal{H}^{N-1} + \int_{\partial E_h^{k_n}} \operatorname{div}_\tau X d\mathcal{H}^{N-1} = \lambda_n \int_{\partial E_h^{k_n}} X \cdot \nu_{E_h^{k_n}} d\mathcal{H}^{N-1}.$$

By Proposition 2.3-ii) and v), we deduce that

$$|\lambda_n| \leq h^{-1} \|d_{E_h^{k_n-1}}\|_{L^\infty(E_h^{k_n})} + \|H_{E_h^{k_n}}\|_{L^\infty(E_h^{k_n})} \leq h^{-1} c_1 + |\bar{\lambda}|.$$

Therefore, by passing to a further subsequence (not relabelled), we can assume that $\lambda_n \rightarrow \lambda$, for some $\lambda \in \mathbb{R}$. Arguing as before, we can localize the Euler-Lagrange equation to each single F_i^n :

$$\int_{\partial F_i^n} \frac{d_{E_h^{k_n-1}}(x + \sigma_i^n)}{h} X \cdot \nu_{F_i^n} d\mathcal{H}^{N-1} + \int_{\partial F_i^n} \operatorname{div}_\tau X d\mathcal{H}^{N-1} = \lambda_n \int_{\partial F_i^n} X \cdot \nu_{F_i^n} d\mathcal{H}^{N-1}.$$

Passing into the limit as $n \rightarrow \infty$ and taking into account Lemma 3.6, we deduce that

$$\int_{\partial \bar{F}_i} \operatorname{div}_\tau X d\mathcal{H}^{N-1} = \lambda \int_{\partial \bar{F}_i} X \cdot \nu_{\bar{F}_i} d\mathcal{H}^{N-1}.$$

In particular, this shows that $R_i = \frac{N-1}{\lambda}$.

Finally, the C^ℓ convergence follows by a classical bootstrap method. The idea is to describe the boundaries ∂F_n (up to changing the reference frame) locally as graphs of suitable functions $f_n : B' \rightarrow \mathbb{R}$, with B' a ball of \mathbb{R}^{N-1} . Then we have

$$\operatorname{div} \left(\frac{\nabla f_n}{\sqrt{1 + |\nabla f_n|^2}} \right) = H_n,$$

with H_n and ∇f_n uniformly bounded in $C^{0,\alpha}$ for every $\alpha \in (0, 1)$. Differentiating the equation in any direction v , we get

$$\operatorname{div}(A(\nabla f_n) \nabla(\partial_v f_n)) = \partial_v H_n,$$

where the coefficients $A(\nabla f_n)$ are uniformly elliptic and uniformly bounded in the $C^{0,\alpha}$ norm. We may therefore apply [13, Theorem 5.19] to conclude that the functions $\partial_v f_n$ are uniformly bounded in the $C^{1,\alpha}$ norm. In order to bound the higher order norms we may now apply a standard bootstrap argument and iteratively use [13, Theorem 5.19]. We leave the details to the reader. \square

We recall the notation introduced before (1.1): $B^{(\mu)}$ stands for the ball of volume μ and $r(\mu)$ denotes its radius.

Lemma 3.8. *Let $\mu > 0$ and $\eta > 0$. There exists $\bar{\delta} > 0$ with the following property: if $f_1, f_2 \in C^1(\partial B^{(\mu)})$ with $\|f_i\|_{C^1} \leq \bar{\delta}$ and $|E_{f_i, \mu}| = \mu$ for $i = 1, 2$ we have*

$$(3.8) \quad r(\mu)^2 (1 - \eta) \frac{\|f_1 - f_2\|_2^2}{2} \leq \mathcal{D}(E_{f_1, \mu}, E_{f_2, \mu}) \leq r(\mu)^2 (1 + \eta) \frac{\|f_1 - f_2\|_2^2}{2},$$

$$(3.9) \quad \frac{1 - \eta}{2} \int_{\partial E_{f_1, \mu}} d_{E_{f_2, \mu}}^2 d\mathcal{H}^{N-1} \leq \mathcal{D}(E_{f_1, \mu}, E_{f_2, \mu}) \leq \frac{1 + \eta}{2} \int_{\partial E_{f_1, \mu}} d_{E_{f_2, \mu}}^2 d\mathcal{H}^{N-1},$$

$$(3.10) \quad |\operatorname{bar}(E_{f_1, \mu}) - \operatorname{bar}(E_{f_2, \mu})| \leq C \sqrt{\mathcal{D}(E_{f_1, \mu}, E_{f_2, \mu})},$$

where $C > 0$ depends only on N and μ .

Proof. We start by observing that for any $\eta' > 0$, if $\bar{\delta}$ is sufficiently small, then for every $p_0 \in \partial E_{f_2, \mu}$ we have

$$(3.11) \quad \partial E_{f_2, \mu} \cap B_{4\bar{\delta}}(p_0) \subset G := \left\{ y \in \mathbb{R}^N : \left| (y - p_0) \cdot \frac{p_0}{|p_0|} \right|^2 \leq \frac{\eta'^2}{1 + \eta'^2} |y - p_0|^2 \right\}.$$

We divide the rest of the proof into two steps.

Step 1. If $\bar{\delta}$ is small enough, for every point $p = \lambda p_0 \in B_{2\bar{\delta}}(p_0)$ ($\lambda > 0$), we have that

$$\frac{1}{1 + \eta'} |p - p_0| \leq \text{dist}(p, \partial E_{f_2, \mu}) \leq |p - p_0|.$$

The second inequality is, indeed, obvious by definition, given that $p_0 \in \partial E_{f_2, \mu}$. Concerning the first one, we notice that $\text{dist}(p, \partial E_{f_2, \mu}) \leq |p - p_0| \leq 2\bar{\delta}$ implies that there exists a point $q \in \partial E_{f_2, \mu} \cap B_{2\bar{\delta}}(p)$ such that $\text{dist}(p, \partial E_{f_2, \mu}) = |p - q|$. In particular, from (3.11) we infer that $q \in G$ and hence we have

$$\text{dist}(p, \partial E_{f_2, \mu}) \geq \text{dist}(p, G) = \frac{1}{\sqrt{1 + \eta'^2}} |p - p_0| \geq \frac{1}{1 + \eta'} |p - p_0|.$$

In particular, if $p_0 = (1 + f_2(s))s \in \partial E_{f_2, \mu}$ with $s \in \partial B^{(\mu)}$ and

$$p_t := p_0 + t \frac{f_1(s) - f_2(s)}{|f_1(s) - f_2(s)|} \frac{s}{|s|} \quad \text{for all } t \in [0, r(\mu)|f_1(s) - f_2(s)|],$$

we deduce that

$$(3.12) \quad \frac{1}{1 + \eta'} t \leq \text{dist}(p_t, \partial E_{f_2, \mu}) \leq t.$$

Then, keeping this same notation and integrating in polar coordinates, we infer that

$$(3.13) \quad \begin{aligned} \mathcal{D}(E_{f_1, \mu}, E_{f_2, \mu}) &= \int_{E_{f_1, \mu} \Delta E_{f_2, \mu}} \text{dist}(x, \partial E_{f_2, \mu}) dx \\ &= \int_{\partial B^{(\mu)}} ds \int_0^{r(\mu)|f_2(s) - f_1(s)|} \text{dist}(p_t, \partial E_{f_2, \mu}) \left(\frac{|p_t|}{r(\mu)} \right)^{N-1} dt. \end{aligned}$$

Recalling that $\left| \frac{|p_t|}{r(\mu)} - 1 \right| \leq \bar{\delta}$ and using (3.12), we get

$$(3.14) \quad \begin{aligned} \mathcal{D}(E_{f_1, \mu}, E_{f_2, \mu}) &\leq (1 + \bar{\delta})^{N-1} \int_{\partial B^{(\mu)}} ds \int_0^{r(\mu)|f_2(s) - f_1(s)|} t dt \\ &= \frac{(1 + \bar{\delta})^{N-1}}{2} r(\mu)^2 \int_{\partial B^{(\mu)}} |f_1(s) - f_2(s)|^2 ds, \end{aligned}$$

from which the second inequality in (3.8) follows by taking $\bar{\delta}$ smaller, if needed. Analogously,

$$(3.15) \quad \begin{aligned} \mathcal{D}(E_{f_1, \mu}, E_{f_2, \mu}) &\geq \frac{(1 - \bar{\delta})^{N-1}}{1 + \eta'} \int_{\partial B^{(\mu)}} ds \int_0^{r(\mu)|f_2(s) - f_1(s)|} t dt \\ &= \frac{(1 - \bar{\delta})^{N-1}}{2(1 + \eta')} r(\mu)^2 \int_{\partial B^{(\mu)}} |f_1(s) - f_2(s)|^2 ds, \end{aligned}$$

from which the first inequality in (3.8) follows by taking η' and $\bar{\delta}$ small enough.

Step 2. The inequalities (3.9) and (3.10) are now easy consequences. Indeed, by (3.12) we have that for every $x = (1 + f_1(s))s \in \partial E_{f_1, \mu}$

$$\frac{r(\mu)}{1 + \eta'} |f_1(s) - f_2(s)| \leq |d_{E_{f_2, \mu}}(x)| \leq r(\mu) |f_1(s) - f_2(s)|.$$

Therefore, (3.9) follows from (3.14) and (3.15), by taking η' and $\bar{\delta}$ smaller if needed, through a simple change of coordinates (recall that the Jacobian of the map $s \mapsto (1 + f_1(s))s$ and its inverse are estimated from above by $1 + C\bar{\delta}$ for a suitable dimensional constant C).

Finally, note that we can write

$$\text{bar}(E_{f_i, \mu}) = \frac{1}{(N + 1)r(\mu)^N \omega_N} \int_{\partial B^{(\mu)}} (1 + f_i(s))^{N+1} s d\mathcal{H}^{N-1}(s).$$

Using the fact that $t \mapsto (1+t)^{N+1}$ is $2(N+1)$ -Lipschitz for t small, we may estimate

$$\begin{aligned} & |\text{bar}(E_{f_1, \mu}) - \text{bar}(E_{f_2, \mu})| \\ & \leq \frac{1}{(N+1)r(\mu)^{N-1}\omega_N} \left| \int_{\partial B(\mu)} ((1+f_1(s))^{N+1} - (1+f_2(s))^{N+1}) d\mathcal{H}^{N-1}(s) \right| \\ & \leq \frac{2}{r(\mu)^{N-1}\omega_N} \int_{\partial B(\mu)} |f_1(s) - f_2(s)| d\mathcal{H}^{N-1}(s) \leq C \|f_1 - f_2\|_2, \end{aligned}$$

where $C > 0$ depends only on N and μ . Thus (3.10) follows from the previous inequality combined with (3.8). \square

Lemma 3.9. *Let $h > 0$, $\mu > 0$. There exists $C = C(h, \mu) > 0$ and $\bar{\delta} = \bar{\delta}(h, \mu) > 0$ with the following property: For any pair of sets $E_{f_1, \mu}$, $E_{f_2, \mu}$ with $f_1, f_2 \in C^2(\partial B(\mu))$, $\|f_i\|_{C^1} \leq \bar{\delta}$, and such that $|E_{f_2, \mu}| = \mu$, $\text{bar}(E_{f_2, \mu}) = 0$ and*

$$(3.16) \quad H_{\partial E_{f_2, \mu}} + \frac{d_{E_{f_1, \mu}}}{h} = \lambda \quad \text{on } \partial E_{f_2, \mu}$$

for some $\lambda \in \mathbb{R}$, we have

$$(3.17) \quad \mathcal{D}(B(\mu), E_{f_2, \mu}) \leq C \mathcal{D}(E_{f_2, \mu}, E_{f_1, \mu}).$$

Proof. By Theorem 1.3 (and Remark 1.4), by choosing $\bar{\delta}$ sufficiently small and using also (3.16), we have

$$(3.18) \quad \begin{aligned} \|f_2\|_{L^2(\partial B(\mu))}^2 & \leq C(\mu) \|H_{\partial E_{f_2, \mu}} - \bar{H}_{\partial E_{f_2, \mu}}\|_{L^2(\partial B(\mu))}^2 \leq C(\mu) \|H_{\partial E_{f_2, \mu}} - \lambda\|_{L^2(\partial B(\mu))}^2 \\ & \leq 2C(\mu) \|H_{\partial E_{f_2, \mu}} - \lambda\|_{L^2(\partial E_{f_2, \mu})}^2 = \frac{2C(\mu)}{h^2} \int_{\partial E_{f_2, \mu}} d_{E_{f_1, \mu}}^2 d\mathcal{H}^{N-1}, \end{aligned}$$

where the third inequality follows by bounding the Jacobian of the change of variables by 2 (which can be done provided $\bar{\delta}$ is small enough). Note now that by (3.8) (with f_1 replaced by 0) and by (3.9) (with the role f_1 and f_2 interchanged), by taking $\bar{\delta}$ smaller if needed, we have

$$\mathcal{D}(B(\mu), E_{f_2, \mu}) \leq \|f_2\|_{L^2(\partial B(\mu))}^2 \quad \text{and} \quad \int_{\partial E_{f_2, \mu}} d_{E_{f_1, \mu}}^2 d\mathcal{H}^{N-1} \leq 4\mathcal{D}(E_{f_2, \mu}, E_{f_1, \mu}).$$

Combining the previous inequalities with (3.18), the conclusion follows. \square

Remark 3.10. *It is clear that the constants C and $\bar{\delta}$ in Lemmas 3.8 and 3.9 are uniform with respect to μ varying on any compact subset of $(0, +\infty)$.*

We are now ready to prove the main result of the paper. The main difficulty is in controlling the translations introduced in Proposition 3.7 and in proving the convergence of the barycenters. A crucial role in such an argument is played by the dissipation/dissipation inequality (3.17), which in turn relies on the quantitative Alexandrov type estimate established in Theorem 1.3.

Proof of Theorem 3.4. We split the proof into several steps.

Step 1. (Exponential decay of dissipations) Recall that by Proposition 3.7 we have

$$P(E_h^n) \rightarrow LP\left(B\left(\frac{m}{L}\right)\right).$$

Thus, summing (3.3) from $n+1 \in \mathbb{N}$ to $+\infty$, we get

$$(3.19) \quad \sum_{k=n+1}^{+\infty} \frac{1}{h} \mathcal{D}(E_h^k, E_h^{k-1}) \leq P(E_h^n) - LP\left(B\left(\frac{m}{L}\right)\right).$$

Recall that again by Proposition 3.7 for n large enough each set E_h^n is made up of L connected components $E_{h,1}^n, \dots, E_{h,L}^n$,

$$(3.20) \quad m_i^n := |E_{h,i}^n| \rightarrow \frac{m}{L}$$

and

$$(3.21) \quad E_{h,i}^n - \xi_i^n \rightarrow B\left(\frac{m}{L}\right) \quad \text{in } C^\ell$$

as $n \rightarrow \infty$, where we set

$$\xi_i^n := \text{bar}(E_{h,i}^n).$$

With Lemma 3.6 in mind, we also get

$$(3.22) \quad E_{h,i}^{n-1} - \xi_i^n \rightarrow B\left(\frac{m}{L}\right) \quad \text{in } C^\ell$$

as $n \rightarrow \infty$. Combining (3.21) and (3.22), we have that for any $\ell \in \mathbb{N}$ and for n large enough there exist functions $f_{1,i}^n, f_{2,i}^n \in C^\ell(\partial B^{(m_i^n)})$ such that (with the notation introduced in (1.1))

$$(3.23) \quad E_{h,i}^n - \xi_i^n = E_{f_{2,i}^n, m_i^n}, \quad E_{h,i}^{n-1} - \xi_i^n = E_{f_{1,i}^n, m_i^n} \quad \text{with } \|f_{1,i}^n\|_{C^\ell}, \|f_{2,i}^n\|_{C^\ell} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Moreover, again by Proposition 3.7 for n large enough we have

$$(3.24) \quad \text{dist}(E_{h,i}^n, E_{h,j}^n) \geq \frac{s_0}{2}, \quad \text{for } i \neq j,$$

with s_0 the constant of Proposition 3.2. Consider now the the admissible competitor for E_h^n given by

$$\mathfrak{B}_n := \bigcup_{i=1}^L \left(\xi_i^{n-1} + B^{(m_i^{n-1})} \right),$$

and note that by (3.20) and (3.21) we also have

$$\text{dist}\left(\xi_i^{n-1} + B^{(m_i^{n-1})}, \xi_j^{n-1} + B^{(m_j^{n-1})}\right) \geq \frac{s_0}{4}$$

for n large enough and $i \neq j$. The above inequality and (3.24), together once again with (3.21) and (3.22), in turn yield that

$$(3.25) \quad \begin{aligned} \mathcal{D}(E_h^n, E_h^{n-1}) &= \sum_{i=1}^L \mathcal{D}(E_{h,i}^n, E_{h,i}^{n-1}) \text{ and} \\ \mathcal{D}(\mathfrak{B}_n, E_h^{n-1}) &= \sum_{i=1}^L \mathcal{D}\left(B^{(m_i^{n-1})}, E_{h,i}^{n-1} - \xi_i^{n-1}\right). \end{aligned}$$

Testing the minimality of $E_{h,i}^n$ with \mathfrak{B}_n and using the second identity in (3.25), we have

$$(3.26) \quad P(E_h^n) + \frac{1}{h} \mathcal{D}(E_h^n, E_h^{n-1}) \leq P(\mathfrak{B}_n) + \frac{1}{h} \sum_{i=1}^L \mathcal{D}\left(B^{(m_i^{n-1})}, E_{h,i}^{n-1} - \xi_i^{n-1}\right).$$

Recall now that by (3.23) and by Lemma 3.9 (see also Remark 3.10) for n large enough we have

$$\begin{aligned} \mathcal{D}\left(B^{(m_i^{n-1})}, E_{h,i}^{n-1} - \xi_i^{n-1}\right) &= \mathcal{D}\left(B^{(m_i^{n-1})}, E_{f_{2,i}^{n-1}, m_i^{n-1}}\right) \\ &\leq C \mathcal{D}\left(E_{f_{2,i}^{n-1}, m_i^{n-1}}, E_{f_{1,i}^{n-1}, m_i^{n-1}}\right) = C \mathcal{D}(E_{h,i}^{n-1}, E_{h,i}^{n-2}). \end{aligned}$$

Thus, from (3.26) and (3.25) we deduce that

$$(3.27) \quad P(E_h^n) - P(\mathfrak{B}_n) \leq \frac{C}{h} \sum_{i=1}^L \mathcal{D}(E_{h,i}^{n-1}, E_{h,i}^{n-2}) = \frac{C}{h} \mathcal{D}(E_h^{n-1}, E_h^{n-2}).$$

Observe now that by concavity

$$\sum_{i=1}^L m_i^{\frac{N-1}{N}} \leq L \left(\frac{m}{L} \right)^{\frac{N-1}{N}} \quad \text{for all } m_1, \dots, m_L \geq 0 \text{ s.t. } \sum_{i=1}^L m_i = m$$

and thus

$$P(\mathfrak{B}_n) \leq LP(B(\frac{m}{L})).$$

Therefore, from (3.27) and (3.19) we get

$$\begin{aligned} & \sum_{k=n-1}^{+\infty} \frac{1}{h} \mathcal{D}(E_h^k, E_h^{k-1}) \\ &= \sum_{k=n+1}^{+\infty} \frac{1}{h} \mathcal{D}(E_h^k, E_h^{k-1}) + \frac{1}{h} \mathcal{D}(E_h^{n-1}, E_h^{n-2}) + \frac{1}{h} \mathcal{D}(E_h^n, E_h^{n-1}) \\ &\leq P(E_h^n) - LP(B(\frac{m}{L})) + \frac{1}{h} \mathcal{D}(E_h^{n-1}, E_h^{n-2}) + \frac{1}{h} \mathcal{D}(E_h^n, E_h^{n-1}) \\ &\leq \frac{C+1}{h} \mathcal{D}(E_h^{n-1}, E_h^{n-2}) + \frac{1}{h} \mathcal{D}(E_h^n, E_h^{n-1}) \leq \frac{C+1}{h} \left(\mathcal{D}(E_h^{n-1}, E_h^{n-2}) + \mathcal{D}(E_h^n, E_h^{n-1}) \right). \end{aligned}$$

We may now apply Lemma 3.11 (with $\ell = 2$) below to conclude

$$(3.28) \quad \mathcal{D}(E_h^n, E_h^{n-1}) \leq \left(1 - \frac{1}{C+1} \right)^{\frac{n}{2}} \left(P(E) - LP(B(\frac{m}{L})) \right),$$

for n large.

Step 2. (Exponential convergence of the barycenters) By (3.23), (3.10) and by (3.28), setting

$$(3.29) \quad b := \left(1 - \frac{1}{C+1} \right)^{\frac{1}{4}} \in (0, 1),$$

we have for n sufficiently large

$$\begin{aligned} |\xi_i^n - \xi_i^{n-1}| &= |\text{bar}(E_{f_{2,i}^n, m_n^i}) - \text{bar}(E_{f_{1,i}^n, m_n^i})| \\ &\leq C \sqrt{\mathcal{D}(E_{f_{2,i}^n, m_n^i}, E_{f_{1,i}^n, m_n^i})} = C \sqrt{\mathcal{D}(E_{h,i}^n, E_{h,i}^{n-1})} \\ &\leq C \left(P(E) - LP(B(\frac{m}{L})) \right)^{\frac{1}{2}} b^n. \end{aligned}$$

In turn, the above estimate implies that $\{\xi_i^n\}_n$ satisfies the Cauchy condition and thus there exist $\xi_i^\infty \in \mathbb{R}^N$, $i = 1, \dots, L$, such that $\xi_i^n \rightarrow \xi_i^\infty$ exponentially fast as $n \rightarrow \infty$; precisely,

$$|\xi_i^n - \xi_i^\infty| \leq \sum_{k=n+1}^{\infty} |\xi_i^k - \xi_i^{k-1}| \leq C \left(P(E) - LP(B(\frac{m}{L})) \right)^{\frac{1}{2}} \frac{b^{n+1}}{1-b}$$

for n large enough and for $i = 1, \dots, L$. Recalling (3.21), we may conclude that for all $\ell \in \mathbb{N}$

$$(3.30) \quad E_{h,i}^n \rightarrow \xi_i^\infty + B(\frac{m}{L}) \quad \text{in } C^\ell \quad \text{as } n \rightarrow \infty \text{ and for } i = 1, \dots, L.$$

Step 3. (Exponential convergence of the sets) By (3.30) we can parametrize the boudaries of the sets $E_{h,i}^n - \xi_i^\infty$ as radial graphs over the limiting ball $B(\frac{m}{L})$. Precisely, again with the notation (1.1), there exist functions g_i^n such that

$$(3.31) \quad E_{h,i}^n - \xi_i^\infty = E_{g_i^n, \frac{m}{L}} \quad \text{and} \quad \|g_i^n\|_{C^\ell(\partial B(\frac{m}{L}))} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

In turn, by Lemma 3.8 (see (3.8)), for n large enough we have that

$$\|g_i^n - g_i^{n-1}\|_{L^2(\partial B(\frac{m}{L}))} \leq C \left(\frac{m}{L} \right) \sqrt{\mathcal{D}(E_{h,i}^n, E_{h,i}^{n-1})}$$

and, thus, recalling (3.28) and arguing as in Step 2, we get

$$\begin{aligned}
 \|g_i^n\|_2 &\leq \sum_{k=n+1}^{\infty} \|g_i^k - g_i^{k-1}\|_2 \leq C \sum_{k=n+1}^{\infty} \sqrt{\mathcal{D}(E_{h,i}^k, E_{h,i}^{k-1})} \\
 (3.32) \qquad &\leq C \left(P(E) - LP(B(\frac{m}{T})) \right)^{\frac{1}{2}} \frac{b^{n+1}}{1-b},
 \end{aligned}$$

where b is as in (3.29). The above estimate yields the exponential decay of the L^2 -norms of the radial graphs. We now recall the following well known interpolation inequality: for every $j \in \mathbb{N}$ there exists $C > 0$ such that if g sufficiently smooth on $\partial B(\frac{m}{T})$, then

$$(3.33) \qquad \|D^\ell g\|_{L^2(\partial B(\frac{m}{T}))} \leq C \|D^{2\ell} g\|_{L^2(\partial B(\frac{m}{T}))}^{\frac{1}{2}} \|g\|_{L^2(\partial B(\frac{m}{T}))}^{\frac{1}{2}},$$

where D^ℓ stands for the collection all ℓ -th order (covariant) derivatives of g , see for instance [3]. Now, using the fact that from (3.31) for every ℓ there exists $n_\ell \in \mathbb{N}$ such that

$$\sup_{n \geq n_\ell} \|D^{2\ell} g_i^n\|_2 \leq 1,$$

we may apply (3.33) to g_i^n to deduce from (3.32) that also $\|D^\ell g_i^n\|_2$ decays exponentially fast for all $\ell \in \mathbb{N}$. This in turn yields the exponential decay in C^ℓ for every ℓ and concludes the proof of the theorem. \square

Lemma 3.11. *Let $\{a_n\}_{n \in \mathbb{N}}$ be a sequence of non-negative numbers. Assume furthermore that there exists $c > 1$, $\ell \in \mathbb{N}$ such that $\sum_{n=k}^{+\infty} a_n \leq c \sum_{j=k}^{k+\ell-1} a_j$ for every $k \in \mathbb{N}$. Then,*

$$a_k \leq \left(1 - \frac{1}{c}\right)^{\frac{k}{\ell}} S$$

for every $k \in \mathbb{N}$, where $S := \sum_{n=1}^{+\infty} a_n$.

Proof. We first consider the case $\ell = 1$. Set $F(k) := \sum_{n=k}^{+\infty} a_n$ and note that by assumption $F(k) \leq c(F(k) - F(k+1))$ for every $k \in \mathbb{N}$. Hence, by iteration we have

$$a_{k+1} \leq F(k+1) \leq \left(1 - \frac{1}{c}\right) F(k) \leq \dots \leq \left(1 - \frac{1}{c}\right)^{k+1} F(0) = \left(1 - \frac{1}{c}\right)^{k+1} S$$

for every $k \in \mathbb{N}$.

In the case $\ell \geq 2$ it is enough to set $b_k := \sum_{j=1}^{\ell} a_{\ell(k-1)+j}$ and to observe that the assumption now reads $\sum_{n=k}^{+\infty} b_n \leq c b_k$ so that we may apply the previous case. \square

ACKNOWLEDGEMENTS

This research was partially supported by the GNAMPA Grant 2018 “Nonlocal geometric flows” and by the GNAMPA Grant 2019 “Variational methods for nonlocal geometric motions”, both funded by INDAM.

E. S. has been supported by the ERC-STG Grant n. 759229 HiCoS “Higher Co-dimension Singularities: Minimal Surfaces and the Thin Obstacle Problem”.

REFERENCES

- [1] E. Acerbi, N. Fusco, M. Morini: Minimality via second variation for a nonlocal isoperimetric problem. *Comm. Math. Phys.* **322** (2013), no. 2, 515–557.
- [2] F. Almgren, J. Taylor, L. Wang: Curvature-driven flows: a variational approach. *SIAM J. Control Optim.* **31** (1993), no. 2, 387–438.
- [3] T. AUBIN, *Some nonlinear problems in Riemannian geometry*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998.

- [4] W. Carter, A. Roosen, J. Cahn, and J. Taylor: Shape evolution by surface diffusion and surface attachment limited kinetics on completely faceted surfaces, *Acta Metallurgica et Materialia*, **43** (1995), pp. 4309–4323.
- [5] G. Bellettini, V. Caselles, A. Chambolle, M. Novaga: The volume preserving crystalline mean curvature flow of convex sets in \mathbb{R}^N . *J. Math. Pure Appl.* **92** (5) (2009), 499–527.
- [6] G. Ciraolo, F. Maggi: On the shape of compact hypersurfaces with almost-constant mean curvature. *Comm. Pure Appl. Math.* **70** (2017), no. 4, 665–716.
- [7] R. Cristoferi: On periodic critical points and local minimizers of the Ohta-Kawasaki functional. *Nonlinear Anal.* **168** (2018), 81–109.
- [8] G. Ciraolo, L. Vezzoni: A sharp quantitative version of Alexandrov’s theorem via the method of moving planes. *J. Eur. Math. Soc. (JEMS)* **20** (2018), 261–299.
- [9] M. Delgadino, F. Maggi: Alexandrov’s theorem revisited. *Anal. PDE* **12** (2019), no. 6, 1613–1642.
- [10] M. Delgadino, F. Maggi, C. Mihaila, R. Neumayer: Bubbling with L^2 -almost constant mean curvature and an Alexandrov-type theorem for crystals. *Arch. Ration. Mech. Anal.* **230** (2018), no. 3, 1131–1177.
- [11] J. Escher, G. Simonett: The volume preserving mean curvature flow near spheres. *Proc. Am. Math. Soc.* **126** (1998), 2789–2796.
- [12] L. Esposito, N. Fusco: A remark on a free interface problem with volume constraint. *J. Convex Anal.* **18** (2011), 417–426.
- [13] M. Giaquinta, L. Martinazzi: An introduction to the regularity theory for elliptic systems, harmonic maps and minimal graphs. Second edition. Appunti. Scuola Normale Superiore di Pisa (Nuova Serie) [Lecture Notes. Scuola Normale Superiore di Pisa (New Series)], 11. Edizioni della Normale, Pisa, 2012. xiv+366 pp.
- [14] I. Kim, D. Kwon: Volume preserving mean curvature flow for star-shaped sets. *Comm. Partial Differential Equations*, to appear.
- [15] B. Krummel, F. Maggi: Isoperimetry with upper mean curvature bounds and sharp stability estimates. *Calc. Var. Partial Differ. Equ.* **56** (2017), Article n. 53.
- [16] S. Luckhaus, T. Sturzenhecker: Implicit time discretization for the mean curvature flow equation. *Calc. Var. Partial Differential Equations* **3** (1995), 253–271.
- [17] F. Maggi: *Sets of finite perimeter and geometric variational problems. An introduction to geometric measure theory.* Cambridge Studies in Advanced Mathematics, 135. Cambridge University Press, Cambridge, 2012.
- [18] U. F. Mayer: A singular example for the average mean curvature flow. *Experimental Mathematics* **10** (1) (2001), 103–107.
- [19] U. F. Mayer, G. Simonett: Self-intersections for the surface diffusion and the volume-preserving mean curvature flow. *Differential and Integral Equations* **13** (79) (2000), 1189–1199.
- [20] L. Mugnai, C. Seis: On the coarsening rates for attachment-limited kinetics. *SIAM J. Math. Anal.* **45** (2013), 324–344.
- [21] L. Mugnai, C. Seis, E. Spadaro: Global solutions to the volume-preserving mean-curvature flow. *Calc. Var.* **55**(2016), Article n. 18.
- [22] G. Huisken: The volume preserving mean curvature flow. *J. Rein. Angew. Math.* **382** (1987), 3548.
- [23] J. Niinikoski: Volume preserving mean curvature flows near strictly stable sets in flat torus. Preprint 2019. <https://arxiv.org/pdf/1907.03618.pdf>
- [24] L. A. Tarshis, J. L. Walker and M. F. X. Gigliotti: Solidification. *Annual Review of Materials Science* **2**(1972), 181–216.
- [25] C. Wagner: Theorie der Alterung von Niederschlägen durch Umlösen (Ostwald-Reifung). *Zeitschrift für Elektrochemie, Berichte der Bunsengesellschaft für physikalische Chemie* **65**, 7-8 (1961), 581–591.

(Massimiliano Morini) DIP. DI SCIENZE MATEMATICHE FISICHE E INFORMATICHE, UNIVERSITÀ DI PARMA, ITALY
E-mail address, M. Morini: massimiliano.morini@unipr.it

(Marcello Ponsiglione) DIP. DI MATEMATICA, UNIVERSITÀ DI ROMA “LA SAPIENZA”, ROMA, ITALY
E-mail address, M. Ponsiglione: ponsigli@mat.uniroma1.it

(Emanuele Spadaro) DIP. DI MATEMATICA, UNIVERSITÀ DI ROMA “LA SAPIENZA”, ROMA, ITALY
E-mail address, E. Spadaro: spadaro@mat.uniroma1.it