

Taxonomic and metabolic development of the human gut microbiome across life stages: a worldwide metagenomic investigation

Leonardo Mancabelli,^{1,2} Christian Milani,^{2,3} Rosita De Biase,³ Fabiana Bocchio,³ Federico Fontana,³ Gabriele Andrea Lugli,³ Giulia Alessandri,³ Chiara Tarracchini,³ Alice Viappiani,⁴ Flora De Conto,¹ Antonio Nouvenne,^{1,2,5} Andrea Ticinesi,^{1,2,5} Ovidio Bussolati,^{1,2} Tiziana Meschi,^{1,2,5} Rossana Cecchi,^{1,2} Francesca Turrone,^{2,3} Marco Ventura^{2,3}

AUTHOR AFFILIATIONS See affiliation list on p. 14.

ABSTRACT The human gut microbiota is a dynamic community of microorganisms that undergo variable changes over the entire life span. To thoroughly investigate the possible fluctuations of the microbiota throughout human life, we performed a pooled analysis of healthy fecal samples across different age groups covering the entire human life span. Our study integrated data from 79 publicly available studies and new stool samples from an Italian cohort, i.e., the Parma Microbiota project, resulting in 6,653 samples processed through the shotgun metagenomic approach. This approach has allowed species-level taxonomic reconstruction of the gut microbiota and investigation of its metabolic potential across the human life span. From a taxonomic point of view, our findings confirmed and detailed at species-level accuracy that the microbial richness of the gut microbiota gradually increases in the first stage of life, becoming relatively stable during adolescence. Moreover, the analysis identified the potential core microbiota representative of distinct age groups, revealing age-related bacterial patterns and the continuous rearrangement of the microbiota in terms of relative abundances across the life span rather than the acquisition and loss of taxa. Furthermore, the shotgun approach provided insights into the functional contribution of the human gut microbiome. The metagenomic analysis revealed functional age-related differences, particularly in carbohydrate and fiber metabolism, suggesting a co-evolution of the microbiome assembly with diet. Additionally, we identified correlations between vitamin synthesis, such as thiamine and niacin, and early life, suggesting a potential role of the microbiome in human physiology, in particular in the functions of the host's nervous and immune systems.

IMPORTANCE In this study, we provided comprehensive insights into the dynamic nature of the human gut microbiota across the human life span. In detail, we analyzed a large data set based on a shotgun metagenomic approach, combining public data sets and new samples from the Parma Microbiota project and obtaining a detailed overview of the possible relationship between gut microbiota development and aging. Our findings confirmed the main stages in microbial richness development and revealed specific core microbiota associated with different age stages. Moreover, the shotgun metagenomic approach allowed the disentangling of the functional changes in the microbiome across the human life span, particularly in diet-related metabolism, which is probably correlated to bacterial co-evolution with dietary habits. Notably, our study also uncovered positive correlations with vitamin synthesis in early life, suggesting a possible impact of the microbiota on human physiology.

Editor Kiran Patil, Medical Research Council Toxicology Unit, United Kingdom

Address correspondence to Marco Ventura, marco.ventura@unipr.it.

The authors declare no conflict of interest.

See the funding table on p. 15.

Received 1 December 2023

Accepted 9 February 2024

Published 5 March 2024

Copyright © 2024 Mancabelli et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

KEYWORDS human gut microbiome, human life span, shotgun metagenomics, human microbiota

The human gut microbiota, a rich and variable consortium of microorganisms residing in the intestinal tract (1–3), plays a crucial role in numerous aspects of human biology, including metabolic health to immune functions (3–5). This complex community, predominantly bacterial, undergoes dynamic changes throughout an individual's life, reflecting the interplay between microbial communities and host development (6, 7). In early life, the gut is gradually colonized, with its microbiota being highly dynamic and influenced by factors such as delivery mode and breastfeeding (8–10). Initially, it is characterized by Actinomycetota and Pseudomonadota (formerly known as Actinobacteria and Proteobacteria, respectively) (2, 11, 12). As children grow, significant shifts occur in the gut microbiota (13, 14), adapting to dietary changes from breastfeeding to solid foods (15).

Despite the current limited research focusing on the gut microbiota during human adolescence, it is widely suggested that, during this crucial window of time, the gut microbial community undergoes significant transformations and evolutions (16, 17). In fact, hormonal changes, dietary preferences, and lifestyle factors may exert influences on the gut microbial ecosystem (18–21). Notably, during adolescence, the microbiota tends to shift toward that of an adult, which is characterized by an increase in microbial genera belonging to the phyla Bacteroidota and Bacillota (formerly known as Bacteroidetes and Firmicutes, respectively), such as *Bacteroides*, *Prevotella*, *Blautia*, and *Faecalibacterium* (22).

Compared with infants and adolescents, adults are endowed with a microbiota characterized by increased stability and resilience (23–25). Its composition is strongly influenced by lifestyle-related factors such as diet (20, 26) and physical activity (27, 28). Numerous studies have revealed that a diverse and stable gut microbiota in adults is associated with improved metabolic and immune health, while an imbalance in composition can be linked to various pathological conditions (1, 29, 30). In fact, the balance between the major phyla and genera of the adult gut microbiota plays a crucial role in ensuring the host's well-being and fulfilling essential metabolic and immunological functions (1, 24).

In the later stages of life, elderly people experience another shift in their gut microbiota composition, which is characterized by a decrease in bacterial richness, particularly in beneficial species such as *Bifidobacterium*, *Akkermansia*, and members of *Clostridium* Clusters IV (31, 32). These changes may affect immune function, nutrient absorption, and susceptibility to age-related conditions, such as frailty and chronic diseases (31, 33, 34).

In this context, comprehensive studies with robust statistical power that investigate the detailed evolution of the gut microbiota across the life span have been notably limited and mainly focused on specific age groups (11, 35–39). Furthermore, a critical knowledge gap remains despite the wealth of knowledge gained in recent years. Currently, most of the existing studies are based on the 16S rRNA gene profiling approach (40), providing insights of the microbiota composition at the genus level but lacking the precision required for species-level analysis (41).

Given these critical gaps of knowledge, our study aims to perform the most extensive gut microbiota analysis based on publicly available shotgun metagenomic data sets regarding studies of the healthy human microbiome across life spans. In detail, we have assembled a very complete data set encompassing 6,653 samples representing various age groups and geographical regions, including 467 samples that originated from a still ongoing local population study, the Parma Microbiota project. This approach enables a more in-depth examination of taxonomic compositions at the species level and the microbiome's functionality, providing a comprehensive and statistically robust exploration of the intricate relationship between age and microbiome composition.

RESULTS

Data set selection

An extensive metagenomic data set search was performed to retrieve the largest number of publicly available shotgun metagenomic studies related to the human microbiome of healthy individuals. In detail, we collected data from 79 publicly available data sets that included healthy human fecal samples based on Illumina shotgun metagenomic methodologies (Fig. S1a; Table S1). In detail, in this pooled analysis, we included only studies in which it was possible to clearly identify the healthy status and the age of the individuals through the reported metadata. Moreover, as part of the Parma Microbiota project, fresh fecal samples from 467 healthy Italian individuals were collected, sequenced, and analyzed (Table S1). Thus, the pooled analysis included a total of 6,653 healthy fecal samples ranging from birth to over 100 years old (Tables S1 and S2) with a robust statistical representation of all the different age groups (see below).

Intra-individual variability across human life

The 6,653 stool samples collected in this pooled analysis were used to assess the microbiota composition through the METAnnotatorX2 software (42) following the standard filtering parameters reported in the manual with *Homo sapiens* reads removal. The downloaded and the sequenced fastq files were processed with the same bioinformatic pipeline to prevent biases, resulting in a total of 110,384,672,113 reads with an average per sample of $14,885,023 \pm 16,560,451$ after quality and human sequence filtering (Table S2). To optimize the taxonomical analysis, following a shallow shotgun metagenomic approach (42, 43), we decided to analyze, after quality and human sequence filtering, a random subset of up to 100,000 reads for each sample, obtaining a total of 424,267,507 classified reads with an average per sample of $63,771 \pm 13,611$ (Table S2). As previously reported, this approach allowed the optimization of the bioinformatic pipeline, ensuring accurate taxonomic profiling. Additionally, it promotes reliable profile comparisons by mitigating disparities resulting from variations in the total number of analyzed reads (42, 43).

The results generated using METAnnotatorX2 software (42) were employed to assess the biodiversity of each sample. In detail, to explore potential variations in species richness throughout the human life span, we categorized the samples into four age groups, that is, G1 (0–4 years), G2 (5–17 years), G3 (18–64 years), and G4 (65 years and older) (Fig. S1), following the guidelines provided by the World Health Organization (WHO) (44). The analysis revealed an increase in bacterial species abundance with age, as highlighted by a pairwise Kruskal-Wallis test ($P < 0.01$). Specifically, there was a substantial difference between G1 (average of 42 ± 23) and G2, G3, and G4 groups (average of 84 ± 20 , 83 ± 20 , and 86 ± 21 , respectively) (Fig. 1A). No significant differences were identified between G2, G3, and G4 groups (Fig. 1A). These results support the notion that the human gut microbiome undergoes developmental changes in the early stages of life, showing increasing complexity in terms of bacterial species until the human host reaches childhood and adolescence (5–17 years) (23). Subsequently, in the later stages of life, the gut microbiome reaches stability, with the number of bacterial species remaining relatively constant. In addition, a subdivision of the samples into further age subgroups (Fig. S1a) showed no significant differences between G1a (0–1 month) and G1b (1–6 months), indicating the presence of a heterogeneous and complexity-varying microbiota (14). Moreover, the G1c (6 months–1 year) and G1d (1–4 years) groups showed a significant trend of increase in species number until reaching the G2a group (5–10 years), where the gut microbiota appears to achieve a stable state that persists in subsequent age groups (pairwise Kruskal-Wallis test $P > 0.05$) (Fig. S1a).

Inter-individual variability between different stages of life

Principal coordinate analysis (PCoA), based on the Bray-Curtis dissimilarity matrix, was used to evaluate the inter-individual differences between age groups. In detail, the




FIG 1 Evaluation of microbial biodiversity. Panel (a) displays the whisker plot representing the species richness identified by subjects of each age group. The x - axis represents the different age groups, while the y - axis indicates the number of species. The 25th and 75th percentiles determine the boxes. The whiskers are determined by the 1.5 interquartile range (IQR). The line in the boxes represents the median, while the square represents the average. Different (Continued on next page)

FIG 1 (Continued)

lowercase letters indicate significant differences at $P < 0.05$ calculated through pairwise Kruskal-Wallis test analyses. In detail, groups with the same letter are not significantly different from each other, while groups with different letters are considered statistically distinct. Panel (b) reports the whisker plot representing the alpha diversity calculated through the Shannon index identified by subjects of each age group. The x - axis represents the different age groups, while the y - axis indicates the Shannon index. The 25th and 75th percentiles determine the boxes. The whiskers are determined by the 1.5 IQR. The line in the boxes represents the median, while the square represents the average. Different lowercase letters indicate significant differences at $P < 0.05$ calculated through pairwise Kruskal-Wallis test analyses. In detail, groups with the same letter are not significantly different from each other, while groups with different letters are considered statistically distinct. Panel (c) shows the pooled analysis of PCoA, subdivided by age groups. The black rows indicate the bacterial species with significant fittings (envit fit $P < 0.005$).

statistical analysis based on PERMANOVA (Table S3) revealed a clear division between the groups (pairwise PERMANOVA $q < 0.01$). Furthermore, the pairwise pseudo- F values (Table S2) evaluation highlighted a clear separation between the G1 group and the others, which exhibited heterogeneity among themselves (Fig. 1B). This division suggested a distinct microbiota composition of the G1 group samples, indicating a significant relationship between age and microbiota structure. Further fitting analyses, considering age and bacterial species as variables, identified *Bifidobacterium longum* as a key microbial taxon (envit fit $P = 0.002$, $r^2 = 0.2751$). This species appeared to exhibit a significant negative relationship with increasing age and is strongly associated with the G1 group. In contrast, other bacterial species with significant fittings (envit fit $P < 0.005$) displayed a positive relationship with increasing age, forming three distinct clusters. A prevalence of species from the *Prevotella* genus, such as *Prevotella copri* (recently classified as *Segatella copri*), characterized the first cluster. *Bacteroides uniformis* and *Alistipes putredinis* characterized the second cluster, while the third cluster exhibited diverse bacterial genera commonly associated with the adult gut microbiota, including *Ruminococcus*, *Roseburia*, and *Faecalibacterium*. These three clusters appeared to reflect the typical enterotypes of adults (45) but provided greater species-level detail. Moreover, the fitting analysis could highlight that enterotype 3, which is associated with the genus *Ruminococcus*, might instead consist of a complex community of bacteria with a less clear dominance of driver species (46, 47).

In addition to the age parameters, we also tested the potential impact of the variables related to the BioProject and nation of the samples on microbiota composition using PERMANOVA analysis. This analysis revealed $q < 0.01$ for both parameters, indicating statistical significance. However, the effect sizes measured by pseudo- F values were 16.7 and 11.4 for BioProject and geographical origin, respectively. These findings suggest that, while these two parameters may have a statistically significant impact on microbiota composition, their effect appears to be relatively modest. This observation could be assigned, in part, to the heterogeneous nature of the data set.

Identification of possible specific bacterial patterns related to different human stages of life

The METAnnotatorX2 software (42) allowed to obtain a detailed taxonomical profile at the species level for each sample. In detail, the sample size employed in this pooled analysis allowed the identification of age-specific core microbiota within each age group (Table S4). Core microbiota members were defined with a prerequisite of a minimum prevalence of 50% and an average relative abundance above 0.1% per age group. These criteria were chosen following the main standards in the field (48) and considering taxonomic complexity at the species level. Additionally, species with a prevalence between 30% and 50%, along with an average relative abundance above 0.1% were classified as accessory taxa. The core microbiota of each age group highlighted that G1 group displayed the simplest core composition, including seven species mainly represented by *B. longum* and *Escherichia coli* species (Table S4). In contrast, G2, G3, and G4 groups displayed larger and more diversified core microbiota, composed of 68, 57, and 63 species, respectively (Table S4). Accessory taxa exhibited a similar upward trend to the core, further validating the hypothesis that the gut microbiota demonstrates

substantial compositional variability during the early developmental stages and then shifts to taxonomic stability with advancing age (2, 23).

In order to determine the mainly representative species that compose the human gut microbiota across the life span, the bacterial species common to both the core and accessory microbiota across all age groups were selected (Table 1). This extensive screening yielded 29 bacterial species, including *B. uniformis*, which was the only species present in all cores, and *Bacteroides fragilis* that was the only ubiquitous species as an accessory taxa (Table 1). Notably, 21 taxa were identified as accessory species exclusively in the G1 group, subsequently transitioning to become core members in the G2, G3, and G4 groups. This pattern suggests that the development of the microbiota is characterized not only by the acquisition and loss of taxa but also by a significant rearrangement in their relative abundances. The early acquisition of certain species, pivotal in later life stages, highlights the critical role of microbiota enrichment during the initial phases of development. Moreover, among the microbial taxa representing the core microbiota of the G1 age group, only *B. longum* remains highly prevalent and abundant even in the G2 age group and then decreases in the subsequent G3 and G4 age groups, constituting one of the accessory taxa. These results suggested that these taxa persisted across the life span but exhibited dynamic interactions, shifting their prevalence over time. This

TABLE 1 Mainly representative species that compose the human gut microbiota across the life span^a

Taxonomy	Prevalence >50% and relative average abundance >0.1%				Relative average abundance			
	G1 (N = 3,100)	G2 (N = 366)	G3 (N = 2,632)	G4 (N = 555)	G1 (N = 3,100)	G2 (N = 366)	G3 (N = 2,632)	G4 (N = 555)
<i>B. longum</i>	69.71%	75.68%	36.17%	41.44%	9.03%	2.53%	0.55%	1.52%
<i>E. coli</i>	64.84%	43.44%	58.51%	54.77%	6.69%	0.87%	1.33%	2.26%
<i>Bacteroides</i> unknown_species	56.97%	96.72%	92.90%	88.47%	0.66%	1.23%	1.23%	1.19%
<i>Clostridium</i> unknown_species	56.84%	96.45%	96.12%	95.86%	0.48%	0.53%	0.60%	0.58%
<i>Blautia</i> unknown_species	53.42%	98.09%	98.52%	98.38%	0.58%	1.57%	2.08%	1.57%
<i>B. uniformis</i>	51.81%	94.54%	89.40%	88.29%	2.99%	4.10%	3.40%	4.10%
Ruminococcus unknown_species	50.58%	97.81%	97.26%	97.48%	0.45%	1.54%	1.68%	1.67%
<i>Blautia wexlerae</i>	48.39%	92.62%	87.42%	84.50%	1.82%	1.77%	1.68%	1.45%
<i>Bifidobacterium</i> unknown_species	46.87%	60.93%	31.08%	42.16%	0.58%	0.29%	0.11%	0.19%
<i>Flavonifractor plautii</i>	46.87%	83.33%	66.53%	78.38%	0.57%	0.37%	0.29%	0.40%
<i>Phocaecicola vulgatus</i>	42.06%	84.43%	72.49%	73.69%	1.31%	1.44%	1.15%	1.12%
<i>Bacteroides thetaiotaomicron</i>	40.90%	84.70%	72.72%	73.87%	0.82%	0.78%	0.59%	0.74%
<i>Phocaecicola dorei</i>	39.45%	80.60%	63.11%	66.13%	2.05%	1.51%	1.29%	1.37%
<i>Eubacterium</i> unknown_species	38.71%	94.54%	95.90%	95.86%	0.26%	0.99%	1.48%	1.29%
<i>Parabacteroides distasonis</i>	37.84%	82.51%	69.38%	77.48%	1.67%	1.15%	0.87%	1.07%
<i>Enterocloster</i> unknown_species	37.23%	87.98%	90.35%	89.55%	0.12%	0.16%	0.22%	0.21%
<i>Roseburia</i> unknown_species	36.74%	93.72%	95.59%	94.05%	0.22%	0.59%	1.14%	0.84%
<i>Faecalibacterium</i> unknown_species	36.65%	93.44%	96.01%	94.77%	1.17%	4.11%	3.23%	3.39%
<i>Faecalibacterium prausnitzii</i>	36.35%	92.62%	96.05%	93.69%	0.97%	3.41%	2.94%	3.12%
<i>Bacteroides xylanisolvens</i>	35.84%	79.78%	56.34%	64.68%	0.52%	0.63%	0.56%	0.55%
<i>Phocaecicola</i> unknown_species	35.68%	79.23%	79.71%	74.59%	0.21%	0.48%	0.65%	0.53%
<i>Anaerostipes hadrus</i>	34.35%	82.51%	63.87%	58.38%	1.11%	1.23%	1.09%	1.09%
<i>B. fragilis</i>	33.58%	43.44%	31.50%	37.48%	2.99%	1.25%	0.43%	0.78%
<i>Coprococcus</i> unknown_species	33.55%	87.43%	88.87%	86.31%	0.14%	0.48%	0.75%	0.58%
<i>Agathobacter rectalis</i> (formerly known as <i>Eubacterium rectale</i>)	31.29%	90.71%	92.86%	87.39%	0.99%	3.23%	4.11%	2.70%
<i>Blautia massiliensis</i>	31.03%	84.70%	75.68%	72.43%	0.41%	1.01%	0.77%	0.62%
<i>Dorea</i> unknown_species	30.77%	89.89%	89.02%	85.95%	0.12%	0.39%	0.58%	0.41%
<i>Bacteroides ovatus</i>	30.35%	68.58%	51.41%	51.53%	0.44%	0.33%	0.22%	0.23%
<i>Roseburia intestinalis</i>	30.06%	80.60%	84.38%	81.26%	0.74%	0.65%	1.01%	0.76%

^aIn detail, only the bacterial species common to both the core and accessory microbiota across all age groups were reported. The bacterial taxa belonging to the core microbiota were highlighted in bold.

dynamic co-existence could imply the formation of a complex ecosystem that could potentially reach a climax condition (23).

In addition, a dedicated correlation analysis was conducted to highlight the bacterial species significantly associated with aging. Based on Spearman's rank correlation coefficient, the correlation analysis revealed a total of 104 taxa with a significant relationship with age ($P < 0.01$) (Table S5). Among these taxa, we focused on those that showed significantly higher relative abundance in at least one of the age groups (multiple comparison analyses Tukey's (honestly significant difference) HSD $P < 0.05$, Fig. 2), resulting in a total of 45 taxa. Within this selection, we identified nine taxa with a negative correlation and 36 taxa with a positive correlation to age (Fig. 2). Notably, the nine taxa with a negative correlation were more abundant in the G1 group and were primarily represented by species belonging to the *Bifidobacterium* genus, such as *Bifidobacterium breve*, *B. longum*, *Bifidobacterium bifidum*, and *Bifidobacterium pseudocatenulatum*, along with species characteristic of the infant microbiota, such as *Veillonella parvula*, *Ruminococcus gnavus*, and *E. coli* (11, 12, 38). Notably, the increase of the correlation value showed a trend related to increasing age, except for the characteristic taxa of the G4 group, which corresponds to elderly individuals. These taxa, such as *Ruthenibacterium lactatiformans*, *Anaerotruncus unknown_species*, and *Butyricicoccus unknown_species*, exhibited a more heterogeneous distribution in terms of correlation values despite being more present in the G4 group, suggesting an increase in variability of the intestinal microbiota composition in this age group.

Exploring microbial functional diversity across the life span

The taxonomical analysis of the 6,653 healthy fecal samples across the life span revealed possible specific bacterial patterns correlated to the different age groups. In this context, in order to explore the genetic features characterizing each microbiome sample, we performed a screening of the microbially driven metabolic enzymatic reactions based on the MetaCyc database (49) and the Enzyme Commission (EC) classification. In detail, the enzymatic reactions that were revealed through the metagenomic analysis were used to perform a correlation analysis with the 104 bacterial species exhibiting a significant relationship with age (see above). Afterward, focusing on the core and accessory bacterial taxa that were statistically associated with each age group and on the main key enzymes involved in the metabolism of the various components of the human diet or in the metabolism of the main microbial metabolic products important for the host (see more detail in Materials and Methods), the correlation analysis seemed to highlight possible specific correlations between the age groups and the metabolic capability of the gut microbiome (Fig. 3). In detail, the functional analysis revealed a different correlation between the EC involved in the metabolism of carbohydrates and the bacterial species characterizing the four age groups, highlighting a greater similarity between groups G2, G3, and G4 compared with group G1 (Fig. 3). Curiously, EC 3.2.1.23, that is, beta-galactosidase, exhibited a significant positive correlation with G1 group compared with groups G2, G3, and G4. This result confirms the possible association between the composition of the intestinal microbiome and the host's milk-based diet. Indeed, group G1, which is associated with infants, is likely influenced by a milk-based diet, where beta-galactosidase is essential for breaking down lactose, the prevalent carbohydrate source in milk (50). Similarly, the bacterial taxa representative of the G1 group showed predominantly negative correlations with EC enzymes involved in fiber metabolisms. This result could probably be linked to the infant's low-fiber diet (51), highlighting the possible relationship between microbiome-EC enzyme and the host's diet.

Furthermore, the analysis of enzymes related to the biosynthesis of B vitamins also revealed a specific correlation between microbiome composition and human life span. Notably, the G1 group showed a positive correlation with enzymes involved in thiamine, i.e., 2.7.1.89, 2.7.1.50, and 3.6.1.27, and niacin, i.e., 2.7.1.23 and 6.3.5.1, biosynthesis in contrast to the G2, G3, and G4 groups. This observation highlighted the importance and




FIG 2 Correlation analysis between the bacterial species and the age of the individuals included in pooled analysis. In detail, only the bacterial taxa that showed a significant Spearman's rank correlation coefficient and significantly higher relative abundance in at least one of the age groups calculated through ANOVA test analysis and multiple comparison analyses Tukey's HSD test were reported.




FIG 3 Correlation analysis between the bacterial species and enzymatic reaction identified in pooled analysis. The red color indicated negative correlations, while the green color represented positive correlations. Only the main key enzymes involved in the human diet resulted in statistical significance; Spearman's rank correlation coefficient were reported.

specificity of microbial groups characterizing the G1 group, such as *B. breve*, *B. longum*, *B. bifidum*, *V. parvula*, and *E. coli*, which could contribute to the biosynthesis of these vitamins impacting the physiological development of the host.


Investigation of the possible impact of geographical origin on the human microbiome across the host life span

In order to identify potential variations in the human gut microbiota based on geographical origin, in particular with continent of origin, we performed an exploratory and preliminary multivariable statistical analysis based on MaAsLin2 software using the most representative population, i.e., the European population, as a reference for each pairwise comparison (52) (see more detail in Materials and Methods). The analysis revealed marked and significant differences in the microbiota profiles (Table S6). In detail, individuals from South America and Africa showed more significant differences than those from Europe (Fig. 4), while samples from North America showed the fewest significant differences (Fig. 4). Notably, South America and Africa exhibited a similar taxonomical correlation trend, revealing a positive correlation with species belonging to *Prevotella*, *Prevotellamassilia*, and *Treponema* genera and a negative correlation with bacterial species belonging to the *Phocaeicola*, *Bacteroides*, and *Alistipes* genera. Curiously, these trend correlations appeared consistent across different age groups, indicating a degree of temporal stability in the microbiota composition. Despite the absence of accurate details about the diet composition, it is tempting to hypothesize that the identified microbial profiles could reflect the lifestyles of different individuals. In fact, individuals originating from Africa and South America, who tend to have a diet characterized more by local and traditional products and less influenced by globalization, seemed to show a microbiota more similar to non-urbanized populations characterized by a high abundance of species belonging to the *Prevotella* and *Treponema* genera (53–56).

Conversely, multivariable analysis, based on the EC class composition of the key enzymes involved in the human diet (Fig. 4; Table S5), showed that the individuals from South America, Africa, and Asia possessed the highest number of negative correlations, i.e., 18, 11, and 11, respectively, compared with Europe, suggesting broader functional diversity or different ecological adaptations in the microbiome of these populations (53, 57). Furthermore, the observed negative correlations remained consistent across different age groups, suggesting a stable pattern in microbiome functionality over time (Fig. 4). These results could corroborate the notion that lifestyle substantially influences the intestinal microbiota, which, in turn, can exert a variable impact on the host, generally maintaining stability in terms of composition and functionality throughout human life. These preliminary results need further investigation using more accurate metagenomic data sets with detailed metadata on lifestyle factors, such as diet composition, use of drugs/antibiotics, and medical treatments. A deeper understanding of these aspects will enable us to better interpret lifestyle impact on the microbiome and its variation across different populations across the human life span.

DISCUSSION

The human intestinal microbiota is widely recognized to play a key role in human health, so it has been the focus of extensive scientific research. However, knowledge regarding the evolution of gut microbiota over an individual's lifetime has been limited. In this context, we decided to perform an extensive pooled analysis based on public and new shotgun metagenomic data sets of the human gut microbiota throughout the human life span, elucidating the dynamic nature of this complex ecosystem. The pooled analysis, encompassing a total of 6,653 fecal samples, identified an increase in microbial diversity in early life, followed by relative stability during adolescence, highlighting the continuous interplay between the microbiome and the host aging. Moreover, the statistical power of this pooled analysis allowed the identification of a potential age-related core microbiota at the bacterial species level. In detail, the samples representing the early



stages of life showed the simplest core microbiota composition, mainly represented by *B. longum* and *E. coli* species. In contrast, adolescent and adult samples showed a more extensive and diversified core microbiota, supporting the hypothesis that the gut microbiota displays considerable changes in composition during the initial stages of development and then shifts to taxonomic stability with advancing age. Intriguingly, taxa, initially classified as accessories in the early stages of life, often become core components in subsequent stages. This transition could highlight a dynamic reorganization within the microbiota, characterized by shifting relative abundances rather than just the progressive acquisition and loss of taxa.

Moreover, the shotgun metagenomic approach allowed to investigate the functional capabilities of the gut microbiome. Specifically, the metagenomic analysis unveiled functional variations linked to age, particularly in the metabolism of carbohydrates and fibers, probably indicating a co-evolution of the microbiome and host influenced by dietary factors. Additionally, the functional analysis revealed possible associations with the biosynthesis of B vitamins and, in particular, with thiamine and niacin metabolisms during early life, suggesting a potential role of the microbiota in shaping human physiology, such as the functions of nervous and immune systems (58, 59).

Furthermore, an exploratory preliminary multivariable analysis investigated the relationship between the human gut microbiome and the geographical origin of the individual. The analysis revealed possible differences in microbiota profiles between continents. In detail, the gut microbiomes from South America and Africa showed distinct microbial compositions compared with other continents, probably related to more traditional diets that are less influenced by globalization. These trends persisted across age groups, indicating temporal microbiota stability. Similarly, the analysis of specific EC classes suggested a possible functional diversity related to geographical origin.

Such findings highlighted the co-evolution of the gut microbiota with the host throughout the human life span, revealing a bacterial adaptation to the host's habits and its potential influence on host physiology. Nevertheless, the uneven distribution of samples across different groups of age and the lack of detailed metadata concerning, among others, diet composition and lifestyle could represent possible limitations of this study that should be overcome through more complex metagenomic analyses, allowing a more comprehensive understanding of the intricate interplay between gut microbiota and human health across the life span. Furthermore, while the current approaches based on genomic databases used for classifying bacterial populations often lack precise species-level identification, they remain a critical tool for microbial analysis. However, the ongoing expansion and enhancement of these genomic databases are expected to significantly improve the accuracy of microbial species identification of the human gut microbiota. Such advancements are crucial for deepening the understanding of the gut microbiota's role in human health across various life stages. Despite these limitations, our study's approach is considered effective for obtaining a comprehensive bacterial profile, particularly compared to other methods based on marker genes (42, 60).

MATERIALS AND METHODS

Selection and collection of samples included in the pooled analysis

In this pooled analysis-based study, we retrieved 79 publicly available data sets from studies regarding the human gut microbiome for a total of 6,186 samples from 37 different nations (Table S1). In particular, we selected shotgun metagenomic data sets obtained by an Illumina sequencing platform to avoid the input data's variability as much as possible. In addition, we included 467 Italian adult healthy individuals collected as part of the Parma Microbiota project (Comitato Etico dell'Area Vasta Emilia Nord, Emilia-Romagna Region, Italy, under the ID 1107/2020/TESS/UNIPR) (Table S1). These Italian fecal samples, once collected, were immediately inactivated with DNA/RNA shield buffer (Zymo Research, USA) and subsequently delivered to the Laboratory of

Probiogenomics of Parma University, where the analysis of bacterial DNA libraries by shotgun metagenomic and the bioinformatic analysis of raw sequencing data were performed.

Shallow shotgun sequencing

According to the manufacturer's instructions, DNA library preparation was performed using the Nextera XT DNA Sample Preparation Kit (Illumina, San Diego, CA, USA). First, 1-ng input DNA from each sample was used for the library preparation, which underwent fragmentation, adapter ligation, and amplification. Then, Illumina libraries were pooled equimolarly, denatured, and diluted to a concentration of 1.5 pM. Next, DNA sequencing was performed on a MiSeq instrument (Illumina) using a 2 × 250-bp Output Sequencing Kit together with a deliberate spike-in of 1% PhiX control library.

Taxonomic classification of sequence reads

Taxonomic profiling of sequenced and downloaded reads was performed employing the METAnnotatorX2 bioinformatic platform (42, 61). In detail, the fastq files were filtered to remove reads with the quality of <25 and to retain reads with a length of >100 bp. Subsequently, human host DNA filtering was performed through Bowtie 2 software (62, 63), following the METAnnotatorX2 manual (42). Afterward, the taxonomic classification of 100,000 reads was achieved by means of MegaBLAST (64) employing a manually curated and pre-processed database of genomes retrieved from the National Center for Biotechnology Information, following the METAnnotatorX2 manual (42).

Functional prediction

Functional profiling of the sequenced reads was performed with the METAnnotatorX2 bioinformatic platform (42, 61). Functional classification of reads was performed to reveal metabolic pathways based on the MetaCyc database (release 24.1) (49) through RAPSearch2 software (65, 66).

Statistical analysis

ORIGIN 2021 (<https://www.originlab.com/2021>) and SPSS software (www.ibm.com/software/it/analytics/spss/) were used to compute statistical analyses. In detail, pairwise Kruskal-Wallis test analyses tested differences in alpha diversity that is calculated through species richness and Shannon index. Moreover, the similarities between samples (beta-diversity) were calculated by the Bray-Curtis dissimilarity matrix based on species abundance, using the "vegdist" function (from `vegan_2.5-7`) on RStudio (<http://www.rstudio.com/>). The range of similarities is calculated between values 0 and 1. Beta-diversity was represented through PCoA using the function "ape" of the `Rsuite` package (67). Moreover, the available metadata and the various detected bacterial species were tested and plotted on the PCoA using the "envfit" and "plot" functions from `vegan` (version 2.5-7), respectively, through RStudio (<http://www.rstudio.com/>). PERMANOVA analyses were performed on RStudio using 999 permutations to estimate *P* values for population differences in PCoA analyses with `adonis2` package (from `vegan_2.5-7`). Furthermore, a correlation analysis between the available metadata and the various detected bacterial species of all samples was performed through Spearman's rank correlation coefficient using "rcorr" function (from `Hmisc_4.6-0`; <https://CRAN.R-project.org/package=Hmisc>), and only significant statistical results were retained. The false discovery rate (FDR) correction based on Benjamini and Hochberg correction (68) and calculated using RStudio through "p.adjust" function (from base package `stats`) was applied to statistically significant results. In detail, correlation analysis was performed between metabolic reactions revealed through the metagenomic analysis and the 104 bacterial species, which exhibited a significant relationship with the host's age. Afterward, we focused our interest on the 46 bacterial taxa that showed significantly higher relative abundance in at least one of the age groups calculated through analysis

of variance (ANOVA) test analysis and multiple comparison analyses Tukey's HSD test and on the main key enzymes involved in the metabolism of the various components of the infant and/or adult human diet, such as human milk oligosaccharides, carbohydrates, and fibers, or in the metabolism of the main microbial products important for the host, such as B vitamins and short-chain fatty acids.

Furthermore, multivariable statistical analysis based on MaAsLin2 software (52) was performed to identify potential variations in the human gut microbiota based on geographical origin. In detail, the multivariable analysis allowed to investigate the possible correlation between the human gut microbiome and the geographical origin of the host. We focused on the continent of origin to reinforce the statistical power of the analysis. The analysis based on MaAsLin2 software was performed separately for each age group, considering the continent of origin, the microbiota composition, and the EC composition of the key enzymes involved in the human diet (see above). Moreover, European individuals were selected as analysis references, primarily due to the higher average number of samples across each age group. Afterward, we focused on the attention on the taxa that exhibit significant statistical correlations across all age groups in at least one continent group.

ACKNOWLEDGMENTS

This study was supported by "Programma Operativo Nazionale Ricerca e Innovazione" 2014–2020 (PON "R&I" 2014–2020) (project ARS01_00530) and by Fondazione Cariparma as part of the Parma Microbiota project. M.V. was funded by the "Fondo per il Programma Nazionale di Ricerca e Progetti di Rilevante Interesse Nazionale (PRIN)," Ministero della Ricerca e dell'Università (20229LEB99). F.T. was supported by PROGETTO Ricerca Finalizzata, Ministero della Salute (RF GR-2018–12365988).

Part of this research is conducted using the high-performance computing facility of the University of Parma.

L.M.: data curation, formal analysis, visualization, and writing—review and editing; C.M.: conceptualization and writing—review and editing.; R.D.B.: investigation, data curation, and formal analysis.; F.B.: investigation and data curation.; F.F.: software.; G.A.L.: software.; G.A.: formal analysis.; C.T.: software.; A.V.: investigation.; F.D.C.: supervision.; A.N.: resources.; A.T.: resources.; O.B.: supervision.; T.M.: resources.; R.C.: resources.; F.T.: conceptualization and supervision.; and M.V.: project administration, conceptualization, and writing—review and editing. All authors have read and approved the final manuscript.

AUTHOR AFFILIATIONS

¹Department of Medicine and Surgery, University of Parma, Parma, Italy

²Interdepartmental Research Centre "Microbiome Research Hub", University of Parma, Parma, Italy

³Laboratory of Probiogenomics, Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Parma, Italy

⁴GenProbio srl, Parma, Italy

⁵Parma University Hospital, Parma, Italy

AUTHOR ORCID*s*

Leonardo Mancabelli  <http://orcid.org/0000-0002-1744-2214>

Christian Milani  <https://orcid.org/0000-0002-5062-3164>

Francesca Turrone  <https://orcid.org/0000-0001-5363-0231>

Marco Ventura  <http://orcid.org/0000-0002-4875-4560>

FUNDING

Funder	Grant(s)	Author(s)
Ministero della Salute (Italy Ministry of Health)	GR-2018-12365988	Francesca Turrone
Ministero della Ricerca e dell'Universita	20229LEB99	Marco Ventura

AUTHOR CONTRIBUTIONS

Leonardo Mancabelli, Data curation, Formal analysis, Visualization, Writing – original draft | Christian Milani, Conceptualization, Writing – review and editing | Rosita De Biase, Data curation, Formal analysis, Investigation | Fabiana Bocchio, Data curation, Investigation | Federico Fontana, software | Gabriele Andrea Lugli, software | Giulia Alessandri, Formal analysis | Chiara Tarracchini, software | Alice Viappiani, Investigation | Flora De Conto, supervision | Antonio Nouvenne, Resources | Andrea Ticinesi, Resources | Ovidio Bussolati, supervision | Tiziana Meschi, Resources | Rossana Cecchi, Resources | Francesca Turrone, Conceptualization, supervision | Marco Ventura, Conceptualization, Project administration, Writing – review and editing

DATA AVAILABILITY

Raw Italian shotgun metagenomic data sequences are accessible through SRA under study accession number [PRJNA1046438](https://www.ncbi.nlm.nih.gov/sra/PRJNA1046438).

ADDITIONAL FILES

The following material is available [online](#).

Supplemental Material

Figure S1 (mSystems01294-23-S0001.tif). Workflow of the pooled analysis performed and species richness identified by subjects of each age subgroup.

Legends (mSystems01294-23-S0002.docx). Supplemental material legends.

Table S1 (mSystems01294-23-S0003.xlsx). List of the public bioprojects included in the pooled analysis.

Table S2 (mSystems01294-23-S0004.xlsx). Metadata of the samples included in the pooled analysis.

Table S3 (mSystems01294-23-S0005.xlsx). PERMANOVA statistical analysis based on the Bray-Curtis dissimilarity matrix calculated the inter-individual differences between age groups.

Table S4 (mSystems01294-23-S0006.xlsx). Core and accessory microbiota calculated on the subjects included in the pooled analysis.

Table S5 (mSystems01294-23-S0007.xlsx). Correlation analysis between the bacterial species and enzymatic reaction identified in pooled analysis.

Table S6 (mSystems01294-23-S0008.xlsx). Multivariate analysis through MaAslin2 software based on bacterial species, age groups, and geographical origin.

REFERENCES

- Hou K, Wu ZX, Chen XY, Wang JQ, Zhang D, Xiao C, Zhu D, Koya JB, Wei L, Li J, Chen ZS. 2022. Microbiota in health and diseases. *Signal Transduct Target Ther* 7:135. <https://doi.org/10.1038/s41392-022-00974-4>
- Milani C, Duranti S, Bottacini F, Casey E, Turrone F, Mahony J, Belzer C, Delgado Palacio S, Arboleya Montes S, Mancabelli L, Lugli GA, Rodriguez JM, Bode L, de Vos W, Gueimonde M, Margolles A, van Sinderen D, Ventura M. 2017. The first microbial colonizers of the human gut: composition, activities, and health implications of the infant gut microbiota. *Microbiol Mol Biol Rev* 81:e00036-17. <https://doi.org/10.1128/MMBR.00036-17>
- de Vos WM, Tilg H, Van Hul M, Cani PD. 2022. Gut microbiome and health: mechanistic insights. *Gut* 71:1020–1032. <https://doi.org/10.1136/gutjnl-2021-326789>
- Zheng D, Liwinski T, Elinav E. 2020. Interaction between microbiota and immunity in health and disease. *Cell Res* 30:492–506. <https://doi.org/10.1038/s41422-020-0332-7>
- Wu HJ, Wu E. 2012. The role of gut microbiota in immune homeostasis and autoimmunity. *Gut Microbes* 3:4–14. <https://doi.org/10.4161/gmic.19320>
- Barreto HC, Gordo I. 2023. Intrahost evolution of the gut microbiota. *Nat Rev Microbiol* 21:590–603. <https://doi.org/10.1038/s41579-023-00890-6>
- Carmody RN, Sarkar A, Reese AT. 2021. Gut microbiota through an evolutionary lens. *Science* 372:462–463. <https://doi.org/10.1126/science.abf0590>
- Milani C, Mancabelli L, Lugli GA, Duranti S, Turrone F, Ferrario C, Mangifesta M, Viappiani A, Ferretti P, Gorfer V, Tett A, Segata N, van

- Sinderen D, Ventura M. 2015. Exploring vertical transmission of bifidobacteria from mother to child. *Appl Environ Microbiol* 81:7078–7087. <https://doi.org/10.1128/AEM.02037-15>
9. Reyman M, van Houten MA, van Baarle D, Bosch A, Man WH, Chu M, Arp K, Watson RL, Sanders EAM, Fuentes S, Bogaert D. 2019. Impact of delivery mode-associated gut microbiota dynamics on health in the first year of life. *Nat Commun* 10:4997. <https://doi.org/10.1038/s41467-019-13014-7>
 10. Ma J, Li Z, Zhang W, Zhang C, Zhang Y, Mei H, Zhuo N, Wang H, Wang L, Wu D. 2020. Comparison of gut microbiota in exclusively breast-fed and formula-fed babies: a study of 91 term infants. *Sci Rep* 10:15792. <https://doi.org/10.1038/s41598-020-72635-x>
 11. Lugli GA, Mancabelli L, Milani C, Fontana F, Tarracchini C, Alessandri G, van Sinderen D, Turrone F, Ventura M. 2023. Comprehensive insights from composition to functional microbe-based biodiversity of the infant human gut microbiota. *NPJ Biofilms Microbiomes* 9:25. <https://doi.org/10.1038/s41522-023-00392-6>
 12. Alessandri G, Fontana F, Mancabelli L, Lugli GA, Tarracchini C, Argentin C, Longhi G, Viappiani A, Milani C, Turrone F, van Sinderen D, Ventura M. 2022. Exploring species-level infant gut bacterial biodiversity by meta-analysis and formulation of an optimized cultivation medium. *NPJ Biofilms Microbiomes* 8:88. <https://doi.org/10.1038/s41522-022-00349-1>
 13. Ronan V, Yeasin R, Claud EC. 2021. Childhood development and the microbiome—the intestinal microbiota in maintenance of health and development of disease during childhood development. *Gastroenterology* 160:495–506. <https://doi.org/10.1053/j.gastro.2020.08.065>
 14. Mei H, Yang S, Peng A, Li R, Xiang F, Zheng H, Tan Y, Zhang Y, Zhou A, Zhang J, Xiao H. 2022. Development of the gut microbiota in healthy twins during the first 2 years of life and associations with body mass index z-score: results from the Wuhan twin birth cohort study. *Front Microbiol* 13:891679. <https://doi.org/10.3389/fmicb.2022.891679>
 15. Fallani M, Amarri S, Uusijarvi A, Adam R, Khanna S, Aguilera M, Gil A, Vieites JM, Norin E, Young D, Scott JA, Doré J, Edwards CA, The Infabio Team. 2011. Determinants of the human infant intestinal microbiota after the introduction of first complementary foods in infant samples from five European centres. *Microbiology (Reading)* 157:1385–1392. <https://doi.org/10.1099/mic.0.042143-0>
 16. Carson MD, Westwater C, Novince CM. 2023. Adolescence and the microbiome: implications for healthy growth and maturation. *Am J Pathol* 193:1900–1909. <https://doi.org/10.1016/j.ajpath.2023.07.004>
 17. Derrien M, Alvarez A-S, de Vos WM. 2019. The gut microbiota in the first decade of life. *Trends Microbiol* 27:997–1010. <https://doi.org/10.1016/j.tim.2019.08.001>
 18. Santos-Marcos JA, Mora-Ortiz M, Tena-Sempere M, Lopez-Miranda J, Camargo A. 2023. Interaction between gut microbiota and sex hormones and their relation to sexual dimorphism in metabolic diseases. *Biol Sex Differ* 14:4. <https://doi.org/10.1186/s13293-023-00490-2>
 19. Ferrario C, Statello R, Carnevali L, Mancabelli L, Milani C, Mangifesta M, Duranti S, Lugli GA, Jimenez B, Lodge S, Viappiani A, Alessandri G, Dall'Asta M, Del Rio D, Sgoifo A, van Sinderen D, Ventura M, Turrone F. 2017. How to feed the mammalian gut microbiota: bacterial and metabolic modulation by dietary fibers. *Front Microbiol* 8:1749. <https://doi.org/10.3389/fmicb.2017.01749>
 20. Milani C, Ferrario C, Turrone F, Duranti S, Mangifesta M, van Sinderen D, Ventura M. 2016. The human gut microbiota and its interactive connections to diet. *J Hum Nutr Diet* 29:539–546. <https://doi.org/10.1111/jhn.12371>
 21. Zhong H, Penders J, Shi Z, Ren H, Cai K, Fang C, Ding Q, Thijs C, Blaak EE, Stehouwer CDA, Xu X, Yang H, Wang J, Wang J, Jonkers D, Masclee AAM, Brix S, Li J, Arts ICW, Kristiansen K. 2019. Impact of early events and lifestyle on the gut microbiota and metabolic phenotypes in young school-age children. *Microbiome* 7:2. <https://doi.org/10.1186/s40168-018-0608-z>
 22. Ou Y, Belzer C, Smidt H, de Weerth C. 2023. Development of the gut microbiota in the first 14 years of life and its relations to Internalizing and externalizing difficulties and social anxiety during puberty. *Eur Child Adolesc Psychiatry*. <https://doi.org/10.1007/s00787-023-02205-9>
 23. Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. 2012. Diversity, stability and resilience of the human gut microbiota. *Nature* 489:220–230. <https://doi.org/10.1038/nature11550>
 24. Sommer F, Anderson JM, Bharti R, Raes J, Rosenstiel P. 2017. The resilience of the intestinal microbiota influences health and disease. *Nat Rev Microbiol* 15:630–638. <https://doi.org/10.1038/nrmicro.2017.58>
 25. Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M, Gill SR, Nelson KE, Relman DA. 2005. Diversity of the human intestinal microbial flora. *Science* 308:1635–1638. <https://doi.org/10.1126/science.1110591>
 26. Kolodziejczyk AA, Zheng D, Elinav E. 2019. Diet-microbiota interactions and personalized nutrition. *Nat Rev Microbiol* 17:742–753. <https://doi.org/10.1038/s41579-019-0256-8>
 27. Fontana F, Longhi G, Tarracchini C, Mancabelli L, Lugli GA, Alessandri G, Turrone F, Milani C, Ventura M. 2023. The human gut microbiome of athletes: metagenomic and metabolic insights. *Microbiome* 11:27. <https://doi.org/10.1186/s40168-023-01470-9>
 28. Monda V, Villano I, Messina A, Valenzano A, Esposito T, Moscatelli F, Viggiano A, Cibelli G, Chieffi S, Monda M, Messina G. 2017. Exercise modifies the gut microbiota with positive health effects. *Oxid Med Cell Longev* 2017:3831972. <https://doi.org/10.1155/2017/3831972>
 29. Vijay A, Valdes AM. 2022. Role of the gut microbiome in chronic diseases: a narrative review. *Eur J Clin Nutr* 76:489–501. <https://doi.org/10.1038/s41430-021-00991-6>
 30. Durack J, Lynch SV. 2019. The gut microbiome: relationships with disease and opportunities for therapy. *J Exp Med* 216:20–40. <https://doi.org/10.1084/jem.20180448>
 31. Ticinesi A, Milani C, Lauretani F, Nouvenne A, Mancabelli L, Lugli GA, Turrone F, Duranti S, Mangifesta M, Viappiani A, Ferrario C, Maggio M, Ventura M, Meschi T. 2017. Gut microbiota composition is associated with polypharmacy in elderly hospitalized patients. *Sci Rep* 7:11102. <https://doi.org/10.1038/s41598-017-10734-y>
 32. Kim S, Jazwinski SM. 2018. The gut microbiota and healthy aging: a mini-review. *Gerontology* 64:513–520. <https://doi.org/10.1159/000490615>
 33. Almeida HM, Sardeli AV, Conway J, Duggal NA, Cavaglieri CR. 2022. Comparison between frail and non-frail older adults' gut microbiota: a systematic review and meta-analysis. *Ageing Res Rev* 82:101773. <https://doi.org/10.1016/j.arr.2022.101773>
 34. Ragonnaud E, Biragyn A. 2021. Gut microbiota as the key controllers of "healthy" aging of elderly people. *Immun Ageing* 18:2. <https://doi.org/10.1186/s12979-020-00213-w>
 35. Biagi E, Franceschi C, Rampelli S, Severgnini M, Ostan R, Turrone S, Consolandi C, Quercia S, Scurti M, Monti D, Capri M, Brigidi P, Candela M. 2016. Gut microbiota and extreme longevity. *Curr Biol* 26:1480–1485. <https://doi.org/10.1016/j.cub.2016.04.016>
 36. David LA, Materna AC, Friedman J, Campos-Baptista MI, Blackburn MC, Perrotta A, Erdman SE, Alm EJ. 2014. Host lifestyle affects human microbiota on daily timescales. *Genome Biol* 15:R89. <https://doi.org/10.1186/gb-2014-15-7-r89>
 37. Mehta RS, Abu-Ali GS, Drew DA, Lloyd-Price J, Subramanian A, Lochhead P, Joshi AD, Ivey KL, Khalili H, Brown GT, DuLong C, Song M, Nguyen LH, Mallick H, Rimm EB, Izard J, Huttenhower C, Chan AT. 2018. Stability of the human faecal microbiome in a cohort of adult men. *Nat Microbiol* 3:347–355. <https://doi.org/10.1038/s41564-017-0096-0>
 38. Mancabelli L, Tarracchini C, Milani C, Lugli GA, Fontana F, Turrone F, van Sinderen D, Ventura M. 2020. Multi-population cohort meta-analysis of human intestinal microbiota in early life reveals the existence of infant community state types (ICSTs). *Comput Struct Biotechnol J* 18:2480–2493. <https://doi.org/10.1016/j.csbj.2020.08.028>
 39. Vandeputte D, De Commer L, Tito RY, Kathagen G, Sabino J, Vermeire S, Faust K, Raes J. 2021. Temporal variability in quantitative human gut microbiome profiles and implications for clinical research. *Nat Commun* 12:6740. <https://doi.org/10.1038/s41467-021-27098-7>
 40. López-Aladid R, Fernández-Barat L, Alcaraz-Serrano V, Bueno-Freire L, Vázquez N, Pastor-Ibáñez R, Palomeque A, Oscanoa P, Torres A. 2023. Determining the most accurate 16S rRNA hypervariable region for taxonomic identification from respiratory samples. *Sci Rep* 13:3974. <https://doi.org/10.1038/s41598-023-30764-z>
 41. Johnson JS, Spakowicz DJ, Hong B-Y, Petersen LM, Demkowicz P, Chen L, Leopold SR, Hanson BM, Agresta HO, Gerstein M, Sodergren E, Weinstock GM. 2019. Evaluation of 16S rRNA gene sequencing for species and strain-level microbiome analysis. *Nat Commun* 10:5029. <https://doi.org/10.1038/s41467-019-13036-1>

42. Milani C, Lugli GA, Fontana F, Mancabelli L, Alessandri G, Longhi G, Anzalone R, Viappiani A, Turrioni F, van Sinderen D, Ventura M. 2021. METAnnotatorX2: a comprehensive tool for deep and shallow metagenomic data set analyses. *mSystems* 6:e0058321. <https://doi.org/10.1128/mSystems.00583-21>
43. Hillmann B, Al-Ghalith GA, Shields-Cutler RR, Zhu Q, Gohl DM, Beckman KB, Knight R, Knights D. 2018. Evaluating the information content of shallow shotgun metagenomics. *mSystems* 3:e00069-18. <https://doi.org/10.1128/mSystems.00069-18>
44. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, Carty C, Chaput J-P, Chastin S, Chou R, et al. 2020. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med* 54:1451–1462. <https://doi.org/10.1136/bjsports-2020-102955>
45. Arumugam M, Raes J, Pelletier E, Le Paslier D, Yamada T, Mende DR, Fernandes GR, Tap J, Bruls T, Batto J-M, et al. 2011. Enterotypes of the human gut microbiome. *Nature* 473:174–180. <https://doi.org/10.1038/nature09944>
46. Lim MY, Rho M, Song YM, Lee K, Sung J, Ko G. 2014. Stability of gut enterotypes in Korean monozygotic twins and their association with biomarkers and diet. *Sci Rep* 4:7348. <https://doi.org/10.1038/srep07348>
47. de Moraes ACF, Fernandes GR, da Silva IT, Almeida-Pititto B, Gomes EP, Pereira A da C, Ferreira SRG. 2017. Enterotype may drive the dietary-associated cardiometabolic risk factors. *Front Cell Infect Microbiol* 7:47. <https://doi.org/10.3389/fcimb.2017.00047>
48. Neu AT, Allen EE, Roy K. 2021. Defining and quantifying the core microbiome: challenges and prospects. *Proc Natl Acad Sci USA* 118:e2104429118. <https://doi.org/10.1073/pnas.2104429118>
49. Caspi R, Billington R, Ferrer L, Foerster H, Fulcher CA, Keseler IM, Kothari A, Krummenacker M, Latendresse M, Mueller LA, Ong Q, Paley S, Subhraveti P, Weaver DS, Karp PD. 2016. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Res* 44:D471–80. <https://doi.org/10.1093/nar/gkv1164>
50. Horner TW, Dunn ML, Eggett DL, Ogden LV. 2011. β -Galactosidase activity of commercial lactase samples in raw and pasteurized milk at refrigerated temperatures. *J Dairy Sci* 94:3242–3249. <https://doi.org/10.3168/jds.2010-3742>
51. Edwards CA, Parrett AM. 2003. Dietary fibre in infancy and childhood. *Proc Nutr Soc* 62:17–23. <https://doi.org/10.1079/PNS2002231>
52. Mallick H, Rahnavard A, McIver LJ, Ma S, Zhang Y, Nguyen LH, Tickle TL, Weingart G, Ren B, Schwager EH, Chatterjee S, Thompson KN, Wilkinson JE, Subramanian A, Lu Y, Waldron L, Paulson JN, Franzosa EA, Bravo HC, Huttenhower C. 2021. Multivariable association discovery in population-scale meta-omics studies. *PLoS Comput Biol* 17:e1009442. <https://doi.org/10.1371/journal.pcbi.1009442>
53. Mancabelli L, Milani C, Lugli GA, Turrioni F, Ferrario C, van Sinderen D, Ventura M. 2017. Meta-analysis of the human gut microbiome from urbanized and pre-agricultural populations. *Environ Microbiol* 19:1379–1390. <https://doi.org/10.1111/1462-2920.13692>
54. Martínez I, Stegen JC, Maldonado-Gómez MX, Eren AM, Siba PM, Greenhill AR, Walter J. 2015. The gut microbiota of rural Papua New Guineans: composition, diversity patterns, and ecological processes. *Cell Rep* 11:527–538. <https://doi.org/10.1016/j.celrep.2015.03.049>
55. De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S, Collini S, Pieraccini G, Lionetti P. 2010. Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci USA* 107:14691–14696. <https://doi.org/10.1073/pnas.1005963107>
56. Porras AM, Shi QJ, Zhou H, Callahan R, Montenegro-Bethancourt G, Solomons N, Brito IL. 2021. Geographic differences in gut microbiota composition impact susceptibility to enteric infection. *Cell Rep* 36:109457. <https://doi.org/10.1016/j.celrep.2021.109457>
57. Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, Magris M, Hidalgo G, Baldassano RN, Anokhin AP, Heath AC, Warner B, Reeder J, Kuczynski J, Caporaso JG, Lozupone CA, Lauber C, Clemente JC, Knights D, Knight R, Gordon JI. 2012. Human gut microbiome viewed across age and geography. *Nature* 486:222–227. <https://doi.org/10.1038/nature11053>
58. Yoshii K, Hosomi K, Sawane K, Kunisawa J. 2019. Metabolism of dietary and microbial vitamin B family in the regulation of host immunity. *Front Nutr* 6:48. <https://doi.org/10.3389/fnut.2019.00048>
59. Calderón-Ospina CA, Nava-Mesa MO. 2020. B vitamins in the nervous system: current knowledge of the biochemical modes of action and synergies of thiamine, pyridoxine, and cobalamin. *CNS Neurosci Ther* 26:5–13. <https://doi.org/10.1111/cns.13207>
60. Lugli GA, Milani C, Mancabelli L, Turrioni F, van Sinderen D, Ventura M. 2019. A microbiome reality check: limitations of in silico-based metagenomic approaches to study complex bacterial communities. *Environ Microbiol Rep* 11:840–847. <https://doi.org/10.1111/1758-2229.12805>
61. Milani C, Casey E, Lugli GA, Moore R, Kaczorowska J, Feehily C, Mangifesta M, Mancabelli L, Duranti S, Turrioni F, Bottacini F, Mahony J, Cotter PD, McAuliffe FM, van Sinderen D, Ventura M. 2018. Tracing mother-infant transmission of bacteriophages by means of a novel analytical tool for shotgun metagenomic datasets: METAnnotatorX. *Microbiome* 6:145. <https://doi.org/10.1186/s40168-018-0527-z>
62. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <https://doi.org/10.1038/nmeth.1923>
63. Langmead B, Wilks C, Antonescu V, Charles R. 2019. Scaling read aligners to hundreds of threads on general-purpose processors. *Bioinformatics* 35:421–432. <https://doi.org/10.1093/bioinformatics/bty648>
64. Chen Y, Ye W, Zhang Y, Xu Y. 2015. High speed BLASTN: an accelerated MegaBLAST search tool. *Nucleic Acids Res* 43:7762–7768. <https://doi.org/10.1093/nar/gkv784>
65. Zhao Y, Tang H, Ye Y. 2012. RAPSearch2: a fast and memory-efficient protein similarity search tool for next-generation sequencing data. *Bioinformatics* 28:125–126. <https://doi.org/10.1093/bioinformatics/btr595>
66. Ye Y, Choi JH, Tang H. 2011. RAPSearch: a fast protein similarity search tool for short reads. *BMC Bioinformatics* 12:159. <https://doi.org/10.1186/1471-2105-12-159>
67. Paradis E, Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 35:526–528. <https://doi.org/10.1093/bioinformatics/bty633>
68. Benjamini Y, Drai D, Elmer G, Kafkafi N, Golani I. 2001. Controlling the false discovery rate in behavior genetics research. *Behav Brain Res* 125:279–284. [https://doi.org/10.1016/s0166-4328\(01\)00297-2](https://doi.org/10.1016/s0166-4328(01)00297-2)