



# UNIVERSITÀ DI PARMA

## ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Antidepressant-like effects of pharmacological inhibition of FAAH activity in socially isolated female rats

This is the peer reviewed version of the following article:

*Original*

Antidepressant-like effects of pharmacological inhibition of FAAH activity in socially isolated female rats / Carnevali, L.; Statello, R.; Vacondio, F.; Ferlenghi, F.; Spadoni, G.; Rivara, S.; Mor, M.; Sgoifo, A.. - In: EUROPEAN NEUROPSYCHOPHARMACOLOGY. - ISSN 0924-977X. - 32:(2020), pp. 77-87. [10.1016/j.euroneuro.2019.12.119]

*Availability:*

This version is available at: 11381/2884825 since: 2024-12-12T08:52:20Z

*Publisher:*

Elsevier B.V.

*Published*

DOI:10.1016/j.euroneuro.2019.12.119

*Terms of use:*

Anyone can freely access the full text of works made available as "Open Access". Works made available

*Publisher copyright*

note finali coverpage

(Article begins on next page)

**Antidepressant-like effects of pharmacological inhibition of FAAH activity in socially  
isolated female rats**

Luca Carnevali<sup>a</sup>, Rosario Statello<sup>a</sup>, Federica Vacondio<sup>b</sup>, Francesca Ferlenghi<sup>b</sup>, Gilberto Spadoni<sup>c</sup>,  
Silvia Rivara<sup>b</sup>, Marco Mor<sup>b</sup>, Andrea Sgoifo<sup>a</sup>

<sup>a</sup> Stress Physiology Lab, Department of Chemistry, Life Sciences and Environmental Sustainability,  
University of Parma, Parma, Italy

<sup>b</sup> Department of Food and Drug, University of Parma, Parma, Italy

<sup>c</sup> Department of Biomolecular Sciences, University of Urbino “Carlo Bo”, Urbino, Italy

**Corresponding author:** Andrea Sgoifo, Stress Physiology Lab, Department of Chemistry, Life  
Sciences and Environmental Sustainability, University of Parma, Parco Area delle Scienze 11/A,  
43124, Parma, Italy. Email address: [andrea.sgoifo@unipr.it](mailto:andrea.sgoifo@unipr.it)

**Total word-count:** 5636

## Abstract

Pharmacological inhibition of the enzyme fatty acid amide hydrolase (FAAH), which terminates signalling of the endocannabinoid N-arachidonylethanolamine (or anandamide, AEA), exerts favourable effects in rodent models of stress-related depression. Yet although depression seems to be more common among women than men and in spite of some evidence of sex differences in treatment efficacy, preclinical development of FAAH inhibitors for the pharmacotherapy of stress-related depression has been predominantly conducted in male animals. Here, adult female rats were exposed to six weeks of social isolation and, starting from the second week, treated with the FAAH inhibitor URB694 (0.3 mg/kg/day, i.p.) or vehicle. Compared to pair-housed females, socially isolated female rats treated with vehicle developed behavioral (mild anhedonia, passive stress coping) and physiological (reduced body weight gain, elevated plasma corticosterone levels) alterations. Moreover, prolonged social isolation provoked a reduction in brain-derived neurotrophic factor (BDNF) and AEA levels within the hippocampus. Together, these changes are indicative of an increased risk of developing a depressive-like state. Conversely, pharmacological inhibition of FAAH activity with URB694 restored both AEA and BDNF levels within the hippocampus of socially isolated rats and prevented the development of behavioral and physiological alterations. These results suggest a potential interplay between AEA-mediated signaling and hippocampal BDNF in the pathogenesis of depression-relevant behaviors and physiological alterations and antidepressant action of FAAH inhibition in socially isolated female rats.

**Keywords:** depression; stress; endocannabinoid; BDNF; females

## 1. Introduction

Prolonged or repeated exposure to stressors of psychosocial nature can act as a precipitating factor for the onset of depression (Cohen et al., 2007; Dinan, 2005). One of the most susceptible brain regions to the effects of psychosocial stress is the hippocampus, a component of the limbic system that regulates emotional and cognitive processes related to psychiatric disorders (Belleau et al., 2019; Sheline et al., 2019). The hippocampus is also a major regulator of the hypothalamic-pituitary-adrenal (HPA) axis (Jacobson and Sapolsky, 1991), the neuroendocrine system responsible for the release of glucocorticoid stress hormones (i.e., cortisol in humans, corticosterone in rodents). In patients with depression, hippocampal volume is decreased (Sapolsky, 2000; Sheline, 1996) and the HPA axis is dysregulated (Stetler and Miller, 2011). Depletion of hippocampal neurogenesis has been implicated as one of the substrates that may explain the hippocampal volume loss seen in depression (Duman and Monteggia, 2006; Levone et al., 2015). Specifically, the neurotrophic hypothesis of depression proposes that stress-induced reductions in the expression of brain-derived neurotrophic factor (BDNF), a member of the neurotrophin family regulating synaptic plasticity (Leal et al., 2017; Lu et al., 2014), occur in key limbic structures, including the hippocampus, to contribute to the pathogenesis of depression (Castren et al., 2007; Duman and Monteggia, 2006). Moreover, several lines of clinical and preclinical evidence indicate that conventional antidepressants (e.g., tricyclics, selective serotonin reuptake inhibitors and norepinephrine reuptake inhibitors) may in part exert their effects through BDNF upregulation (Hayley and Anisman, 2013; Pittenger and Duman, 2008; Tardito et al., 2006).

The past two decades have witnessed a driven focus on the identification of novel therapeutic targets for depression, in an attempt to overcome the notable limitations of conventional antidepressant treatments, poor efficacy being perhaps the most critical (Connolly and Thase, 2012). For example, substantial evidence has accumulated implicating a deficit in endocannabinoid (eCB) neurotransmission in the etiology of depression (for a comprehensive review see Gorzalka and Hill, 2011). At the preclinical level, a deficiency in the signaling mediated by the eCB N-arachidonylethanolamine (or anandamide, AEA) has been noted in the hippocampus, hypothalamus, ventral striatum, and prefrontal cortex of rats exposed to several stressors (i.e.,

67 chronic unpredictable stress and social defeat stress) and presenting a "depressive-like" phenotype  
68 (reviewed in Carnevali et al., 2017b). These findings have triggered significant interest in the  
69 development of eCB-interacting drugs, including direct-acting receptor ligands and catabolism  
70 inhibitors for the pharmacotherapy of depression (Micale et al., 2013). Specifically, within preclinical  
71 models, facilitation of AEA signaling through pharmacological inhibition of its degrading enzyme (i.e.,  
72 fatty acid amide hydrolase (FAAH)) can enhance monoaminergic transmission, increase cellular  
73 plasticity and neurotrophin expression within the hippocampus, dampen HPA axis activity, and evoke  
74 antidepressant-like behavioral effects (reviewed in Carnevali et al., 2017b). However, while the  
75 literature has been unequivocal in showing that women experience depression at twice the rate of  
76 men (e.g., Grigoriadis and Robinson, 2007), very few preclinical studies have been conducted on  
77 female experimental animals (Beery, 2018; Kokras and Dalla, 2014). Moreover, despite the  
78 existence of sex differences in response to antidepressant treatment (Sloan and Kornstein, 2003),  
79 preclinical research on the antidepressant action of FAAH inhibitors has been predominantly  
80 conducted in male rodents (Carnevali et al., 2017b; Fowler, 2015). Therefore, there is a clear need  
81 to use female animals in preclinical models of stress to either confirm and generalize to females the  
82 previously obtained male animal-based findings or underscore potential sex differences in the  
83 etiology of depression and/or in the efficacy of new treatments.

84 Based on this background, the purpose of the current study was two-fold. First, we aimed at  
85 documenting the development of behavioral (passive stress coping, anhedonia) and biological  
86 (reduced hippocampal BDNF levels, HPA axis hyperactivity, body weight loss) alterations in adult  
87 female rats exposed to prolonged social isolation, a mild chronic social stressor that has been widely  
88 used to model symptoms that are often associated with an increased risk of developing a depressive-  
89 like state in rodents (Carnevali et al., 2017a). Second, we tested the hypothesis that pharmacological  
90 inhibition of FAAH activity would correct the alterations associated with prolonged social isolation.  
91 To this aim, we employed the FAAH inhibitor URB694 (6-hydroxy-[1,1'-biphenyl]-3-yl-  
92 cyclohexylcarbamate) which was shown to exhibit higher selectivity and more prolonged and  
93 profound access to the brain than the standard inhibitor URB597 (Clapper et al., 2009).

94

## 2. Experimental procedures

### 2.1. Animals and housing conditions

Four-month-old female wild-type Groningen rats were used in this study. This rat population, originally derived from the University of Groningen (the Netherlands) and currently bred in our laboratory under standard conditions, shows considerable individual differences in trait-like patterns of behavioral and physiological responses to environmental challenges (Carnevali et al., 2014; de Boer et al., 2017). After weaning, female animals were housed in same-sex sibling pairs and kept in rooms with controlled temperature ( $22 \pm 2$  °C) and humidity ( $50 \pm 10$  %), under a reversed light-dark cycle (light on from 19:00 to 7:00 h), with food and water ad libitum except when required for the sucrose preference test (see below). A total of 40 pairs were included in the study, but only one female rat from each pair was submitted to the experimental procedures described below. Experiments were performed in accordance with the European Community Council Directive 2010/63/UE and approved by the Italian legislation on animal experimentation (D.L. 04/04/2014, n. 26, authorization n. 449/2017-PR). All efforts were made to reduce sample size and minimize animal suffering.

### 2.2. Experimental design

The experimental timeline is depicted in Figure 1. Specific procedures and data analysis are described in the following sections. On day 0, animals were randomly divided in socially isolated (SI) and paired-housed (PH) groups. Female rats from the SI group were separated from their respective sibling and individually housed in a soundproof room for 6 weeks to avoid any sensory (visual, olfactory, and acoustic) contact with their conspecifics. On the contrary, female rats from the PH group were continually housed with their respective sibling and kept in the same room with other pairs. Handling and cage cleaning were matched between the two groups. Starting from the beginning of the third week of the social isolation/pair-housing condition, animals received daily i.p. injection of either the FAAH inhibitor URB694 or vehicle (VEH). Thus, four experimental subgroups emerged: (i) SI + VEH (n = 10), (ii) SI + URB694 (n = 10), (iii) PH + VEH (n = 10), and (iv) PH + URB694 (n = 10). Experiments were conducted on separate cohorts of 8 experimental animals each

(n = 4 SI and n = 4 PH rats), starting with the VEH-treated animals. Experimental animals were tested four times in the sucrose preference test and once in the forced swim test during the dark phase of the daily cycle between 10.00 and 12.00 h. At sacrifice (day 42), trunk blood, adrenal glands, and hippocampus were harvested. Body weight was measured weekly throughout the study. Moreover, the estrous cycle phase of female rats was determined immediately after each behavioral test and before sacrifice using vaginal smear cytology. Vaginal smears were collected by gently introducing a moistened (0.9% NaCl) cotton swab in the rat's vagina. The sample was transferred to a glass slide and examined microscopically following Giemsa staining. The phase of the cycle (metaestrous, diestrous, pro-estrous or estrous) was determined based upon the presence of leukocytes, nucleated epithelial or cornfield epithelial cells (Marcondes et al., 2002).

133

### 2.3. Drug treatment

URB694 is a carbamate FAAH inhibitor that irreversibly carbamoylates the nucleophile catalytic serine in FAAH active site (Tarzia et al., 2006). URB694 is a second generation inhibitor with improved metabolic stability and selectivity for FAAH (Clapper et al., 2009). URB694 was freshly dissolved in VEH containing 5% PEG, 5% Tween 80, and 90% saline. VEH (vol:1 ml/kg) or URB694 (0.3 mg/kg, i.p.) were injected i.p. between 11.00 and 13.00 h and, on the days of the sucrose solution and forced swim tests, at least 1 h after the completion of the test. URB694 dose was chosen based on our previous studies (Carnevali et al., 2015a; Carnevali et al., 2015b), and a pilot study showing that FAAH activity in the brain of female wild-type Groningen rats was substantially inhibited 24 h after administration of this drug dose (Supplemental Figure S1).

144

### 2.4. Sucrose preference test

*Ad libitum* 2% sucrose solution was available for 5 days before the beginning of the experimental procedures to allow adaptation to its taste. Food and water were removed from the cage for 16 hours before each sucrose preference test; moreover, one hour before the test, all experimental animals (paired and isolated) were moved into individual cages to ensure accurate fluid intake measurements of paired animals. Water and 2% sucrose solution were placed in premeasured bottles in the

individual cage, and fluid intake was monitored for 1 hour. Animals were returned to their respective home cages immediately after the test (Grippe et al., 2007). Sucrose preference tests were conducted in baseline conditions (day -3) and after 11, 25, and 39 days of social isolation (Figure 1). Sucrose solution intake was expressed as the relative percentage of the total liquid intake and was taken as an operational index of anhedonia, defined as reduced sucrose preference relative to control animals and baseline values (Grippe et al., 2007).

157

## 158 2.5. Forced swim test

159 An adapted version of the forced swim test originally described by Porsolt (Porsolt et al., 1977) was  
160 used. On day 35 (Fig. 1), female rats were forced to swim individually for 5 min in a Plexiglas cylinder  
161 (height: 40 cm, diameter: 30 cm) filled with water (temperature:  $24 \pm 2^{\circ}\text{C}$ ; depth: 30 cm). During the  
162 test, rats' behavior was video-taped. The overall time spent in immobility (floating and making only  
163 those movements necessary to keep the head above water) was scored by a trained experimenter  
164 blind to animals' condition and treatment. Immobility during the single session of the forced swim  
165 test was used as an index of passive stress coping (Commons et al., 2017).

166

## 167 2.6. Measurements at sacrifice

168 Twenty-four hours after the last administration of URB694 or VEH (i.e., at 11.00 h; day 42, Figure  
169 1), female rats were euthanized by decapitation under isoflurane anesthesia (2% in 100% oxygen).  
170 Trunk blood was collected in EDTA-coated tubes (Sarsted AG, Numbrecht, Germany) and plasma  
171 was separated by centrifugation (2600 g,  $4^{\circ}\text{C}$ , 10 min). Brains were immediately removed and the  
172 hippocampus rapidly dissected and snap-frozen in nitrogen. All samples were stored at  $-80^{\circ}\text{C}$  until  
173 further analysis, as described below. Adrenal glands were also removed and weighed.

### 174 2.6.1. Plasma corticosterone levels

175 Plasma was deproteinized by addition of two volumes of organic solvent (ice-cold acetonitrile),  
176 containing the internal standard dexamethasone (structural analog of corticosterone, 75 nmol/L).  
177 After centrifugation (14000 g,  $4^{\circ}\text{C}$ , 10 min), the supernatant was directly injected in the liquid  
178 chromatography/tandem mass spectrometry system (HPLC/MS/MS) for quantification of



179 corticosterone levels, in accordance with previously published analytical methods (Plenis et al.,  
180 2011). A detailed description of the HPLC/MS/MS analytical method and related MS instrumentation  
181 is reported in the Supplemental Material.

#### 182 2.6.2. *BDNF hippocampal content*

183 BDNF content in the hippocampus was measured using a commercially available sandwich enzyme-  
184 linked immune sorbent assay (ELISA) kit (Quantikine ®ELISA-Total BDNF, R&D Systems,  
185 Minneapolis, MN, USA) according to the manufacturer's instructions. A detailed description of the  
186 experimental procedure is reported in the Supplemental Material. BDNF tissue content was  
187 expressed as a percentage of the control group (PH+VEH rats).

#### 188 2.6.3. *AEA hippocampal levels*

189 AEA was extracted from 10% w/v hippocampal tissue homogenates employing two volumes of ice-  
190 cold acetonitrile containing the deuterated internal standard AEA-d<sub>4</sub> and quantified by HPLC/MS/MS  
191 as previously reported (Carnevali et al., 2015a) The analytical standards AEA and AEA-d<sub>4</sub> were  
192 purchased from Cayman Chemical (Ann Arbor, MI, USA) as stock solutions in ethanol. AEA levels  
193 were expressed as pmol/g wet weight of tissue. A detailed description of the HPLC/MS/MS analytical  
194 method and related MS instrumentation is reported in the Supplemental Material.

#### 195 2.6.4. *FAAH activity in the hippocampus*

196 For ex vivo determination of FAAH activity, frozen hippocampi were thawed and homogenized in ice-  
197 cold Tris buffer (10 volumes, 50 mM, pH 7.5) containing 0.32 M sucrose. The homogenates were  
198 centrifuged (1000 g, 10 min, 4°C) and total protein content was quantified in the supernatant by the  
199 bicinchoninic acid (BCA) protein kit (Pierce Biotechnology, Rockford, IL, USA). FAAH activity was  
200 measured at 37°C for 30 min in 0.5 mL Tris buffer (50 mM, pH 7.5) containing fatty acid-free bovine  
201 serum albumin (BSA) (0.05%, w/v), 50 µg of protein from brain homogenates, 10 µM AEA and [<sup>3</sup>H]-  
202 AEA (10000 disintegrations per minute) as previously described (Clapper et al., 2009). Briefly, the  
203 reactions were stopped with 1 mL chloroform:methanol (1:1). After centrifugation (2000 g, 10 min,  
204 4°C), [<sup>3</sup>H]-ethanolamine was measured in the aqueous phase by liquid scintillation counting. [<sup>3</sup>H]-  
205 AEA (specific activity: 60 Ci/mmol), employed as a substrate for ex vivo FAAH assay, was purchased  
206 from American Radiolabeled Chemicals (St. Louis, MI, USA).

## 207 2.7. Statistical analysis

208 All statistical analyses were performed using SPSS v. 25 (IBM software package). Data are  
209 presented as mean  $\pm$  standard error of the mean (SEM). The influence of the estrous cycle phase  
210 on behavioral and biochemical measurements was controlled in all statistical analyses. A three-way  
211 ANOVA for repeated measures with “condition” (2 levels: isolation, pair-housing) and “treatment” (2  
212 levels: VEH, URB694) as the between subject factors, and “time” as the within subject factor (3  
213 levels: days 11, 25, and 39) was applied on delta changes in sucrose solution preference with respect  
214 to baseline. All other data were analyzed with 2 (factor “condition”: isolation or pair-housing) x 2  
215 (factor “treatment”: URB694 or VEH) factorial design ANOVAs. Follow-up analyses were conducted  
216 using Student’s “t” tests, with a Bonferroni correction for multiple comparisons. Pearson’s  $r$   
217 correlations were performed to assess the correlation between plasma corticosterone levels, BDNF  
218 hippocampal content and AEA hippocampal levels. Statistical significance was set at  $p < 0.05$ .

219

## 220 3. Results

### 221 3.1. Body weight

222 There were no significant differences in body weight among groups at the start of the experiment  
223 (i.e., when animals were assigned to the different housing conditions) (PH + VEH =  $230 \pm 2$  g; IS +  
224 VEH =  $237 \pm 5$  g; PH + URB694 =  $231 \pm 4$  g; IS + URB694 =  $226 \pm 8$  g). However, a significant time  
225 x condition interaction emerged on body weight gain calculated as the difference between weight at  
226 the end (i.e., immediately before animals were euthanized) and at the start of the experiment ( $F =$   
227  $7.1$ ,  $p = .012$ ). As shown in Figure 2, socially isolated female rats treated with VEH gained  
228 significantly less weight compared with their respective pair-housed counterparts ( $p = .002$ ). This  
229 effect of social isolation was prevented by URB694 treatment (SI + URB694 vs SI + VEH,  $p = .012$ ).

### 230 3.2. Sucrose preference test

231 Total fluid intake did not differ among groups at each assessment point (Supplemental Table S1).  
232 Also, there were no significant differences among groups in their baseline preference for the  
233 consumption of the sucrose solution (PH + VEH =  $85 \pm 2$  %; IS + VEH =  $88 \pm 2$  %; PH + URB694 =  
234  $83 \pm 3$  %; IS + URB694 =  $82 \pm 3$  %). Of note, the estrous cycle phase had no effect on baseline

sucrose solution preference ( $F = 0.3$ ,  $p = .543$ ). However, factorial ANOVA yielded a significant time x condition interaction ( $F = 5.1$ ,  $p = .028$ ) on preference changes during the social isolation period (calculated as the difference between each assessment point and the baseline), with no significant effects of the estrous cycle phase ( $F = 0.4$ ,  $p = .497$ ). Specifically, as shown in Figure 3, no group differences were observed on day 11. However, on day 25, socially-isolated female rats treated with VEH showed a significantly larger reduction in the preference for sucrose solution consumption compared with their respective pair-housed counterparts ( $p = .025$ ). This effect was prevented by URB694 treatment (SI + URB694 vs SI + VEH,  $p = .003$ ). A similar trend was observed on day 39, although differences did not reach full statistical significance (SI + VEH vs PH + VEH,  $p = .056$ ; SI + VEH vs SI + URB694,  $p = .067$ ).

### 3.3. Forced swim test

Behavior during the forced swim test is illustrated in Figure 4. Factorial ANOVA yielded a significant effect of treatment ( $F = 4.9$ ,  $p = .033$ ) and a strong trend for condition x treatment interaction ( $F = 3.5$ ,  $p = .071$ ) on immobility time, with no significant effects of estrous cycle phase ( $F = 0.2$ ,  $p = .632$ ). Specifically, socially isolated female rats treated with VEH spent significantly more time in immobility compared with their respective pair-housed counterparts ( $p = .024$ ). This behavioral effect of social isolation was significantly corrected by URB694 treatment (SI + URB694 vs SI + VEH,  $p = .007$ ).

### 3.4. Measurements at sacrifice

#### 3.4.1. Plasma corticosterone levels and adrenal weight

Factorial ANOVA yielded a significant condition x treatment interaction ( $F = 7.1$ ,  $p = .012$ ) on plasma corticosterone levels at the end of the experimental protocol, with no significant effects of the estrous cycle phase ( $F = 0.6$ ,  $p = .430$ ). As depicted in Figure 5, socially isolated female rats treated with VEH had significantly higher plasma corticosterone levels than their respective pair-housed counterparts ( $p = .016$ ). URB694 treatment prevented the effect of social isolation on plasma corticosterone levels (SI + URB694 vs SI + VEH,  $p = .003$ ).

There were no significant effects of condition and/or treatment on adrenal weight corrected for body weight at the end of the experiment (PH + VEH =  $0.021 \pm 0.002$  mg/g; IS + VEH =  $0.027 \pm 0.003$  mg/g; PH + URB694 =  $0.027 \pm 0.002$  mg/g; IS + URB694 =  $0.026 \pm 0.002$  mg/g).

### 263 3.4.2. *BDNF hippocampal content*

264 Factorial ANOVA yielded a significant effect of treatment ( $F = 7.3$ ,  $p = .012$ ) and a significant  
265 condition x treatment interaction ( $F = 6.9$ ,  $p = .014$ ) on BDNF content in the hippocampus at the end  
266 of the experimental protocol. As illustrated in Figure 6A, socially isolated female rats treated with  
267 VEH showed a significantly lower BDNF hippocampal content compared with their respective pair-  
268 housed counterparts ( $p = .023$ ). This effect of social isolation was prevented by URB694 treatment  
269 (SI + URB694 vs SI + VEH,  $p = .001$ ). Moreover, we found a negative, although not significant,  
270 correlation between plasma corticosterone levels and BDNF hippocampal content (Table 1).

### 271 3.4.3. *AEA hippocampal levels*

272 Factorial ANOVA yielded significant effects of condition ( $F = 19.7$ ,  $p < .001$ ) and treatment ( $F = 27.6$ ,  
273  $p < .001$ ), and a significant condition x treatment interaction ( $F = 5.3$ ,  $p = .028$ ) on AEA hippocampal  
274 levels at the end of the experimental protocol. As shown in Figure 6B, socially isolated female rats  
275 treated with VEH showed significantly lower AEA hippocampal levels compared with their respective  
276 pair housed counterpart ( $p < .001$ ). As expected, URB694-treated groups showed significantly  
277 greater AEA levels than corresponding VEH-treated groups, both in the social isolation ( $p < .001$ )  
278 and pair-housing ( $p = .040$ ) condition. Moreover, we found a significant positive correlation between  
279 AEA levels and BDNF content within the hippocampus (Table 1), as well as a strong trend for a  
280 negative correlation between AEA hippocampal levels and plasma corticosterone levels (Table 1).

### 281 3.4.4. *FAAH activity*

282 Factorial ANOVA yielded a significant effect of treatment ( $F = 456.0$ ,  $p < .001$ ) on FAAH activity in  
283 the hippocampus, being, as expected, significantly lower in URB694-treated than VEH-treated rats  
284 in both the social isolation ( $p < .001$ ) and pair-housing ( $p < .001$ ) condition (Figure 6C).

285

## 286 4. Discussion

287 The major findings of the current investigation are the following. Compared to pair-housed females,  
288 socially isolated female rats developed behavioral (mild anhedonic state, passive stress coping) and  
289 physiological (reduced body weight gain, elevated plasma corticosterone levels) changes, and  
290 showed a reduction in BDNF and AEA levels within the hippocampus. Together, these changes are

291 indicative of an increased risk of developing a depressive-like state. Notably, pharmacological  
292 inhibition of FAAH activity with URB694 restored AEA and BDNF hippocampal levels, and prevented  
293 the development of behavioral and physiological alterations following prolonged social isolation.

#### 294 4.1. Depressive-like changes in socially isolated female rats

295 Psychiatric disorders in humans have been linked prevalently with social stress and/or reduced  
296 social interaction (Bjorkqvist, 2001; Heinrich and Gullone, 2006). Within preclinical models, the social  
297 defeat paradigm has been shown to have a substantial impact on depression-relevant behavioral  
298 and physiological parameters in adult male rats, while solitary housing is particularly effective in  
299 precipitating depressive-like symptoms in previously group-housed female rats (Beery and Kaufer,  
300 2015; Carnevali et al., 2017a). Of note, the social isolation protocol adopted in this study included  
301 both solitary housing and long-term deprivation of sensory stimuli originating from the surrounding  
302 social environment. Therefore, it is likely that the described effects are due to a combination of both.  
303 Specifically, female rats showed a reduction in body weight gain, signs of a mild anhedonic-like state  
304 (i.e., reduced preference for the consumption of a sucrose solution), passive coping (i.e., increased  
305 immobility in the forced swim test), and elevated plasma corticosterone levels. Deficits in body weight  
306 gain in isolated rats may be explained by reduced food intake, as previously demonstrated in  
307 individually housed mice and rats (Izadi et al., 2018; Sun et al., 2014), particularly around light-dark  
308 phase transitions (Sun et al., 2014). Interestingly, reductions in heat production and in the respiratory  
309 exchange ratio were also found during light-dark transitions in individually housed mice (Sun et al.,  
310 2014), suggesting that metabolic functions may have been affected also in our socially isolated rats.  
311 Moreover, the mild reduction in the preference for the consumption of a palatable solution observed  
312 only after 25 days of social isolation resembles the time course of changes reported in female Wistar  
313 rats exposed to chronic mild stress (Grippe et al., 2005) and in socially isolated female prairie voles  
314 (Grippe et al., 2007). However, we acknowledge that the interpretation of this result is limited by the  
315 difference, albeit not statistically significant, between the two stressed groups on day 11 (i.e., before  
316 the start of the pharmacological treatment). Notably, the estrous cycle phase did not seem to have  
317 any effect on any of the behavioral and biological variables assessed in the current study, although  
318 our analysis is limited by the small sample size given that four different stages were considered.

319 Nevertheless, this is in line with empirical research across multiple rodent species demonstrating  
320 that estrous cyclicity is not a major source of variability in females or, at least, is not greater than  
321 intrinsic variability in males (Beery, 2018; Finnell et al., 2018; Kokras et al., 2015).

322 Animal and human studies have provided support for the role of stress in the pathogenesis of  
323 depression via alterations in BDNF-mediated signaling (Hashimoto, 2010; Stepanichev et al., 2014),  
324 a neurotrophin that primarily regulates synaptic plasticity (Leal et al., 2017; Lu et al., 2014). In line  
325 with these findings, we found that BDNF content was reduced in the hippocampus of socially isolated  
326 female rats with depressive-like symptoms. Remarkably, such downregulation of hippocampal BDNF  
327 was paralleled by a decrease in AEA hippocampal levels. Converging lines of evidence support the  
328 possibility that AEA signaling at the cannabinoid receptor 1 (CB1R) may be an important mediator  
329 of neuroplastic phenomena within the hippocampus (Aguado et al., 2005; Hashimotodani et al.,  
330 2007; Hill et al., 2010; Scarante et al., 2017; Burstein et al., 2018). Particularly relevant for the current  
331 results are findings of decreased BDNF levels in the hippocampus of CB1R knockout mice (Aso et  
332 al., 2008). Thus, we hypothesize that a deficiency in AEA-mediated signaling at the CB1R might be  
333 implicated in the downregulation of BDNF hippocampal content observed in socially isolated female  
334 rats. Moreover, the positive correlation found here between AEA levels and BDNF content further  
335 supports a role for the eCB system in adult hippocampal neurogenesis (Scarante et al., 2017).

336 Notably, while one study reported a similar decrease in AEA levels in the hippocampus of chronically  
337 stressed male rats (Hill et al., 2008), other studies showed no changes in AEA hippocampal levels  
338 upon chronic stress exposure (Bortolato et al., 2007; Carnevali et al., 2015a; Hill et al., 2005). Of  
339 note, our data suggest that reduced AEA levels in the hippocampus of socially isolated rats were not  
340 due to an upregulation of FAAH enzymatic activity. This is in line with previous studies showing that  
341 FAAH activity is not affected by chronic stress exposure in rats (Bortolato et al., 2007; Hill et al.,  
342 2008), suggesting that the stress-induced decline in the hippocampal pool of AEA might be due to  
343 diminished biosynthetic mechanisms. Empirical evidence indicates the eCB system may be a  
344 biochemical effector of glucocorticoids in the brain (Hill and McEwen, 2010). Notably, the  
345 hippocampus itself is particularly sensitive to the action of glucocorticoid stress hormones due the  
346 rich concentration of receptor sites for glucocorticoids (De Kloet et al., 1998). The negative, although

only marginally significant, correlation found between plasma corticosterone levels and AEA hippocampal levels prompts further investigation into the specific mechanisms underlying the effects of stress exposure on AEA metabolism and their causal relationship with BDNF hippocampal downregulation. Interestingly, sex-specific mechanisms of eCB-mediated synaptic modulation within the hippocampus have been proposed to partly explain sex disparities in prevalence of depression (Huang and Woolley, 2012; Tabatadze et al., 2015). Decreased levels of BDNF may contribute to the atrophy of the hippocampus that has been observed in patients with depression (Sheline, 1996; Sheline et al., 2019). Recently, Belleau and colleagues (Belleau et al., 2019) proposed a model according to which chronic life stress can trigger the initial development of hippocampal volume reduction. However, this reduction would be neither necessary nor sufficient to produce a major depressive episode (Belleau et al., 2019). On the other hand, stress also initiates a set of neurotoxic processes (HPA axis dysregulation, inflammation, and neurotransmitter disturbances) that interact and may drive the development of a more chronic type of depression marked by further hippocampal volume reduction (Belleau et al., 2019). Although hippocampal volume was not assessed in the current study, we speculate that AEA–BDNF interactions might be implicated in the development of depressive symptoms and hippocampal volume decline under chronic life stress. Future longitudinal studies in rodent models of social stress may be informative in this regard.

#### 4.2. Antidepressant-like effects of the FAAH inhibitor URB694

In an attempt to replicate findings of our previous study demonstrating antidepressant-like effects of the FAAH inhibitor URB694 in chronically stressed male rats (Carnevali et al., 2015a), pharmacological treatment with URB694 started after two weeks of social isolation (i.e., we anticipated that depressive-like behaviors would already have begun to manifest by then). However, contrary to our expectations, we failed to conclusively demonstrate the onset of an anhedonic-like state before the start of the treatment. Thus, the fact that URB694-treated females did not show depressive-like behavioral and biological symptoms after a prolonged period of social isolation suggests, more cautiously, that inhibition of FAAH activity represents an effective preventive measure in this animal model. These results are in line with a growing body of evidence demonstrating that pharmacological inhibition of FAAH activity produces an antidepressant response

375 in chronically stressed male rodents (Carnevali et al., 2017b). Interestingly, FAAH inhibitors have  
376 been shown to increase hippocampal neurogenesis in adult rats (Goncalves et al., 2008; Hill et al.,  
377 2006; Marchalant et al., 2009) and prevent stress-induced BDNF downregulation in the brain  
378 (Burstein et al., 2018), supposedly via facilitation of CB1R-mediated activation of the extracellular  
379 signal-regulated kinase signaling pathway (Derkinderen et al., 2003; Rubino et al., 2006). Therefore,  
380 given that CB1Rs are highly abundant in the rodent (and human) hippocampus (Mackie, 2005), we  
381 hypothesize that the antidepressant-like action of the FAAH inhibitor URB694 in socially isolated  
382 female rats may be partly mediated by a preservation of hippocampal BDNF content via  
383 enhancement of AEA signaling at the CB1R. However, the antidepressant-like effects of URB694  
384 may also be interpreted in light of experimental evidence showing that AEA-signaling enhancement  
385 at the CB1R facilitates adaptive stress coping behaviors (Haller et al., 2013) and attenuates the  
386 neuroendocrine response to psychological stressors (Gorzalka et al., 2008). Moreover, given that  
387 FAAH inhibitors also increase the levels of other fatty acid amines with activity at peroxisome  
388 proliferator activated receptor- $\alpha$  (N-oleoylethanolamine (OEA) and N-palmitoylethanolamine (PEA)),  
389 the possibility of other non-cannabinoid receptor-mediated mechanisms cannot be completely ruled  
390 out. For example, a growing body of preclinical evidence suggests that PEA could have  
391 antidepressant-like activity (De Gregorio et al., 2019). On the other hand, increases in the  
392 endogenous levels of OEA may reduce food intake by regulating systems that control hunger and  
393 satiety in the brain (Romano et al., 2015). However, these compounds might also prolong and  
394 enhance AEA biological activity by competing with AEA for FAAH-mediated degradation (Petrosino  
395 et al., 2009). Of note, the current drug regimen had no effects on control animals, suggesting that  
396 the FAAH inhibitor did not affect normal biological processes and behavioral responses.

397

#### 398 4.3. Conclusion

399 The results of this study suggest a potential interplay between AEA-mediated signaling and BDNF  
400 at the level of the hippocampus in the development of depression-relevant behaviors and  
401 physiological changes in female rats exposed to prolonged social isolation. Moreover, the current  
402 results document the ability of the FAAH inhibitor URB694 to correct the alterations associated with



403 prolonged social isolation. One should note, however, that the current results are merely suggestive,  
404 and their translational implications for depression should be interpreted within the context of their  
405 limitations. First, the behavioral and biological changes described here after social isolation are often  
406 associated with an increased risk of developing major depression, but are not clinical symptoms of  
407 major depression per se. Second, we adopted a rodent model of prolonged social isolation, which  
408 should not be intended as a diagnostic model, but rather as a model of risk and vulnerability factor  
409 of stress-related depression. Moreover, we must acknowledge that, at present, clinical research on  
410 FAAH inhibitors has been slowed down by the serious adverse effects caused by the FAAH inhibitor  
411 BIA 10-2474 for the treatment of pain (von Schaper, 2016), which displayed both intrinsic toxic  
412 effects at high doses and off-targets effects (van Esbroeck et al., 2017). Investigations conducted by  
413 a Temporary Specialist Scientific Committee concluded that the toxicity of BIA 10-2474 is unlikely  
414 due to FAAH inhibition (Temporary Specialist Scientific Committee, 2016). A communication from  
415 the U.S. Food and Drug Administration also reported that the unique toxicity of this drug does not  
416 extend to other FAAH inhibitors (Food and Drug Administration, 2016), which are well tolerated by  
417 patients enrolled in clinical trials, and remarkably lack of the common adverse events elicited by  
418 exogenous cannabinoid-like compounds, including impairment in cognition, motor coordination, and  
419 psychoses (Mallet et al., 2016). The disorders for which these agents are being tested are mostly  
420 neuropsychiatric, such as pain conditions, depression, anxiety disorders, and phobias (Mallet et al.,  
421 2016). Nevertheless, the current results in female rats and previous research in male rodents using  
422 the carbamate FAAH inhibitors URB597 (e.g. Bortolato et al., 2007) and URB694 (Carnevali et al.,  
423 2015a) warrant more translational studies to examine the mood-modulating properties of this class  
424 of FAAH inhibitors (Gururajan et al., 2019). Recently, sex differences in hippocampal response to  
425 pharmacological inhibition of FAAH activity have been reported in rats after acute intense stress  
426 (Zer-Aviv and Akirav, 2016). This suggests that preclinical development of FAAH inhibitors for the  
427 pharmacotherapy of stress-related depression should aim at comparing the underlying  
428 neurobiological mechanisms between males and females.

429 **Table 1** Correlation matrix between plasma corticosterone levels, brain-derived neurotrophic factor  
 430 (BDNF) hippocampal content, and anandamide (AEA) hippocampal levels at the end of the  
 431 experimental protocol.

		Corticosterone	BDNF	AEA
Corticosterone	r	-		
	p			
BDNF	r	-.32	-	
	p	.082		
AEA	r	-.31	.44	-
	p	.068	.015	

432

433

434 **Figure legends**

435 **Figure 1.** Timeline of experimental procedures.

436

437 **Figure 2.** Body weight gain of paired-housed (PH) and socially isolated (SI) female rats treated with  
438 vehicle (VEH) or URB694, calculated as the difference between weight at the end (immediately  
439 before animals were euthanized) and at the start (when animals were assigned to the different  
440 housing conditions) of the experiment (n = 10 per group). Data are expressed mean $\pm$ SEM. \* =  
441 significantly different from corresponding PH + VEH group; # = significantly different from  
442 corresponding SI + VEH group (p values are reported in the text).

443

444 **Figure 3.** Changes in sucrose solution preference in paired-housed (PH) and socially isolated (SI)  
445 female rats treated with vehicle (VEH) or URB694, calculated as the difference between each  
446 assessment point during the social isolation period and the baseline (n = 10 per group). Data are  
447 expressed mean $\pm$ SEM. \* = significantly different from corresponding PH + VEH group; # =  
448 significantly different from corresponding SI + VEH group (p values are reported in the text).

449

450 **Figure 4.** Time spent in immobility during the forced swim test by paired-housed (PH) and socially  
451 isolated (SI) female rats treated with vehicle (VEH) or URB694 (n = 10 per group). Data are  
452 expressed mean $\pm$ SEM. \* = significantly different from corresponding PH + VEH group; # =  
453 significantly different from corresponding SI + VEH group (p values are reported in the text).

454

455 **Figure 5.** Plasma corticosterone levels at the end of the experimental protocol in paired-housed (PH)  
456 and socially isolated (SI) female rats treated with vehicle (VEH) or URB694 (n = 10 per group). Data  
457 are expressed mean $\pm$ SEM. \* = significantly different from corresponding PH + VEH group; # =  
458 significantly different from corresponding SI + VEH group (p values are reported in the text).

459

460 **Figure 6.** Brain-derived neurotrophic factor (BDNF; panel A) and anandamide (panel B) levels, and  
461 fatty acid amide hydrolase (FAAH) activity (panel C) in the hippocampus of paired-housed (PH)  
462 and socially isolated (SI) female rats treated with vehicle (VEH) or URB694 (n = 10 per group).

463 Data are expressed mean $\pm$ SEM. BDNF values are expressed as a percentage of the control group  
464 (PH+VEH rats). \* = significantly different from corresponding PH + VEH group; # = significantly  
465 different from corresponding VEH group (p values are reported in the text).

466

## 467   **References**

- 468   Aguado, T., Monory, K., Palazuelos, J., Stella, N., Cravatt, B., Lutz, B., Marsicano, G., Kokaia, Z.,  
469   Guzman, M., Galve-Roperh, I., 2005. The endocannabinoid system drives neural progenitor  
470   proliferation. *Faseb J.* 19, 1704-1706.
- 471   Aso, E., Ozaita, A., Valdizan, E.M., Ledent, C., Pazos, A., Maldonado, R., Valverde, O., 2008. BDNF  
472   impairment in the hippocampus is related to enhanced despair behavior in CB1 knockout mice. *J.*  
473   *Neurochem.* 105, 565-572.
- 474   Beery, A.K., 2018. Inclusion of females does not increase variability in rodent research studies. *Curr.*  
475   *Opin. Behav. Sci.* 23, 143-149.
- 476   Beery, A.K., Kaufer, D., 2015. Stress, social behavior, and resilience: insights from rodents.  
477   *Neurobiol. Stress* 1, 116-127.
- 478   Belleau, E.L., Treadway, M.T., Pizzagalli, D.A., 2019. The Impact of Stress and Major Depressive  
479   Disorder on Hippocampal and Medial Prefrontal Cortex Morphology. *Biol Psychiatry* 85, 443-453.
- 480   Bjorkqvist, K., 2001. Social defeat as a stressor in humans. *Physiol. Behav.* 73, 435-442.
- 481   Bortolato, M., Mangieri, R.A., Fu, J., Kim, J.H., Arguello, O., Duranti, A., Tontini, A., Mor, M., Tarzia,  
482   G., Piomelli, D., 2007. Antidepressant-like activity of the fatty acid amide hydrolase inhibitor URB597  
483   in a rat model of chronic mild stress. *Biol. Psychiatry* 62, 1103-1110.
- 484   Burstein, O., Shoshan, N., Doron, R., Akirav, I., 2018. Cannabinoids prevent depressive-like  
485   symptoms and alterations in BDNF expression in a rat model of PTSD. *Prog.*  
486   *Neuropsychopharmacol. Biol. Psychiatry*, 84, 129–139.
- 487   Carnevali, L., Montano, N., Statello, R., Sgoifo, A., 2017a. Rodent models of depression-  
488   cardiovascular comorbidity: Bridging the known to the new. *Neurosci. Biobehav. Rev.* 76, 144-153.
- 489   Carnevali, L., Nalivaiko, E., Sgoifo, A., 2014. Respiratory patterns reflect different levels of  
490   aggressiveness and emotionality in Wild-type Groningen rats. *Respir. Physiol. Neurobiol.* 204, 28-  
491   35.
- 492   Carnevali, L., Rivara, S., Nalivaiko, E., Thayer, J.F., Vacondio, F., Mor, M., Sgoifo, A., 2017b.  
493   Pharmacological inhibition of FAAH activity in rodents: A promising pharmacological approach for  
494   psychological-cardiac comorbidity? *Neurosci. Biobehav. Rev.* 74, 444-452.

495 Carnevali, L., Vacondio, F., Rossi, S., Callegari, S., Macchi, E., Spadoni, G., Bedini, A., Rivara, S.,  
 496 Mor, M., Sgoifo, A., 2015a. Antidepressant-like activity and cardioprotective effects of fatty acid  
 497 amide hydrolase inhibitor URB694 in socially stressed Wistar Kyoto rats. *Eur.*  
 498 *Neuropsychopharmacol.* 25, 2157-2169.

499 Carnevali, L., Vacondio, F., Rossi, S., Macchi, E., Spadoni, G., Bedini, A., Neumann, I.D., Rivara,  
 500 S., Mor, M., Sgoifo, A., 2015b. Cardioprotective effects of fatty acid amide hydrolase inhibitor  
 501 URB694, in a rodent model of trait anxiety. *Sci. Rep.* 5, 18218.

502 Castren, E., Voikar, V., Rantamaki, T., 2007. Role of neurotrophic factors in depression. *Curr. Opin.*  
 503 *Pharmacol.* 7, 18-21.

504 Clapper, J.R., Vacondio, F., King, A.R., Duranti, A., Tontini, A., Silva, C., Sanchini, S., Tarzia, G.,  
 505 Mor, M., Piomelli, D., 2009. A second generation of carbamate-based fatty acid amide hydrolase  
 506 inhibitors with improved activity in vivo. *ChemMedChem* 4, 1505-1513.

507 Cohen, S., Janicki-Deverts, D., Miller, G.E., 2007. Psychological stress and disease. *Jama* 298,  
 508 1685-1687.

509 Commons, K.G., Cholanians, A.B., Babb, J.A., Ehlinger, D.G., 2017. The Rodent Forced Swim Test  
 510 Measures Stress-Coping Strategy, Not Depression-like Behavior. *ACS Chem. Neurosci.* 8, 955-960.

511 Connolly, K.R., Thase, M.E., 2012. Emerging drugs for major depressive disorder. *Expert. Opin.*  
 512 *Emerg. Drugs* 17, 105-126.

513 de Boer, S.F., Buwalda, B., Koolhaas, J.M., 2017. Untangling the neurobiology of coping styles in  
 514 rodents: Towards neural mechanisms underlying individual differences in disease susceptibility.  
 515 *Neurosci. Biobehav. Rev.* 74(Pt B), 401-422.

516 De Gregorio, D., Manchia, M., Carpinello, B., Valtorta, F., Nobile, M., Gobbi, G., Comai, S., 2019.  
 517 Role of palmitoylethanolamide (PEA) in depression: Translational evidence: Special Section on  
 518 "Translational and Neuroscience Studies in Affective Disorders". Section Editor, Maria Nobile MD,  
 519 PhD. This Section of JAD focuses on the relevance of translational and neuroscience studies in  
 520 providing a better understanding of the neural basis of affective disorders. The main aim is to briefly  
 521 summaries relevant research findings in clinical neuroscience with particular regards to specific  
 522 innovative topics in mood and anxiety disorders. *J. Affect. Disord.* 255, 195-200.

523 De Kloet, E.R., Vreugdenhil, E., Oitzl, M.S., Joels, M., 1998. Brain corticosteroid receptor balance in  
524 health and disease. *Endocr. Rev.* 19, 269-301.

525 Derkinderen, P., Valjent, E., Toutant, M., Corvol, J.C., Enslen, H., Ledent, C., Trzaskos, J., Caboche,  
526 J., Girault, J.A., 2003. Regulation of extracellular signal-regulated kinase by cannabinoids in  
527 hippocampus. *J. Neurosci.* 23, 2371–2382.

528 Dinan, T.G., 2005. Stress: the shared common component in major mental illnesses. *Eur. Psychiatry*  
529 20 Suppl 3, S326-328.

530 Duman, R.S., Monteggia, L.M., 2006. A neurotrophic model for stress-related mood disorders. *Biol.*  
531 *Psychiatry* 59, 1116-1127.

532 Finnell, J.E., Muniz, B.L., Padi, A.R., Lombard, C.M., Moffitt, C.M., Wood, C.S., Wilson, L.B.,  
533 Reagan, L.P., Wilson, M.A., Wood, S.K., 2018. Essential Role of Ovarian Hormones in Susceptibility  
534 to the Consequences of Witnessing Social Defeat in Female Rats. *Biol. Psychiatry* 84, 372-382.

535 Food and Drug Administration, Drug Safety and Availability report, 2016. FDA finds drugs under  
536 investigation in the U.S. related to French BIA 10-2474 drug do not pose similar safety risks.  
537 [https://www.fda.gov/drugs/drug-safety-and-availability/fda-finds-drugs-under-investigation-us-](https://www.fda.gov/drugs/drug-safety-and-availability/fda-finds-drugs-under-investigation-us-related-french-bia-10-2474-drug-do-not-pose-similar-safety)  
538 [related-french-bia-10-2474-drug-do-not-pose-similar-safety.](https://www.fda.gov/drugs/drug-safety-and-availability/fda-finds-drugs-under-investigation-us-related-french-bia-10-2474-drug-do-not-pose-similar-safety)

539 Fowler, C.J., 2015. The potential of inhibitors of endocannabinoid metabolism as anxiolytic and  
540 antidepressive drugs--A practical view. *Eur. Neuropsychopharmacol.* 25, 749-762.

541 Goncalves, M.B., Suetterlin, P., Yip, P., Molina-Holgado, F., Walker, D.J., Oudin, M.J., Zentar, M.P.,  
542 Pollard, S., Yanez-Munoz, R.J., Williams, G., Walsh, F.S., Pangalos, M.N., Doherty, P., 2008. A  
543 diacylglycerol lipase-CB2 cannabinoid pathway regulates adult subventricular zone neurogenesis in  
544 an age-dependent manner. *Mol. Cell. Neurosci.* 38, 526-536.

545 Gorzalka, B.B., Hill, M.N., 2011. Putative role of endocannabinoid signaling in the etiology of  
546 depression and actions of antidepressants. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 35, 1575-  
547 1585.

548 Gorzalka, B.B., Hill, M.N., Hillard, C.J., 2008. Regulation of endocannabinoid signaling by stress:  
549 implications for stress-related affective disorders. *Neurosci. Biobehav. Rev.* 32, 1152-1160.

550 Grigoriadis, S., Robinson, G.E., 2007. Gender issues in depression. *Ann. Clin. Psychiatry* 19, 247-  
551 255.

552 Grippo, A.J., Cushing, B.S., Carter, C.S., 2007. Depression-like behavior and stressor-induced  
553 neuroendocrine activation in female prairie voles exposed to chronic social isolation. *Psychosom.*  
554 *Med.* 69, 149-157.

555 Grippo, A.J., Sullivan, N.R., Damjanoska, K.J., Crane, J.W., Carrasco, G.A., Shi, J., Chen, Z.,  
556 Garcia, F., Muma, N.A., Van de Kar, L.D., 2005. Chronic mild stress induces behavioral and  
557 physiological changes, and may alter serotonin 1A receptor function, in male and cycling female rats.  
558 *Psychopharmacology (Berl)* 179, 769-780.

559 Gururajan, A., Reif, A., Cryan, J.F., Slattery, D.A., 2019. The future of rodent models in depression  
560 research. *Nat. Rev. Neurosci.*, in press.

561 Haller, J., Goldberg, S.R., Pelczar, K.G., Aliczki, M., Panlilio, L.V., 2013. The effects of anandamide  
562 signaling enhanced by the FAAH inhibitor URB597 on coping styles in rats. *Psychopharmacology*  
563 *(Berl)* 230, 353-362.

564 Hashimoto, K., 2010. Brain-derived neurotrophic factor as a biomarker for mood disorders: an  
565 historical overview and future directions. *Psychiatry Clin. Neurosci.* 64, 341-357.

566 Hashimotodani, Y., Ohno-Shosaku, T., Kano, M., 2007. Endocannabinoids and synaptic function in  
567 the CNS. *Neuroscientist* 13, 127-137.

568 Hayley, S., Anisman, H., 2013. Neurotrophic paths in the treatment of depression. *J. Psychiatry*  
569 *Neurosci.* 38, 291-293.

570 Heinrich, L.M., Gullone, E., 2006. The clinical significance of loneliness: a literature review. *Clin.*  
571 *Psychol. Rev.* 26, 695-718.

572 Hill, M.N., Carrier, E.J., McLaughlin, R.J., Morrish, A.C., Meier, S.E., Hillard, C.J., Gorzalka, B.B.,  
573 2008. Regional alterations in the endocannabinoid system in an animal model of depression: effects  
574 of concurrent antidepressant treatment. *J. Neurochem.* 106, 2322-2336.

575 Hill, M.N., Kambo, J.S., Sun, J.C., Gorzalka, B.B., Galea, L.A., 2006. Endocannabinoids modulate  
576 stress-induced suppression of hippocampal cell proliferation and activation of defensive behaviours.  
577 *Eur. J. Neurosci.* 24, 1845-1849.



578 Hill, M.N., McEwen, B.S., 2010. Involvement of the endocannabinoid system in the neurobehavioural  
579 effects of stress and glucocorticoids. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 34, 791-797.

580 Hill, M.N., Patel, S., Carrier, E.J., Rademacher, D.J., Ormerod, B.K., Hillard, C.J., Gorzalka, B.B.,  
581 2005. Downregulation of endocannabinoid signaling in the hippocampus following chronic  
582 unpredictable stress. *Neuropsychopharmacology* 30, 508-515.

583 Hill, M.N., Titterness, A.K., Morrish, A.C., Carrier, E.J., Lee, T.T., Gil-Mohapel, J., Gorzalka, B.B.,  
584 Hillard, C.J., Christie, B.R., 2010. Endogenous cannabinoid signaling is required for voluntary  
585 exercise-induced enhancement of progenitor cell proliferation in the hippocampus. *Hippocampus* 20,  
586 513-523.

587 Huang, G.Z., Woolley, C.S., 2012. Estradiol acutely suppresses inhibition in the hippocampus  
588 through a sex-specific endocannabinoid and mGluR-dependent mechanism. *Neuron* 74, 801-808.

589 Izadi, M.S., Radahmadi, M., Ghasemi, M., Rayatpour, A., 2018. Effects of Isolation and Social  
590 Subchronic Stresses on Food Intake and Levels of Leptin, Ghrelin, and Glucose in Male Rats. *Adv.*  
591 *Biomed. Res.* 7, 118.

592 Jacobson, L., Sapolsky, R., 1991. The role of the hippocampus in feedback regulation of the  
593 hypothalamic-pituitary-adrenocortical axis. *Endocr. Rev.* 12, 118-134.

594 Kokras, N., Antoniou, K., Mikail, H.G., Kafetzopoulos, V., Papadopoulou-Daifoti, Z., Dalla, C., 2015.  
595 Forced swim test: What about females? *Neuropharmacology* 99, 408-421.

596 Kokras, N., Dalla, C., 2014. Sex differences in animal models of psychiatric disorders. *Br. J.*  
597 *Pharmacol.* 171, 4595-4619.

598 Leal, G., Bramham, C.R., Duarte, C.B., 2017. BDNF and Hippocampal Synaptic Plasticity. *Vitam.*  
599 *Horm.* 104, 153-195.

600 Levone, B.R., Cryan, J.F., O'Leary, O.F., 2015. Role of adult hippocampal neurogenesis in stress  
601 resilience. *Neurobiol. Stress* 1, 147-155.

602 Lu, B., Nagappan, G., Lu, Y., 2014. BDNF and synaptic plasticity, cognitive function, and dysfunction.  
603 *Handb. Exp. Pharmacol.* 220, 223-250.

604 Mackie, K., 2005. Distribution of cannabinoid receptors in the central and peripheral nervous system.  
605 *Handb. Exp. Pharmacol.*, 299-325.

606 Mallet, C., Dubray, C., Dualé, C., 2016. FAAH inhibitors in the limelight, but regrettably. *Int. J. Clin.*  
607 *Pharmacol. Ther.* 54, 498-501.

608 Marchalant, Y., Brothers, H.M., Wenk, G.L., 2009. Cannabinoid agonist WIN-55,212-2 partially  
609 restores neurogenesis in the aged rat brain. *Mol. Psychiatry* 14, 1068-1069.

610 Marcondes, F.K., Bianchi, F.J., Tanno, A.P., 2002. Determination of the estrous cycle phases of rats:  
611 some helpful considerations. *Braz. J. Biol.* 62, 609-614.

612 Micale, V., Di Marzo, V., Sulcova, A., Wotjak, C.T., Drago, F., 2013. Endocannabinoid system and  
613 mood disorders: priming a target for new therapies. *Pharmacol. Ther.* 138, 18-37.

614 Petrosino, S., Ligresti, A., Di Marzo, V., 2009. Endocannabinoid chemical biology: a tool for the  
615 development of novel therapies. *Curr. Opin. Chem. Biol.* 13, 309-320.

616 Pittenger, C., Duman, R.S., 2008. Stress, depression, and neuroplasticity: a convergence of  
617 mechanisms. *Neuropsychopharmacology* 33, 88-109.

618 Plenis, A., Konieczna, L., Oledzka, I., Kowalski, P., Baczek, T., 2011. Simultaneous determination  
619 of urinary cortisol, cortisone and corticosterone in parachutists, depressed patients and healthy  
620 controls in view of biomedical and pharmacokinetic studies. *Mol. Biosyst.* 7, 1487-1500.

621 Porsolt, R.D., Le Pichon, M., Jalfre, M., 1977. Depression: a new animal model sensitive to  
622 antidepressant treatments. *Nature* 266, 730-732.

623 Romano, A., Tempesta, B., Provensi, G., Passani, M.B., Gaetani, S., 2015. Central mechanisms  
624 mediating the hypophagic effects of oleoylethanolamide and N-acylphosphatidylethanolamines:  
625 different lipid signals? *Front. Pharmacol.* 6, 137.

626 Rubino, T., Forlani, G., Vigano, D., Zippel, R., Parolaro, D., 2005. Ras/ERK signalling in cannabinoid  
627 tolerance: from behaviour to cellular aspects. *J. Neurochem.* 93, 984–991.

628 Sapolsky, R.M., 2000. The possibility of neurotoxicity in the hippocampus in major depression: a  
629 primer on neuron death. *Biol. Psychiatry* 48, 755-765.

630 Scarante, F.F., Vila-Verde, C., Detoni, V.L., Ferreira-Junior, N.C., Guimaraes, F.S., Campos, A.C.,  
631 2017. Cannabinoid Modulation of the Stressed Hippocampus. *Front. Mol. Neurosci.* 10, 411.

632 Sheline, Y.I., 1996. Hippocampal atrophy in major depression: a result of depression-induced  
633 neurotoxicity? *Mol. Psychiatry* 1, 298-299.

634 Sheline, Y.I., Liston, C., McEwen, B.S., 2019. Parsing the Hippocampus in Depression: Chronic  
635 Stress, Hippocampal Volume, and Major Depressive Disorder. *Biol. Psychiatry* 85, 436-438.

636 Sloan, D.M., Kornstein, S.G., 2003. Gender differences in depression and response to  
637 antidepressant treatment. *Psychiatr. Clin. North. Am.* 26, 581-594.

638 Stepanichev, M., Dygalo, N.N., Grigoryan, G., Shishkina, G.T., Gulyaeva, N., 2014. Rodent models  
639 of depression: neurotrophic and neuroinflammatory biomarkers. *Biomed. Res. Int.* 2014, 932757.

640 Stetler, C., Miller, G.E., 2011. Depression and hypothalamic-pituitary-adrenal activation: a  
641 quantitative summary of four decades of research. *Psychosom. Med.* 73, 114-126.

642 Sun, M., Choi, E.Y., Magee, D.J., Stets, C.W., During, M.J., Lin, E.J., 2014. Metabolic Effects of  
643 Social Isolation in Adult C57BL/6 Mice. *Int. Sch. Res. Notices* 2014, 690950.

644 Tabatadze, N., Huang, G., May, R.M., Jain, A., Woolley, C.S., 2015. Sex Differences in Molecular  
645 Signaling at Inhibitory Synapses in the Hippocampus. *J. Neurosci.* 35, 11252-11265.

646 Tardito, D., Perez, J., Tiraboschi, E., Musazzi, L., Racagni, G., Popoli, M., 2006. Signaling pathways  
647 regulating gene expression, neuroplasticity, and neurotrophic mechanisms in the action of  
648 antidepressants: a critical overview. *Pharmacol Rev.* 58, 115-134.

649 Tarzia, G., Duranti, A., Gatti, G., Piersanti, G., Tontini, A., Rivara, S., Lodola, A., Plazzi, P.V., Mor,  
650 M., Kathuria, S., Piomelli, D., 2006. Synthesis and structure-activity relationships of FAAH inhibitors:  
651 cyclohexylcarbamic acid biphenyl esters with chemical modulation at the proximal phenyl ring.  
652 *ChemMedChem* 1, 130-139.

653 Temporary Specialist Scientific Committee, 2016. FAAH (Fatty Acid Amide Hydrolase), on the  
654 causes of the accident during a Phase 1 clinical trial in Rennes in January 2016.  
655 [https://ansm.sante.fr/var/ansm\\_site/storage/original/application/744c7c6daf96b141bc9509e2f85c2](https://ansm.sante.fr/var/ansm_site/storage/original/application/744c7c6daf96b141bc9509e2f85c227e.pdf)  
656 [27e.pdf](https://ansm.sante.fr/var/ansm_site/storage/original/application/744c7c6daf96b141bc9509e2f85c227e.pdf)

657 van Esbroeck, A.C.M., Janssen, A.P.A., Cognetta, A.B., Ogasawara, D., Shpak, G., van der Kroeg,  
658 M., Kantae, V., Baggelaar, M.P., de Vrij, F.M.S., Deng, H., Allarà, M., Fezza, F., Lin, Z., van der Wel,  
659 T., Soethoudt, M., Mock, E.D., den Dulk, H., Baak, I.L., Florea, B.I., Hendriks, G., De Petrocellis, L.,  
660 Overkleeft, H.S., Hankemeier, T., De Zeeuw, C.I., Di Marzo, V., Maccarrone, M., Cravatt, B.F.,


661 Kushner, S.A., van der Stelt, M., 2017. Activity-based protein profiling reveals off-target proteins of  
662 the FAAH inhibitor BIA 10-2474. *Science* 356, 1084–1087.

663 von Schaper, E., 2016. Bial incident raises FAAH suspicions. *Nat. Biotechnol.* 34, 223.

664 Zer-Aviv, T.M., Akirav, I., 2016. Sex differences in hippocampal response to endocannabinoids after  
665 exposure to severe stress. *Hippocampus* 26, 947-957.

666

667




668

669

670

FIGURE 1

671



672

673

FIGURE 2

674





FIGURE 3

678



679

680

FIGURE 4

681






FIGURE 5




FIGURE 6