

University of Parma Research Repository

Sharp and Fast Bounds for the Celis-Dennis-Tapia Problem

This is the peer reviewd version of the followng article:

Original

Sharp and Fast Bounds for the Celis-Dennis-Tapia Problem / Consolini, Luca; Locatelli, Marco. - In: SIAM JOURNAL ON OPTIMIZATION. - ISSN 1052-6234. - 33:2(2023), pp. 868-898. [10.1137/21M144548X]

Availability: This version is available at: 11381/2971032 since: 2024-02-15T16:57:40Z

Publisher: SIAM PUBLICATIONS

Published DOI:10.1137/21M144548X

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

Author queries:

- Q1: Should this be defined?
- Q2: Should this be defined?
- Q3: Should this be defined?
- Q4: Should "being" and "H" be transposed?
- Q5: Should this be defined?
- Q6: Delete [17] since it is a duplicate of [16]?

2

1 SHARP AND FAST BOUNDS FOR THE CELIS-DENNIS-TAPIA PROBLEM*

LUCA CONSOLINI[†] AND MARCO LOCATELLI[†]

3 **Abstract.** In the Celis-Dennis-Tapia (CDT) problem a quadratic function is minimized over a region defined by two strictly convex quadratic constraints. In this paper we rederive a necessary 4 a region defined by two strictly convex quadratic constraints. In this paper we rederive a necessary 5 and sufficient optimality condition for the exactness of the dual Lagrangian bound (equivalent to the 6 Shor relaxation bound in this case). Starting from such a condition, we propose strengthening the 7 dual Lagrangian bound by adding one or two linear cuts to the Lagrangian relaxation. Such cuts are 8 obtained from supporting hyperplanes of one of the two constraints. Thus, they are redundant for
9 the original problem, but they are not for the Lagrangian relaxation. The computational experiments 9 the original problem, but they are not for the Lagrangian relaxation. The computational experiments 10 show that the new bounds are effective and require limited computing times. In particular, one of the show that the new bounds are effective and require limited computing times. In particular, one of the 11 proposed bounds is able to solve all but one of the 212 hard instances of the CDT problem presented
12 in [S. Burer and K. M. Anstreicher, {\it SIAM J. Optim.}, 23 (2013), pp. 432–451]. in $[S. Burer and K. M. Anstreicher, \{\it{SIAM J. Optim.}\}$, 23 (2013), pp. 432-451.

13 Key words. CDT problem, dual Lagrangian bound, linear cuts

14 MSC codes. 90C20, 90C22, 90C26

DOI. 10.1137/21M144548X

15 **1. Introduction.** The Celis-Dennis-Tapia problem (CDT problem in what fol-16 lows) is defined as follows:

(1.1)
$$
p^* = \min \mathbf{x}^\top \mathbf{Q} \mathbf{x} + \mathbf{q}^\top \mathbf{x} \n\mathbf{x}^\top \mathbf{x} \le 1 \n\mathbf{x}^\top \mathbf{A} \mathbf{x} + \mathbf{a}^\top \mathbf{x} \le a_0,
$$

17 where $\mathbf{Q}, \mathbf{A} \in \mathbb{R}^{n \times n}$, $\mathbf{q}, \mathbf{a} \in \mathbb{R}^n$, $a_0 \in \mathbb{R}$, while **A** is assumed to be positive definite. We 18 will denote by

$$
H = \{ \mathbf{x} \in \mathbb{R}^n : \mathbf{x}^\top \mathbf{A} \mathbf{x} + \mathbf{a}^\top \mathbf{x} \le a_0 \}
$$

19 the ellipsoid defined by the second constraint, by ∂H its border, and by $int(H)$ its interior. The CDT problem was originally proposed in [13] and has attracted a lot of attention in the last two decades. For some special cases a convex reformulation is available. For instance, in [26] it is shown that a semidefinite reformulation is 23 possible when no linear terms are present, i.e., when $q = a = 0$. However, up to now no tractable convex reformulation of general CDT problems has been proposed in the literature. In spite of that, recently three different works [9, 14, 22] independently proved that the CDT problem is solvable in polynomial time. More precisely, in [14, 22] polynomial solvability is proved by identifying all KKT points through the solution 28 of a bivariate polynomial system with polynomials of degree at most $2n$. The two unknowns are the Lagrange multipliers of the two quadratic constraints. Instead, in [9] an approach based on the solution of a sequence of feasibility problems for systems of quadratic inequalities is proposed. The systems are solved by a polynomial-time

^{*} Received by the editors September 9, 2021; accepted for publication (in revised form) November 29, 2022; published electronically DATE.

<https://doi.org/10.1137/21M144548X>

[†] Dipartimento di Ingegneria e Architettura, Università di Parma, Parma, 43121, Italy [\(luca.consolini@unipr.it,](mailto:luca.consolini@unipr.it) [marco.locatelli@unipr.it\)](mailto:marco.locatelli@unipr.it).

 algorithm based on Barvinok's construction [6]. Though polynomial, all of these approaches are computationally demanding since the degree of the polynomial is quite large. Conditions guaranteeing that the classical Shor SDP relaxation or, equivalently in this case, the dual Lagrangian bound is exact, are discussed in [2, 7]. In particular, in [2] a necessary and sufficient condition is presented. It is shown that the lack of exactness is related to the existence of KKT points with the same Lagrange multipliers but two distinct primal solutions, both active at one of the two constraints but one violating and the other one fulfilling the other constraint. In [10] necessary and sufficient conditions for local and global optimality are discussed based on copositivity. In [11] an exactness condition is given for a copositive relaxation, also for the case with additional linear constraints. A trajectory following method to solve the CDT problem has been discussed in [26], while different branch-and-bound solvers are tested in [19].

 Recently, different papers proposed valid bounds for the CDT problem. In [12] the Shor relaxation bound is strengthened by adding all RLT constraints obtained by $\mathbf{AQ1}$
47 supporting hyperplanes of the two ellipsoids. By fixing the supporting hyperplane for supporting hyperplanes of the two ellipsoids. By fixing the supporting hyperplane for one ellipsoid, the RLT constraints obtained with all the supporting hyperplanes of the 49 other can be condensed into a single $SOC-RLT$ constraint. Varying the supporting hy- $AQ2$
50 perplane of the first ellipsoid gives rise to an infinite number of $SOC-RLT$ constraints perplane of the first ellipsoid gives rise to an infinite number of SOC-RLT constraints which, however, can be separated in polynomial time. The addition of these con- straints does not allow one to close the duality gap, but it is computationally shown that many instances which are not solved via the SDP bound, are solved with the addition of these SOC-RLT cuts. The authors generate 1000 random test instances 55 for each $n = 5, 10, 20$, following a procedure described in [18] to generate trust-region problems with one local and nonglobal minimizer. The proposed bound based on 57 SOC-RLT cuts allows for solving most instances except for 212 (38 for $n = 5$, 70 for $n = 70$, and 104 for $n = 20$. Such unsolved instances are considered as hard ones in subsequent works. In [25] lifted-RLT cuts are introduced and it is shown that the 60 new constraints allow one to derive an exact bound for $n = 2$ but also to improve the 61 bounds of [12] over the hard instances for $n > 2$. In [27] it is proved that the duality gap can be reduced to 0 by solving two subproblems with SOC constraints when the second constraint is the product of two linear functions and an exactness result is also provided for the case of problems with two variables. Due to its relations with the approach proposed in this work, we will further discuss the approach proposed in [27] 66 at the end of section 3. In [3] cuts are introduced. These are Kronecker product con- \angle AQ3 straints which generalize both the classical RLT constraints obtained from two linear straints which generalize both the classical RLT constraints obtained from two linear inequality constraints, and the SOC-RLT constraints obtained from one linear inequal- ity constraint and a SOC constraint. Further hard instances from [12] are solved with the addition of these cuts. In the very recent paper [4] a branch and bound approach is proposed. The main feature of this approach is eigenvector branching, i.e., a branch- ing rule based on the use of the eigenvector corresponding to the maximal eigenvalue 73 of $X^* - x^* x^{*T}$, where (X^*, x^*) is the optimal solution of an SDP relaxation.

 In this paper we investigate ways to strengthen the dual Lagrangian bound through the addition of one or two linear cuts. In particular, the paper is structured as follows. In section 2 we derive some theoretical results for a class of problems with two constraints which includes the CDT problem as a special case. We develop a bi- section technique to solve the dual Lagrangian relaxation for such class of problems. In the following sections we apply the results of section 2 to the CDT problem. In particular, in section 3 we introduce some results through which it will be possible to rederive the necessary and sufficient exactness condition discussed in [2] and we

 discuss how to improve the dual bound for the CDT problem by the addition of a linear cut. Next, in sections 4 and 5 we discuss techniques to further improve the bound. More precisely, in section 4 we still present a bound based on the addition of a linear cut, but we develop a technique to locally adjust a given linear cut, while in section 5 we consider a bound based on the addition of two linear cuts. Finally, in section 6 we present some computational experiments which show that the newly pro- posed bounds, in particular those based on two linear cuts, are both computationally cheap and effective. In particular, one of the bounds will be able to solve all but one of the hard instances from [12]. We also investigate which are the most challenging instances for the proposed bounds and, as we will see, the difficulties are related to the existence of multiple solutions of Lagrangian relaxations.

 It is also worthwhile to highlight the contribution of this paper under another perspective. As previously discussed, while there is no known convex relaxation of the CDT problem, there are several problems, related to CDT, which do have exact SDP relaxations. These include the trust-region subproblem (TRS), the TRS with a single linear constraint (TRS1, see [12, 23]), and the TRS with two linear constraints, at least one of which is tight (TRS2eq, see [26]). This paper shows that such special cases, for which an exact convex relaxation exists, can be used to help solve the general CDT problem. Indeed, we first observe that the subproblems to be solved in section 3 to improve the dual Lagrangian bound, obtained by adding a linear cut corresponding to a supporting hyperplane for the second ellipsoidal constraint, turn out to be TRS1 problems. In section 4 we also discuss how to pick a ""good"" supporting hyperplane, i.e., one which leads to a good SDP relaxation and, in fact, we also provide a necessary and sufficient condition under which we can guarantee that the supporting hyperplane is the best one. Next, in section 5 we observe that the bound can be further improved by adding two linear cuts, one of which must be active, so that the subproblems to be solved in this case are TRS2eq problems.

 2. Lower bounds obtained from the Lagrangian relaxation. The CDT 110 problem (1.1) is a specific instance of the following, more general, one:

(2.1)
$$
p^* = \min_{\mathbf{x} \in \mathbb{R}^n}, \quad f(\mathbf{x}),
$$

$$
g(\mathbf{x}) \le 0,
$$

$$
h(\mathbf{x}) \le 0.
$$

 In this section, we discuss a class of lower bounds on the solution of problem (2.1) that can be obtained from its Lagrangian relaxation. In the next sections, we will apply these bounds to the specific case of the CDT problem (1.1). Throughout this and the following sections, we make the following assumptions.

- $11\overline{6}$ Assumption 2.1. In problem (2.1)
- 117 (a) q, h are continuous;
- 118 (b) the set $\{ x \in \mathbb{R}^n : g(x) \leq 0\}$ is bounded;
- (c) it holds that

(2.2)
$$
h_0 = \min_{\mathbf{x} \; : g(\mathbf{x}) \le 0} h(\mathbf{x}) < 0;
$$

(d) the solution set of problem (2.1) without the last constraint, that is

$$
\bar{P} = \arg \min_{\mathbf{x} \in \mathbb{R}^n}, \quad f(\mathbf{x}),
$$

$$
g(\mathbf{x}) \le 0,
$$

121 is such that $(\forall x \in \overline{P}) h(x) > 0$.

 Note that if the last condition in Assumption 2.1 is violated, we can find the solution of problem (2.1) by removing the last constraint and the relaxation discussed 124 in this section is useless. Now, let $G = \{ x \in \mathbb{R}^n : g(x) \leq 0 \}$ and $H = \{ x \in \mathbb{R}^n : g(x) \leq 0 \}$ $h(\mathbf{x}) \leq 0$. Let $X \supset H$ be a closed subset of $\Bbb R^n$ and for $\lambda \in \Bbb R$, with $\lambda \geq 0$, define the Lagrangian relaxation

(2.3)
$$
p_X(\lambda) = \min_{\mathbf{x} \in X \cap G} f(\mathbf{x}) + \lambda h(\mathbf{x}),
$$

127 and the corresponding solution set

$$
P_X(\lambda) = \arg \min_{\mathbf{x} \in X \cap G} f(\mathbf{x}) + \lambda h(\mathbf{x}).
$$

128 Note that $P_X(\lambda)$ is compact, since $G \cap X$ is nonempty (in view of part (c)) 129 of Assumption 2.1) and compact (in view of the compactness of G which follows 130 from parts (a) and (b) of Assumption 2.1), and $f + \lambda h$ is continuous. Due to well-131 known properties of the Lagrangian relaxation, we have that function p_X is such that 132 $(\forall \lambda \geq 0) p_X(\lambda) \leq p^*$, and is concave (it is the pointwise minimum of a set of functions 133 linear in λ). The best bound that can be obtained as the solution of (2.3) is given by

(2.4)
$$
\bar{p}_X = \max_{\lambda \geq 0} p_X(\lambda),
$$

134 and corresponds to the solution of the dual Lagrangian problem. Note that function 135 p_X depends on the choice of set X.

136 Now, we recall that the supergradient of a function $q : \mathbb{R} \rightarrow \mathbb{R}$ at $x \in \mathbb{R}$ is defined 137 as

$$
\partial q(x) = \{ z \in \mathbb{R} : (\forall y \in \mathbb{R}) q(y) - q(x) \le z(y - x) \}.
$$

138 Since p_X is concave, for any $\lambda \in \mathbb{R}$, the supergradient $\partial p_X(\lambda)$ is nonempty. 139 For $A \subset \mathbb{R}^n$ define the following subset of \mathbb{R} :

$$
h(A) = \{h(\mathbf{x}) \ : \ \mathbf{x} \in A\}
$$

140 For $X \subset \mathbb{R}^n$, define a (set-valued) function $Q_X : \mathbb{R}_+ \rightarrow \mathcal{P} (\Bbb R)$,

$$
(2.5) \tQ_X(\lambda) = h(P_X(\lambda))
$$

141 (\mathbb{R}_+ denotes the set of nonnegative reals and $\mathcal{P}(\mathbb{R})$ is the power set of the set of real 142 numbers). Also set $h_X^{\min}(\lambda) = \min Q_X(\lambda)$ and $h_X^{\max}(\lambda) = \max Q_X(\lambda)$. The following 143 proposition shows that function Q_X is monotone nonincreasing (see Definition 3.5.1) 144 of [5]) and upper semicontinuous (see Definition 1.4.1 of [5]). These two properties 145 will play an important role in the computation of a lower bound for problem (2.1). 146 Moreover, this proposition characterizes the supergradient of p_X at each $\lambda \geq 0$. In the 147 proof of the proposition we will make use of Berge's maximum theorem (see [8]). In 148 particular, we will consider the slightly different formulation presented as the corollary 149 to Theorem 3 on page 30 of [15].

150 COROLLARY 2.1. Let the correspondence (i.e., the set-valued function) β of S 151 into T be compact-valued and continuous, and let $\phi : S \times T \rightarrow \Bbb R$ be a continuous 152 function. Then, we have the following:

153 (a) The function $z \mapsto m(z) := max\{ \phi (z, y)| y \in \beta (z)\}$ is continuous.

154 (b) The correspondence $z \mapsto \{ y \in \beta (z)| \phi (z, y) = m(z)\}$ is nonempty and compact-155 valued and upper semicontinuous.

- 156 PROPOSITION 2.2. For any $X \subset \mathbb{R}^n$,
- 157 (i) Q_X is monotone non-increasing, that is if $\lambda_1 \geq \lambda_2, y_1 \in Q_X(\lambda_1), y_2 \in Q_X(\lambda_2),$ 158 then $y_1 \leq y_2$.
- 159 (ii) Q_X is upper semicontinuous, that is, if $Q_X(\lambda) \subset U$, where U is an open subset 160 of $\Bbb R$, then there exists a neighborhood V of λ such that $(\forall z \in V) Q_X(z) \subset U$.
- 161 (iii) $\partial p_X(\lambda) = [\min Q_X(\lambda) , \max Q_X(\lambda)].$

162 Proof. (i) Let $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{R}^n$ be such that $y_1 = h(\mathbf{x}_1)$ and $y_2 = h(\mathbf{x}_2)$, then $f(\mathbf{x}_1)$ + 163 $\lambda_1 h(\mathbf{x}_1) \leq f(\mathbf{x}_2) +\lambda_1 h(\mathbf{x}_2)$ and $f(\mathbf{x}_2) +\lambda_2 h(\mathbf{x}_2) \leq f(\mathbf{x}_1) +\lambda_2 h(\mathbf{x}_1)$. By adding up the 164 previous inequalities, it follows that $(\lambda_1 - \lambda_2)(h(\mathbf{x}_1) - h(\mathbf{x}_2)) \leq 0$.

- 165 (ii) Apply Corollary 2.1 with $T = G \cap X$, $S = \mathbb{R}_+$, constant function $(\forall \lambda \in \mathbb{R})$ 166 S) $\beta (\lambda) = G\cap X, \phi (\lambda ,x) = - f(x) - \lambda \cdot h(x)$. Since ϕ is continuous, set-valued function 167 $P_X(\lambda) = \{ \mathbf{x} \in G \cap X : \phi(\lambda , \mathbf{x}) = \max_{\mathbf{y} \in G \cap X} \phi (\lambda , \mathbf{y})\}$ is upper semicontinuous. Hence, 168 also Q_X is upper semicontinuous, since it is obtained as the composition of P_X with 169 h, which is continuous (see Theorem 1' on page 113 of $[8]$).
- 170 (iii) It is a consequence of Theorem 4.4.2 in [16], being G compact.
- 171 The next proposition characterizes the optimal solution of the dual Lagrangian prob-172 lem (2.4).
- 173 Proposition 2.3. Under Assumption 2.1, the optimal value of the dual La-174 grangian problem (2.4) is: (i) either attained at $\lambda_X = 0$ in case $\partial p_X(0) \cap \mathbb{R}_- \neq \emptyset$, 175 where \mathbb{R}_- denotes the set of nonpositive real numbers; (ii) or is attained at $\lambda_X > 0$ 176 such that $0 \in \partial p_X(\lambda_X)$. In the former case, $\bar{p}_X = p^*$ holds.

177 Proof. The proposition, apart from the last statement, is a direct consequence of 178 the optimality conditions for the maximum of concave functions (see Theorem 1.1.1 179 in Chapter 7 of [16]). To prove the last statement, namely that if (i) holds $\bar{p}_X = p^*$, 180 note that, if $\partial p_X(0) \cap \mathbb{R}_- \neq \emptyset$, then, in view of part (iii) of Proposition 2.2, $h_X^{\min}(0) \leq 0$ 181 and, thus, there exists an optimal solution \mathbf{x}^* of $\min_{\mathbf{x} \in X \cap G} f(\mathbf{x})$ such that $h(\mathbf{x}^*) \leq 0$. 182 This implies that \mathbf{x}^* is also an optimal solution of the original problem (2.1), so that 183 $\bar{p}_X = p^*$ holds. Л

184 The following property shows that it is always possible to find a sufficiently high 185 value of λ such that $P_X(\lambda) \subset H$, that is, the elements of $P_X(\lambda)$ are feasible solutions 186 of problem (2.1).

187 Lemma 2.4. If

(2.6)
$$
\lambda \geq \hat{\lambda} = \frac{\max_{\mathbf{x} \in G \cap X} f(\mathbf{x}) - \min_{\mathbf{x} \in G \cap X} f(\mathbf{x})}{|h_0|},
$$

188 where h_0 is defined in (2.2), then $P_X(\lambda) \subset H$.

189 Proof. By contradiction, assume that there exists $\mathbf{x} \in P_X(\lambda)$ such that $h(\mathbf{x}) > 0$, 190 and let $\mathbf{x}_0 \in G \cap H$ be such that $h(\mathbf{x}_0) = h_0 < 0$; then $f(\mathbf{x}) + \lambda h(\mathbf{x}) \leq f(\mathbf{x}_0) + \lambda h(\mathbf{x}_0)$. Since $h(\mathbf{x}) > 0$, it follows that $\lambda \leq \frac{f(\mathbf{x}_0) - f(\mathbf{x})}{|h(\mathbf{x}_0)| + h(\mathbf{x})} < \frac{\max_{\mathbf{x} \in G \cap X} f(\mathbf{x}) - \min_{\mathbf{x} \in G \cap X} f(\mathbf{x})}{|h(\mathbf{x}_0)|}$ 191 Since $h(\mathbf{x}) > 0$, it follows that $\lambda \leq \frac{f(\mathbf{x}_0) - f(\mathbf{x})}{|h(\mathbf{x}_0)| + h(\mathbf{x})} < \frac{\max_{\mathbf{x} \in G \cap X} f(\mathbf{x}) - \min_{\mathbf{x} \in G \cap X} f(\mathbf{x})}{|h(\mathbf{x}_0)|}$, which 192 contradicts the assumption on λ . П

- 193 The following proposition shows that if $0 \in Q_X(\lambda)$, then $p_X(\lambda)$ is equal to the 194 optimal value of problem (2.1).
- 195 Proposition 2.5. Under Assumption 2.1, the following statements are equiva-196 lent for $\lambda > 0$:
- 197 (i) $0 \in Q_X(\lambda)$,
- 198 $\text{(ii)} \ p^* = p_X(\lambda) \ \text{and there exists } \bar{\mathbf{x}} \in \arg \min_{\mathbf{x} \in G \cap H} f(\mathbf{x}) \ \text{such that } h(\bar{\mathbf{x}}) = 0.$

 \Box

199 Proof. (i) \Rightarrow (ii). Let $\bar{\mathbf{x}}$ be such that $h(\bar{\mathbf{x}}) \in Q_X(\lambda)$ and $h(\bar{\mathbf{x}}) = 0$. Let \mathbf{x}^* be a 200 solution of (2.1). Then, $p_X(\lambda) = f(\bar{\mathbf{x}}) + \lambda h(\bar{\mathbf{x}}) = f(\bar{\mathbf{x}}) \leq f(\mathbf{x}^*) + \lambda h(\mathbf{x}^*) \leq f(\mathbf{x}^*)$, hence 201 $p_X(\lambda) \leq p^*$. Moreover, $p_X(\lambda) = f(\bar{\mathbf{x}}) \geq \min_{\mathbf{x} \in G \cap H} f(\mathbf{x}) = p^*$.

202 (ii) \Rightarrow (i). Assume that $p_X(\lambda) = p^*$, and let $\mathbf{x} \in P_X(\lambda)$. Then, by (ii), $f(\mathbf{x})$ + 203 $\lambda h(\mathbf{x}) = f(\bar{\mathbf{x}}) = f(\bar{\mathbf{x}}) + \lambda h(\bar{\mathbf{x}})$. It follows that $\bar{\mathbf{x}} \in P_X(\lambda)$ and $Q_X(\lambda) \ni h(\bar{\mathbf{x}}) = 0$. \Box

204 Remark 2.6. If $0 \in Q_X(\lambda)$, by point (iii) of Proposition 2.2, $\partial p_X(\lambda) \ni 0$, so that 205 λ corresponds to a maximizer of the dual Lagrangian. Note that equation $0 \in Q_X(\lambda)$ 206 always admits a solution if Q_X is continuous. However, in the general case, Q_X is 207 only upper semicontinuous. In this case, the value of λ for which $\partial p_X(\lambda) \ni 0$ may not 208 satisfy $0 \in Q_X(\lambda)$. Thus, the optimal value of the dual Lagrangian (2.4) is not equal 209 to the optimal value of (2.1) but it represents a lower bound of it.

- 210 In order to evaluate a numerical solution algorithm, we define the following weak 211 solution of (2.1).
- 212 DEFINITION 2.7. x is an η -solution of (2.1) if $\mathbf{x} \in G \cap H$ and $f(\mathbf{x}) p^* \leq \eta$.
- 213 The following proposition presents a bound on the error committed on the esti-214 mation of p^* .

215 PROPOSITION 2.8. For any $\lambda \geq 0$ such that $P_X(\lambda) \cap H \neq \emptyset$, and for any $\mathbf{x} \in \mathbb{R}$ 216 $P_X(\lambda) \cap H$, it holds that $f(x) - p^* \leq \lambda | h(x)|$, i.e., x is an η -solution of problem of 217 (2.1) with $\eta = \lambda | h(\mathbf{x}) |$.

218 Proof. Since $\mathbf{x} \in P_X(\lambda)$ and observing that $\mathbf{x}^* \in G \cap X$ for any $X \supset H$, $f(\mathbf{x})$ + 219 $\lambda h(\mathbf{x}) \leq f(\mathbf{x}^*) + \lambda h(\mathbf{x}^*) \leq f(\mathbf{x}^*)$, from which $f(\mathbf{x}) - f(\mathbf{x}^*) \leq \lambda | h(\mathbf{x})|$. \Box

220 Now we introduce Algorithm 2.1 which is based on a binary search through differ-221 ent λ values and is able to return the solution of the dual Lagrangian problem, i.e., the 222 maximum of function $p_X(\lambda)$ and, in some cases, even the solution of problem (2.1). 223 The algorithm also returns a point $\mathbf{z}_1(\lambda^{\max}) \in H$ and (possibly) a point $\mathbf{z}_2(\lambda^{\min}) \notin H$. 224 Note that according to Proposition 2.8, point $z_1(\lambda^{\max})$ is an η -solution of problem 225 (2.1) with $\eta = \lambda | h(\mathbf{z}_1(\lambda^{\text{max}}))|$.

226 The algorithm starts with an initial interval of λ values $[\lambda^{\min}, \lambda^{\max}] = [0, \lambda^{\text{init}}],$ 227 where λ ^{init} is a suitably large value and can be set equal to $\hat{\lambda}$ as defined in Lemma 228 2.4. At each iteration the algorithm halves such interval by evaluating the set Q_X^{λ} 229 at $\lambda = (\lambda^{\max} + \lambda^{\min})/2$. Then, the algorithm sets- are $\lambda^{\min} = \lambda$, if $h_X^{\min}(\lambda) > 0$; 230 $\lambda^{\max} = \lambda$ if $h_X^{\max}(\lambda) < 0$. Instead, if $0 \in \partial p_X(\lambda) = [h_X^{\min}(\lambda), h_X^{\max}(\lambda)]$, the algorithm 231 sets $\lambda^{\max} = \lambda^{\min} = \lambda$ and exits the loop.

232 The following proposition characterizes Algorithm 2.1.

233 Proposition 2.9. (i) Algorithm 2.1 terminates in a finite number of iterations: 234 (ii) at each iteration $\lambda^{\min} \leq \lambda_X \leq \lambda^{\max}$,

235 (iii) at termination $|\lambda^{\max} - \lambda_X| \leq \epsilon$,

236 (iv) at each iteration, if $\lambda^{\min} < \lambda_X < \lambda^{\max}$, then $[h_X^{\max}(\lambda^{\max}), h_X^{\min}(\lambda^{\min})] \supset$ 237 $\qquad \partial p_X(\lambda_X),$

238 (v) point $\mathbf{z}_1(\lambda^{\max}) \in P_X(\lambda^{\max}) \cap H$ is an η -solution of (2.1) with $\eta = \lambda_{\max}$ 239 $h(\mathbf{z}_1(\lambda^{\max}))$.

240 Proof. (i) At each iteration the length of the interval $[\lambda^{\min}, \lambda^{\max}]$ is halved. Hence, 241 in a sufficient large number of iterations, the termination condition of the main loop 242 will be satisfied.

243 (ii) At the beginning of the algorithm we have that $\lambda^{\min} \leq \lambda_X \leq \lambda^{\max}$. Every time 244 λ^{\min} is updated, we set $\lambda^{\min} = \lambda$ if condition $h_X^{\min}(\lambda) > 0$ holds. Since $h_X^{\min}(\lambda_X) \leq 0$,

Algorithm 2.1 Binary search algorithm for the solution of the dual Lagrangian problem for (1.1).

 $\overline{\text{DualLagrangian}(X, \, \lambda_{\text{init}})}$ Set $\lambda_{\text{min}} = 0, \ \lambda^{\text{max}} = \lambda^{\text{init}}$ while $\lambda^{\max} - \lambda^{\min} > \varepsilon$ do Set $\lambda = (\lambda^{\max} + \lambda^{\min})/2$ Solve problem (2.3), and let $P_X(\lambda)$ be its set of optimal solutions Compute the set $Q_X(\lambda)$ and the values $h^{\min}_X(\lambda), h^{\max}_X(\lambda)$ if $h_X^{\min}(\lambda) > 0$ then Set $\lambda_{\min} = \lambda$ else if $h_X^{\max}(\lambda) < 0$ then Set $\lambda^{\max} = \lambda$ else Set $\lambda^{\max} = \lambda^{\min} = \lambda$ end if end while Set $Lb = p_X(\lambda^{\max})$, and let $\mathbf{z}_1(\lambda^{\max})$ be some point in $P_X(\lambda^{\max}) \cap H$ and $\mathbf{z}_2(\lambda^{\min})$ be some point (if any) in $P_X(\lambda^{\min}) \setminus H$ $\textbf{return }Lb, \lambda^{\max}, \mathbf{z}_1(\lambda^{\max}), \mathbf{z}_2(\lambda^{\min})$

245 by the monotonicity of function h_X^{\min} , which is a consequence of the monotonicity of 246 function Q_X , condition $\lambda^{\min} \leq \lambda_X$ is maintained. The same reasoning can be used to 247 prove that $\lambda^{\max} \geq \lambda_X$.

248 (iii) It is a consequence of (ii) and the termination condition.

249 $\text{(iv) } \partial p_X(\lambda_X) = [h_X^{\min}(\lambda_X), h_X^{\max}(\lambda_X)] \subset [h_X^{\max}(\lambda^{\max}), h_X^{\min}(\lambda^{\min})],$ due to point 250 (ii) and the monotonicity of functions h_X^{max} and h_X^{min} , which is a consequence of the 251 monotonicity of function Q_X .

252 (v) It is a consequence of Proposition 2.8.

 \Box

 $\frac{253}{254}$ The following property is a direct consequence of the upper semicontinuity of Q_X .

255 PROPOSITION 2.10. Let $X \supset H$ be such that $\sup Q_X(\lambda) < 0$; then there exists a 256 neighborhood U of λ such that $(\forall \eta \in U)$ max $Q_X(\eta) < 0$.

257 As a consequence of the previous proposition, it is possible to improve the lower 258 bound on problem (2.1) , obtained as the solution of (2.3) , by replacing set X with a 259 different set $Y \supset H$ fulfilling a given condition.

260 PROPOSITION 2.11. Let $Y \supset H$ be such that $\max Q_Y (\lambda_X) \leq 0$ or, equivalently, 261 $P_Y(\lambda_X) \setminus H = \emptyset$, and assume that $\bar{p}_X = p_X(\lambda_X) < p^*$. Then $\bar{p}_Y = p_Y (\lambda_Y) > \bar{p}_X$.

262 Proof. Note that, by Proposition 2.3, $\bar{p}_X = p_X(\lambda_X) < p^*$ implies $\lambda_X > 0$. Now, 263 in case max $Q_Y (\lambda_X) = 0$, then $0 \in Q_Y (\lambda_X)$ and, by Proposition 2.5, $\bar{p}_Y = p^* > \bar{p}_X$. 264 Thus, we only consider the case max $Q_Y(\lambda_X) < 0$. In such case, by Proposition 2.10, 265 $\lambda_Y < \lambda_X$. If $\lambda_Y = 0$, by Proposition 2.3 we have that $\bar{p}_Y = p^* > \bar{p}_X$ and we are done. 266 Otherwise, if $\lambda_Y > 0$, again by Proposition 2.3 we have that $0 \in [h_Y^{\min}(\lambda_Y), h_Y^{\max}(\lambda_Y)],$ 267 and, consequently, there exists $y \in P_Y (\lambda Y)$ such that $h(y) \leq 0$. Note that $\bar{p}_Y =$ 268 $f(\mathbf{y}) + \lambda_Y h(\mathbf{y})$. If $h(\mathbf{y}) = 0$, then, by Proposition 2.5, $p_Y(\lambda_Y) = p^*$, so that the thesis 269 is satisfied in view of $\bar{p}_X < p^*$. Otherwise, if $h(\mathbf{y}) < 0$, let $\mathbf{x} \in \mathbb{R}^n$ be such that $\bar{p}_X =$ 270 $f(\mathbf{x}) + \lambda_X h(\mathbf{x})$. Then $\bar{p}_X = f(\mathbf{x}) + \lambda_X h(\mathbf{x}) \leq f(\mathbf{y}) + \lambda_X h(\mathbf{y}) < f(\mathbf{y}) + \lambda_Y h(\mathbf{y}) = \bar{p}_Y$, 271 where we used the facts that $h(\mathbf{y}) < 0$ and that $\lambda_Y < \lambda_X$. \Box 272 The following proposition deals with the special case of the previous result when 273 $Y \subset X$.

274 PROPOSITION 2.12. Let $X \supset Y \supset H$ be such that $Y \cap (P_X(\lambda_X) \setminus H) = \emptyset$, and 275 assume that $\bar{p}_X = p_X(\lambda_X) < p^*$. Then $\bar{p}_Y = p_Y (\lambda_Y) > \bar{p}_X$.

276 Proof. Since $h_X^{\min}(\lambda_X) < 0$ we have that $P_X(\lambda_X) \cap H \neq \emptyset$ and, consequently, since 277 $Y \supset H$, also $Y \cap P_X(\lambda_X) \neq \emptyset$. Then, $Y \subset X$ implies $P_Y (\lambda_X) = Y \cap P_X(\lambda_X)$. Moreover, 278 if $Y \cap (P_X(\lambda_X)\setminus H) = \emptyset$, then the condition max $Q_Y (\lambda_X) \leq 0$ is satisfied and the result 279 follows from Proposition 2.11. Л

280 Stated in another way, the previous propositions show that, in case the lower 281 bound \bar{p}_X is not exact, we are able to improve (increase) it if we are able to replace 282 set X with a new set Y which cuts away all members of $P_X(\lambda_X)$ outside H.

283 Remark 2.13. Up to now we have not discussed the difficulty of computing the 284 values of function p_X or, equivalently, the difficulty of solving problem (2.3). Such 285 difficulty is strictly related to the specific problem (i.e., to the specific functions f, g, h), 286 and also to the specific set X . In the next sections we apply the general theory 287 developed in this section to the CDT problem. We show that for suitably defined sets 288 X (defined by one or two linear cuts), the computation of function p_X can be done 289 efficiently, and, moreover, the corresponding lower bounds \bar{p}_X improve the standard 290 dual Lagrangian bound, corresponding to the case $X = \mathbb{R}^n$.

291 Remark 2.14. In principle one could also define a cutting algorithm where a 292 sequence of sets $\{ X_k\}$ is generated such that (i) $X_k \supset X_{k+1} \supset H$ for all k; (ii) 293 $X_{k+1} \cap (P_{X_k}(\lambda_{X_k}) \setminus H) = \emptyset;$ (iii) $\bigcap_{k=1}^{\infty} X_k = H$. The corresponding sequence of lower 294 bounds $\{\bar{p}_{X_k}\}\$ is strictly increasing in view of Proposition 2.12, and converges to p^* . 295 However, the difficulty related to such an algorithm is that forcing (ii) may not be 296 trivial and, moreover, as already commented in Remark 2.13, computing p_{X_k} may be 297 computationally demanding.

 The following algorithm, Algorithm 2.2, in principle, is able to always find an approximate solution of (2.1). The algorithm is based on an iterative reduction of set X, in order to eliminate its elements in which function h is positive. In practice, Algorithm 2.2 could be unimplementable. Indeed, it may require a large number of cuts on set X and each added cut may increase the complexity of the optimization problem that we need to solve to evaluate DualLagrangian. In section 4, we will see that, to refine the lower bound on the solution of the CDT problem, it is computa- tionally more convenient to adjust existing cuts instead of adding new ones. We stress that we will not actually use Algorithm 2.2 for the solution of the CDT problem. We present this algorithm just as a theoretical contribution.

308 PROPOSITION 2.15. Algorithm 2.2 terminates and $\bar{\mathbf{x}}$ is such that $h(\bar{\mathbf{x}}) \leq \frac{\eta}{\lambda^{\max}}$ 309 $\qquad \text{and } |\bar{f} - f^*| \leq \eta.$

310 Proof. By contradiction, assume that the algorithm does not terminate. Let l_i 311 be the value of λ^{\min} returned by the *i*th call to **DualLagrangian**. Sequence l_i is 312 monotone nonincreasing; moreover, the domain of the sequence is a subset of finite 313 cardinality of interval $[0, \lambda^{\text{init}}]$ (its maximum cardinality depends on λ^{init} and ϵ). 314 Indeed, the termination condition of function DualLagrangian allows only for a 315 finite number of divisions of the interval $[0, \lambda^{\text{init}}]$. Hence, sequence l_i converges in 316 a finite number of iterations to its limit $l_\infty = \lim_{i\to \infty} l_i$ and there exists $\bar{i} \in \mathbb{N}$ such 317 that $(\forall i \geq \overline{i})$ $l_i = l_\infty$. By (iv) of Proposition 2.9, $h^{\max}(l_\infty) \geq 0$ and, since the

Algorithm 2.2 Bound improvement through redefinition of set X .

1: Set $X = \mathbb{R}^n$ 2: Set $\lambda^{\max} = \lambda^{\text{init}}$

- 3: repeat
-
- 4: Let
- $[Lb, \lambda^{\min}, \lambda^{\max}, \mathbf{z}_1(\lambda^{\max}), \mathbf{z}_2(\lambda^{\min}), h_X^{\min}, h_X^{\max}] = \textbf{DualLagrangian}(X, \lambda^{\text{init}})$ 5: Set $Z = \{ \mathbf{x} \in P_X(\lambda^{\min}) : h(\mathbf{x}) > 0\}$
- 6: Redefine $X = Y$, where Y is such that $X \supset Y \supset H$ and $Z \cap Y = \emptyset$.
- 7: **until** min $\{ h_X^{\max}(\lambda^{\min}), -h_X^{\min}(\lambda^{\max}) \} \lambda^{\max} \leq \eta$
- 8: return $\bar{\mathbf{x}} \in P_X(\lambda_X^{min}) \cup P_X(\lambda_X^{max})$ with $|h(\bar{\mathbf{x}})| \leq \eta$, $\bar{f} = f(\bar{\mathbf{x}})$.

318 algorithm does not terminate, $h^{\max}(l_{\infty}) \geq \eta$. At the $\bar{i} + 1$ -iteration, the algorithm 319 calls **DualLagrangian** (X, l_∞) , which returns the value $\lambda^{\min} = l_\infty$. Anyway, at the 320 previous iteration \overline{i} , the elements $P_X(\lambda^{\min})$ at which function h is positive had already 321 been removed from X. This implies that **DualLagrangian** (X, l_∞) cannot return the 322 strictly positive value $\lambda^{\min} = l_{\infty}$, leading to a contradiction. Hence, the algorithm 323 terminates and the stated bounds hold because of the termination condition and by 324 Proposition 2.8. \Box

 3. Dual Lagrangian bound and a possible improvement. In this section, we apply the general properties presented in section 2 to the CDT problem (1.1). In 327 fact, the CDT problem is a specific instance of (2.1) in which $f(\mathbf{x}) = \mathbf{x}^\top \mathbf{Q} \mathbf{x} + \mathbf{q}^\top \mathbf{x}$, $g(\mathbf{x}) = \mathbf{x}^\top \mathbf{x} - 1$, $h(\mathbf{x}) = \mathbf{x}^\top \mathbf{A} \mathbf{x} + \mathbf{a}^\top \mathbf{x} - a_0$.

329 Note that the first two requirements of Assumption 2.1 are satisfied; in order to 330 satisfy the third one we assume that

(3.1)
$$
h_0 = \min_{\mathbf{x} \; : \; \mathbf{x}^\top \mathbf{x} \leq 1} \mathbf{x}^\top \mathbf{A} \mathbf{x} + \mathbf{a}^\top \mathbf{x} - a_0 < 0,
$$

331 i.e., the feasible region of (1.1) has a nonempty interior. Note that the assumption can 332 be checked in polynomial time by the solution of a trust region problem. As before, 333 we denote by $X \subseteq \mathbb{R}^n$ a closed set such that $X \supset H$, i.e., it contains the ellipsoid 334 defined by the second constraint. For each $\lambda \geq 0$, the Lagrangian relaxation (2.3) 335 takes on the form

(3.2)
$$
p_X(\lambda) = \min_{\mathbf{x} \in X} \mathbf{x}^\top (\mathbf{Q} + \lambda \mathbf{A}) \mathbf{x} + (\mathbf{q} + \lambda \mathbf{a})^\top \mathbf{x} - \lambda a_0
$$

$$
\mathbf{x}^\top \mathbf{x} \le 1.
$$

336 If $X = \mathbb{R}^n$, this is the standard Lagrangian relaxation of problem (1.1) and it can be 337 solved efficiently since it is a trust region problem. Following the notation of section 338 2, let

$$
P_X(\lambda) = arg \min_{\mathbf{x} \in X \ : \ \mathbf{x}^\top \mathbf{x} \le 1} \mathbf{x}^\top (\mathbf{Q} + \lambda \mathbf{A}) \mathbf{x} + (\mathbf{q} + \lambda \mathbf{a})^\top \mathbf{x}
$$

339 be the set of optimal solutions of (3.2). To apply Algorithm 2.1 to the CDT problem 340 with $X = \mathbb{R}^n$, we need to characterize the set of optimal solutions $P_{\mathbb{R}^n} (\lambda)$ of problem 341 (3.2) with $X = \mathbb{R}^n$, which is a trust region problem. The set of optimal solutions of a 342 trust region problem has been derived, e.g., in [1, 20, 21]. Here we briefly recall the 343 different cases. For simplicity, let $S_\lambda = Q + \lambda A$ and $s_\lambda = q + \lambda a$. We distinguish the ³⁴⁴ following cases: ³⁴⁵

346 Case 1 If $S_\lambda \succ O$ and $\left\| -\frac{1}{2} S_\lambda^{-1} s_\lambda \right\| \leq 1$, then $\left| -\frac{1}{2} S_\lambda^{-1} s_\lambda \right|$ is the unique optimal solution 347 of (3.2);

348 **Case 2** Let \mathbf{u}_j be the orthonormal eigenvectors of matrix \mathbf{S}_λ , and let γ_j be the cor-349 responding eigenvalues. Let $\gamma_{\min} = \min_j \gamma_j$ and $J_\lambda = arg \min_j \gamma_j$. For each γ 350 such that $(\forall j) \gamma \neq \gamma_j$, let

$$
\mathbf{y}(\gamma) = \mathbf{y}_1(\gamma) + \mathbf{y}_2(\gamma),
$$

351 where

$$
\mathbf{y}_1(\gamma) = -\sum_{j\not\in J_{\lambda}} \frac{\mathbf{s}_\lambda^\top \mathbf{u}_j}{\gamma_j - \gamma} \mathbf{u}_j, \quad \mathbf{y}_2(\gamma) = -\sum_{j\in J_{\lambda}} \frac{\mathbf{s}_\lambda^\top \mathbf{u}_j}{\gamma_j - \gamma} \mathbf{u}_j.
$$

³⁵² Then, we have the following subcases. ³⁵³

- 354 **Case 2.1** It holds that $\mathbf{s}_{\lambda}^{\top} \mathbf{u}_{j} \neq 0$ for some $j \in J_{\lambda}$. Then, there exists a unique 355 $\gamma^* \in (-\gamma_{\min},+\infty)$ such that $\|\mathbf{y}(\gamma^*)\| = 1$ and $\mathbf{y}(\gamma^*)$ is the unique 356 optimal solution of (3.2).
- 357 **Case 2.2** It holds that $\mathbf{s}_\lambda^\top \mathbf{u}_j = 0$ for all $j \in J_\lambda$ but $\|\mathbf{y}_1(\gamma_{\min})\| \geq 1$. In this 358 case there exists a unique $\gamma^* \in [-\gamma_{\min},+\infty)$ such that $\|\mathbf{y}_1(\gamma^*)\| = 1$ 359 and $\mathbf{y}_1(\gamma^*)$ is the unique optimal solution of (3.2).
- 360 **Case 2.3** It holds that $\mathbf{s}_\lambda^\top \mathbf{u}_j = 0$ for all $j \in J_\lambda$ and $\|\mathbf{y}_1(\gamma_{\min})\| < 1$. In this 361 case we have that $P_{\mathbb{R}^n}(\lambda)$ is not a singleton and is made up by the 362 following points:

$$
(3.3) \quad P_{\mathbb{R}^n}(\lambda) = \left\{ \mathbf{y}_1(\gamma_{\min}) + \sum_{j \in J_{\lambda}} \xi_j \mathbf{u}_j \; : \; \sum_{j \in J_{\lambda}} \xi_j^2 = 1 - ||\mathbf{y}_1(\gamma_{\min})||^2 \right\}.
$$

³⁶³ Thus, we recognize two further subcases. ³⁶⁴

- 365 **Case 2.3.1** $|J_{\lambda}| = 1$, in which case $P_{\mathbb{R}^n}(\lambda)$ contains exactly two distinct 366 points.
- 367 **Case 2.3.2** $|J_{\lambda}| \geq 2$, in which case the set $P_{\mathbb{R}^n} (\lambda)$ contains an infinite num-368 ber of points and is a connected set.
- 369 Note that in Cases 2.3.1 and 2.3.2 we can compute the two values $h_{\mathbb{R}^n}^{\min}(\lambda), h_{\mathbb{R}^n}^{\max}(\lambda)$ 370 by solving a trust region problem over the border of a $|J_\lambda|$ -dimensional ball. More 371 precisely, we need to solve the following problems:

(3.4)
$$
\min/\max_{\xi} \mathbf{w}(\xi)^{\top} \mathbf{A} \mathbf{w}(\xi) + \mathbf{a}^{\top} \mathbf{w}(\xi) - a_0
$$

$$
\|\mathbf{w}(\xi)\|^2 = 1,
$$

372 where $\mathbf{w}(\boldsymbol{\xi}) = \mathbf{y}_1(\gamma_{\min}) + \sum_{j\in J_{\lambda}} \xi_j \mathbf{u}_j$. In these cases, where $P_{\mathbb{R}^n}(\lambda)$ is not a singleton, 373 we also set

(3.5)
$$
\mathbf{z}_1(\lambda) = \mathbf{w}(\xi_1), \quad \xi_1 \in \arg \min_{\xi : \|\mathbf{w}(\xi)\| = 1} \mathbf{w}(\xi)^\top \mathbf{A} \mathbf{w}(\xi) + \mathbf{a}^\top \mathbf{w}(\xi) - a_0, \n\mathbf{z}_2(\lambda) = \mathbf{w}(\xi_2), \quad \xi_2 \in \arg \max_{\xi : \|\mathbf{w}(\xi)\| = 1} \mathbf{w}(\xi)^\top \mathbf{A} \mathbf{w}(\xi) + \mathbf{a}^\top \mathbf{w}(\xi) - a_0,
$$

374 while in all other cases, when $P_{\mathbb{R}^n} (\lambda) = \{ \mathbf{z}^* (\lambda)\}$ is a singleton, we set

(3.6)
$$
\mathbf{z}_1(\lambda) = \mathbf{z}_2(\lambda) = \mathbf{z}^*(\lambda).
$$

375 The following statement is a direct consequence of Proposition 2.5.

 $\frac{376}{377}$ PROPOSITION 3.1. In the CDT problem (1.1), if $\lambda > 0$ and

378 \bullet $h_{\mathbb{R}^n}^{\min}(\lambda) = 0;$

- 379 \bullet or if $h_{\mathbb{R}^n}^{\max}(\lambda) = 0;$
- 380 \bullet or $h_{\mathbb{R}^n}^{\min}(\lambda) < 0 < h_{\mathbb{R}^n}^{\max}(\lambda)$ and $|J_{\lambda}| \geq 2$ (i.e., we are in Ccase 2.3.2);
- 381 $then \ p_{\mathbb{R}^n} (\lambda) = p^*$.

382 Proof. Since $\{ h_{\mathbb{R}^n}^{\min}(\lambda) , h_{\mathbb{R}^n}^{\max}(\lambda) \} \in Q_X(\lambda)$, in the first two cases $0 \in Q_X(\lambda)$ and 383 the thesis is a consequence of Proposition 2.5. If $h_{\mathbb{R}^n}^{\min}(\lambda) < 0 < h_{\mathbb{R}^n}^{\max}(\lambda)$ and $|J_{\lambda}| \geq 2$, 384 we observed that $P_{\mathbb{R}^n}(\lambda)$ is a connected set. Then, there exists $\mathbf{x}^* \in P_{\mathbb{R}^n}(\lambda)$ such 385 that $\mathbf{x}^* \in \partial H$. More precisely, \mathbf{x}^* is a point along the curve in $P_{\mathbb{R}^n}(\lambda)$ connecting 386 points $\mathbf{z}_1(\lambda)$ and $\mathbf{z}_2(\lambda)$, defined in (3.5). Thus, the lower bound $p_{\mathbb{R}^n}(\lambda)$ is equal to the 387 optimal value of problem (1.1). \Box

 Note that the first two conditions of Proposition 3.1 imply exactness of the bound 389 also for generic regions $X \supset H$, while the last condition is specific to the case $X = \mathbb{R}^n$. The following result is related to the necessary and sufficient condition under which the dual Lagrangian bound is not exact discussed in [2].

392 PROPOSITION 3.2. In the CDT problem (1.1), $p_{\mathbb{R}^n}(\lambda_{\mathbb{R}^n}) \neq p^*$ if and only if $\lambda_{\mathbb{R}^n} >$ 393 0, $P_{\mathbb{R}^n}(\lambda_{\mathbb{R}^n})$ contains exactly two points (Case 2.3.1), and $0 \in (h_{\mathbb{R}^n}^{\min}(\lambda_{\mathbb{R}^n}), h_{\mathbb{R}^n}^{\max}(\lambda_{\mathbb{R}^n}))$.

394 Proof. First note that, in view of Proposition 2.3, the dual Lagrangian bound is 395 always exact when $\lambda \Bbb B^n = 0$. When $\lambda \Bbb B^n > 0$, the result is a consequence of Proposition 396 3.1 and the fact that for $|J_{\lambda_{\Bbb R} n} |= 1$ it holds that $Q_{\Bbb R^n} (\lambda_{\Bbb R^n}) = \{ h_{\Bbb R^n}^{\min} (\lambda_{\Bbb R^n}) , h_{\Bbb R^n}^{\max} (\lambda_{\Bbb R^n}) \} \not \ni$ 397 0. \Box

398 Now, we introduce an example where $p_{\mathbb{R}^n} (\lambda_{\mathbb{R}^n}) \neq p^*$, that is the dual Lagrangian 399 bound is not exact, which will also be helpful in the following sections.

400 Example 3.3. Let us consider the following example taken from [12]:

$$
\mathbf{Q} = \begin{pmatrix} -4 & 1 \\ 1 & -2 \end{pmatrix}, \quad \mathbf{A} = \begin{pmatrix} 3 & 0 \\ 0 & 1 \end{pmatrix}, \quad \mathbf{q} = (1\ 1) \quad \mathbf{a} = (0\ 0), \quad a_0 = 2.
$$

401 Such an instance has optimal value -4 attained at points $\Big(\frac{\sqrt{2}}{2}, -\frac{\sqrt{2}}{2}\Big)$ and $\Big(-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2}\Big)$. 402 The maximizer of $p_{\mathbb{R}^2} (\lambda)$ is $\lambda_{\mathbb{R}^2} = 1$ for which we have

$$
h_{\mathbb{R}^2}^{\min}\approx -0.66<0<0.66\approx h_{\mathbb{R}^2}^{\max},
$$

and, moreover,
$$
|J_{\lambda_{\mathbb{R}^2}}| = 1
$$
, so that we have exactly two optimal solutions of (3.2), one violating the second constraint, namely $\mathbf{z}_2(\lambda_{\mathbb{R}^2}) = (-0.911, 0.4114)$, point x_1 in Figure 1, displayed as \circ , the other in $int(H)$, point z_1 in Figure 1, displayed as \times . The lower bound is $p_{\mathbb{R}^2}(1) = -4.25$, which is not exact.

407 Now, let us assume that the dual Lagrangian bound is not exact, i.e., as previously 408 stated in Proposition 3.2

$$
0 \in \left(h^{\min}_{\mathbb{R}^n}(\lambda_{\mathbb{R}^n}), h^{\max}_{\mathbb{R}^n}(\lambda_{\mathbb{R}^n})\right), \quad |J_{\lambda_{\mathbb{R}^n}}| = 1.
$$

409 Recall that, by Proposition 3.2, in this case, there exists a single point $\mathbf{z}_1(\lambda_{\mathbb{R}^n}) \in$ $P_{\mathbb{R}^n} (\lambda_{\mathbb{R}^n})\cap H$ (actually $\mathbf{z}_1(\lambda_{\mathbb{R}^n}) \in int(H)$), and a single point $\mathbf{z}_2(\lambda_{\mathbb{R}^n}) \in P_{\mathbb{R}^n} (\lambda_{\mathbb{R}^n})\setminus H$. Now we show that the dual Lagrangian bound can be strictly improved through the addition of a linear cut. We first observe that the optimal value of problem (1.1) does not change if we add constraints which are implied by the second one.

414 In the following proposition, we define a projection $\Pi_{A,a} : \mathbb{R}^n \setminus H \rightarrow \partial H$ that 415 maps $\mathbf{x} \notin H$ to the element of ∂H located on the segment that joins \mathbf{x} to the center 416 of the ellipsoid H (given by $\boldsymbol{\alpha} = -\frac{1}{2}\mathbf{A}^{-1}\mathbf{a}$).

417 PROPOSITION 3.4. For $\mathbf{x} \notin H$, set $\Pi_{\mathbf{A},\mathbf{a}}(\mathbf{x}) = \sqrt{\frac{-h(\boldsymbol{\alpha})}{h(\mathbf{x}) - h(\boldsymbol{\alpha})}}(\mathbf{x} - \boldsymbol{\alpha}) + \boldsymbol{\alpha}$, where 418 $\alpha = -\frac{1}{2}\mathbf{A}^{-1}\mathbf{a}$ is the center of the ellipsoid. Then $h(\Pi_{\mathbf{A},\mathbf{a}}(\mathbf{x})) = 0$.

FIG. 1. Optimal solutions of the dual Lagrangian bound outside H (x_1) and in int(H) (z_1) , denoted by \circ and \times , respectively. The continuous red curve is the border of the unit ball, while the dotted blue curve is the border of the ellipsoid H. (Figure in color online.)

419 *Proof.* Note that
$$
(\forall \beta \in \mathbb{R}) h(\beta(\mathbf{x} - \alpha) + \alpha) - h(\alpha) = \beta^2(h(\mathbf{x}) - h(\alpha))
$$
 (it is a

420 consequence of the fact that function h is quadratic and it can be verified by direct substitution). Then $h(\Pi_{\mathbf{A},\mathbf{a}}(\mathbf{x})) = h\left(\sqrt{\frac{-h(\boldsymbol{\alpha})}{h(\mathbf{x}) - h(\boldsymbol{\alpha})}}(\mathbf{x} - \boldsymbol{\alpha}) + \boldsymbol{\alpha}\right) = \frac{-h(\boldsymbol{\alpha})}{h(\mathbf{x}) - h(\boldsymbol{\alpha})}$ 421 substitution). Then $h(\Pi_{\mathbf{A},\mathbf{a}}(\mathbf{x})) = h\left(\sqrt{\frac{-h(\alpha)}{h(\mathbf{x})-h(\alpha)}}(\mathbf{x}-\alpha)+\alpha\right) = \frac{-h(\alpha)}{h(\mathbf{x})-h(\alpha)}(h(\mathbf{x})-$

422
$$
h(\alpha) + h(\alpha) = 0.
$$

423 Given any $\bar{\mathbf{x}} \in \mathbb{R}^n$, it holds, by convexity, that

$$
\mathbf{x}^\top \mathbf{A} \mathbf{x} + \mathbf{a}^\top \mathbf{x} \ge \bar{\mathbf{x}}^\top \mathbf{A} \bar{\mathbf{x}} + \mathbf{a}^\top \bar{\mathbf{x}} + (2\mathbf{A} \bar{\mathbf{x}} + \mathbf{a})^\top (\mathbf{x} - \bar{\mathbf{x}}).
$$

 \Box

424 Thus, the following linear constraint is implied by the second constraint in (1.1):

(3.7)
$$
(2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a})^{\top}\mathbf{x} - \bar{\mathbf{x}}^{\top}\mathbf{A}\bar{\mathbf{x}} \leq a_0,
$$

425 and, consequently, it can be added to problem (1.1) without modifying its feasible 426 region. In particular, if $\bar{\mathbf{x}} \in \partial H$, being $\bar{\mathbf{x}}^T \mathbf{A} \bar{\mathbf{x}} + \mathbf{a}^T \bar{\mathbf{x}} = a_0$, the linear constraint is

(3.8)
$$
(2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a})^{\top}(\mathbf{x} - \bar{\mathbf{x}}) \leq 0.
$$

427 Due to the redundancy of the linear constraint for problem (1.1) , we can define, for a 428 given $\bar{\mathbf{x}} \in \partial H$, the new Lagrangian problem

(3.9)
$$
p_X(\lambda) = \min_{\mathbf{x}} \quad \mathbf{x}^\top (\mathbf{Q} + \lambda \mathbf{A}) \mathbf{x} + (\mathbf{q} + \lambda \mathbf{a})^\top \mathbf{x} - \lambda a_0
$$

$$
\mathbf{x}^\top \mathbf{x} \le 1
$$

$$
(2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a})^\top (\mathbf{x} - \bar{\mathbf{x}}) \le 0,
$$

429 where

(3.10)
$$
X = \Omega_{\bar{\mathbf{x}}} = {\mathbf{x} : (2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a})^{\top}(\mathbf{x} - \bar{\mathbf{x}}) \leq 0} \supset H.
$$

430 If we set $\bar{\mathbf{x}} = \Pi_{\mathbf{A},\mathbf{a}}(\mathbf{z}_2(\lambda_{\mathbb{R}^n}))$, i.e., $\bar{\mathbf{x}}$ is the projection over ∂H of the single point in 431 $P_{\mathbb{R}^n}(\lambda_{\mathbb{R}^n}) \setminus H$, then $\Bbb R^n \supset X \supset H$ and, moreover, $X \cap (P_{\Bbb R^n}(\lambda_{\mathbb{R}^n}) \setminus H) = \emptyset$, so that, 432 by Proposition 2.12, $\bar{p}_X > \bar{p}_{\mathbb{R}^n}$. Then, if we run again Algorithm 2.1 with input 433 $X = \Omega_{\overline{\mathbf{x}}}$ defined in (3.10) and λ _{init} $= \lambda_{\mathbb{R}^n}$ (or $\lambda_{\text{init}} = \lambda_{\mathbb{R}^n}$), we are able to improve 434 strictly the dual Lagrangian bound. Note that problem (3.9), needed to compute 435 function $p_{\Omega_{\bar{x}}}$, can be solved in polynomial time according to the results proved in 436 [12, 23]. But we also discuss an alternative way to solve problem (3.9), based on the 437 solution of a trust region problem. For $\lambda = \lambda_{\mathbb{R}^n}$, after the addition of the linear cut, 438 a unique optimal solution exists, lying in $int(H)$ and, consequently, in $int(\Omega_{\bar{x}})$, since also the linear constraint in (3.9) is not active at it, being H a subset of the region $\mathbf{AQ4}$ defined by the linear cut. By continuity, for λ values smaller than but close to $\lambda_{\mathbb{R}^n}$, defined by the linear cut. By continuity, for λ values smaller than but close to $\lambda_{\mathbb{R}^n}$, 441 the unique optimal solution of (3.9) also lies in $int(H)$, i.e., $P_{\Omega_{\bar{x}}}(\lambda) = {\bf z}_1(\lambda)$ with 442 $\mathbf{z}_1(\lambda) \in int(H)$. Thus, such optimal solution must be a local and nonglobal optimal 443 solution of the trust region problem (3.2) with $X = \mathbb{R}^n$. Indeed, the globally optimal 444 solutions of this trust region problem always violate the second constraint in (1.1) for 445 all $\lambda < \lambda_{\mathbb{R}^n}$. Now, for all $\lambda \in [0, \lambda_{\mathbb{R}^n})$, we first check whether a local and nonglobal 446 optimal solution of problem (3.2) with $X = \mathbb{R}^n$ exists by exploiting the necessary and 447 sufficient condition stated in [24]. Also recall that, if it exists, the local and nonglobal 448 minimizer is unique. If it does not exist, then we set $f_1 = +\infty$. Otherwise, if it exists, 449 we denote it by $\mathbf{z}_1(\lambda)$. If $\mathbf{z}_1(\lambda) \notin \Omega_{\overline{\mathbf{x}}}$, then we set again $f_1 = +\infty$, otherwise we denote 450 by f_1 the value of the objective function of (3.9) evaluated at $z_1(\lambda)$. If some globally 451 optimal solution of the trust region problem (3.2) with $X = \mathbb{R}^n$ belongs to $\Omega_{\bar{\mathbf{x}}}\setminus H$, then 452 it is also a solution of (3.9) and we set f_2 equal to the optimal value of this problem. 453 Note that in this case $f_2 < f_1$, since f_1 is the function value at a local and nonglobal 454 solution of the trust region problem. Then, Algorithm 2.1 sets $\lambda^{\min} = \lambda$. Instead, 455 if all globally optimal solutions of the trust region problem do not belong to $\Omega_{\bar{x}}$, we 456 proceed as follows. We consider the best feasible solutions of problem (3.9) for which 457 the linear constraint is imposed to be active. The resulting problem is converted into 458 a trust region problem, after the change of variable $\mathbf{x} = \bar{\mathbf{x}} + \mathbf{V}\mathbf{z}$, where $\mathbf{V} \in \mathbb{R}^{n \times (n-1)}$ 459 is a matrix whose columns form a basis for the null space of vector $2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a}$. The 460 resulting (trust region) problem is (3.11)

$$
\min_{\mathbf{w} \in \mathbb{R}^{n-1}} \quad \mathbf{w}^\top \mathbf{V}^\top (\mathbf{Q} + \lambda \mathbf{A}) \mathbf{V} \mathbf{w} + \left[2\bar{\mathbf{x}}^\top (\mathbf{Q} + \lambda \mathbf{A}) \mathbf{V} + (\mathbf{q} + \lambda \mathbf{a})^\top\right] \mathbf{w} + \ell(\bar{\mathbf{x}}, \lambda) \|\bar{\mathbf{x}} + \mathbf{V} \mathbf{w}\|^2 \le 1,
$$

461 where $\ell(\bar{\mathbf{x}}, \lambda) = \bar{\mathbf{x}}^\top (\mathbf{Q} + \lambda \mathbf{A})\bar{\mathbf{x}} + (\mathbf{q} + \lambda \mathbf{a})^\top \bar{\mathbf{x}} - \lambda a_0$ is constant with respect to the 462 vector of variables **w**. Let $W^*(\lambda)$ be the set of optimal solutions of problem (3.11) 463 and

$$
P_1^{\star}(\lambda) = \{ \bar{\mathbf{x}} + \mathbf{V} \mathbf{w}^{\star} \; : \; \mathbf{w}^{\star} \in W^{\star}(\lambda) \}.
$$

464 Note that the set $W^*(\lambda)$ can be computed through the procedure presented in section 465 3 with the different cases (namely, Cases 1, 2.1, 2.2, 2.3.1, 2.3.2) after rewriting it as a 466 classical trust region problem. Moreover, let $f_2 < +\infty$ be the optimal value of problem 467 (3.11). Now, after comparing f_1 and f_2 , we are able to define the set $P_{\Omega_{\bar{x}}}(\lambda)$ of optimal 468 solutions for problem (3.9). More precisely, if $f_2 > f_1$, then $P_{\Omega_{\bar{x}}}(\lambda) = {\bf z}_1(\lambda)$, i.e., 469 $\mathbf{z}_1(\lambda)$ is the unique optimal solution of problem (3.9). In this case

$$
h_{\Omega_{\bar{\mathbf{x}}}}^{\min}(\lambda) = h_{\Omega_{\bar{\mathbf{x}}}}^{\max}(\lambda) = \mathbf{z}_1(\lambda)^{\top} \mathbf{A} \mathbf{z}_1(\lambda) + \mathbf{a}^{\top} \mathbf{z}_1(\lambda) - a_0.
$$

470 Instead, if $f_2 < f_1$, which always holds, e.g., if $f_1 = +\infty$, then $P_{\Omega_{\bar{x}}}(\lambda) = P_1^{\star}(\lambda)$. Since 471 all points in $P_1^{\star}(\lambda)$ lie over a supporting hyperplane of H, we must have that

$$
h_{\Omega_{\bar{\mathbf{x}}}}^{\min}(\lambda) = \min_{\mathbf{x} \in P_1^{\star}(\lambda)} \mathbf{x}^{\top} \mathbf{A} \mathbf{x} + \mathbf{a}^{\top} \mathbf{x} - a_0 \ge 0,
$$

472 and equality holds only if $\bar{\mathbf{x}} \in P_1^*(\lambda)$. In the latter case, the bound is exact, otherwise 473 Algorithm 2.1 sets $\lambda^{\min} = \lambda$. Finally, if $f_1 = f_2$, then $P_{\Omega_{\bar{\mathbf{x}}}}(\lambda) = P_1^{\star}(\lambda) \cup \{ \mathbf{z}_1(\lambda) \}$ and 474 in this case $0 \in [h_{\Omega_{\bar{\mathbf{x}}}}^{\min}(\lambda), h_{\Omega_{\bar{\mathbf{x}}}}^{\max}(\lambda)]$ and the algorithms exits the loop. The following 475 result is a straightforward consequence of Proposition 2.12.

476 PROPOSITION 3.5. Algorithm 2.1 with $\varepsilon = 0$ will stop after a finite number of 477 iterations or will converge to some $\lambda_{\Omega_{\bar{x}}} < \lambda_{\mathbb{R}^n}$ with a new lower bound $\bar{p}_{\Omega_{\bar{x}}} > \bar{p}_{\mathbb{R}^n}$.

478 Proof. Strict inequalities hold in view of Proposition 2.12 with $X = \mathbb{R}^n$ and 479 $Y = \Omega_{\bar{\mathbf{x}}}$, since, as already observed, $\Omega_{\bar{\mathbf{x}}} \cap (P_{\mathbb{R}^n} (\lambda_{\mathbb{R}^n}) \setminus H) = \emptyset$. О

480 If the final bound is not exact, i.e., $\bar{p}_{\Omega_{\bar{x}}} = p_{\Omega_{\bar{x}}} (\lambda_{\Omega_{\bar{x}}}) \lt p^*$, at $\lambda_{\Omega_{\bar{x}}}$ we have $f_1 = f_2$ and 481 $P_{\Omega_{\bar{x}}}(\lambda_{\Omega_{\bar{x}}})$ contains multiple optimal solutions, in particular, one in $int(H)$ and the 482 other(s) outside H, more precisely on $\partial \Omega_{\bar{x}} \setminus H$. We illustrate all this on Example 3.3.

484 Example 3.6. The optimal solution of (3.2) with $X = \mathbb{R}^n$ for $\lambda_{\mathbb{R}^n} = 1$ which 485 violates the second constraint is $\mathbf{z}_2(\lambda_{\mathbb{R}^n}) = (-0.911, 0.4114)$. The lower bound is 486 p $\mathbb{R}^n (1) = -4.25$. After the addition of the linear inequality (3.7) obtained with $\bar{\mathbf{x}} =$ 487 $\Pi_{\mathbf{A},\mathbf{a}}(\mathbf{z}_2(\lambda \Bbb B^n))$, equal to the projection of $\mathbf{z}_2(\lambda \Bbb B^n)$ over the boundary of the second 488 constraint, we can run again Algorithm 2.1 with $X = \Omega_{\bar{\mathbf{x}}}$ and we get to $\lambda_{\Omega_{\bar{\mathbf{x}}} \approx 0.726$ 489 and $p_{\Omega_{\bar{x}}}(\lambda_{\Omega_{\bar{x}}} \approx -4.097$, which improves the previous lower bound. In Figure 2 we 490 show the linear cut and the two new optimal solutions outside H and in $int(H)$ (x_2) 491 and z_2 , respectively) obtained at $\lambda_{\Omega_{\bar{x}}}$. In the same figure we also display the previous 492 pair of optimal solutions in order to show the progress of the algorithm.

493 It is worthwhile to discuss at this point the relations between the approach pro-494 posed in this work and the one proposed in [27], where the classical SDP relaxation

FIG. 2. First linear cut and the two optimal solutions lying outside H (x_2) and in int(H) (z_2) , denoted by \circ and \times , respectively.

483

 of problem (1.1) is considered. Both approaches stem from the necessary and suf- ficient condition under which the dual Lagrangian bound is not exact discussed in 497 [2], namely the existence of two distinct optimal solutions, one belonging to $int(H)$ 498 and the other outside H. In both cases it is observed that, in order to improve the bound, it is necessary to separate such optimal solutions. But the way the separation is carried on in the two approaches is different. Following the terminology employed in Integer Programming, in [27] the separation is performed through a branching op- eration, while in this work it is performed through the addition of a cutting plane. 503 Indeed, in [27] first, a hyperplane $\mathbf{w}^T \mathbf{x} = v$ separating the two optimal solutions is introduced; then, two distinct subproblems are solved, one by adding the inequality $\mathbf{w}^T \mathbf{x} \leq v$ (converted into an SOCP constraint) to the SDP relaxation, the other by 506 adding the inequality $\mathbf{w}^T \mathbf{x} \geq v$ to the SDP relaxation; finally, the new bound is set equal to the minimum of the bounds over the two subregions into which the original feasible region has been split. Note that one of the two subregions may be empty, in 509 which case its corresponding lower bound is set equal to $+\infty$ and the linear inequality is a separating hyperplane between H and the optimal solution outside H. In this paper the separation is performed through the addition of a linear cut and a single subproblem is solved. Moreover, in [27] it is observed that one could search for an 513 "optimal' hyperplane separating the two optimal solutions, namely one which leads to the best possible bound. In that paper such a hyperplane is derived in the special case when the function h is the product of two affine functions and an exactness result is also provided for the case of problems with two variables, but the question about how to characterize an '`optimal`' affine function is left open in the general case. In the next section we will be able to provide a necessary and sufficient condition for a linear cut to be the one delivering the best bound (Proposition 4.2). Based on this condition, we will also be able to propose a procedure to improve the bound by local adjustments of the linear cut. Finally, in this paper we will also show in section 5 that the bound can be further improved through the addition of a second linear cut, possibly followed by a local adjustment of the two linear cuts. The experiments in section 6 will show that the bound obtained by the addition of two linear cuts is quite a good one, allowing one to solve all except one of the 212 hardest instances intro- duced in [12]. As a final remark, we observe that the approach presented in [27] and the one discussed in this paper could actually be combined by performing a branching operation (as in [27]) followed by the addition of a linear cut (as in this work) in each branch. Borrowing again from the terminology of Integer Programming, this can be viewed as a branch-and-cut approach.

 4. Improving the bound by local adjustments of the linear cut. In the 532 previous section we proposed to set $\bar{\mathbf{x}}$ equal to the projection over ∂H of $\mathbf{z}_2(\lambda \mathbb{R}^n)$, the 533 optimal solution of problem (3.2) with $X = \mathbb{R}^n$ lying outside H. However, this point can be improved by some local adjustment. We first give a necessary and sufficient 535 condition under which the current point $\bar{\mathbf{x}}$ cannot be improved. The proof will also suggest how to improve the point (and the bound) when the condition is not fulfilled. Let

(4.1)
$$
r(\mathbf{w}, \mathbf{x}) = (2\mathbf{A}\mathbf{w} + \mathbf{a})^\top (\mathbf{x} - \mathbf{w}) + \mathbf{w}^\top \mathbf{A} \mathbf{w} + \mathbf{a}^\top \mathbf{w} - a_0 = (2\mathbf{A}\mathbf{w} + \mathbf{a})^\top \mathbf{x} - \mathbf{w}^\top \mathbf{A} \mathbf{w} - a_0
$$

 be the linearization of the ellipsoid constraint at w. Note that constraint (3.7) can be 539 written as $r(\bar{\mathbf{x}}, \mathbf{x}) \leq 0$. Also note that for each \mathbf{x}, r is a concave function with respect to w. Next, we set

$$
p(\lambda, \mathbf{w}) = p_{\Omega_{\mathbf{w}}}(\lambda),
$$

541 in order to highlight the dependency of the bound not only on λ but also on **w**. Then, 542 in order to maximize the lower bound, we need to solve the following problem:

$$
\max_{\lambda \geq 0, \mathbf{w}} p(\lambda, \mathbf{w}).
$$

543 As before, we denote by $P_{\Omega_{\bf w}}(\lambda)$ the optimal set of problem (3.9) with $\bar{\bf x} = {\bf w}$, while we 544 denote by $P^1_{\Omega_\mathbf{w}}(\lambda) = P_{\Omega_\mathbf{w}}(\lambda) \setminus int(H)$ the set of optimal solutions of the same problem 545 lying outside the interior of the ellipsoid H . We will need the following lemma.

546 LEMMA 4.1. Set-valued functions $\mathbf{w} \sim P_{\Omega_{\mathbf{w}}}(\lambda)$ and $\mathbf{w} \sim P_{\Omega_{\mathbf{w}}}^1(\lambda)$ are upper semi-547 continuous for any $\lambda \geq 0$.

548 Proof. Upper semicontinuity of $\mathbf{w} \sim P_{\Omega_{\mathbf{w}}}(\lambda)$ follows from the maximum theorem 549 (see, for instance, Theorem 1.4.16 of [5]), while upper semicontinuity of $\mathbf{w} \sim P_{\Omega}^{1}(\lambda)$ 549 (see, for instance, Theorem 1.4.16 of [5]), while upper semicontinuity of $\mathbf{w} \sim P_{\Omega_{\mathbf{w}}}^1(\lambda)$ 550 follows from the fact that $P_{\Omega_{\mathbf{w}}}^1(\lambda)$ is obtained by intersecting the upper semicontinuous 551 function $\mathbf{w} \sim P_{\Omega_{\mathbf{w}}}(\lambda)$ with the compact set $\{\mathbf{x} : ||\mathbf{x}|| \leq 1\} \ \int \int \mathbf{g}$ (see, for instance, Proposition 1.4.9 of [5]). Proposition 1.4.9 of $[5]$).

553 Now, the following proposition characterizes the maxima of p.

554 PROPOSITION 4.2. Let $(\lambda^*, \mathbf{w}^*)$ be such that $\mathbf{w}^* \in \partial H, \ \lambda^* > 0, \ 0 \in (h_{\Omega_{\mathbf{w}^*}}^{\min}(\lambda^*),$ 555 $h_{\Omega_{\mathbf{w^*}}}^{\max}(\lambda^*))$, and $0 \notin Q_{\Omega_{\mathbf{w^*}}}(\lambda^*)$. Assume also that $(\forall \mathbf{v} \in P_{\Omega_{\mathbf{w^*}}}^1(\lambda^*))$ $r(\mathbf{w^*}, \mathbf{v}) = 0$. Then, 556 the following statements are equivalent:

557 (i)
$$
(\lambda^*, \mathbf{w}^*) = \operatorname{argmax}_{(\lambda \ge 0, \mathbf{w})} p(\lambda, \mathbf{w}).
$$

558 (ii)
$$
(\forall \mathbf{w} \in \mathbb{R}^n) P^1_{\Omega_{\mathbf{w}}}(\lambda^*) \neq \emptyset
$$
.

559 (iii)

(4.2)
$$
(\forall \mathbf{d} \in \mathbb{R}^n) (\exists \mathbf{v} \in P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*)) \; : \; -\mathbf{d}^\top \mathbf{A} \mathbf{d} + 2\mathbf{d}^\top \mathbf{A} (\mathbf{v} - \mathbf{w}^*) \leq 0.
$$

560 Proof. Before proving the result we make some remarks. First, note that $0 \in$ $(h_{\Omega_{\mathbf{w^*}}}^{\min}(\lambda^*), h_{\Omega_{\mathbf{w^*}}}^{\max}(\lambda^*))$ implies that $\lambda^* = \lambda_{\Omega_{\mathbf{w^*}}}$. Moreover, since $0 \notin Q_{\Omega_{\mathbf{w^*}}}(\lambda^*)$ and $\lambda^* > 0$, by Propositions 2.3 and 2.5, $p_{\Omega_{\mathbf{w}^*}}(\lambda^*) < p^*$ (i.e., the bound is not exact). If 563 the bound were exact, the current pair $(\mathbf{w}^*, \lambda^*)$ would obviously be optimal. Also 564 note that $h_{\Omega_{\mathbf{w}^*}}^{\min}(\lambda^*) < 0 < h_{\Omega_{\mathbf{w}^*}}^{\max}(\lambda^*)$ implies that $P_{\Omega_{\mathbf{w}^*}}^1(\lambda^*) \neq \emptyset$. Finally, condition $(\forall \mathbf{v} \in P^1_{\Omega_{\mathbf{w}^*}}(\mathbf{\lambda}^*))$ $r(\mathbf{w}^*, \mathbf{v}) = \mathbf{0}$ means that $P^1_{\Omega_{\mathbf{w}^*}}(\mathbf{\lambda}^*)^{\sim} \subset \partial \Omega_{\mathbf{w}^*}$, which, according to the discussion about the optimal solutions of problem (3.9), holds true provided that $\mathbf{z}_2(\lambda \Bbb{R}^n) \notin \Omega_{\mathbf{w}^*}.$

568 (i) \rightarrow (ii) By contradiction, let **w** be such that $P_{\Omega_{\mathbf{w}}}^1(\lambda^*) = \emptyset$. Then, $P_{\Omega_{\mathbf{w}}}(\lambda^*) \subset int(H)$. Therefore, function $\lambda \sim p_{\Omega_{\mathbf{w}}}(\lambda)$ is strictly decreasing at λ^* . As a consequence, 570 there exists $0 \leq \bar{\lambda} < \lambda^*$ such that $p(\bar{\lambda}, \mathbf{w}) = p_{\Omega_{\mathbf{w}}}(\bar{\lambda}) > p_{\Omega_{\mathbf{w}}}(\lambda^*) = p(\lambda^*, \mathbf{w})$. More-571 over, $p(\lambda^*, \mathbf{w}^*) = p(\lambda^*, \mathbf{w})$. Indeed, since $0 \in (h_{\Omega_{\mathbf{w}^*}}^{\min}(\lambda^*), h_{\Omega_{\mathbf{w}^*}}^{\max}(\lambda^*))$, we have that 572 $h_{\Omega_{\mathbf{w^*}}}^{\min}(\lambda^*) \leq 0$, so that $P_{\Omega_{\mathbf{w^*}}}(\lambda^*) \cap int(H)$ is not empty and is equal to $P_{\Omega_{\mathbf{w}}}(\lambda^*)$. 573 Hence, $p(\bar{\lambda}, \mathbf{w}) > p(\lambda^*, \mathbf{w}^*)$, which contradicts (i).

574 (ii) \rightarrow (iii) By contradiction, let us assume that there exists $\mathbf{d} \in \mathbb{R}^n$ such that

(4.3)
$$
(\forall \mathbf{v} \in P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*)) - \mathbf{d}^\top \mathbf{A} \mathbf{d} + 2\mathbf{d}^\top \mathbf{A} (\mathbf{v} - \mathbf{w}^*) > 0.
$$

575 By continuity of the left-hand side of the inequality in (4.3) with respect to v, there 576 exists a neighborhood B_1 of $P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*)$ such that

$$
(\forall \mathbf{v} \in B_1) \mathbf{-d}^\top \mathbf{A} \mathbf{d} + 2\mathbf{d}^\top \mathbf{A} (\mathbf{v} - \mathbf{w}^*) > 0,
$$

577 which implies that

(4.4)
$$
(\forall \mathbf{v} \in B_1) 2\mathbf{d}^\top \mathbf{A} (\mathbf{v} - \mathbf{w}^*) > 0.
$$

578 By upper semicontinuity of set-valued function $\mathbf{w} \sim P_{\Omega_{\mathbf{w}}}^1(\lambda^*)$ (see Lemma 4.1), 579 there exists a neighborhood B_2 of \mathbf{w}^* such that $(\forall \mathbf{w} \in B_2)$ $P^1_{\Omega_{\mathbf{w}}}(\lambda^*) \subset B_1$. Let $\bar{\eta} > 0$ 580 be such that $\mathbf{w}^* + \bar{\eta} \mathbf{d} \in B_2$ and consider function $\rho : [0, \bar{\eta}] \times \Bbb R^n \rightarrow \Bbb R, \rho (\eta , \mathbf{v}) =$ 581 $r(\mathbf{w}^* + \eta \mathbf{d}, \mathbf{v})$. Then, by definition of r in (4.1),

$$
\rho(\eta, \mathbf{v}) = -\eta^2 \mathbf{d}^\top \mathbf{A} \mathbf{d} + 2\eta \mathbf{d}^\top \mathbf{A} (\mathbf{v} - \mathbf{w}^*) + \rho(0, \mathbf{v}).
$$

582 By (4.4), $(\forall v \in B_1) \partial_{\eta} \rho (0,v) > 0$, where ∂_{η} denotes the partial derivative with respect 583 to η . By continuity of ρ , there exists a continous function $\hat{\eta} : B_1 \rightarrow (0, \bar{\eta}]$ such that

(4.5)
$$
(\forall \mathbf{v} \in B_1, \eta \in (0, \hat{\eta}(\mathbf{v})]) \rho(\eta, \mathbf{v}) > \rho(\mathbf{0}, \mathbf{v}).
$$

584 Hence, since B_1 is a compact set and $\hat{\eta}$ is continuous and strictly positive, setting

585 $\tilde{\eta} = \min_{\mathbf{v} \in B_1} \hat{\eta}(\mathbf{v})$, it follows that $(\forall \eta \in [0, \tilde{\eta}]) P^1_{\Omega_{\mathbf{w}^* + \eta \mathbf{d}}}(\lambda^*) \subseteq P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*)$. Moreover, 586 since, by assumption, $(\forall \mathbf{v} \in P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*)) r(\mathbf{w}^*, \mathbf{v}) = 0$,

$$
(\forall \mathbf{v} \in P_{\Omega_{\mathbf{w}^*}}^1(\lambda^*)) \,\rho(0,\mathbf{v}) = 0.
$$

587 Hence, (4.5) implies that

(4.6)
$$
(\forall \mathbf{v} \in P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*)) \rho(\tilde{\eta}, \mathbf{v}) > 0.
$$

588 Being $P^1_{\Omega_{\mathbf{w}^*} + \bar{\eta} \mathbf{d}}(\lambda^*) \subseteq P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*), (4.6)$ implies that $P^1_{\Omega_{\mathbf{w}^*} + \bar{\eta} \mathbf{d}}(\lambda^*) = \emptyset$, which contradicts 589

590 (iii) \rightarrow (i) By contradiction, there exists a couple $(\bar{\lambda}, \bar{\mathbf{w}})$ such that $p(\bar{\lambda}, \bar{\mathbf{w}}) > p(\lambda^*, \mathbf{w}^*)$. 591 . In particular, we can take $\overline{\lambda} = \lambda_{\Omega_{\overline{\mathbf{w}}}}$. In case $\overline{\lambda} = 0$, by assumption $\overline{\lambda} < \lambda^*$. Now we show that the same inequality holds true also when $\bar{\lambda} > 0$. If $\bar{\lambda} > 0$, then $\bar{\lambda} = \lambda_{\Omega_{\bar{\mathbf{w}}}}$ 592 593 implies that $0 \in [h_{\Omega_{\mathbf{w}}}^{\min}(\bar{\lambda})]$, $h_{\Omega_{\mathbf{w}}}^{\max}(\bar{\lambda})]$. Since, by assumption, $0 \in (h_{\Omega_{\mathbf{w}}^{*}}^{\min}(\lambda^{*}), h_{\Omega_{\mathbf{w}}^{*}}^{\max}(\lambda^{*}))$, 594 we have that both $P_{\Omega_{\mathbf{w}}}(\bar{\lambda}) \cap H \neq \emptyset$ and $P_{\Omega_{\mathbf{w}^*}}(\lambda^*) \cap H \neq \emptyset$ (i.e., the minimum values 595 $p(\lambda^*, \mathbf{w}^*)$ and $p(\bar{\lambda}, \bar{\mathbf{w}})$ are both attained in H). Hence, $p(\bar{\lambda}, \bar{\mathbf{w}}) > p(\lambda^*, \mathbf{w}^*)$ implies 596 bat $\bar{\lambda} < \lambda^*$. Indeed, let us assume that $\bar{\lambda} \geq \lambda^*$ and let $\mathbf{z} \in P_{\Omega_{\mathbf{w}^*}}(\lambda^*) \cap H$. Then,

$$
p(\lambda^*, \mathbf{w}^*) = \mathbf{z}^\top \mathbf{Q} \mathbf{z} + \mathbf{q}^\top \mathbf{z} + \lambda^* (\mathbf{z}^\top \mathbf{Q} \mathbf{z} + \mathbf{q}^\top \mathbf{z} - a_0)
$$

\n
$$
\geq \mathbf{z}^\top \mathbf{Q} \mathbf{z} + \mathbf{q}^\top \mathbf{z} + \bar{\lambda} (\mathbf{z}^\top \mathbf{Q} \mathbf{z} + \mathbf{q}^\top \mathbf{z} - a_0) \geq p(\bar{\lambda}, \bar{\mathbf{w}}),
$$

597 which is a contradiction. Then, function $p(\lambda^*, \bar{\mathbf{w}})$ must be decreasing at λ^* or, equiv-598 alently, $P_{\Omega_{\bar{\mathbf{w}}}}(\lambda^*) \subset int(H)$ and $P_{\Omega_{\bar{\mathbf{w}}}}(\lambda^*) \neq \emptyset$. Since $h_{\Omega_{\mathbf{w}^*}}^{\min}(\lambda^*) < 0$, then $P_{\Omega_{\bar{\mathbf{w}}}}(\lambda^*)$ 599 and $P_{\Omega_{\mathbf{w}}*}(\lambda^*)$ have a common nonempty intersection within H and, consequently, 600 $p(\lambda^*, \bar{\mathbf{w}}) = p(\lambda^*, \mathbf{w}^*)$ holds. This implies that $P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*) \cap \Omega_{\bar{\mathbf{w}}} = \emptyset$. Indeed, assume 601 there exists $\mathbf{v} \in P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*) \cap \Omega_{\bar{\mathbf{w}}}$. Note that $\mathbf{v} \notin int(\tilde{H})$ and, since $P^1_{\Omega_{\bar{\mathbf{w}}}}(\lambda^*) \subset P_{\Omega_{\bar{\mathbf{w}}}}(\lambda^*),$ 602 v would also belong to $P_{\Omega_{\bar{\mathbf{w}}}}(\lambda^*)$ which, however, contradicts $P_{\Omega_{\bar{\mathbf{w}}}}(\lambda^*) \subset int(H)$.

603 Condition $P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*) \cap \Omega_{\bar{\mathbf{w}}} = \emptyset$ is equivalent to

$$
\left(\forall {\bf v}\in P^1_{\Omega_{{\bf w}^*}}(\lambda^*)\right)r(\bar{\bf w},{\bf v})>0.
$$

604 Note that, by assumption, $\mathbf{v} \in P^1_{\Omega_{\mathbf{w}^*}}(\lambda^*)$ implies $\mathbf{v} \in \partial \Omega_{\mathbf{w}^*}$. Moreover,

$$
r(\bar{\mathbf{w}}, \mathbf{v}) = r((\bar{\mathbf{w}} - \mathbf{w}^*) + \mathbf{w}^*, \mathbf{v}) = -(\bar{\mathbf{w}} - \mathbf{w}^*)^\top \mathbf{A} (\bar{\mathbf{w}} - \mathbf{w}^*) + 2(\bar{\mathbf{w}} - \mathbf{w}^*)^\top \mathbf{A} (\mathbf{v} - \mathbf{w}^*) + r(\mathbf{w}^*, \mathbf{v}) > 0.
$$

605 Being $\mathbf{w}^* \in \partial H$ and $\mathbf{v} \in \partial \Omega_{\mathbf{w}^*}$, we have that $r(\mathbf{w}^*, \mathbf{v}) = 0$. Then, by taking $\mathbf{d} = \bar{\mathbf{w}} - \mathbf{w}^*$, 606 (iii) is contradicted. \Box

607 Given the current point $\bar{\mathbf{x}}$ with $\lambda_{\Omega_{\bar{\mathbf{x}}}} > 0$, the question now is either to find a direction 608 d fulfilling

(4.7)
$$
(\forall \mathbf{v} \in P_1^{\star}(\lambda_{\Omega_{\bar{\mathbf{x}}}})) - \mathbf{d}^{\top} \mathbf{A} \mathbf{d} + 2 \mathbf{d}^{\top} \mathbf{A} (\mathbf{v} - \bar{\mathbf{x}}) > 0
$$

609 or to establish that it does not exist. In case it does not exist,

$$
p(\bar{\mathbf{x}}, \lambda_{\Omega_{\bar{\mathbf{x}}}}) = \max_{\lambda \geq 0, \mathbf{w}} p(\lambda, \mathbf{w}).
$$

610 Otherwise, direction $(d, -1)$ is an increasing direction for function p. We discuss 611 different cases depending on the cardinality of $P_1^*(\lambda_{\Omega_{\bar{x}}})$ (see the cases discussed in 612 section 3 for the trust region problem).

613 4.1. $|P_1^{\star}(\lambda_{\Omega_{\bar{x}}})|=1$. In this case, let v be the unique point in $P_1^{\star}(\lambda_{\Omega_{\bar{x}}})$. Then 614 we need to solve the following convex optimization problem:

$$
\max_{\mathbf{d}\in\mathbb{R}^n} -\mathbf{d}^\top\mathbf{A}\mathbf{d} + 2\mathbf{d}^\top\mathbf{A}(\mathbf{v} - \bar{\mathbf{x}}),
$$

615 whose optimal solution is $\mathbf{d} = \mathbf{v} - \bar{\mathbf{x}}$ and its optimal value is $(\mathbf{v} - \bar{\mathbf{x}})^{\top} \mathbf{A}(\mathbf{v} - \bar{\mathbf{x}}) > 0$.

616 Therefore, if $|P_1^{\star}(\lambda_{\Omega_{\bar{x}}})|=1$, we are always able to locally adjust the current point \bar{x} 617 in such a way that the bound can be improved.

618 4.2. $|P_1^{\star}(\lambda_{\Omega_{\bar{x}}})|=2$. In this case, let v_1 and v_2 be the two optimal points in 619 $P_1^{\star}(\lambda_{\Omega_{\bar{x}}})$. Then, we need to solve the following optimization problem:

(4.8)
$$
\max_{\mathbf{d}\in\mathbb{R}^n}\min\{-\mathbf{d}^\top\mathbf{A}\mathbf{d}+2\mathbf{d}^\top\mathbf{A}(\mathbf{v}_1-\bar{\mathbf{x}}),-\mathbf{d}^\top\mathbf{A}\mathbf{d}+2\mathbf{d}^\top\mathbf{A}(\mathbf{v}_2-\bar{\mathbf{x}})\},
$$

620 or, equivalently

$$
\begin{array}{cc} v \\ v \leq & -\bm{d}^\top \bm{A}\bm{d} + 2\bm{d}^\top \bm{A}(v_1 - \bar{x}) \\ v \leq & -\bm{d}^\top \bm{A}\bm{d} + 2\bm{d}^\top \bm{A}(v_2 - \bar{x}). \end{array}
$$

621 This is a convex optimization problem, whose solution can be obtained in closed form.

622 Indeed, by imposing the KKT conditions, it can be seen that the optimal solution has 623 the following form:

(4.9)
$$
\mathbf{d} = \beta(\mathbf{v}_1 - \bar{\mathbf{x}}) + (1 - \beta)(\mathbf{v}_2 - \bar{\mathbf{x}}), \quad \beta \in [0, 1].
$$

 $m\epsilon$

624 Now, let

$$
a = (\mathbf{v}_1 - \bar{\mathbf{x}})^{\top} \mathbf{A} (\mathbf{v}_1 - \bar{\mathbf{x}}) > 0,
$$

\n
$$
b = (\mathbf{v}_2 - \bar{\mathbf{x}})^{\top} \mathbf{A} (\mathbf{v}_2 - \bar{\mathbf{x}}) > 0,
$$

\n
$$
c = (\mathbf{v}_1 - \bar{\mathbf{x}})^{\top} \mathbf{A} (\mathbf{v}_2 - \bar{\mathbf{x}}).
$$

625 By replacing (4.9) in the objective function of (4.8) , we have that (4.8) can be rewritten 626 as

$$
\max_{\beta \in [0,1]} \min \left\{ (-\beta^2 + 2\beta)a - (1 - \beta)^2 b + 2(1 - \beta)^2 c, -\beta^2 a + (1 - \beta^2)b + 2\beta^2 c \right\}.
$$

627 The optimal solution of this problem is

$$
\beta^* = \begin{cases} 0 & \text{if } b \leq c, \\ 1 & \text{if } a \leq c, \\ \frac{b-c}{a+b-2c} & \text{otherwise.} \end{cases}
$$

628 Then, the optimal value is

$$
\begin{cases} b & \text{if } b \leq c, \\ a & \text{if } a \leq c, \\ \frac{ab-c^2}{a+b-2c} & \text{otherwise.} \end{cases}
$$

629 We notice that $a, b > 0$,

$$
a+b-2c = (\mathbf{v}_1 - \mathbf{v}_2)^{\top} \mathbf{A} (\mathbf{v}_1 - \mathbf{v}_2) > 0,
$$

630 and, by the Cauchy-Schwarz inequality,

$$
ab - 2c^2 \ge 0,
$$

631 and equality holds if and only if $(v_1 - \bar{x})$ and $(v_2 - \bar{x})$ are linearly dependent. Thus, 632 the optimal value of (4.8) is always strictly positive unless the two vectors $(v_1 - \bar{x})$ 633 and $(v_2 - \bar{x})$ lie along the same direction. More precisely, the optimal value is null 634 only if the two vectors have the same direction but opposite sign. Indeed, let

$$
\mathbf{v}_1 - \bar{\mathbf{x}} = \gamma (\mathbf{v}_2 - \bar{\mathbf{x}}).
$$

635 Then, we have $b = \gamma^2 a$ and $c = \gamma a$. If γ is positive, then either $b \leq c$ (if $\gamma \leq 1$), or $a \leq c$ 636 (if $\gamma \geq 1$) occurs, so that the optimal value is equal to a or b and is, thus, positive. If 637 ($v_1 - \bar{x}$) is not a negative multiple of $(v_2 - \bar{x})$, we are able to locally adjust \bar{x} along 638 direction

$$
\mathbf{d} = \beta^{\star}(\mathbf{v}_1 - \bar{\mathbf{x}}) + (1 - \beta^{\star})(\mathbf{v}_2 - \bar{\mathbf{x}}).
$$

639 **4.3.** $P_1^{\star}(\lambda_{\Omega_{\bar{x}}})$ is an infinite connected set. In this case we need to solve the 640 following optimization problem:

(4.10)
$$
\max_{\mathbf{d}\in\mathbb{R}^n}\min_{\mathbf{v}\in P_1^{\star}(\lambda_{\Omega_{\bar{\mathbf{x}}}})} -\mathbf{d}^{\top}\mathbf{A}\mathbf{d} + 2\mathbf{d}^{\top}\mathbf{A}(\mathbf{v} - \bar{\mathbf{x}}).
$$

 An improving direction exists if and only if the optimal value of this problem is strictly positive (note that the optimal value is always nonnegative since the inner 643 minimization problem has optimal value 0 for $d = 0$). We first remark that the problem is convex. Indeed, for each fixed \mathbf{v} , we have a concave function with respect to d, and the minimum of an infinite set of concave functions is itself a concave function (to be maximized, so that the problem is convex). The inner minimization problem can be solved in closed form. After removing the terms which do not depend on v, the inner problem to be solved is

$$
\min_{\mathbf{v}\in P_1^{\star}(\lambda_{\Omega_{\bar{\mathbf{x}}}})} 2\mathbf{d}^\top \mathbf{A} \mathbf{v}.
$$

649 • According to Case 2.3.2 in section 3, $P_1^{\star}(\lambda_{\Omega_{\bar{x}}})$ can be written as in (3.3) and the 650 minimization problem can be reduced to the computation of the minimum of a linear 651 function over the unit sphere:

$$
\min_{\pmb{\xi} \in \mathbb{R}^q \; : \; \|\pmb{\xi}\|^2 = 1} \bar{\mathbf{c}}(\mathbf{d})^\top \pmb{\xi},
$$

$$
\xi^\star = -\frac{\bar{\mathbf{c}}(\mathbf{d})}{\|\bar{\mathbf{c}}(\mathbf{d})\|},
$$

655 while the optimal value is $- \|\bar{\mathbf{c}}(\mathbf{d})\|$.

656 4.4. An algorithm for the refinement of the bound. Let $\bar{\mathbf{x}}$ and $\lambda_{\mathbb{R}^n}$ be 657 defined as in section 3. We propose Algorithm 4.1 for a bound based on successive 658 local adjustments of the linear cut. In line 2, Algorithm 2.1 is run with input $X = \Omega_{\bar{\mathbf{x}}}$ 659 and $\lambda \Bbb R^n$. Note that with a slight abuse here we are assuming that the algorithm 660 returns $\lambda \Omega$ _x and the related points z_1 and z_2 , while in practice close approximations 661 of these quantities are returned, namely λ^{\max} , $z_1(\lambda^{\max})$, and $z_2(\lambda^{\min})$. In line 3, z is 662 initialized with the input point $\bar{\mathbf{x}}$ itself and the direction \mathbf{d}^* , following the discussion 663 in section 4.1, is set equal to the difference between $\mathbf{z}_2(\lambda \Omega_{\bar{x}})$, the point outside H 664 returned by Algorithm 2.1, and $\bar{\mathbf{x}}$. The outer while loop of the algorithm (lines 4-665 20) is repeated until the bound is improved by at least a tolerance value tol. Inside 666 this loop, in line 5 the initial step size $\eta = 1$ is set and a new incumbent $y \in \partial H$ 667 is computed. The inner while loop (lines 7--15) computes the step size: until the 668 optimal value of problem (3.9) with $\bar{\mathbf{x}} = \mathbf{y}$ and $\lambda = \lambda \Omega_z$, denoted by *opt*, is lower 669 than the current lower bound Lb, we need to decrease the step size and recompute a 670 new incumbent y (lines $10-11$). If the step size falls below a given tolerance value, 671 we exit the inner loop and also the outer one. Otherwise, we have identified a new 672 valid incumbent and we set to 1 the exit flag stop for the inner loop (line 13), so 673 that, later on, a new linear inequality (3.7) with $\bar{\mathbf{x}} = \mathbf{y}$ will be computed. Then, at 674 line 17 we run Algorithm 2.1 with input $X = \Omega_{\mathbf{y}}$ and $\lambda_{\Omega_{\mathbf{z}}}$. Finally, in line 18, we 675 update point **z** and the direction \mathbf{d}^* . We remark that at each iteration $\mathbf{z}_2(\lambda_{\Omega_z})$ is 676 one optimal solution of the current subproblem (3.9) with $\lambda = \lambda_{\Omega_z}$ lying outside H 677 and at which the linear cut of the subproblem is active, i.e., $\mathbf{z}_2(\lambda_{\Omega_z}) \in P_1^{\star}(\lambda_{\Omega_z})$. As 678 seen in section 4.1, if $|P_1^{\star}(\lambda_{\Omega_z})| = 1$, i.e., $z_2(\lambda_{\Omega_z})$ is the unique optimal solution of the 679 current subproblem (3.9) with $\lambda = \lambda_{\Omega_z}$ lying outside H, then, in view of Proposition 680 2.11, the local adjustment employed in Algorithm 4.1 is guaranteed to improve the 681 bound. However, as seen in sections 4.2 and 4.3, if $P_1^{\star}(\lambda_{\Omega_z})$ contains more than one 682 point, than the proposed local adjustment is not guaranteed to improve the bound. 683 Sections 4.2 and 4.3 suggest how to define perturbing directions which still allow 684 one to improve the bound, in case they exist. However, as we will see through the 685 computational experiments, Algorithm 4.1 turns out to be time-consuming, and it 686 is more convenient to improve the bound by adding a further linear cut, as we do 687 in section 5, rather than further locally adjusting the current linear cut. In order to 688 clarify this point, we can make a comparison with Integer Linear Programming (ILP). 689 In ILP problems, once a linear relaxation is solved, a valid cut removes one optimal 690 solution of the relaxation. If the optimal solution is unique, then after the addition 691 of the valid cut, the bound improves. But if the linear relaxation has got multiple 692 solutions, then the valid cut is not guaranteed to remove all of them and, thus, the bound may not improve. It is possible to try to strengthen the valid cut in such a way

Algorithm 4.1 Bound improvement through a local adjustment of the linear cut.

Input: $\bar{\mathbf{x}}, \lambda \mathbb{R}^n$ 1: Set $Lb_{old} = -\infty$ $2: \;\;\; \mathrm{Let}\; [Lb, \lambda_{\Omega_{\bar{\mathbf{x}}}}, \mathbf{z}_1(\lambda_{\Omega_{\bar{\mathbf{x}}}}), \mathbf{z}_2(\lambda_{\Omega_{\bar{\mathbf{x}}}})] = \mathbf{DualLagrangian}(\Omega_{\bar{\mathbf{x}}}, \lambda_{\mathbb{R}^n})$ 3: Set $\mathbf{z} = \bar{\mathbf{x}}$ and $\mathbf{d}^* = \mathbf{z}_2(\lambda_{\Omega_{\bar{\mathbf{x}}}}) - \bar{\mathbf{x}}$ 4: while $Lb - Lb_{old} > tol$ do 5: Set $Lb_{old} = Lb, \eta = 1$ and $\mathbf{y} = \Pi_{\mathbf{A},\mathbf{a}}(\mathbf{z} + \mathbf{d}^*) \in \partial H$ 6: Set $stop = 0$ 7: while $stop = 0$ and $\eta > \varepsilon$ do 8: Solve problem (3.9) with $\bar{\mathbf{x}} = \mathbf{y}$ and $\lambda = \lambda \Omega_z$, and let *opt* be its optimal value 9: if $opt < Lb$ then 10: Set $\eta = \eta /2$ 11: Set $\mathbf{y} = \Pi_{\mathbf{A},\mathbf{a}}(\mathbf{z} + \eta \mathbf{d}^*) \in \partial H$ 12: else 13: Set $stop = 1$ 14: end if 15: end while 16: if $stop = 1$ then 17: Let $[Lb, \lambda \Omega$ _y, $z_1(\lambda \Omega$ _y $), z_2(\lambda \Omega$ _y $)$] = **DualLagrangian**(Ω _y, $\lambda \Omega$ _z) 18: Set $\mathbf{z} = \mathbf{y}, \ \dot{\mathbf{d}}^* = \mathbf{z}_2(\lambda_{\Omega_{\mathbf{y}}}) - \mathbf{z}$ 19: end if 20: end while 21: return Lb

- 693 that all optimal solutions of the linear relaxations are removed. But, more commonly, 694 new linear cuts are added.
- 695 Now we apply Algorithm 4.1 to our example.

696 Example 4.3. We have that **z** is initialized with $(-0.7901, 0.3565)$ and Lb with 697 -4.0971 . During the execution of Algorithm 4.1, **z** and Lb are updated as indicated 698 in Table 1.

 Interestingly, the best bound obtained in the example is exactly the one obtained for the same problem by the approach proposed in [12], based on the addition of SOC-RLT constraints. Figure 3 displays the situation at the last iteration of Algorithm 702 4.1. Problem (3.9) has three optimal solutions, one in $int(H)$ and two outside H. The

Iteration	z	Lh
1	$(-0.7204, 0.6658)$	-4.0850
2	$(-0.7742, 0.4493)$	-4.0638
3	$(-0.7481, 0.5665)$	-4.0477
4	$(-0.7607, 0.5136)$	-4.0416
5	$(-0.7556, 0.5361)$	-4.0378
6	$(-0.7571, 0.5296)$	-4.0364
7	$(-0.7568, 0.5309)$	-4.0362

TABLE 1 Iterations of Algorithm 4.1 over the example.

Fig. 3. Final linear cut after running Algorithm 4.1. Problem (3.9) has three optimal solutions, one in $int(H)$ and two outside H . The latter solutions are opposite to each other with respect to the final vector z.

 703 two optimal solutions outside H are opposite to each other with respect to the final 704 vector z, so that, as discussed in section 4.2, no further local adjustment is possible 705 to improve the bound in this case.

 5. Bound improvement through the addition of a further linear cut. Another possible way to improve the bound is by adding a further linear cut to (3.9). 708 Let $\bar{\mathbf{x}}$ and $\lambda_{\mathbb{R}^n}$ be defined as in section 3. In line 2 of Algorithm 4.1, we compute $[Lb, \lambda \Omega_{\bar{x}}, z_1(\lambda \Omega_{\bar{x}}), z_2(\lambda \Omega_{\bar{x}})] = \textbf{DualLagrangian}(\Omega_{\bar{x}}, \lambda_{\mathbb{R}^n})$, and, later on, we try to 710 locally adjust \bar{x} . Rather than doing that, we can add a further linear cut, cutting $\mathbf{z}_2(\lambda \Omega_{\bar{x}}) \notin H$ away. In particular, we add the one obtained through the projection over ∂H of $\mathbf{z}_2(\lambda \Omega_{\bar{\mathbf{x}}})$. Let $\tilde{\mathbf{x}} = \Pi_{\mathbf{A},\mathbf{a}}(\mathbf{z}_2(\lambda \Omega_{\bar{\mathbf{x}}})) \in \partial H$ be such projection. Then, we define the following problem:

(5.1)
\n
$$
\min_{\mathbf{x}} \quad \mathbf{x}^{\top}(\mathbf{Q} + \lambda \mathbf{A})\mathbf{x} + (\mathbf{q} + \lambda \mathbf{a})^{\top}\mathbf{x} - \lambda a_0
$$
\n
$$
\mathbf{x}^{\top}\mathbf{x} \le 1
$$
\n
$$
(2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a})^{\top}(\mathbf{x} - \bar{\mathbf{x}}) \le 0
$$
\n
$$
(2\mathbf{A}\tilde{\mathbf{x}} + \mathbf{a})^{\top}(\mathbf{x} - \tilde{\mathbf{x}}) \le 0,
$$

714 which is equivalent to problem (3.2) where

$$
X = \Omega_{\bar{\mathbf{x}}} \cap \Omega_{\tilde{\mathbf{x}}} = \{ \mathbf{x} : (2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a})^\top (\mathbf{x} - \bar{\mathbf{x}}) \le 0, (2\mathbf{A}\tilde{\mathbf{x}} + \mathbf{a})^\top (\mathbf{x} - \tilde{\mathbf{x}}) \le 0 \} \supset H.
$$

 A convex reformulation as the one proposed in [12, 23] for problem (3.9) is not available in this case (unless the two linear inequalities do not intersect in the interior of the unit ball). But in this case the alternative procedure discussed in section 3 turns out 718 to be useful. As before, for each value λ in the while loop of Algorithm 2.1 we can first 719 check whether a local and nonglobal optimal solution of problem (3.2) with $X = \mathbb{R}^n$ exists, by exploiting the necessary and sufficient condition stated in [24]. If it exists,

721 and belongs to $\Omega_{\bar{\mathbf{x}}} \cap \Omega_{\bar{\mathbf{x}}}$, we denote it by $\mathbf{z}_1(\lambda)$. Next, we need to compute the optimal 722 value of (5.1) when at least one of the two linear constraints is active, i.e., we need to 723 solve the following problem:

$$
\min_{\mathbf{x}} \quad \mathbf{x}^{\top}(\mathbf{Q} + \lambda \mathbf{A})\mathbf{x} + (\mathbf{q} + \lambda \mathbf{a})^{\top}\mathbf{x} - \lambda a_0
$$
\n
$$
\mathbf{x}^{\top}\mathbf{x} \le 1
$$
\n
$$
(2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a})^{\top}(\mathbf{x} - \bar{\mathbf{x}}) \le 0
$$
\n
$$
(2\mathbf{A}\tilde{\mathbf{x}} + \mathbf{a})^{\top}(\mathbf{x} - \tilde{\mathbf{x}}) \le 0
$$
\n
$$
[(2\mathbf{A}\bar{\mathbf{x}} + \mathbf{a})^{\top}(\mathbf{x} - \bar{\mathbf{x}})]^{\top}[(2\mathbf{A}\tilde{\mathbf{x}} + \mathbf{a})^{\top}(\mathbf{x} - \tilde{\mathbf{x}})] = 0.
$$

724 A convex reformulation of this problem has been proposed in [26]. Alternatively, one 725 can solve two distinct problems, each imposing that one of the two linear inequalities 726 is active. Each of these problems can be converted into a trust region problem with 727 an additional linear inequality, which can be solved in polynomial time through the 728 already mentioned convex reformulation proposed in [12, 23]. Thus, we compute the 729 set $P_1^*(\lambda) \subseteq \partial \Omega_{\bar{\mathbf{x}}} \cap \Omega_{\bar{\mathbf{x}}}$ of optimal solutions of (5.1) for which the first linear cut is 730 active, and then the set $P_2^{\star}(\lambda) \subseteq \Omega_{\bar{\mathbf{x}}} \cap \partial \Omega_{\bar{\mathbf{x}}}$ of optimal solutions of (5.1) for which 731 the second linear cut is active. Finally, the optimal values of these problems are 732 compared with the value of the local and nonglobal minimizer (if it exists) in order 733 to identify the set $P_X(\lambda)$ of optimal solutions of (5.1). At this point we are able to 734 compute $h_X^{\min}(\lambda), h_X^{\max}(\lambda)$ and update λ^{\min} and λ^{\max} accordingly. If for some λ we 735 have that $\mathbf{z}_1(\lambda) \in P_X(\lambda)$ and $P_X(\lambda) \cap [P_1^*(\lambda) \cup P_2^*(\lambda)] \neq \emptyset$, i.e., problem (5.1) has 736 an optimal solution in $int(H)$ and (at least) one optimal solution outside H , then **738** $0 \in [h_X^{\min}(\lambda), h_X^{\max}(\lambda)]$ and Algorithm 2.1 stops. We illustrate all this on Example 3.3.

 Example 5.1. We add a second linear cut obtained through the projection over ∂H of the optimal solution of problem (3.9) with $\lambda_{\Omega_{\bar{x}}} = 0.726$ outside H. This leads 741 bo a further improvement with $\lambda_{\Omega_{\bar{x}} \cap \Omega_{\bar{x}}} \approx 0.39$ and $p_{\Omega_{\bar{x}} \cap \Omega_{\bar{x}}} (\lambda_{\Omega_{\bar{x}} \cap \Omega_{\bar{x}}}) \approx -4.005$, which almost closes the gap. In Figure 4 we show the two linear cuts and the two new optimal

FIG. 4. Two linear cuts and the two optimal solutions outside H (x_3) and in int(H) (z_3) , denoted by \circ and \times , respectively.

743 solutions, one outside H and one belonging to $int(H)$ (x₃ and z₃, respectively). Again, 744 we also report the previous pairs of optimal solutions in order to show the progress.

745 Now, assume that the returned bound is not exact. Also in this case $\bar{\mathbf{x}}$ and $\tilde{\mathbf{x}}$ can be 746 locally adjusted. One can combine the techniques presented in section 4 and in the 747 current section, by using a technique similar to the one described in the former section 748 to improve the pair of points $\bar{\mathbf{x}}$ and $\tilde{\mathbf{x}}$. In particular, at $\lambda \Omega_{\bar{\mathbf{x}}} \cap \Omega_{\bar{\mathbf{x}}}$ we have one optimal 749 solution of problem 5.1 belonging to $int(H)$, namely the local and nonglobal optimal 750 solution of problem (3.2) with $X = \mathbb{R}^n$, and at least another one outside H. We denote 751 the latter by v and we observe that at least one of the two linear cuts is active at this 752 point, i.e., either $\mathbf{v} \in \partial \Omega_{\bar{\mathbf{x}}}$ or $\mathbf{v} \in \partial \Omega_{\bar{\mathbf{x}}}$ (or both). Then, if only the first cut is active at 753 v, we update $\bar{\mathbf{x}}$ as follows: $\bar{\mathbf{x}}' = \Pi_{\mathbf{A},\mathbf{a}}(\bar{\mathbf{x}}+\eta (\mathbf{v} - \bar{\mathbf{x}}))$ for a sufficiently small η value, while 754 $\tilde{\mathbf{x}}' = \tilde{\mathbf{x}}$. If only the second cut is active, we update $\tilde{\mathbf{x}}$ as follows: $\tilde{\mathbf{x}}' = \Pi_{\mathbf{A},\mathbf{a}}(\tilde{\mathbf{x}} + \eta (\mathbf{v} - \tilde{\mathbf{x}}))$ for a sufficiently small η value, while $\bar{\mathbf{x}}' = \bar{\mathbf{x}}$. Finally, if both are active we select one 756 of the two cuts and perturb it. After the perturbation, we run again Algorithm 2.1 757 with input $X = \Omega_{\tilde{\mathbf{x}}'} \cap \Omega_{\tilde{\mathbf{x}}'}$ and $\lambda_{\Omega_{\tilde{\mathbf{x}}} \cap \Omega_{\tilde{\mathbf{x}}}}$, and we repeat this procedure until there is 758 a significant reduction of the bound. Note, however, that it might happen that no 759 improvement is possible. In case $|P_1^{\star}(\lambda_{\Omega_{\bar{\mathbf{x}}}\cap \Omega_{\bar{\mathbf{x}}}})|=1$ and $P_2^{\star}(\lambda_{\Omega_{\bar{\mathbf{x}}}\cap \Omega_{\bar{\mathbf{x}}}})=\emptyset$ (similar 760 for $|P_2^{\star}(\lambda_{\Omega_{\bar{\mathbf{x}}}\cap \Omega_{\bar{\mathbf{x}}}})| = 1$ and $P_1^{\star}(\lambda_{\Omega_{\bar{\mathbf{x}}}\cap \Omega_{\bar{\mathbf{x}}}}) = \emptyset$, then the proposed perturbation $\bar{\mathbf{x}}' =$ 761 $\Pi_{\mathbf{A},\mathbf{a}}(\bar{\mathbf{x}} + \eta (\mathbf{v} - \bar{\mathbf{x}}))$ for η sufficiently small allows improvement of the bound. Indeed, 762 in such cases the local adjustment is able to cut the unique solution outside H away. 763 In order to illustrate other different cases we employ Figures 5a--5c. As usual, in these 764 figures the point in $int(H)$ is denoted by \times , while the others (outside H) are denoted 765 by \circ . If $|P_1^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}})| = |P_2^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}})| = 1$ and $P_1^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}}) \cap P_2^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}}) = \emptyset$ (see 766 Figure 5a), or $|P_1^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}})|=2, |P_2^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}})|=1$, and $P_1^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}})\cap P_2^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}})\neq \emptyset$ 767 (see Figure 5b), then it is not possible to remove all the solutions outside H by 768 perturbing a single linear cut. Indeed, in both cases the perturbation of a single 769 linear cut is able to remove just one of the two optimal solutions outside H. But it is 770 possible to remove both by perturbing both linear cuts. Instead, Figure 5c illustrates 771 a case where $|P_1^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}})| = |P_2^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}})| = 2$ and $P_1^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}}) \cap P_2^{\star}(\lambda_{\Omega_{\tilde{\mathbf{x}}}\cap\Omega_{\tilde{\mathbf{x}}}}) \neq \emptyset$. 772 In this case even the perturbation of both linear cuts is unable to remove all three 773 solutions outside H. The only way to remove all three solutions outside H is through 774 the addition of a further linear cut, but, of course, this leads to a more complex 775 problem with one trust region constraint and three linear inequalities.

776 Example 5.2. In our example, this refinement is finally able to close the gap 777 and return the exact optimal value - 4. In Figure 6 we report the result of the first 778 perturbation of the linear cuts. Since only the second linear cut is active at x_3 , in 779 this case the second linear cut is slightly perturbed and becomes equivalent to the 780 tangent to H at the optimal solution $(-\sqrt{2}/2, \sqrt{2}/2)$ of the original problem (1.1). It 781 is interesting to note that the new optimal solution outside H , indicated by x_4 , lies 782 in a different region with respect to the previous ones and is further from ∂H with 783 respect to x_2 and x_3 (the reduction of λ reduces the penalization of points outside 784 H). Such a solution is cut by the new linear inequality, obtained by a (not so small) 785 perturbation of the first linear cut, displayed in Figure 7, together with the two new 786 optimal solutions $(x_5 \text{ and } z_5)$, now corresponding to the two optimal solutions of 787 problem (1.1).

788 6. Computational experiments. In this section we report the computational 789 results for the proposed bounds over the set of hard instances selected from the random 790 ones generated in [12] and inspired by [18]. More precisely, in [12] 1000 random

(a) Three optimal solutions, none with both linear cuts active.

(b) Three optimal solutions, one with both linear cuts active.

(c) Four optimal solutions.

Fig. 6. Perturbation of the second linear cut and the two new optimal solutions outside H (x_4) and in $int(H)$ (z_4) , denoted by \circ and \times , respectively.

FIG. 7. Perturbation of the first linear cut and the two optimal solutions outside $H(x_5)$ and in $int(H)$ (z₅), denoted by \circ and \times , respectively.

791 instances were generated for each size $n = 5, 10, 20$. Some of these instances have been 792 declared hard ones, namely those for which the bound obtained by adding SOC-RLT 793 constraints was not exact. In particular, these are 38 instances with $n = 5$, 70 instances 794 with $n = 10$, and 104 instances with $n = 20$. Such instances have been made available 795 in GAMS, AMPL, and COCOUNT formats in [19]. We tested our bounds on such instances.

796 All tests have been performed on an Intel Core i7 running at 1.8 GHz with 16 GB of 797 RAM. All bounds have been coded in MATLAB.

- 798 We computed the following bounds:
- 800 \bullet LbDual, the dual Lagrangian bound computed through Algorithm 2.1 with 801 input $X = \mathbb{R}^n;$
- 802 LbOneCut, the bound obtained by adding a single linear cut and computed 803 through Algorithm 2.1 with input $X = \Omega_{\bar{x}}$;
- 804 \bullet LbOneAdj, the bound obtained by local adjustments of the added linear cut 805 as indicated in Algorithm 4.1;
- 806 \bullet LbTwoCut, the bound obtained by adding two linear cuts;
- 807 \bullet LbTwoAdj, the bound obtained by adjusting the two linear cuts.

808 According to what was done in [3, 12, 25], an instance is considered to be 'solved' 809 when the relative gap between the lower bound, say LB, and the upper bound, say 810 UB , is not larger than 10^{-4} , i.e.,

$$
\frac{UB - LB}{|UB|} \le 10^{-4}.
$$

811 We set UB equal to the lowest value obtained by running, after the addition of the 812 first linear cut, two local searches for the original problem (1.1), one from the optimal 813 solution $\mathbf{z}_1(\lambda_{\Omega_{\bar{\mathbf{x}}}}) \in int(H)$ of (3.9) returned at the end of Algorithm 2.1, and the other 814 from an optimal solution of the same problem outside H . In Tables 2-4 we report the 815 average and maximum relative gaps for each bound, and the average and maximum 816 computing times for $n = 5, 10, 20$, respectively. In the last line of the tables we also report the same values for the $SOC-RLT$ bound presented in [12], computed by Mosek.¹ 817 818 Note that the average gap is taken only over the instances which were not solved (in 819 the sense specified above) by a given bound. Moreover, the average computing time 820 for bound LbTwoAdj is computed only over the instances (87 overall, as we will see) 821 which are *not* solved by bound $LbTwoCut$.

822 We remark that the bound LbTwoCut is computed by adding the first cut as in 823 bound LbOneCut, i.e., the supporting hyperplane at $\bar{\mathbf{x}} \in \partial H$, and then adding a further 824 linear cut through the projection of an optimal solution outside H obtained when com-825 puting bound LbOneCut, i.e., point $\mathbf{z}_2(\lambda_{\min})$ returned by procedure DualLagrangian 826 with input $X = \Omega_{\bar{x}}$. We could as well choose the adjusted cut computed by bound 827 LbOneAdj as the first cut for bound LbTwoCut, but we observed that with this choice 828 no improvement over LbOneAdj is obtained. This is related to what already observed

TABLE 2

Average and maximum relative gaps and computing times (in seconds) for the instances with $n = 5$.

Bound	Average relative gap $(\%)$	Max relative gap $(\%)$	Average time Max time	
LbDual	0.90%	2.97%	0.013	0.015
LbOneCut	0.31%	1.27%	0.035	0.040
LbOneAdj	0.130%	0.548%	0.266	0.388
LbTwoCut	0.07%	0.21%	0.089	0.108
LbTwoAdj	0%	0%	0.146	0.281
$SOC-RLT$	0.131%	0.548%	1.435	2.080

¹The authors are very grateful to Professor Samuel Burer for providing the MATLAB code for the computation of the $\texttt{SOC-RLT}$ bound.

TABLE 3

Average and maximum relative gaps and computing times (in seconds) for the instances with $n = 10$

TABLE 4

Average and maximum relative gaps and computing times (in seconds) for the instances with $n = 20$.

 in Figure 3: bound LbOneAdj cannot be improved any more when there are (at least) 830 two optimal solutions outside H (besides the one in $int(H)$). Thus, the second cut is able to remove one of such optimal solutions but not the other, so that the bound cannot be improved. Similarly, for bound LbTwoAdj the two initial cuts are the ones computed for bound LbTwoCut.

 For what concerns the computing times, we observe that these are lower than those reported in [25] for the bound obtained by adding lifted-RLT cuts (around 836 92s for an instance with $n = 20$). They are also lower than those reported in [3] 837 for the bound obtained by adding KSOC cuts (up to 2s for $n = 20$ instances). For the sake of correctness, we point out that the computing times reported in those papers have been obtained with different processors. However, such processors have comparable performance with respect to the one employed for the computational experiments in this paper. In general, the proposed bounds are very cheap. Only 842 for two instances with $n = 20$, LbTwoAdj required times above 1s (around 1.5s in both cases). Usually the computing times are (largely) below 1s. Both the dual Lagrangian bound and the bound obtained by a single linear cut are pretty cheap but with poorer performance in terms of relative gap. The bound obtained by Algorithm 4.1 with a local adjustment of the linear cut is better than the two previous ones in terms of gap but is also more expensive (although still cheap). The bound LbTwoCut offers a good combination between quality and cheap computing time. But a more careful choice of the two linear cuts, through a local adjustment, improves the quality without compromising the computing times. This is confirmed by the results reported 851 for LbTwoAdj. Although this bound is more expensive than the others, the additional search for adjusted linear cuts further increases the quality of the bound. In Table 5 we report the number of solved instances for LbTwoCut and LbTwoAdj. According to what was reported in [3], the total number of unsolved instances out of the 212 hard

TABLE 5 Number of solved instances for the bounds LbTwoCut and LbTwoAdj.

			Bound $n = 5$ (out of 38) $n = 10$ (out of 70) $n = 20$ (out of 104)
LbTwoCut			
LbTwoAdj	38	70	103

TABLE 6 Minimum, average, and maximum $PercDiff$ values, where $PercDiff$ is defined in (6.1).

855 instances is equal to the following: 133 for the bound proposed in [25] (18 with $n = 5$, 856 49 with $n = 10$, and 66 with $n = 20$; 85 for the bound proposed in [3] (18 with $n = 5$, 857 22 with $n = 10$, and 45 with $n = 20$; 56 by considering the best bound between the 858 one in [25] and the one in [3] (10 with $n = 5$, 15 with $n = 10$, and 31 with $n = 20$). 859 For bound LbTwoCut the total number of unsolved instances reduces to 87 (24, 29, 860 and 34 for $n = 5$, $n = 10$, and $n = 20$, respectively). Finally, for bound LbTwoAdj 861 we have the remarkable outcome that there is just one unsolved instance (namely, 862 instance 20.628). For the sake of correctness, we should warn that the value UB in [3, 863 25] is not computed by running two local searches as done in this paper. It is instead 864 computed from the final solution of the relaxed problem, so that it could be slightly 865 worse and justify the larger number of unsolved instances. All the same, the quality 866 of the proposed bounds appears to be quite good.

 We still need to compare our bounds with the SOC-RLT bound (last line in Ta- bles 2-4). In terms of computing times we notice that both the average and the maximum computing times of the SOC-RLT bound are larger than those of all the pro- posed bounds. But we believe that the most interesting observation is that, in terms of average and maximum gap, the SOC-RLT bound is almost identical to the LbOneAdj bound. In order to better investigate the relation between the two bounds, in Table 6 we report the minimum, average, and maximum percentages difference between the two bounds, i.e., the quantity

(6.1)
$$
PercDiff = 100 * \frac{\text{LbOneAdj} - \text{SOC} - \text{RLT}}{|\text{LbOneAdj}|} \%
$$

 We notice that the difference is sometimes positive and sometimes negative, suggesting that none of the two bounds dominate the other. But the differences are also so small 877 (below the tolerance value under which an instance is declared to be "solved" by a given bound) that they could also be numerical differences due to the tolerance values employed in the solvers. We believe that an interesting question for future research is to establish whether these two bounds are, in fact, equivalent, which would lead to a new interpretation of the SOC-RLT bound proposed in [12].

882 6.1. Investigating the hardest instance. As a final experiment, we investi-883 gate the behavior of bound LbTwoAdj over the hardest instance with $n = 20$, the one

884 for which the relative error is above 10^{-4} . For this instance, at the last iteration we recorded the following objective function values, corresponding to values of local 886 minimizers of problem (5.1) , which certainly include the global minimizer(s) of such

888 a problem:

- 889 \bullet the value at the optimal solution of problem (5.1) belonging to $int(H)$;
- 890 \bullet the value at a globally optimal solution of the trust region problem obtained by fixing in problem (5.1) the first linear cut to an equality, in case such solution fulfills the second linear cut, or, alternatively, the value at the local and nonglobal solution of the same problem, in case such solution exists and fulfills the second linear cut (if the global minimizer does not fulfill the second linear cut and the local and nonglobal minimizer does not exist or does not fulfill the second linear cut, then the value is left undefined);
- \bullet the same value as above but after fixing the second linear cut to an equality in problem (5.1);
- 899 \bullet the value at a globally optimal solution of the trust region problem obtained by fixing both cuts to equalities in problem (5.1).

 Note that two of the four values must be equal. In particular, one of the two 902 equal values is always the first one, attained in $int(H)$. But for the hardest instance we observed that all four values are very close to each other and all of them are lower 904 than the UB value. Thus, it appears that for this instance a situation like the one displayed in Figure 5c occurs. In this case even the perturbation of both linear cuts 906 is unable to remove all of the three solutions outside H .

 7. Conclusions. In this paper we discussed the CDT problem. First, we derived some theoretical results for a class of problems which includes the CDT problem as a special case. Then, from the theory developed for such class, we have rederived a necessary and sufficient condition for the exactness of the Shor relaxation and of the equivalent dual Lagrangian bound for the CDT problem. The condition is based on the existence of multiple solutions for a Lagrangian relaxation. Based on such con- dition, we proposed to strengthen the dual Lagrangian bound by adding one or two linear cuts. These cuts are based on supporting hyperplanes of one of the two qua- dratic constraints, and they are, thus, redundant for the original CDT problem (1.1). However, the cuts are not redundant for the Lagrangian relaxation and their addition allows one to improve the bound. We ran different computational experiments over the 212 hard test instances selected from the three thousand ones randomly gener- ated in [12], reporting gaps and computing times. We have shown that the bounds are computationally cheap and are quite effective. In particular, one of them, based on the addition of two linear cuts, is able to solve all but one of the hard instances. We have also investigated more in detail such hardest instance for which the bound is not exact (though quite close to the optimal value). An interesting topic for fu- ture research could be that of establishing the relations between the bounds proposed in this work and those presented in the recent literature (in particular, as already 926 mentioned, it would be interesting to establish whether bound LbOneAdj is equivalent to the SOC-RLT bound introduced in [12]). Moreover, it would also be interesting to develop procedures which are able to generate CDT instances for which the bound LbTwoAdj is unable to return the optimal value. Finally, it would be interesting to 930 see if the results presented in this work could be extended to QP problems with more $\mathbf{A}\mathbf{Q5}$
931 than two constraints. Some preliminary studies, which will appear elsewhere, show than two constraints. Some preliminary studies, which will appear elsewhere, show that for such problems it is sometimes possible to improve the dual Lagrangian bound with the addition of a linear cut, but it may be hard to identify it and it is not even guaranteed to exist.

REFERENCES

- 935 [1] S. ADACHI, S. IWATA, Y.NAKATSUKASA, AND A. TAKEDA, Solving the trust-region subprob-
936 *lem by a generalized eigenvalue problem.* SIAM J. Optim., 27 (2017). pp. 269–291. lem by a generalized eigenvalue problem, SIAM J. Optim., 27 (2017), pp. 269-291, 937 [https://doi.org/10.1137/16M1058200.](https://doi.org/10.1137/16M1058200)
938 [2] W. AI AND S. ZHANG, *Strong duality for th*
- 938 [2] W. AI AND S. ZHANG, *Strong duality for the CDT subproblem: A necessary and sufficient con-dition*, SIAM J. Optim., 19 (2009), pp. 1735–1756, https://doi.org/10.1137/07070601X. 939 dition, SIAM J. Optim., 19 (2009), pp. 1735–1756, [https://doi.org/10.1137/07070601X.](https://doi.org/10.1137/07070601X)
940 [3] K. M. ANSTREICHER, Kronecker product constraints with an application to the two-tru
- 940 [3] K. M. ANSTREICHER, Kronecker product constraints with an application to the two-trust region subproblem. SIAM J. Optim., 27 (2017), pp. 368–378, https://doi.org/10.1137/ 941 region subproblem, SIAM J. Optim., 27 (2017), pp. 368–378, [https://doi.org/10.1137/](https://doi.org/10.1137/16M1078859)
942 16M1078859. [16M1078859.](https://doi.org/10.1137/16M1078859)
- 943 [4] K. M. ANSTREICHER, Solving two-trust-region subproblems using semidefinite optimization with 944 eigenvector branching, J. Optimi. Theory Appl., (2022), to appear.
- 945 [5] J. P. AUBIN AND H. FRANKOWSKA, Set-Valued Analysis, Modern Birkhäuser Classics, 946 Birkhäuser Boston, Boston, 2009.
- 947 [6] A. I. BARVINOK, Feasibility testing for systems of real quadratic equations, Discrete Comput. 948 Geom., 10 (1993), pp. 1–13.
949 [7] A. BECK AND Y. C. ELDAR,
- 949 [7] A. BECK AND Y. C. ELDAR, *Strong duality in nonconvex quadratic optimization with* 950 two *quadratic constraints*. SIAM J. Optim., 17 (2006), pp. 844–860, https://doi.org/ two quadratic constraints, SIAM J. Optim., 17 (2006), pp. 844-860, [https://doi.org/](https://doi.org/10.1137/050644471) 951 [10.1137/050644471.](https://doi.org/10.1137/050644471)
952 [8] C. BERGE, Topological S
- 952 [8] C. Berge, Topological Spaces: Including a Treatment of Multi-valued Functions, Vector Spaces and Convexity, Dover Publications, Mineola, New York, 1963.
- 954 [9] D. BIENSTOCK, A note on polynomial solvability of the CDT problem, SIAM J. Optimi., 26
955 (2016), pp. 488–498, https://doi.org/10.1137/15M1009871. 955 (2016), pp. 488--498, [https://doi.org/10.1137/15M1009871.](https://doi.org/10.1137/15M1009871)
- 956 [10] I. M. BOMZE AND M. L. OVERTON, Narrowing the difficulty gap for the Celis-Dennis-Tapia 957 [10] $problem$. Math. Program., 151 (2015), pp. 459–476. $problem, Math. Program., 151 (2015), pp. 459–476.$
- 958 [11] I. M. BOMZE, V. JEYAKUMAR, AND G. LI, *Extended trust-region problems with one or two balls:* 959 Exact copositive and lagrangian relaxations, J. Global Optim., 71 (2018), pp. 551–569.
960 [12] S. BURER AND K. M. ANSTREICHER, Second-oder-cone constraints for extended trust-req.
- 960 [12] S. Burer and K. M. Anstreicher, Second-oder-cone constraints for extended trust-region 961 subproblems, SIAM J. Optim., 23 (2013), pp. 432--451, [https://doi.org/10.1137/110826862.](https://doi.org/10.1137/110826862)
- 962 [13] M. R. CELIS, J. E. DENNIS, AND R. A. TAPIA, A trust region strategy for nonlinear equality constrained optimization, IN Numerical Optimization 1984, P. T. BOGGS, R. H. BYRD, constrained optimization, IN Numerical Optimization 1984, P. T. Boggs, R. H. BYRD, 964 AND R. B. SCHNABEL, eds., SIAM, Philadelphia, 1985, pp. 71-82.
- 965 [14] L. CONSOLINI AND M.LOCATELLI, On the complexity of quadratic programming with two qua-966 *dratic constraints*, Math. Program., 164 (2017), pp. 91–128.
967 [15] W. HILDENBRAND, *Core and Equilibria of a Large Economy.* (*P.*
- [15] W. HILDENBRAND, Core and Equilibria of a Large Economy. (PSME-5), Princeton University 968 Press, Princeton, NJ, 1974.
- 969 [16] J. B. HIRIART-URRUTY AND C. LEMARECHAL, Convex Analysis and Minimization Algorithms 970 [16] J. Fundamentals, Springer-Verlag, Berlin, 1993. 970 I, Fundamentals, Springer-Verlag, Berlin, 1993.
971 [17] J.-BA HIRIART-URRUTY AND C. LEMARECHAL, Con-
	- [17] J.-BA HIRIART-URRUTY AND C. LEMARECHAL, Convex Analysis and Minimization Algorithms
- 972 *I*, Springer-Verlag, 1996.
973 [18] J. M. MARTINEZ, *Local minimizers of quadratic functions on Euclidean balls and spheres*, SIAM 974 J. Optim., 4 (1994), pp. 159–176, [https://doi.org/10.1137/0804009.](https://doi.org/10.1137/0804009)
- 975 [19] T. MONTAHNER, A. NEUMAIER, AND F. DOMES, A computational study of global optimization
976 **Solvers** on two trust region subproblems, J. Global Optim., 71 (2018), pp. 915–934. solvers on two trust region subproblems, J. Global Optim., 71 (2018), pp. 915-934.
- 977 [20] J. J. Moré AND D. C. Sorensen, *Computing a trust region step*, SIAM J. Sci. Stat. Comput., 978 4 (1983), pp. 553–572, https://doi.org/10.1137/0904038. 4 (1983), pp. 553-572, [https://doi.org/10.1137/0904038.](https://doi.org/10.1137/0904038)
- 979 [21] D. C. SORENSEN, Newton's method with a model trust region modification, SIAM J. Numer. 980 Anal., 19 (1982), pp. 409–426, [https://doi.org/10.1137/0719026.](https://doi.org/10.1137/0719026)
981 [22] S. SAKAUE, Y. NAKATSUKASA, A. TAKEDA, AND S. IWATA, Solvine
- [22] S. SAKAUE, Y. NAKATSUKASA, A. TAKEDA, AND S. IWATA, Solving generalized CDT prob-982 lems via two-parameter eigenvalues, SIAM J. Optim., 26 (2016), pp. 1669--1694, 983 [https://doi.org/10.1137/15100624X.](https://doi.org/10.1137/15100624X)
984 [23] J. F. STURM AND S. ZHANG, On cones o
- [23] J. F. STURM AND S. ZHANG, On cones of nonnegative quadratic functions, Math. Oper. Res., 985 28 (2003), pp. 246–267.
986 24 J. WANG AND Y. XIA. Clos
- [24] J. WANG AND Y. XIA, Closing the gap between necessary and sufficient conditions for local 987 nonglobal minimizer of trust region subproblem, SIAM J. Optim., 30 (2020), pp. 1980– 988 1995, [https://doi.org/10.1137/19M1294459.](https://doi.org/10.1137/19M1294459)
989 [25] B. YANG AND S. BURER, A two-variable approac
- [25] B. YANG AND S. BURER, A two-variable approach to the two-trust region subproblem, SIAM J. 990 Optim., 26 (2016), pp. 661–680, [https://doi.org/10.1137/130945880.](https://doi.org/10.1137/130945880)
991 [26] Y. YE AND S. ZHANG, New results on quadratic minimization, SIAM.
- 991 [26] Y. YE AND S. ZHANG, New results on quadratic minimization, SIAM J. Optim., 14 (2003), pp. 245-267, https://doi.org/10.1137/S105262340139001X. 992 pp. 245–267, [https://doi.org/10.1137/S105262340139001X.](https://doi.org/10.1137/S105262340139001X)
993 [27] J. YUAN, M. WANG, W. AI, AND T. SHUAI, New results on
- [27] J. YUAN, M. WANG, W. AI, AND T. SHUAI, New results on narrowing the duality gap on 994 the extended Celis-Dennis-Tapia problem, SIAM J. Optim., 27 (2017), pp. 890-909, 995 [https://doi.org/10.1137/16M1080082.](https://doi.org/10.1137/16M1080082)