

BIBLIOGRAFIA

Adham N, Borden LA, Schechter LE et al.: Cell-specific coupling of the cloned human 5-HT_{1F} receptor to multiple signal transduction pathways. *Naunyn Schmiedeberg's Arch. Pharmacol.* 348:566-75, 1993

Albert PR, Zhou QY, Vantol HHM et al.: Cloning, functional expression and messenger RNA tissue distribution of the rat 5-hydroxytryptamine_{1A} receptor gene. *J. Biol. Chem.* 265:5825-32, 1990

Anderson GM, Horne WC, Chatterjee D, Cohen DJ: The hyperserotonemia of autism. *Ann. NY Acad. Sci.* 600:331-40, discussion 341-2, 1990

Andrade R, Malenka RC, Nicoll RA: A G-protein couples serotonin and GABA_B receptors to the same channels in hippocampus. *Science* 234:1261-5, 1986

Andrews PLR, Bhandari P: The 5-hydroxytryptamine receptor antagonists as antiemetics: preclinical evaluation and mechanism of action. *Eur. J. Cancer* 29A (Suppl 1):511-6, 1993

Andrews PLR, Rapeport WG, Sanger GJ: Neuropharmacology of emesis induced by anticancer therapy. *Trends Pharmacol. Sci.* 9:334-41, 1988

Anguelova M, Benkelfat C, Turecki G: A systematic review of association studies investigating genes coding for serotonin receptors and the serotonin transporter: II. Suicidal behavior. *Mol. Psychiatry* 8:646-53, 2003

Arias B, Catalan R, Gasto C, et al.: 5-HTTLPR polymorphism of the serotonin transporter gene predicts non-remission in major depression patients treated with citalopram in a 12-weeks follow up study. *J Clin Psychopharmacol*, 23:563-7, 2003

Atkinson W, Lockhart S, Whorwell PJ et al.: Altered 5-hydroxytryptamine signalling in patients with constipation- and diarrhea-predominant irritable bowel syndrome. *Gastroenterology* 130, 34-43, 2006

Azpiroz F, Enck P, Whitehead WE: Anorectal functional testing: review of collective experience. *Am. J. Gastroenterol.* 97:232-40, 2002

Baig MK, Zhao RH, Woodhouse SL et al.: Variability in serotonin and enterochromaffin cells in patients with colonic inertia and idiopathic diarrhoea as compared to normal controls. *Colorectal Dis.* 4:348-54, 2002

Baker DE: Rationale for using serotonergic agents to treat irritable bowel syndrome. *Am. J. Health-Syst. Pharm.* 62: 700-711, 2005

Bardhan KD, Bodemar G, Geldof H et al.: A double-blind, randomized, placebo-controlled dose-ranging study to evaluate the efficacy of alosetron in the treatment of irritable bowel syndrome. *Aliment. Pharmacol. Ther.* 14:23-34, 2000

Barnes JM, Barnes NM, Cooper SJ: Behavioural pharmacology of 5-HT₃ receptor ligands. *Neurosci. Biobehav. Rev.* 16:107-13, 1992

Battersby S, Ogilvie AD, Blackwood DH et al.: Presence of multiple functional polyadenylation signals and a single nucleotide polymorphism in the 3' untranslated region of the human Serotonin transporter gene. *J. Neurochem.* 72:1384-8, 1999

Bax WA, Renzenbrink GJ, Van Heuven-Nolsen D et al.: 5-HT receptors mediating contractions of the isolated human coronary artery. *Eur. J. Pharmacol.* 239:203-10, 1993

Beer MS, Stanton JA, Bevan Y et al.: An investigation of the 5-HT_{1D} receptor binding affinity of 5-hydroxytryptamine, 5-carboxamidotryptamine and sumatriptan in the central nervous system of seven species. *Eur. J. Pharmacol.* 213:193-7, 1992

Bellini M, Rappelli L, Blandizzi C et al.: Platelet serotonin transporter in patients with diarrhea-predominant irritable bowel syndrome both before and after treatment with alosetron. *Am. J. Gastroenterol.* 98:2704-9, 2003

Bellivier F, Szoke A, Henry C et al.: Possible association between serotonin transporter gene polymorphism and violent suicidal behavior in mood disorders. *Biol. Psychiatry.* 48:319-22, 2000

Bertrand PP, Kunze WAA, Furness JB, Bornstein JC: The terminals of myenteric intrinsic primary afferent neurons of the guinea-pig ileum are excited by 5-HT acting at 5-HT₃ receptors. *Neuroscience* 101:459-69, 2000

Bezprozvanny I, Scheller RH, Tsien RW: Functional impact of syntaxin on gating of N-type and Q-type calcium channels. *Nature* 378:623-6, 1995

Bharucha AE, Camilleri M, Haydock S et al.: Effects of a serotonin 5-HT₄ receptor antagonist SB-207266 on gastrointestinal motor and sensory function in humans. *Gut* 47:667-74, 2000

Billett EA, Richter MA, King N et al.: Obsessive compulsive disorder, response to serotonin reuptake inhibitors and the serotonin transporter gene. *Mol. Psychiatry* 2:403-6, 1997

Blakely RD, De Felice LJ, Hartzell HC: Molecular physiology of norepinephrine and serotonin transporters. *J. Exp. Biol.* 196:263-81, 1994

Blakely RD, Ramamoorthy S, Schroeter S et al.: Regulated phosphorylation and trafficking of antidepressant-sensitive serotonin transporter proteins. *Biol. Psychiatry* 44:169-78, 1998

Blandina P, Goldfarb J, Craddock-Royal B, Green JP: Release of endogenous dopamine by stimulation of 5-hydroxytryptamine receptors in rat striatum. *J. Pharmacol. Exp. Ther.* 251:803-9, 1989

Blauw GJ, Van Brummelen P, Van Zwieten PA: Serotonin induced vasodilatation in the human forearm is antagonised by the selective 5-HT₃ receptor antagonist ICS 205930. *Life Sci.* 43:1441-9, 1988

Blier P and de Montigny C: Possible serotonergic mechanisms underlying the antidepressant and anti-obsessive-compulsive disorder responses. *Biol. Psychiatry* 44:313-23, 1998

Bockaert J, Fozard JR, Dumuis A, Clarke DE: The 5-HT₄ receptor: a place in the sun. *Trends Pharmacol. Sci.* 13:141-5, 1992

Bradley CC, Blakely RD: Alternative splicing of the human serotonin transporter gene. *J. Neurochem.* 69:1356-67, 1997

Bueno L, De Ponti F, Fried M et al.: Serotonergic and non-serotonergic targets in the pharmacotherapy of visceral hypersensitivity. *Neurogastroenterol Motil*, 19(1 Suppl): 89-119, 2007

Burns CM, Chu H, Rueter SM et al.: Regulation of serotonin-2C receptor G-protein coupling by RNA editing. *Nature* 387: 303-8, 1997

Camilleri M, Heading RC, Thompson WG: Consensus report: clinical perspectives, mechanisms, diagnosis and management of irritable bowel syndrome. *Aliment Pharmacol Ther*, 16: 1407-1430, 2002

Caspi A, Sugden K, Moffitt TE, et al.: Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science*, 301:368–89, 2003

Catalano M: This challenges of psychopharmacogenetics. *Am. J. Hu. Genet.* 65: 606-10, 1999

Chang AS, Chang SM, Starnes DM et al.: Cloning and expression of the mouse serotonin transporter. *Mol. Brain Res.* 43:185-92, 1996

Chang AS, Lam DM: Mechanistic analyses of ion dependences in a high-affinity human serotonin transport system in transfected murine fibroblast cells. *J. Physiol.* 510:903-13, 1998

Chen JJ, Zhishan L, Pan H et al.: Maintenance of serotonin in the intestinal mucosa and ganglia of mice that lack the high affinity serotonin transporter (SERT): abnormal intestinal motility and the expression of cation transporters. *J. Neurosci.* 21:6348-61, 2001

Chen JX, Pan H, Rothman TP, Wade PR et al.: Guinea-pig 5-HT transporter: cloning, expression, distribution and function in intestinal sensory reception. *Am. J. Physiol.* 275:G433-48, 1998

Cho HJ, Meira-Lima I, Cordeiro Q, et al.: Population-based and family-based studies on the serotonin transporter gene polymorphisms and bipolar disorder: a systematic review and meta-analysis. *Mol Psychiatry*. 10(8):771-81. 2005

Coates MD, Moses PL, Mahoney OR, et al.: Serotonin signaling in the mucosa of the human colon. *Gastroenterol.* 124: A341, 2003

Coates MD, Mahoney CR, Linden DR et al.: Molecular defects in mucosal serotonin content and decreased serotonin reuptake transporter in ulcerative colitis and irritable bowel syndrome. *Gastroenterology* 126, 1657–1666, 2004

Coccaro EF, Kavoussi RJ: Fluoxetine and impulsive aggressive behavior in personality-disordered subjects. *Arch. Gen. Psychiatry* 54:1081-8, 1997

Cohen ML, Wittenauer LA: Serotonin receptor activation of phosphoinositide turnover in uterine, fundal, vascular and tracheal smooth muscle. *J. Cardiovasc. Pharmacol.* 10:176-81, 1987

Colucci R, Blandizzi C, Bellini M et al.: The genetics of the serotonin transporter and irritable bowel syndrome. *Trends Mol Med.*;14(7):295-304, 2008

Cool DR, Leibach FH, Ganapathy V: Modulation of serotonin uptake kinetics by ions and ion gradients in human placental brush-border membrane vesicles. *Biochemistry* 29:1818-22, 1990

Costall B, Kelly ME, Naylor RJ et al.: Neuroanatomical sites of action of 5-HT₃ receptor agonist and antagonists for alteration of aversive behaviour in the mouse. *Br. J. Pharmacol.* 96:325-32, 1989

Costall B, Naylor RJ: 5-Hydroxytryptamine: new receptors and novel drugs for gastrointestinal motor disorders. *Scand. J. Gastroenterol.* 25:769-87, 1990

Coto E, Reguero JR, Alvarez V et al.: 5-Hydroxytryptamine 5-HT_{2A} receptor and 5-hydroxytryptamine transporter polymorphisms in acute myocardial infarction. *Clin. Sci. (Lond).* 104:241-5, 2003

Coyle N, Jones I, Robertson E et al.: Variation at the serotonin transporter gene influences susceptibility to bipolar affective puerperal psychosis. *Lancet* 356:1490-1, 2000

Craig DA, Eglen RM, Walsh LKM, et al.: 5-Methoxytryptamine and 2-methyl-5-hydroxytryptamine-induced desensitisation as a discriminative tool for the 5-HT₃ and putative 5-HT₄ receptors in guinea-pig ileum. *Naunyn Schmiedeberg's Arch. Pharmacol.* 342:9-16, 1990

Curzon G, Kennett GA: m-CPP: a tool for studying behavioural responses associated with 5-HT_(1C) receptors. *Trends Pharmacol. Sci.* 11:181-2, 1990

De Ponti F: Pharmacology of serotonin: what a clinician should know. *Gut*; 53(10):1520-35, 2004

De Ponti F, Tonini M: Irritable bowel syndrome: new agents targeting serotonin receptor subtypes. *Drugs* 61:317-32, 2001

Delbruck SJ, Wendel B, Grunewald I et al.: A novel allelic variant of the human serotonin transporter gene regulatory polymorphism. *Cytogenet. Cell. Genet.* 79:214-20, 1997

Derkach V, Surprenant AM, North RA: 5-HT₃ receptors are membrane ion channels. *Nature (Lond.)* 339:706-9, 1989

Di Bella D, Erzegovesi S, Cavallini MC, et al. Obsessive-compulsive disorder, 5-HTTLPR polymorphism and treatment response. *Pharmacogenomics J.* 2:176-81, 2002

Drossmann DA: Do psychosocial factors define symptom severity and patient status in irritable bowel syndrome? *Am. J. Med.* 107:415-505, 1999

Duan J, Wainwright MS, Comeron JM et al.: Synonymous mutations in the human dopamine receptor D2 (DRD2) affect mRNA stability and synthesis of the receptor. *Hum. Mol. Genet.* 12:205-16, 2003

Duncan GE, Little KY, Kirkman JA et al.: Autoradiographic characterisation of [3H]imipramine and [3H]citalopram binding in rat and human brain: species differences and relationships to serotonin innervation patterns. *Brain Res.* 591:181-97, 1992

Eddahibi S, Humbert M, Fadel E et al.: Serotonin transporter overexpression is responsible for pulmonary artery smooth muscle hyperplasia in primary pulmonary hypertension. *J. Clin. Invest.* 108:1141-50, 2001

El-Salhy M, Norrgard O, Spinnell S: Abnormal colonic endocrine cells in patients with chronic idiopathic slow-transit constipation. *Scand. J. Gastroenterol.* 34:1007-11, 1999

Erlander MO, Lovenberg TW, Baron BM et al.: Two members of a distinct subfamily of 5-hydroxytryptamine receptors differentially expressed in rat brain. *Proc. Natl. Acad. Sci. USA* 90:3452-6, 1993

Fayyaz M, Lackner JM: Serotonin receptor modulators in the treatment of irritable bowel syndrome. *Ther Clin Risk Manag.* 4(1):41-8, 2008

Ferguson SS, Barak LS, Zhang J et al.: G-protein-coupled receptor regulation: role of G-protein-coupled receptor kinases and arrestins. *Can. J. Physiol. Pharmacol.* 74:1095-110, 1996

Fiskerstrand CE, Lovejoy EA, Quinn JP: An intronic polymorphic domain often associated with susceptibility to affective disorders has allele dependent differential enhancer activity in embryonic stem cells. *FEBS Lett.* 458:171-4, 1999

Flattem NL, Blakely RD: Modified structure of the human serotonin transporter promoter. *Mol. Psychiatry* 5:110-5, 2000

Flory JD, Manuck SB, Ferrell RE et al.: Neuroticism is not associated with the serotonin transporter (5-HTTLPR) polymorphism. *Mol. Psychiatry* 4:93-6, 1999

Flynn DL, Zabrowski DL, Becker DP et al.: SC-53116: the first selective agonist at the newly identified serotonin 5-HT₄ receptor subtype. *J. Med. Chem.* 35:1486-9, 1992

Foguet M, Hoyer D, Pardo LA et al.: Cloning and functional characterisation of the rat stomach fundus serotonin receptor. *Embo J.* 11:3481-7, 1992

Ford APDW, Clarke DE: The 5-HT₄ receptor. *Med. Res. Rev.* 13:633-62, 1993

Ford AC, Talley NJ, Schoenfeld PS et al.: Efficacy of Antidepressants and Psychological Therapies in Irritable Bowel Syndrome: Systematic Review and Meta-analysis. *Gut.* 2008 (in press)

Fozard JR, Gray JA: 5-HT_{1C} receptor activation: a key step in the initiation of migraine? *Trends Pharmacol. Sci.* 10:307-9, 1989

Fozard JR: Role of 5-HT₃ receptors in nociception. In 5-HT₃ Receptor Antagonists, edited by B. Jones, F.D. King, and O.J. Sanger, CRC Inc., Boca Raton, FL, pp. 241-53, 1993

Francis CY, Morris J, Whorwell PJ: The irritable bowel severity scoring system: a simple method of monitoring irritable bowel syndrome and its progress. *Aliment Pharmacol Ther* 11: 395-402 1997

Fukumoto S, Tatewaki M, Yamada T et al.: Short-chain fatty acids stimulate colonic transit via intraluminal 5-HT release in rats. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 284:R1269-76, 2003

Fumeron F, Betoulle D, Nicaud V et al.: Serotonin transporter gene polymorphism and myocardial infarction: Etude Cas-Temoins de l'Infarctus du Myocarde (ECTIM). *Circulation* 105:2943-5, 2002

Furman BL, Waton NG: 5-Hydroxytryptamine and peripheral secretory mechanisms. *In* The Peripheral Actions of 5-Hydroxytryptamine, edited by J. R. Fozard, pp. 274-300, Oxford University Press, Oxford, UK, 1989

Galligan JJ: Ligand-gated ion channels in the enteric nervous system. *Neurogastroenterol. Motil.* 14:611-23, 2002

Galzin AM, Langer SZ: Modulation of 5-HT release by presynaptic inhibitory and facilitatory 5-HT receptors in brain slices. *Adv. Biosci.* 82:59-62, 1991

Gelernter J, Kranzler H, Cubells JF: Serotonin transporter protein (SLC6A4) allele and haplotype frequencies and linkage disequilibria in African- and European-American and Japanese populations and in alcohol-dependent subjects. *Hum. Genet.* 101, 243-6, 1997

Gershon MD: Serotonin: its role and receptors in enteric neurotransmission. *Adv Exp Med Biol*, 294, 221-230, 1991

Gershon MD: Serotonin and its implication for the management of irritable bowel syndrome. *Rev Gastroenterol Disorders*, 3 (Suppl.2): S25-S34, 2003

Gershon MD, Wade PR, Florica-Howells E et al.: 5-HT_{1P} receptors in the bowel: G-protein coupling, localisation and function. *In* Serotonin: Biology, Receptors and Functional Effects, edited by J. R. Fozard and P. R. Sazena, pp. 133-143, Birkhauser Verlag, Basel, Switzerland, 1991

Gershon MD, Tack J: The serotonin signaling system: from basic understanding to drug development for functional GI disorders. *Gastroenterology*.132(1):397-414, 2007

Gilbert DG: *In smoking: Individual difference, Psychopathology, and Emotion*; Baker K.P., Seltzer H. Ends.: Taylor & Francis: Washington DC pp.149-76, 1995

Gill R.K., Pant N., Saksena S., et al.: Function, expression and characterization of the serotonin transporter (SERT) in the native human intestine. *Am. J. Physiol. Gastrointest. Liver Physiol.* 294, G254–G262, 2008

Gillman PK: Tricyclic antidepressant pharmacology and therapeutic drug interactions updated. *Br J Pharmacol*, 151(6): 737-48, 2007

Givens RC, Watkins PB: Pharmacogenetics and clinical gastroenterology *Gastroenterol.* 12:240-8, 2003

Glatt CE, Reus VI: Pharmacogenomics of monoamine transporters. *Pharmacogenomics.* 4:583-96, 2003

Goldberg PA, Kamm MA, Setti-Carraro P et al.: Modification of visceral sensitivity and pain in irritable bowel syndrome by 5-HT₃ antagonism (ondansetron). *Digestion* 57:478-83, 1996

Gore S, Gilmore IT, Haigh CG, et al.: Colonic transit in man is slowed by ondansetron (GR38032F), a selective 5-hydroxytryptamine receptor (type 3) antagonist. *Aliment. Pharmacol. Ther.* 4:139-44, 1990

Gorwood P, Batel P, Ades J, et al.: Serotonin transporter gene polymorphisms, alcoholism, and suicidal behavior. *Biol. Psychiatry* 48:259-64, 2000

Greenberg BD, Li Q, Lucas FR, et al.: Association between the serotonin transporter promoter polymorphism and personality traits in a primarily female population sample. *Am. J. Med. Genet.* 96:202-16, 2000

Grider JR, Foxx-Orenstein AE, Ji-Guang J: 5-Hydroxytryptamine₄ receptor agonists initiate the peristaltic reflex in human, rat, and guinea-pig intestine. *Gastroenterol.* 115:370-80, 1998

Grider JR, Kuemmerle JF, Jin JG: 5-HT released by mucosal stimuli initiates peristalsis by activating 5-HT₄/5-HT_{1P} receptors on sensory CGRP neurons. *Am. J. Physiol.* 270:G778-82, 1996

Gripenberg J: Inhibition of reserpine, guanethidine and imipramine of the uptake of 5-hydroxytryptamine by rat peritoneal mast cells in vitro. *Acta Physiol. Scand.* 96:407-16, 1976

Hahn MK, Blakely RD: Monoamine transporter gene structure and polymorphisms in relation to psychiatric and other complex disorders. *Pharmacogenomics J.* 2:217-35, 2002

Hamon M, Gallissot MC, Ménard F et al.: 5-HT₃ receptor binding sites are on capsaicin-sensitive fibres in the rat spinal cord. *Eur. J. Pharmacol.* 164:315-22, 1989

Hamon M, Lanfumey LEL, Mestikawi S et al.: The main features of central 5-HT₁ receptors. *Neuropsychopharmacol.* 3:349-60, 1990

Hannon J, Hoyer D: Molecular biology of 5-HT receptors. *Behavioural Brain Research*; 195: 198–213, 2008

Hansen MB: The enteric nervous system III: a target for pharmacological treatment. *Pharmacol. Toxicol.* 93:1-13, 2003

Harraf F, Schmulson M, Saba L et al.: Subtypes of constipation predominant irritable bowel syndrome based on rectal perception. *Gut* 43:388-94, 1998

Heils A, Mossner R, Lesch KP: The human serotonin transporter gene polymorphism-basic research and clinical implications. *J. Neural Transm.* 104:1005-14, 1997

Hernandez I, Sokolov BP: Abnormal expression of serotonin transporter mRNA in the frontal and temporal cortex of schizophrenics. *Mol. Psychiatry* 2:57-64, 1997

Houghton LA, Foster JM, Whorwell PJ: Alosetron, a 5-HT₃ receptor antagonist, delays colonic transit in patients with irritable bowel syndrome and healthy volunteers. *Aliment. Pharmacol. Ther.* 14:775-82, 2000

Hrdina PD, Foy B, Hepner A et al.: Antidepressant binding sites in brain: autoradiographic comparison of [3H]paroxetine and [3H]imipramine localization and relationship to serotonin transporter. *J. Pharmacol. Exp. Ther.* 252:410-8, 1990

Hu S, Brody CL, Fisher C et al.: Interaction between the serotonin transporter gene and neuroticism in cigarette smoking behavior. *Mol. Psychiatry* 5:181-8, 2000

Hu X, Oroszi G, Chun J et al.: An expanded evaluation of the relationship of four alleles to the level of response to alcohol and the alcoholism risk. *Alcohol. Clin. Exp. Res.* 29, 8–16, 2005

Humprey PPA, Hartig P, Hoyer D: A proposed new nomenclature for 5-HT receptors. *Trends Pharmacol. Sci.* 14:233-6, 1993

Ingelman-Sundberg M: Pharmacogenetics on opportunity for safer and more efficient pharmacotherapy. *Inter. Med. J.* 250:186-200, 2001

Jarrett ME, Kohen R, Cain KC et al.: Relationship of SERT polymorphisms to depressive and anxiety symptoms in irritable bowel syndrome. *Biol. Res. Nurs.* 9, 161–169, 2007

Jin H, Oksenberg D, Ashkenazi A et al.: Characterization of the human 5-hydroxytryptamine_{1B} receptor. *J. Biol. Chem.* 267:5735-8, 1992

Jin JG, Foxx-Orenstein AE, Grider JR: Propulsion in guinea-pig colon induced by 5-hydroxytryptamine via 5-HT₄ and 5-HT₃ receptors. *J. Pharmacol. Exp. Ther.* 288:93-7, 1999

Johanson JF: Options for patients with irritable bowel syndrome: contrasting traditional and novel serotonergic therapies. *Neurogastroenterol. Motil.* 16: 701-711, 2004

Jonnakuty C, Gragnoli C: What do we know about serotonin? *J Cell Physiol*, 217(2):301-6, 2008

Kaiser R, Tremblay PB, Schmider J et al.: Serotonin transporter polymorphisms: no association with response to antipsychotic treatment, but associations with the schizoparanoid and residual subtypes of schizophrenia. *Mol. Psychiatry* 6:179-85, 2001

Kannan R, Grant NJ, Aunis D, Langley K: SNAP-25 is differentially expressed by noradrenergic and adrenergic chromaffin cells. *FEBS Lett.* 385:159-64, 1996

Kaumann AJ, Murray KJ, Brown AM et al.: A receptor for 5-HT in human atrium. *Br. J. Pharmacol.* 98:664P, 1989

Kaumann AJ, Sanders L, Brown AM et al.: A 5-hydroxytryptamine receptor in human atrium. *Br. J. Pharmacol.* 100:879-85, 1990

Kerckhoffs APM, Ter Linde JJ, Akkermans LM et al.: Trypsinogen IV, serotonin transporter transcript levels and serotonin content are increased in small intestine of irritable bowel syndrome patients. *Neurogastroenterol. Motil.* 20(8):900-7, 2008

Kim DK, Lim SW, Lee S et al.: Serotonin transporter gene polymorphism and antidepressant response. *Neuroreport* 11:215-9, 2000

Kim DY, Camilleri M: Serotonin: a mediator of the brain-gut connection. *Am. J. Gastroenterol.* 95:2698-709, 2000

Kim HJ, Camilleri M, Carlson PJ et al.: Association of distinct alpha(2) adrenoceptor and serotonin transporter polymorphisms with constipation and somatic symptoms in functional gastrointestinal disorders. *Gut* 53:829-37, 2004

Kim M, Cooke HJ, Javed NH et al.: D-glucose releases 5-hydroxytryptamine from human BON cells as a model of enterochromaffin cells. *Gastroenterol.* 121:1400-6, 2001

Kirchgessner AL, Liu MT, Tamir H et al.: Identification and localisation of 5-HT_{1P} receptors in the guinea-pig pancreas. *Am. J. Physiol.* 262: G553-66, 1992

Kirov G, Rees M, Jones I et al.: Bipolar disorder and the serotonin transporter gene: a family-based association study. *Psychol. Med.* 29:1249-54, 1999

Kozlowski CM, Green A, Grundy D et al.: The 5-HT(3) receptor antagonist alosetron inhibits the colorectal distention induced depressor response and spinal c-fos expression in the anaesthetised rat. *Gut* 46:474-80, 2000

Kraft JB, Slager SL, McGrath PJ et al.: Sequence analysis of the serotonin transporter and associations with antidepressant response. *Biol. Psychiatry* 58:374-381, 2005

Kunze WA, Furness JB: The enteric nervous system and regulation of intestinal motility. *Ann. Rev. Physiol.* 61:117-42, 1999

Kursar JD, Nelson DL, Wainscott DB et al.: Molecular cloning, functional expression and pharmacological characterisation of a novel serotonin receptor (5-hydroxytryptamine_{2F}) from rat stomach fundus. *Mol. Pharmacol.* 42:549-57, 1992

Laporte AM, Kidd EJ, Verge D et al.: Autoradiographic mapping of central 5-HT₃ receptors. *In* Centrale and Peripheral 5-HT₃ Receptors, edited by M.Hamon pp.157-87, Accademic Press, London UK, 1992

Leabman MK, Huang CC, DeYoung J et al.: Pharmacogenetics of membrane transporters investigators. Natural variation in human membrane transporter genes reveals evolutionary and functional constraints. *Proc. Natl. Acad. Sci. USA* 100:5896-901, 2003

Lee DY, Park H, Kim WH et al.: Serotonin transporter gene polymorphism in healthy adults and patients with irritable bowel syndrome. *Korean J. Gastroenterol.* 43:18-22, 2004

Lefebvre H, Contesse V, Delarue C et al.: Serotonin-induced stimulation of cortisol secretion from human adrenocortical tissue is mediated through activation of a serotonin₄ receptor subtype. *Neuroscience* 47:999-1007, 1992

Lesch KP, Balling U, Gross J et al.: Organisation of the human serotonin transporter gene. *J. Neural Transm. Gen. Sect.* 95:157-62, 1994

Lesch KP, Bengel D, Heils A et al.: Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science* 274:1527-31, 1996

Lesch KP, Jatzke S, Meyer J et al.: Mosaicism for a serotonin transporter gene promoter-associated deletion: Decreased recombination in depression. *J. Neural. Transm.* 106:1223-30, 1999

Lesch KP, Moessner R: Genetically driven variation in serotonin uptake: Is there a link to affective spectrum, neurodevelopmental, and neurodegenerative disorders? *Biol. Psychiatry* 44:179-92, 1998

Lesch KP, Wolozin BL, Murphy DL et al. Primary structure of the human platelet serotonin uptake site: identity with the brain serotonin transporter. *J. Neurochem.* 60:2319-22, 1993

Levy FO, Gudermann T, Peres-Reyes E et al.: Molecular cloning of a human serotonin receptor (S12) with a pharmacological profile resembling that of the 5-HT_{1D} subtype. *J. Biol. Chem.* 267:7553-62, 1992 a

Levy FO, Gudermann T, Birbaumer M et al.: Molecular cloning of a human gene (S31) encoding a novel serotonin receptor mediating inhibition of adenylyl cyclase. *FEBS Lett.* 296:201-6, 1992 b

Li Y, Nie Y, Xie J et al.: The Association of Serotonin Transporter Genetic Polymorphisms and Irritable Bowel Syndrome and Its Influence on Tegaserod Treatment in Chinese Patients. *Dig Dis Sci* .52: 2942–2949, 2007

Lim JE, Papp A, Pinsonneault J et al.: Allelic expression of serotonin transporter (SERT) mRNA in human pons: lack of correlation with the polymorphism SERTLPR. *Mol. Psychiatry* 11: 649–662, 2006

Linden DR, Foley KF, McQuoid C et al.: Serotonin transporter function and expression are reduced in mice with TNBS-induced colitis. *Neurogastroenterol. Motil.* 17: 565–574, 2005

Longstreth GF, Thompson WG, Chey WD et al.: Functional bowel disorders. *Gastroenterology*, 130(5):1480–91, 2006

Lotrich FE, Pollock BG, Ferrell RE: Polymorphism of the serotonin transporter: implications for the use of selective serotonin reuptake inhibitors. *Am. J. Pharmacogenomics.* 1:153-64, 2001

Lowther S, De Paermentier F, Rufus Crompton M et al.: The distribution of 5-HT_(1D) and 5-HT_(1E) binding sites in human brain. *Eur. J. Pharmacol.* 222:137-42, 1992

Lucki I: 5-HT₁ receptors and behaviour. *Neurosci. Biobehav. Rev.* 16:83-93, 1992

MacKenzie A, Quinn J: A serotonin transporter gene intron 2 polymorphic region, correlated with affective disorders, has allele-dependent differential enhancer-like properties in the mouse embryo. *Proc. Natl. Acad. Sci. USA* 96:15251-5, 1999

Maestrini E, Lai C, Marlow A et al.: Serotonin transporter (5-HTT) and gamma-aminobutyric acid receptor subunit beta3 (GABA β 3) gene polymorphisms are not associated with autism in the IMGSA families. The International Molecular Genetic Study of Autism Consortium. *Am J Med Genet.* 88:492-6, 1999

Manji HK, Quiroz JA, Spom J et al.: Enhancing neuronal plasticity and cellular resilience to develop novel, improved therapeutics for difficult-to-treat depression. *Biol. Psychiatry* 53:707-42, 2003

Marcinkiewicz M, Vergé D, Gozlan H et al.: Autoradiographic evidence for the heterogeneity of 5-HT₁ sites in the rat brain. *Brain Res.* 291:159-63, 1984

Matthes M, Boschert U, Amlaiky N et al.: Mouse 5-hydroxytryptamine_{5A} and 5-hydroxytryptamine_{5B} receptors define a new family of serotonin receptors: cloning, functional expression and chromosomal localisation. *Mol. Pharmacol* 43:313-9, 1993

Mawe GM, Coates MD, Moses PL: Review article: intestinal serotonin signaling in irritable bowel syndrome. *Aliment Pharmacol Ther*; 23, 1067-76, 2006

McAllister G, Charlesworth A, Snodin C et al.: Molecular cloning of a serotonin receptor from human brain (5HT1E): a fifth 5HT1-like subtype. *Proc. Natl. Acad. Sci. USA* 89:5517-21, 1992

McDougle CJ, Epperson CN, Price LH et al.: Evidence for linkage disequilibrium between serotonin transporter protein gene (SLC6A4) and obsessive compulsive disorder. *Mol. Psychiatry* 3:270-3, 1998

Melikian HE, Ramamoorthy S, Tate CG et al.: Inability to N-glycosylate the human norepinephrine transporter reduces protein stability, surface trafficking, and transport activity but not ligand recognition. *Mol. Pharmacol.* 50:266-76, 1996

Meltzer CC, Smith G, DeKosky ST et al.: Serotonin in aging, late-life depression, and Alzheimer's disease: the emerging role of functional imaging. *Neuropsychopharmacol.* 18:407-30, 1998

Meneses A: 5-HT system and cognition. *Neurosci. Biobehav. Rev.* 23:1111-25, 1999

Merz H: Role of the brain and sensory pathways in gastrointestinal sensory disorders in humans. *Gut* 51:129-33, 2002

Michaelovsky E, Frisch A, Rockah R et al.: A novel allele in the promoter region of the human serotonin transporter gene. *Mol. Psychiatry* 4:97-9, 1999

Miwa J, Echizen H, Matsueda K, Umeda N: Patients with constipation-predominant irritable bowel syndrome (IBS) may have elevated serotonin concentrations in colonic mucosa as compared with diarrhea-predominant patients and subjects with normal bowel habits. *Digestion* 63:188-94, 2001

Mohammad-Zadeh LF, Moses L, Gwaltney-Brant SM: Serotonin: a review. *J Vet Pharmacol Ther*, 31(3):187-99, 2008

Molderings GJ, Werner K, Likungu J, Gothert M: Inhibition of noradrenaline release from the sympathetic nerves of the human saphenous vein via presynaptic 5-HT receptor similar to the 5-HT_{1D} subtype. *Naunyn Schmiedebergs Arch. Pharmacol.* 342:371-7, 1990

Monferini E, Gaetani P, Rodriguez Y et al.: Pharmacological characterisation of the 5-hydroxytryptamine receptor coupled to adenylyl cyclase stimulation in human brain. *Life Sci.* 52:61-5, 1993

Monro RL, Bertrand PP, Bornstein JC: ATP and 5-HT are the principal neurotransmitters in the descending excitatory reflex pathway of the guinea-pig ileum. *Neurogastroenterol. Motil.* 14:255-64, 2002

Mortensen OV, Thomassen M, Larsen MB et al.: Functional analysis of a novel human serotonin transporter gene promoter in immortalized raphe cells. *Mol. Brain Res.* 68, 141-148, 1999

Murphy DL, Lerner A, Rudnick G, Lesch KP: Serotonin transporter: gene, genetic disorders and pharmacogenetics. *Mol. Interventions* 4:109-123, 2004

Murphy DL, Fox MA, Timpano KR et al.: How the serotonin story is being rewritten by new gene-based discoveries principally related to SLC6A4, the serotonin transporter gene, which functions to influence all cellular serotonin systems. *Neuropharmacology*, 55(6):932-60, 2008

Mylecharane EJ: Agonists and antagonists of 5-HT₂ receptors. *In* Cardiovascular Pharmacology of 5-Hydroxytryptamine, edited by P. R. Saxena, D. I. Wallis, W. Wouters, and P. Bevan, pp. 81-100, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1990

Nakamura M, Ueno S, Sano A, Tanabe H: The human serotonin transporter gene linked polymorphism (5-HTTLPR) shows ten novel allelic variants. *Mol. Psychiatry* 5:32-8, 2000

Nakamura T, Matsushita S, Nishiguchi N et al.: Association of a polymorphism of the 5HT2A receptor gene promoter region with alcohol dependence. *Mol. Psychiatry* 4:85-8, 1999

Naren AP, Nelson DJ, Xie W et al.: Regulation of CFTR chloride channels by syntaxin and Munc18 isoforms. *Nature* 390:302-5, 1997

Naylor L, Dean B, Opeskin K et al.: Changes in the serotonin transporter in the hippocampus of subjects with schizophrenia identified using [3H]paroxetine. *J. Neural Transm.* 103:749-57, 1996

Nichols DE, Nichols CD: Serotonin receptors. *Chem Rev*, 108: 1614-1641, 2008

Ogilvie AD, Battersby S, Bubb VJ et al.: Polymorphism in serotonin transporter gene associated with susceptibility to major depression. *Lancet* 347:731-3, 1996

Olivier B, Soudijn W, Van Wijngaarden I: Serotonin, dopamine and norepinephrine transporters in the central nervous system and their inhibitors. *Prog Drug Res.* 54:59-119, 2000

Orwin JM, Fozard JR: Blockade of the flare response to intradermal 5-hydroxytryptamine in man by MDL 72222, a selective antagonist at neuronal 5-hydroxytryptamine receptors. *Eur. J. Clin. Pharmacol.* 30:209-12, 1986

Ozsarac N, Santha E, Hoffman BJ: Alternative non-coding exons support serotonin transporter mRNA expression in the brain and gut. *J. Neurochem.* 82:336-44, 2002

Pae CU, Masand PS, Ajwani N et al.: Irritable bowel syndrome in psychiatric perspectives: a comprehensive review. *Int. J. Clin. Pract.* 61, 1708-1718, 2007

Palacios JM, Waeber C, Mengod G, Pompeiano M: Molecular neuroanatomy of 5-HT receptors. *In* Serotonin: Molecular Biology, Receptors and Functional Effects, edited by J.R. Fozard and P.R. Saxena, pp. 5-20, Birkhauser Verlag, Basel, Switzerland, 1991

Pan H, Galligan JJ: 5-HT_{1A} and 5-HT₄ receptors mediate inhibition and facilitation of fast synaptic transmission in enteric neurons. *Am. J. Physiol.* 266:G230-8, 1994

Pan H, Gershon MD: Activation of intrinsic afferent pathways in submucosal ganglia of the guinea-pig small intestine. *J. Neurosci.* 20:3295-309, 2000

Park MJ, Choi MG, Park JA et al.: Serotonin transporter gene polymorphism and irritable bowel syndrome. *Neurogastroenterol Motil.* 18, 995-1000, 2006

Pata C, Erdal ME, Derici E et al.: Serotonin transporter gene polymorphism in irritable bowel syndrome. *Am. J. Gastroenterol.* 97:1780-4, 2002

Paudice P, Raiteri M: Cholecystokinin release mediated by 5-HT₃ receptors in rat cerebral cortex and nucleus accumbens. *Br. J. Pharmacol.* 103:1790-4, 1991

Pazos A, Palacios JM: Quantitative autoradiographic mapping of serotonin receptors in the rat brain. I. Serotonin-1 receptors. *Brain Res.* 346:205-30, 1985

Pazos A, Probst A, Palacios JM: Serotonin receptors in the human brain. III. Autoradiographic mapping of serotonin-1 receptors. *Neuroscience* 21:97-122, 1987

Perlis RH, Mischoulon D, Smoller JW et al.: Serotonin transporter polymorphisms and adverse effects with fluoxetine treatment. *Biol. Psychiatry* 54:879-83, 2003

Persico AM, Militerni R, Bravaccio C et al.: Lack of association between serotonin transporter gene promoter variants and autistic disorder in two ethnically distinct samples. *Am. J. Med. Genet.* 96:123-7, 2000

Peters JA, Malone HM, Lambert JJ: Characterisation of 5-HT₃ receptor-mediated electrical responses in nodose ganglion neurones and clonal neuroblastoma cells maintained in culture. *In* Serotonin: Molecular Biology, Receptors and Functional Effects, edited by J.R. Fozard and P.R. Sazena, pp. 84-94, Birkhauser Verlag, Basel, Switzerland, 1991

Plassat JL, Boschert U, Amlaiky N, Hen R: The mouse 5HT₅ receptor reveals a remarkable heterogeneity within the 5HT_{1D} receptor family. *Embo J.* 11:4779-86, 1992

Pollock BG, Ferrell RE, Mulsant BH et al.: Allelic variation in the serotonin transporter promoter affects onset of paroxetine treatment response in late-life depression. *Neuropsychopharmacology.* 23: 587-90, 2000

Pratt GD, Bowery NG, Kilpatrick GJ et al.: Consensus meeting agrees distribution of 5-HT₃ receptors in mammalian hindbrain. *Trends Pharmacol. Sci.* 11:135-7, 1990

Prins NH, Briejer MR, Van Bergen PJ, et al.: Evidence for 5-HT₇ receptors mediating relaxation of human colonic circular smooth muscle. *Br J Pharmacol.*; 128(4): 849-52, 1999

Prior A, Maxton DG, Whorwell PJ: Anorectal manometry in irritable bowel syndrome: differences between diarrhoea and constipation predominant subjects. *Gut* 31:458-62, 1990

Qian Y, Melikian HE, Rye DB et al.: Identification and characterization of antidepressant-sensitive serotonin transporter proteins using site-specific antibodies. *J. Neurosci.* 15:1261-74, 1995

Quick MW: Regulating the conducting states of a mammalian serotonin transporter. *Neuron* ,40(3):537-49, 2003

Racke K, Reimann A, Schworer H, Kilbinger H: Regulation of 5-HT release from enterochromaffin cells. *Behav. Brain Res.* 73:83-7, 1996

Radja F, Laporte AM, Daval G et al.: Autoradiography of serotonin receptor subtypes in the central nervous system. *Neurochem. Int.* 18:1-15, 1991

Ragnarsson G, Bodemar G: Division of the irritable bowel syndrome into subgroups on the basis of daily recorded symptoms in two outpatients samples. *Scand. J. Gastroenterol.* 34:993-1000, 1999

Ramamoorthy S, Bauman AL, Moore KR et al.: Antidepressant and cocaine sensitive human serotonin transporter: molecular cloning, expression, and chromosomal localization. *Proc. Natl. Acad. Sci. USA.* 90:2542-6, 1993 a

Ramamoorthy S, Cool DR, Mahesh VB et al.: Regulation of the human serotonin transporter. Cholera toxin-induced stimulation of serotonin uptake in human placental choriocarcinoma cells is accompanied by increased serotonin transporter mRNA levels and serotonin transporter-specific ligand binding. *J. Biol. Chem.* 268:21626-31, 1993 b

Raybould HE, Glatzle J, Robin C et al.: Expression of 5-HT₃ receptors by extrinsic duodenal afferents contribute to intestinal inhibition of gastric emptying. *Am. J. Physiol. Gastrointest. Liver Physiol.* 284:G367-72, 2003

Risch NJ: Searching for genetic determinants in the new millennium. *Nature* 405:847-56, 2000

Robinson MB: Regulated trafficking of neurotransmitter transporters: common notes but different melodies. *J. Neurochem.* 80:1-11, 2002

Roses AD: "Pharmacogenetics and the practice of medicine" *Nature* 405:857-65, 2000

Routledge C, Middlemiss DN: The 5-HT hypothesis of depression revisited. *Mol. Psychiatry* 1:437, 1996

Ruat M, Traiffort E, Arrang JM et al.: A novel rat serotonin (5-HT₅) receptor molecular cloning, localisation and stimulation of cAMP accumulation. *Biochem. Biophys. Res. Commun.* 193:268-76, 1993

Rudnick G: Mechanisms of biogenic amine neurotransmitter transporters in "Neurotransmitters transporters – structure, function and regulation"; Reith E.A. ed. Humana Press, New Jersey. 73-100, 1997

Rudnick G, Clark J: From synapse to vesicle: the reuptake and storage of biogenic amine neurotransmitters. *Biochim. Biophys. Acta.* 1144:249-63, 1993

Rudnick G: Serotonin transporters-Structure and function. *J Membrane Biol* 213, 101-110, 2006

Rudnick G: Active transport of 5-hydroxytryptamine by plasma membrane vesicles isolated from human blood platelets. *J. Biol. Chem.* 252:2170-4, 1977

Saito YA, Locke GR, Zimmerman JM et al.: A genetic association study of 5-HTT LPR and GNb3 C825T polymorphisms with irritable bowel syndrome. *Neurogastroenterol Motil.* 19: 465–470, 2007

Saito YA: Genes and irritable bowel syndrome: is there a link? *Curr Gastroenterol Rep*; 10(4):355-62. 2008

Sakai K, Nakamura M, Ueno S et al.: The silencer activity of the novel human serotonin transporter linked polymorphic regions. *Neurosci. Lett.* 327, 13–16, 2002

Sauer WH, Berlin JA, Kimmel SE: Effect of antidepressants and their relative affinity for the serotonin transporter on the risk of myocardial infarction. *Circulation* 108:32-6, 2003

Saxena PR, Villalòn CM: 5-Hydroxytryptamine: a chameleon in the heart. *Trends Pharmacol. Sci.* 12:223-7, 1991

Schikowski A, Thewissen M, Mathis C et al.: Serotonin type-4 receptors modulate the sensitivity of intramural mechanoreceptive afferents of the cat rectum. *Neurogastroenterol. Motil.* 14:221-7, 2002

Schmuck K, Lubbert C, Engels P, Lubbert H: Cloning and functional characterization of the human 5-HT_{2B} serotonin receptor. *FEBS Lett.* 342:85-90, 1994

Schoemaker RG, Du XY, Bax WA, Saxena PR: 5-Hydroxytryptamine stimulates human isolated atrium but not ventricle. *Eur. J. Pharmacol.* 239:103-5, 1993

Seeger G, Schloss P, Schmidt MH: Marker gene polymorphisms in hyperkinetic disorder-predictors of clinical response to treatment with methylphenidate? *Neurosci. Lett.* 313:45-8, 2001

Serretti A, Lilli A, Smeraldi E: Pharmacogenetics in affective disorders. *Eur. J. Pharmacol.* 438:117-28, 2002a

Serretti A, Lilli R, Lorenzi C et al.: Serotonin transporter gene (5-HTTLPR) and major psychoses. *Mol. Psychiatry* 7:95-9, 2002b

Shih JC: Molecular basis of human MAO-A and B. *Neuropsychopharmacol.* 4:1-7, 1991

Shioe K, Ichimiya T, Suhara T et al.: No association between genotype of the promoter region of serotonin transporter gene and serotonin transporter binding in human brain measured by PET. *Synapse* 48, 184–188, 2003

Simrén M, Abrahamsson H, Bjornsson ES: An exaggerated sensory component of the gastrocolonic response in patients with irritable bowel syndrome. *Gut* 48:20-7, 2001

Slater BJ, Plusa SM, Smith AN, Varma JS: Rectal hypersensitivity in the irritable bowel syndrome. *Int. J. Colorectal Dis.* 12:29-32, 1997

Spiller R: Serotonin and GI clinical disorders. *Neuropharmacology.* 55(6):1072-80. 2008

Spiller RC, Jenkins D, Thornley JP et al.: Increased rectal mucosal enteroendocrine cells, T lymphocytes, and increased gut permeability following acute Campylobacter enteritis and in post-dysenteric irritable bowel syndrome. *Gut* 47:804-11, 2000

Steadman CJ, Talley NJ, Phillips SF, Zinsmeister AR: Selective 5-hydroxytryptamine type 3 receptor antagonism with ondansetron as treatment for diarrhea-predominant irritable bowel syndrome: a pilot study. *Mayo Clin. Proc.* 67:732-8, 1992

Steens J, Van Der Schaar PJ, Penning C, Brussee J, Masclee AA: Compliance, tone and sensitivity of the rectum in different subtypes of irritable bowel syndrome. *Neurogastroenterol. Motil.* 14:241-7, 2002

Steiner JA, Carneiro AM, Blakely RD: Going with the flow: trafficking-dependent and -independent regulation of serotonin transport. *Traffic*, 9(9):1393-402, 2008

Stober G, Jatzke S, Heils A et al.: Susceptibility for schizophrenia is not influenced by a functional insertion/deletion variant in the promoter of the serotonin transporter gene. *Eur. Arch. Psychiatry Clin. Neurosci.* 248:82-6, 1998

Sudhof TC: The synaptic vesicle cycle: a cascade of protein-protein interactions. *Nature* 375:645-53, 1995

Takaki M, Branchek T, Tamir H, Gershon MD: Specific antagonism of enteric neural serotonin receptors by dipeptides of 5-hydroxytryptophan: evidence that serotonin is a mediatory of slow synaptic excitation in the myenteric plexus. *J. Neurosci.* 5:1769-80, 1985

Takeshi I, Ichiro K, Itsuhiro T: Serotonin transporters. *Current Drug Targets-CNS & Neurological Disorders.* 1:519-29, 2002

Tam FSF, Hiller K, Bunce K: Investigation of the 5-HT receptor type involved in inhibition of spontaneous activity of human colonic circular muscle. *Br. J. Pharmacol.* 106:25P, 1992

Tang WJ and Gilman AG: Type-specific regulation of adenylyl cyclase by G-protein $\beta\gamma$ subunits. *Science* (Washington DC) 254:1500-3, 1991

Taylor CB: Treatment of anxiety disorders. In. *American Psychiatric Press Textbook of Psychopharmacology*, 2nd ed. Washington DC, American Psychiatric Press 775-89, 1998

Tonini M, Vicini R, Cervio E et al.: 5-HT₇ receptors modulate peristalsis and accommodation in the guinea pig ileum. *Gastroenterology*, 129(5):1557-66, 2005

Tsai SJ, Hong CJ, Yu YW et al.: Association study of a functional serotonin transporter gene polymorphism with schizophrenia, psychopathology and clozapine response. *Schizophr. Res.* 44:177-81, 2000

Turconi M, Schiantarielli P, Borsini F et al.: Azabicycloalkyl benzimidazolones: interaction with serotonergic 5-HT₃ and 5-HT₄ receptors and potential therapeutic implications. *Drugs Future* 16:1011-26, 1991

Van Kerkhoven LAS., Laheij RJF. and Jansen J.B.M.J.: Meta-analysis: a functional polymorphism in the gene encoding for activity of the serotonin

transporter protein is not associated with the irritable bowel syndrome *Aliment Pharmacol Ther*; 26: 979–986, 2007

Vane JR: The relative activities of some tryptamine analogues on the isolated rat stomach strip preparation. *Br. J. Pharmacol.* 14:87-98, 1959

Verkes RJ, Van der Mast RC, Hengeveld MW et al.: Reduction by paroxetine of suicidal behavior in patients with repeated suicide attempts but not major depression. *Am. J. Psychiatry.* 155:543-7, 1998

Videlock EJ, Chang L: Irritable bowel syndrome: current approach to symptoms, evaluation, and treatment. *Gastroenterol Clin North Am.* 36(3):665-85, 2007

Viramontes BE, Camilleri M, McKinzie S et al.: Gender-related differences in slowing colonic transit by a 5-HT₃ antagonist in subjects with diarrhea-predominant irritable bowel syndrome. *Am. J. Gastroenterol.* 96: 2671-6, 2001

Wade PR, Chen J, Jaffe B et al.: Localisation and function of a 5-HT transporter in crypt epithelia of the gastrointestinal tract. *J. Neurosci.* 16:2352-64, 1996

Waeber C, Hoyer D, Palacios JM: 5-HT₃ receptors in the human brain: autoradiographic visualisation. *Neuroscience* 31:393-400, 1989

Waeber C, Schoeffer P, Hoyer D, Palacios JM: The serotonin 5-HT_{1D} receptor: a progress review. *Neurochem. Res.* 15:567-82, 1990

Wallis DI, Elliott P: The electrophysiology of 5-HT. *In Serotonin: Molecular Biology, Receptors and Functional Effects*, edited by J.R. Fozard and P.R. Saxena, pp. 203-219, Birkhauser Verlag, Basel, Switzerland, 1991

Wallis DI: Interaction of 5-hydroxytryptamine with autonomic and sensory neurones. *In The Peripheral Actions of 5-Hydroxytryptamine*, edited by J.R. Fozard, pp. 220-246, Oxford University Press, Oxford, UK, 1989

Wang L, Erlandsen H, Haavik J et al.: Three-dimensional structure of human tryptophan hydroxylase and its implications for the biosynthesis of the neurotransmitters serotonin and melatonin. *Biochemistry.* 41:12569-74, 2002

Wang BM et al.: Serotonin transporter gene polymorphism in irritable bowel syndrome. *Zhonghua Nei Ke Za Zhi.* 43: 439–441, 2004

Weinshank RL, Zgombick JM, Macchi MJ et al.: Human serotonin 1D receptor is encoded by a subfamily of two distinct genes: 5-HT1D α and 5-HT1D β . *Proc. Natl. Acad. Sci. USA* 89:3630-4, 1992

Wendland JR et al.: Simultaneous genotyping of four functional loci of human SLC6A4, with a reappraisal of 5-HTTLPR and rs25531. *Mol. Psychiatry* 11, 224-226, 2006

Whitehead WE, Engel BT, Schuster MM: Irritable bowel syndrome: physiological and psychological differences between diarrhea-predominant and constipation-predominant patients. *Dig. Dis. Sci.* 25:404-13, 1980

Yagaloff KA, Hartig PR: [¹²⁵]Lysergic acid diethylamide binds to a novel serotonin site on rat choroid plexus epithelial cells. *J. Neurosci.* 5:3178-83, 1985

Yakel JL, Jackson MB: 5-HT₃ receptors mediate rapid responses in cultured hippocampus and a donal cell line. *Neuron.* 1:615-21, 1988.

Yeo A, Boyd P, Lumsden S et al.: Association between a functional polymorphism in the serotonin transporter gene and diarrhoea predominant irritable bowel syndrome in women. *Gut* 53:1452-8, 2004

Yirmiya N, Pilowsky T, Nemanov L et al.: Evidence for an association with the serotonin transporter promoter region polymorphism and autism. *Am. J. Med. Genet.* 105:381-6, 2001

Yoshida K, Ito K, Sato K et al.: Influence of the serotonin transporter gene-linked polymorphic region on the antidepressant response to fluvoxamine in Japanese depressed patients. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 26:383-6, 2002

Yu PL, Fujimura M, Hayashi N et al.: Mechanisms in regulating the release of serotonin from the perfused rat stomach. *Am J Physiol Gastrointest. Liver Physiol.* 280:G1099-105, 2001

Zhao R, Baig MK, Wexner SD et al.: Enterochromaffin and serotonin cells are abnormal for patients with colonic inertia. *Dis. Colon Rectum* 43:858-63, 2000

Zhou FC, Tao-Cheng JH, Segu L et al.: Serotonin transporters are located on the axons beyond the synaptic junctions: anatomical and functional evidence. *Brain Res.* 805:241-54, 1998

Zhou SF, Di YM, Chan E et al.: Clinical pharmacogenetics and potential application in personalized medicine. *Curr Drug Metab.*; 9(8):738-84, 2008