



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Forensic application of two short tandem repeat systems: HUMTH01 and HUMvWA31

This is the peer reviewed version of the following article:

Original

Forensic application of two short tandem repeat systems: HUMTH01 and HUMvWA31 / Tagliabracci, A.; Buscemi, L.; Cucurachi, Nicola; Ferrara, S. D.. - STAMPA. - XLIV:(1994), pp. 96-98. (Intervento presentato al convegno XVIth Congress of the International Academy of Legal Medicine tenutosi a Strasbourg, France nel 31 may - 2 June 1994).

Availability:

This version is available at: 11381/2786072 since: 2015-02-24T16:44:27Z

Publisher:

Springer

Published

DOI:

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

24 April 2024

**ACTA
MEDICINÆ
LEGALIS**

Volume XLIV

Proceedings of the
XVIth Congress of the
International Academy
of Legal Medicine



Springer

FORENSIC APPLICATION OF TWO SHORT TANDEM REPEAT SYSTEMS: HUMTH01 AND HUMvWA31.

*A. Tagliabracci, L. Buscemi, N. Cucurachi and S.D. Ferrara

*Istituto di Medicina Legale e delle Assicurazioni, Università di Ancona, Italy.

Istituto di Medicina Legale e delle Assicurazioni, Università di Parma, Italy.

INTRODUCTION

Over the past few years many laboratories worldwide have introduced analysis of DNA polymorphisms as a routine method for forensic purposes. In particular, VNTRs, investigated by PCR, have reached an important position after considerable development. At present laboratories are studying a further series of highly polymorphic loci whose polymorphism derives from a dimeric-trimeric-tetrameric tandem repeated core sequence, named short tandem repeats (STRs) (Weber and May 1989; Edwards et al. 1991). The aim of the present study was to investigate two of these STRs, HUMTH01 (Edwards et al. 1992) and HUMvWA31 (Kimpton et al. 1992) to evaluate: 1) allele frequencies in an Italian population sample, and 2) their efficiency for personal identification on aged bloodstains from various substrates.

MATERIALS AND METHODS

Stain preparation : a series of stains prepared from 3, 6, 10 and 20 μ l of whole fresh blood obtained from six subjects was collected on paper, white cotton and glass and were stored for 3 years at room temperature.

DNA extraction : extracted from the fresh blood of unrelated healthy subjects (HUMTH01 n= 288; HUMvWA31 n=211) and from 72 bloodstains using phenol-chloroform (Budowle and Baetchel 1990) and chelex (Walsh et al. 1991) methods.

PCR amplification : according to Brinkmann et al. (1993a) for HUMTH01 and Wiegand et al. (1993) for HUMvWA31 with the primers proposed by Edwards et al. (1991) for HUMTH01 and by Kimpton et al. (1992) for HUMvWA31.

Electrophoresis : separation by high-resolution PAGE (Budowle et al. 1991) for HUMTH01 and denaturing PAGE for HUMvWA31 using a discontinuous buffer (Allen et al. 1989). Bands were visualised by silver staining (Budowle et al. 1991) and phenotyped by side-to-side comparison with an allelic ladder (Fig. 1 a,b,c).




Fig. 1 (a,b,c) Amplified DNA fragments in HUMTH01 and HUMvWA31 systems using polyacrylamide gels and silver staining.

a) HUMTH01: from left to right: allele ladder is in lanes 1,5,6; HUMTH01 phenotypes: lane 2: 8-9,3; lane 3: 6-7; lane 4: 8-10; lane 7: 6-9; lane 8: 8-9,3.

b) HUMvWA31: from left to right: allele ladder is in lanes 1,6; HUMvWA31 phenotypes: lane 2: 17-18; lane 3: 15-19; lane 4: 11-17; lane 5: 14-16; lane 7: 18-20; lane 8: 17-19.

c) HUMTH01: from left to right: allele ladder is in lanes 1,4,7; lanes 2 and 3: 6-9,3 phenotypes from 3 and 20 μ l aged bloodstains; lanes 5, 6: 8-9,3 phenotypes from 3 and 20 μ l aged bloodstains.

RESULTS AND DISCUSSION

Six alleles for HUMTH01 (183-207 bp) and 7 for HUMvWA31 (138-162 bp) were observed. A new allele located on the anodic side of allele HUMvWA31*14 was found (temporarily designated as HUMvWA*11). The frequency distribution of HUMTH01 and HUMvWA31 alleles is shown in figure 2 a,b. Observed and expected

genotypes were compared by assuming the Hardy-Weinberg law (HUMTH01 $\chi^2 = 9.40$ $0.80 < P < 0.90$; 15

d.f. - HUMvWA31 $\chi^2 = 22.166$ $0.75 < P < 0.90$; 28 d.f.). The power of discrimination (PD) and chance of exclusion were, respectively, 0.92 and 0.57 for HUMTH01 and 0.93 and 0.61 for HUMvWA31, suggesting that these STR systems may be a powerful tool in forensics. Table 1 summarizes the results expressed as percentages of amplification and typing from aliquoted bloodstains on various substrates, using two extraction methods (phenol-chloroform vs chelex). The results show that: a) 3 μ l are sufficient to obtain positive amplification; b) paper and glass are better substrates than white cotton; c) in our bloodstain storage conditions, phenol-isoamyl alcohol was better than the chelex method.




Fig. 2 (a,b) Frequency distribution of HUMTH01 (a) and HUMvWA31(b) system alleles in an Italian population sample.

SIZE	PAPER		WHITE COTTON		GLASS	
	HUMTH01	HUMvWA31	HUMTH01	HUMvWA31	HUMTH01	HUMvWA31
	a % b	a % b	a % b	a % b	a % b	a % b
20 mcl	100 (33.3)	100 (16.7)	16.7 (66.7)	66.7 (66.7)	100 (66.7)	100 (33.3)
10 mcl	100 (33.3)	100 (16.7)	16.7 (66.7)	50 (50)	100 (66.7)	100 (33.3)
6 mcl	100 (33.3)	83.3 (16.7)	16.7 (66.7)	50 (50)	83.3 (50)	83.3 (16.7)
3 mcl	83.3 (33.3)	66.7 (16.7)	83.3 (50)	16.7 (16.7)	66.7 (33.3)	83.3 (16.7)

a) phenol-chloroform; b) chelex

Table 1. Amplification and typing of HUMTH01 and HUMvWA 31 from bloodstains of different sizes on various substrates using two extraction methods

REFERENCES

- Allen RC, Graves G, Budowie B (1989) Polymerase chain reaction amplification products separated on rehydratable polyacrylamide gels and stained with silver. *Bio Tech* 7: 736-744
- Brinkmann B, Rand S, Bajanowsky T (1992) Forensic identification of urine samples. *Int J Leg Med* 105: 59-61
- Budowie B, Baechtel FS (1990) Modifications to improve effectiveness of restriction fragment length polymorphism typing. *Appl Theor Electrophor* 1:181-187
- Budowie B, Chakraborty R, Giusti AM, Eisenberg AJ, Allen RC (1991) Analysis of the variable number of tandem repeats locus D1S80 by the polymerase chain reaction followed by high resolution polyacrylamide gel electrophoresis. *Am J Hum Genet* 48: 137-144
- Edwards A, Civitello A, Hammond HA, Caskey CT (1991) DNA typing and genetic mapping with trimeric and tetrameric tandem repeats. *Am J Hum Genet* 49: 746-756
- Edwards A, Hammond HA, Jin L, Caskey CT, Chakraborty R (1992) Genetic variation at five trimeric and tetrameric tandem repeat loci in four human population groups. *Genomics* 12: 241-253
- Kimpton C, Walton A, Gill P (1992) A further tetranucleotide repeat polymorphism in the vWF gene. *Hum Mol Genet* 1:287
- Walsh PS, Metzger DA, Higuchi R (1991) Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *Biotechniques* 10: 506-513
- Weber J L and May P E (1989) Abundant class of human DNA polymorphisms which can be typed using the polymerase chain reaction. *Am J Hum Genet* 44: 388-396
- Wiegand P, Bajanowski T, Brinkmann B (1993) DNA typing of debris from fingernails. *Int J Leg Med* 106: 81-83