

Review

# PdI<sub>2</sub>-Based Catalysis for Carbonylation Reactions: A Personal Account

Raffaella Mancuso <sup>1</sup>, Nicola Della Ca' <sup>2</sup>, Lucia Veltri <sup>1</sup>, Ida Ziccarelli <sup>1</sup> and Bartolo Gabriele <sup>1,\*</sup>

<sup>1</sup> Laboratory of Industrial and Synthetic Organic Chemistry (LISOC), Department of Chemistry and Chemical Technologies, University of Calabria, Via P. Bucci 12/C, 87036 Arcavacata di Rende (CS), Italy

<sup>2</sup> Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale Parco Area delle Scienze, Università di Parma, 17/a, 43124 Parma, Italy

\* Correspondence: bartolo.gabriele@unical.it; Tel.: +39-0984-492-815

Received: 2 July 2019; Accepted: 13 July 2019; Published: 18 July 2019



**Abstract:** In this account, we review our efforts in the field of carbonylation reactions promoted by palladium iodide-based catalysts, which have proven to be particularly efficient in diverse kinds of carbonylation processes (oxidative carbonylations as well as additive and substitutive carbonylations). Particularly in the case of oxidative carbonylations, more emphasis has been given to the most recent results and applications.

**Keywords:** carbon monoxide; carbonylation; heterocycles; homogeneous catalysis; palladium; palladium iodide

## 1. Introduction



Metal-catalyzed carbonylation reactions are very important processes in industrial chemistry and organic synthesis, since they allow the direct incorporation of the simplest C-1 unit (carbon monoxide) into a substrate, with the direct formation of high value-added carbonylated compounds [1–30].

In this field, our research group has been active for many years. In particular, by employing palladium iodide-based catalysts, many carbonylation reactions have been developed, under oxidative as well as non-oxidative conditions. We have previously reviewed our efforts on oxidative carbonylations and carbonylative synthesis of heterocycles [23–30]. In this personal account, we will describe our achievements on PdI<sub>2</sub>-catalyzed carbonylations leading to various carbonyl compounds under oxidative as well as non-oxidative conditions. Particularly in the case of oxidative carbonylations, more emphasis will be given to the most recent results and applications.

## 2. PdI<sub>2</sub>-Catalyzed Oxidative Carbonylations


### 2.1. Background: PdI<sub>2</sub>-Catalyzed Oxidative Dialkoxycarbonylation of Alkynes

According to the literature [26], in organic synthesis, oxidative carbonylation is a process in which carbon monoxide (CO) is inserted into (an) organic substrate(s) (SH<sub>2</sub>) to yield a carbonylated product [S(CO)], under the promoting action of a metal species [M(X)], which is reduced to M(X-2) at the end of the process (Scheme 1a). A catalytic version of this reaction is achieved when operating in the presence of a suitable external oxidant (OX), able to reoxidize M(X-2) back to M(X), with the formation of OXH<sub>2</sub> (Scheme 1b) [26].




**Scheme 1.** (a) Oxidative carbonylation of organic substrate  $\text{SH}_2$  promoted by the metal species  $\text{M(X)}$ ; (b)  $\text{M(X)}$ -catalyzed oxidative carbonylation of  $\text{SH}_2$  in the presence of the external oxidant  $\text{OX}$  [26].

At the beginning of the nineties, we reported that a very simple palladium-based catalytic system, consisting of  $\text{PdI}_2$  in conjunction with an excess of  $\text{KI}$ , was a very efficient catalyst for promoting the oxidative dialkoxycarbonylation of terminal alkynes, carried out with oxygen as the external oxidant, with the formation of maleic diesters and their cyclic tautomers in high yields and excellent TONs (Scheme 2) [31,32]. Later on, we found that, using water as the nucleophile under suitable conditions, it was also possible to effectively synthesize maleic anhydrides or acids [33]. The excess of  $\text{KI}$  was necessary to solubilize  $\text{PdI}_2$  in the alcoholic solvent and to stabilize the catalytically active species  $\text{PdI}_4^{2-}$ . The main reason for the high efficiency shown by this complex was related to a very efficient reoxidation of  $\text{Pd(0)}$ , through a mechanism involving the initial oxidation of the 2 equiv of  $\text{HI}$  (ensuing from the carbonylation process) to  $\text{I}_2$  followed by the oxidative addition of the latter to  $\text{Pd(0)}$  (Scheme 3; in this and in all the following Schemes in this account, unreactive iodide ligands are omitted for clarity). The formation of maleic diesters can be rationalized by triple bond *syn* insertion into an alkoxyacetylpalladium iodide complex (formed by the reaction between  $\text{PdI}_2$ ,  $\text{R}'\text{OH}$ , and  $\text{CO}$ ) followed by carbon monoxide insertion and nucleophilic displacement by  $\text{R}'\text{OH}$  (Scheme 3) [32,34–36].



**Scheme 2.** Synthesis of maleic diesters by  $\text{PdI}_2/\text{KI}$ -catalyzed oxidative dialkoxycarbonylation of terminal alkynes (DMA = *N,N*-dimethylacetamide) [32].




**Scheme 3.** Mechanism of the formation of maleic diesters by  $\text{PdI}_2/\text{KI}$ -catalyzed oxidative dialkoxycarbonylation of terminal alkynes [32].

## 2.2. $\text{PdI}_2$ -Catalyzed Oxidative Cyclocarbonylation-Alkoxyacetylation of Functionalized Alkynes


The  $\text{PdI}_2$ -catalyzed oxidative carbonylation of the triple bond described in Section 2.1 was successively exploited to synthesize a variety of carbonylated heterocycles (such as  $\beta$ -lactones [37,38],  $\gamma$ -lactones [38],  $\beta$ -lactams [39], indolones [40], and pyridinones [41]), starting from suitably

functionalized alkynes bearing a nucleophilic group in appropriate position for cyclization. In these reactions, heterocyclization occurred with CO incorporation into the cycle (cyclocarbonylation) and was accompanied by exocyclic alkoxy carbonylation, as outlined in Scheme 4. Thus, the formation of a triple bond-stabilized complex  $[-Nu(CO)PdI]$  from the reaction between the substrate,  $PdI_2$ , and CO takes place, followed by cyclization by *syn* intramolecular triple bond insertion, CO insertion, and nucleophilic displacement by the external alcohol (Scheme 4). This kind of reactivity usually occurs with substrates bearing a terminal triple bond (because the intramolecular triple bond insertion step is quite sensitive to steric hindrance) and with relatively strong nucleophiles (such as amino groups), able to readily afford a carbamoylpalladium iodide species  $[-N(R)(CO)PdI]$ . On the other hand, with weaker nucleophilic groups (such as the hydroxyl), the cyclocarbonylation-alkoxy carbonylation mechanism is generally observed when the direct nucleophilic attack to the triple bond coordinated to palladium (see Section 2.3) is not allowed by Baldwin's rules.




**Scheme 4.** Mechanism of the formation of carbonylated heterocycles by  $PdI_2/KI$ -catalyzed oxidative cyclocarbonylation-alkoxy carbonylation of terminal alkynes bearing a suitably placed nucleophilic group.

Recently,  $\zeta$ -lactams with antitumor activity against breast cancer cells were also successfully synthesized, as shown in Scheme 5 [42]. In this reaction, the formation of a carbamoylpalladium iodide intermediate is followed by a remarkable intramolecular *syn* 8-*exo-dig* triple bond insertion and nucleophilic displacement by an alcohol (Scheme 5) [42].



**Scheme 5.** Synthesis of  $\zeta$ -lactams by  $PdI_2/KI$ -catalyzed oxidative cyclocarbonylation-alkoxy carbonylation of 2-(2-ethynylphenoxy)anilines [42].


The nucleophile may also be bonded to a heterocyclic ring, leading to fused heterocycles, as shown in the synthesis of benzimidazopyrimidinones from propynylbenzimidazolamines (Scheme 6) [43].



**Scheme 6.** Synthesis of benzimidazopyrimidinones by PdI<sub>2</sub>/KI-catalyzed oxidative cyclocarbonylation-alkoxycarbonylation of propynylbenzimidazolamines [43].

### 2.3. PdI<sub>2</sub>-Catalyzed Oxidative Heterocyclization-Alkoxycarbonylation of Functionalized Alkynes


With alkynes bearing less nucleophilic groups (such as hydroxyl or amido groups), an initial *exo* or *endo* heterocyclization without CO incorporation into the cycle generally occurs, as long as it is allowed by Baldwin's rules. Cyclization, which may take place with the terminal as well as with internal triple bonds, is then followed by alkoxy-carbonylation of the ensuing vinylpalladium moiety, to afford the final monocarbonylated heterocycle (Scheme 7). This mechanism is particularly favored when the initially formed heterocyclic ring may further undergo aromatization. Furans [44,45], tetrahydrofurans [46–48], oxazolidinones [49,50], oxazolines [39,47,51,52], pyrroles [41,53], dihydroisobenzofurans [48,54], isochromenes [54,55], isoquinolines [55], benzoxazines [56,57], quinolines [58], and indoles [59] were synthesized by this approach.



**Scheme 7.** Mechanism of the formation of carbonylated heterocycles by PdI<sub>2</sub>/KI-catalyzed oxidative heterocyclization-alkoxycarbonylation of terminal alkynes bearing a suitably placed nucleophilic group.


More recently, we have applied this kind of reactivity to the synthesis of other important heterocyclic derivatives. Thus, indole-3-carboxylic esters were prepared in good yields

(50–84%) from 2-alkynylanilines bearing an internal triple bond and a secondary amino group (Scheme 8a) [60]. With a primary amino group, and in the presence of trimethyl orthoformate, 1-(dimethoxymethyl)indole-3-carboxylates were obtained, which could be easily converted under acidic conditions into the corresponding unprotected indoles (Scheme 8b) [60]. It is worth noting that the latter could not be obtained by the direct oxidative carbonylation of 2-alkynylanilines bearing a primary amino group under the usual conditions (i.e., in the absence of trimethyl orthoformate) [60].



**Scheme 8.** Synthesis of alkyl indole-3-carboxylates by PdI<sub>2</sub>-catalyzed oxidative *endo* heterocyclization-alkoxycarbonylation of 2-alkynylanilines bearing a secondary (a) or a primary (b) amino group [60].


Using an ROH/HC(OR)<sub>3</sub> mixture as the solvent, 2-alkynylbenzamides were also subjected to PdI<sub>2</sub>-catalyzed oxidative carbonylation conditions and were selectively converted into isobenzofuranimine derivatives through 5-*exo-dig* *O*-heterocyclization followed by alkoxy carbonylation (Scheme 9) [61]. These derivatives have shown a promising herbicidal activity [62].




**Scheme 9.** Synthesis of 3-[(alkoxycarbonyl)methylene]isobenzofuran-1(3*H*)imines by PdI<sub>2</sub>-catalyzed oxidative *exo* *O*-heterocyclization-alkoxycarbonylation of 2-alkynylbenzamides [61,62].

Pyrrole-3-carboxylates were synthesized starting from *N*-Boc-1-amino-3-yn-2-ols [63]. In this case, the 5-*endo-dig* heterocyclization-alkoxycarbonylation process was accompanied by dehydrative aromatization, and led to a mixture of Boc-protected and *N*-unsubstituted pyrrole derivatives. This mixture could be easily converted into *N*-unsubstituted pyrrole-3-carboxylic esters by a basic treatment with NaOR (Scheme 10a) [63]. Starting from substrates bearing geminal alkynyl groups, 4-acyl-3-pyrrolecarboxylates were directly obtained, from *N*-deprotection in situ and regioselective water addition to the triple bond of the second alkynyl moiety (Scheme 10b) [63].

In a similar manner, 3-yne-1,2-diol derivatives were smoothly converted into furan-3-carboxylates (Scheme 11a) [64,65] or 4-methylene-4,5-dihydrofuran-3-carboxylates (Scheme 11b) [65], depending on whether dehydration from the alcoholic function at C-2 occurred in an endocyclic (with substrates bearing a secondary alcoholic group at C-1) or exocyclic (tertiary alcoholic group at C-1) way, respectively.




**Scheme 10.** Synthesis of pyrrole-3-carboxylates (a) and 4-acyl-3-pyrrolecarboxylates (b) by PdI<sub>2</sub>-catalyzed *endo* heterocyclization-alkoxycarbonylation of *N*-Boc-1-amino-3-yn-2-ols and *N*-Boc-2-alkynyl-1-amino-3-yn-2-ols, respectively [63].



**Scheme 11.** Synthesis of furan-3-carboxylates (a) [64,65] or 4-methylene-4,5-dihydrofuran-3-carboxylates (b) [65] by PdI<sub>2</sub>-catalyzed oxidative *endo*-heterocyclization-alkoxycarbonylation of 3-yn-1,2-diol derivatives.


Very recently, we found that 2-(ethynyl)benzoic acids (obtained in situ by desilylation of 2-[(trimethylsilyl)ethynyl]benzoic acids) can be stereoselectively converted into (*Z*)-2-[oxoisobenzofuran-1-(3*H*)-ylidene]acetates in high to excellent yields (70–98%) (Scheme 12) [66]. In this case, to account for product stereochemistry (which was confirmed by XRD analysis), the formation of a palladium carboxylate can be assumed, followed by 5-*exo-dig* intramolecular insertion of the triple bond and alkoxylation (Scheme 12) [66].




**Scheme 12.** Synthesis of (*Z*)-2-[oxoisobenzofuran-1-(3*H*)-ylidene]acetates by PdI<sub>2</sub>-catalyzed oxidative heterocyclization-alkoxycarbonylation of 2-(ethynyl)benzoic acids [66].

#### 2.4. PdI<sub>2</sub>-Catalyzed Oxidative Carbonylative Double Cyclization of Functionalized Alkynes

With acetylenic substrates bearing two suitably placed nucleophilic groups, a cascade double cyclization process may take place, ensuing from an initial heterocyclization followed by intramolecular cyclocarbonylation. This reactivity was observed with 2-(hydroxypropyn-1-yl)anilines, which were converted into 3,4-dihydrofuro[3,4-*b*]indol-1-ones by 5-*endo-dig* *N*-cyclization followed by carbon monoxide insertion and intramolecular nucleophilic displacement by the alcoholic group (Scheme 13) [67]. In a similar way, furobenzofuranone derivatives were synthesized in 74–86% yields starting from 2-(hydroxypropyn-1-yl)phenols, in BmimBF<sub>4</sub> (1-butyl-3-methylimidazolium tetrafluoroborate) as unconventional solvent (Scheme 14) [68].




**Scheme 13.** Synthesis of 3,4-dihydrofuro[3,4-*b*]indol-1-ones by PdI<sub>2</sub>-catalyzed oxidative carbonylative double cyclization of 2-(hydroxypropyn-1-yl)anilines: 5-*endo-dig* *N*-cyclization followed by cyclocarbonylation [67].



**Scheme 14.** Synthesis of furo[3,4-*b*]benzofuran-1(3*H*)-ones by PdI<sub>2</sub>-catalyzed oxidative carbonylative double cyclization of 2-(hydroxypropyn-1-yl)phenols in BmimBF<sub>4</sub> [68].


In the case of 5-trimethylsilyl-4-yne-1,3-diols, an initial 5-*exo-dig* *O*-cyclization was followed by cyclocarbonylation to afford dihydrofurofuranone derivatives, which have shown significant antitumor activity *in vitro* against breast cancer cells, including the most aggressive MDA-MB-231 and MDA-MD-468 cells (Scheme 15) [69].



**Scheme 15.** Synthesis of 6,6*a*-dihydrofuro[3,2-*b*]furan-2(5*H*)-one derivatives by PdI<sub>2</sub>-catalyzed oxidative carbonylative double cyclization of 5-trimethylsilyl-4-yne-1,3-diols: 5-*exo-dig* O-cyclization followed by cyclocarbonylation [69].

### 2.5. PdI<sub>2</sub>-Catalyzed Oxidative Monoaminocarbonylation of Terminal Alkyne Derivatives


When carried out in the presence of a nucleophilic secondary amine as external nucleophile instead of an alcohol, the PdI<sub>2</sub>-catalyzed oxidative carbonylation of terminal alkynes selectively led to 2-ynamides, corresponding to a monocarbonylation of the triple bond ensuing from *sp* C-H activation (Scheme 16) [70]. In fact, under these conditions, the amine may initially act as a base, promoting the formation of an alkynylpalladiumiodide species. Carbon monoxide insertion is then followed by nucleophilic displacement by the same amine to afford the final product (Scheme 16) [70].



**Scheme 16.** PdI<sub>2</sub>-catalyzed oxidative monoaminocarbonylation of terminal alkynes leading to 2-ynamides [70].

This reactivity may lead to the formation of heterocyclic derivatives when applied to suitably substituted terminal alkynes, because the initially formed 2-ynamide compounds present a triple bond conjugated to the amido group and are therefore susceptible to nucleophilic attack (either intermolecular or intramolecular). Thus, furanone [71,72] and pyrrolone [73] derivatives were formed starting from

propargyl alcohols or amines, respectively, ensuing from intermolecular conjugate addition by the secondary amine and intramolecular alcoholysis or aminolysis [71–73]. In other cases, the nucleophilic function in the 2-ynamide intermediate was already placed in a suitable position for the occurrence of an intramolecular conjugate addition, in agreement with Baldwin's rules. By this latter approach, different kinds of carbonylated heterocycles, including tetrahydrofurans [72,74], dihydrobenzodioxines [75], dihydrobenzoxazines [75], and furans [72,76] were synthesized, as exemplified in Scheme 17.




**Scheme 17.** Synthesis of various functionalized heterocycles by PdI<sub>2</sub>-catalyzed oxidative monoaminocarbonylation of terminal alkynes followed by intramolecular conjugate addition [72,74–76].


More recently, isoindolinones were conveniently prepared by sequential oxidative aminocarbonylation-intramolecular conjugate addition of 2-ethynylbenzamides, as shown in Scheme 18a [61,77]. The heterocyclic derivatives thus obtained were used for preparing new spiro[isoindole-1,5'-isoxazolidin]-3(2*H*)ones with antitumor activity against neuroblastoma SH-SY5Y, colorectal HT-29, and hepatocellular HepG2 cell lines (Scheme 18b) [78].

Propynylbenzimidazolamines, already successfully employed for the synthesis of benzimidazopyrimidinones under oxidative alkoxy carbonylation conditions (Scheme 6, Section 2.2) [43], were also used as substrates under aminocarbonylation conditions, to produce benzimidazoimidazoles in good to excellent yields after a final isomerization process (Scheme 19) [79]. In a similar way, benzimidazothiazoles were synthesized from propynylsulfanylbenzimidazoles, obtained in situ from the corresponding benzimidazolium bromide salts (Scheme 20) [80]. In this latter case, the internal heteronucleophile corresponded to the nitrogen of an imidazole ring.


Aminocarbonylation followed by carbocyclization is also possible, with substrates leading in situ to stabilized carbonucleophiles (by carbon deprotonation under the basic reaction conditions) in a suitable position with respect to the triple bond. Thus, 2-(2-ethynylbenzyl)malonates were converted into functionalized indanes in one single procedure, as shown in Scheme 21 [81].




**Scheme 18.** (a) Synthesis of isoindolinones by sequential  $\text{PdI}_2$ -catalyzed oxidative aminocarbonylation-*N*-cyclization of 2-ethynylbenzamides [61,77] and (b) their use for the preparation of spiro[isoindole-1,5'-isoxazolidin]-3(2H)ones with antitumor activity [78].



**Scheme 19.** Synthesis of benzimidazoimidazoles by sequential oxidative  $\text{PdI}_2$ -catalyzed aminocarbonylation-*N*-cyclization-isomerization of propynylbenzimidazolamines [79].




**Scheme 20.** Synthesis of benzimidazothiazoles by sequential  $\text{PdI}_2$ -catalyzed oxidative aminocarbonylation-*N*-cyclization-isomerization of propynylsulfanylbenzimidazoles [80].




**Scheme 21.** Synthesis of indanes by  $\text{PdI}_2$ -catalyzed by oxidative aminocarbonylation-carbocyclization of 2-(2-ethynylbenzyl)malonates (EWG =  $\text{CO}_2\text{Me}$ , EWG' =  $\text{CO}_2\text{Me}$ , COMe, CN) [81].

In some cases, and under suitable conditions, the initial  $\text{PdI}_2$ -catalyzed aminocarbonylation of the terminal triple bond is followed by  $\text{PdI}_2$ -catalyzed cyclocarbonylation through the concatenation of two different catalytic cycles, both catalyzed by the same species (auto-tandem catalysis). We observed this reactivity with propargyl amines, when the oxidative carbonylation reactions were carried out in the presence of water as a promoter [82]. The cyclocarbonylation process started with palladation of the nitrogen of the 2-ynamide intermediate (deriving from the aminocarbonylation process) and was followed by CO insertion, O-cyclization triggered by water attack, and  $\beta$ -H elimination from the HO-C-Pd moiety (Scheme 22) [82]. More recently, we also found that it is possible to carry out the process in the ionic liquid (IL) EmimEtSO<sub>4</sub> (1-ethyl-3-methylimidazolium ethyl sulfate) as a nonconventional solvent, with the possibility to recycle the catalyst-IL medium [83].



**Scheme 22.** Synthesis of oxazolidinones from propargylic amines by  $\text{PdI}_2$ -promoted auto-tandem catalysis:  $\text{PdI}_2$ -catalyzed oxidative aminocarbonylation followed by  $\text{PdI}_2$ -catalyzed, water-promoted oxidative cyclocarbonylation [82].


Aminocarbonylation followed by cyclocarbonylation was also observed in the case of monosubstituted propynylthioimidazoles, which led to imidazothiazinones in one step and in satisfactory yields. In this case, in the cyclocarbonylation cycle, nitrogen palladation was followed by CO insertion, intramolecular triple bond insertion, protonolysis, and isomerization (Scheme 23) [84].



**Scheme 23.** Synthesis to imidazothiazinones by PdI<sub>2</sub>-catalyzed oxidative aminocarbonylation-cyclocarbonylation of propynylthioimidazoles [84].

### 2.6. PdI<sub>2</sub>-Catalyzed Oxidative Monoalkoxycarbonylation of $\alpha$ -Olefins


In combination with CuI as co-catalyst, PdI<sub>2</sub>-based catalysts are also able to promote the monoalkoxycarbonylation of  $\alpha$ -olefins to give (*E*)- $\alpha,\beta$ -unsaturated esters, under relatively mild conditions (120 °C and under 2 atm of CO and 35 atm of air). Pd/C was used as the catalytic precursor in this process, leading in situ to PdI<sub>2</sub> by the combined action of CuI and oxygen. Interestingly, the direct use of PdI<sub>2</sub>/KI or PdI<sub>2</sub>/CuI catalytic systems led to less satisfactory results (both in terms of product yield and selectivity) with respect to Pd/C-CuI, perhaps also suggesting the involvement, in the last case, of Pd-Cu bimetallic species, more active toward olefin carbonylation. The formation of an alkoxycarbonylpalladium iodide species was then followed by regioselective double bond insertion and stereoselective  $\beta$ -H elimination to give the final product (Scheme 24). The reaction worked nicely with several arenes, leading to the corresponding cinnamates in yields up to 95% (Scheme 24). In the case of aliphatic  $\alpha$ -olefins, however, a mixture of isomeric  $\alpha,\beta$ - and  $\beta,\gamma$ -unsaturated esters were obtained [85].



**Scheme 24.** Synthesis of alkyl cinnamates by PdI<sub>2</sub>/CuI-catalyzed oxidative monoalkoxycarbonylation of arenes [85].

### 2.7. PdI<sub>2</sub>-Catalyzed Oxidative Carbonylation of Amines, β-Amino Alcohols, and 2-Aminophenols to Ureas, Oxamides, Oxazolidinones, and Benzoxazolones


We have seen in Sections 2.2 and 2.5 that PdI<sub>2</sub> can easily react with amino groups to form carbamoylpalladiumiodide intermediates. This reactivity can be conveniently exploited for the synthesis of ureas, when applied to simple primary amines. In fact, in this case, the initially formed RNH(CO)PdI intermediate may undergo β-H elimination from the HN-CO-Pd moiety, with the formation of an isocyanate and palladium(0). While the latter is readily reoxidized to PdI<sub>2</sub> in the presence of oxygen, the isocyanate may be attacked by a second molecule of amine (either the same primary amine or a secondary amine, used as additional substrate) to yield a symmetrical 1,3-disubstituted or a trisubstituted urea, respectively (Scheme 25) [86,87]. We originally developed this process using 1,2-dimethoxyethane (DME) as the solvent, using, in some cases, an excess of CO<sub>2</sub> to ensure better catalytic performances [87]. In the synthesis of symmetrically 1,3-disubstituted ureas, carbon dioxide could even be used as the reaction solvent, reaching, in the conversion of aniline to 1,3-diphenylurea (DPU), an unprecedented catalytic efficiency of 43,500 mol of DPU per mol of palladium employed [88,89].




**Scheme 25.** Formation of symmetrically 1,3-disubstituted and trisubstituted ureas by PdI<sub>2</sub>-catalyzed oxidative carbonylation of amines [86–89].

More recently, we found that it is also possible to carry out this transformation in the ionic liquid (IL) BmimBF<sub>4</sub> (1-butyl-3-methylimidazolium tetrafluoroborate) as an unconventional solvent, with the possibility to recycle the catalyst-IL system (Scheme 26) [90]. Interestingly, while aromatic primary amines led to 1,3-diarylureas (Scheme 26a), aliphatic primary amines could be divergently converted into 1,3-dialkylureas (Scheme 26b) or *N,N'*-dialkyloxamides (deriving from a double carbonylation process; Scheme 26c), depending on reactions conditions [90]. On the other hand, secondary amines (normally unreactive under the “classical” conditions [86–88]) selectively afforded tetrasubstituted oxamides (Scheme 26d) [90].

The isocyanate intermediate deriving from a primary amine under PdI<sub>2</sub>-catalyzed oxidative carbonylation conditions can also be trapped intramolecularly by a second nucleophilic group. Thus, cyclic ureas [87], 2-oxazolidinones [90–93], and benzoxazolones [90,92] were obtained in excellent yields from aromatic diamines, β-amino alcohols, and 2-aminophenols, respectively, as exemplified in Scheme 27.




**Scheme 26.** Synthesis of (a) 1,3-diaryloxamides; (b) 1,3-dialkylureas; (c) *N,N'*-dialkylureas; and (d) tetraalkylureas by PdI<sub>2</sub>-catalyzed oxidative carbonylation of amines in BmimBF<sub>4</sub> [90].



**Scheme 27.** Synthesis of (a) benzimidazolone [87]; (b) 2-oxazolidinones [92]; and (c) benzoxazolones (Y = CH,N) [90] by PdI<sub>2</sub>-catalyzed oxidative carbonylation of 1,2-benzenediamine, β-amino alcohols, and 2-aminophenols, respectively.


### 2.8. PdI<sub>2</sub>-Catalyzed Oxidative Carbonylation of 1,2- and 1,3-Diols to Cyclic Carbonates

Under PdI<sub>2</sub>-catalyzed oxidative carbonylation conditions 1,2- and 1,3-diols, were also reactive and led to the corresponding cyclic carbonates in good to excellent yields (66–94%, Scheme 28) [94,95]. In some cases, the use of a dehydrating agent, such as trimethyl orthoacetate, was necessary to achieve more satisfactory product selectivities. In this case, the alkoxyacetyl palladium iodide intermediate deriving from the reaction between the first hydroxyl group, PdI<sub>2</sub> and CO, was intramolecularly trapped by nucleophilic displacement by the second hydroxyl (Scheme 28) [94,95].



**Scheme 28.** Synthesis of 5-membered and 6-membered cyclic carbonates by PdI<sub>2</sub>-catalyzed oxidative cyclocarbonylation of 1,2- and 1,3-diols, respectively [94,95].

Notably, the reaction could also be applied to glycerol, affording, under suitable conditions, glycerol carbonate (a very important derivative, used in many applicative fields, including industrial, pharmaceutical, and material chemistry) in 62% yield (Scheme 29a) [95]. More complex polyols, such as D-glucose, underwent a double cyclocarbonylative process leading to bicyclic carbonates (Scheme 29b) [95].




**Scheme 29.** Synthesis of (a) glycerol carbonate and (b)  $\alpha$ -D-glucofuranose 1,2:5,6-dicarbonate by PdI<sub>2</sub>-catalyzed oxidative cyclocarbonylation of glycerol and D-glucose, respectively [95].

## 3. PdI<sub>2</sub>-Catalyzed Non-Oxidative Carbonylations

### 3.1. Combination Between Oxidative and Reductive Carbonylation and Additive Carbonylation of Terminal Alkynes


As we have seen in Section 2.1, the palladium-iodine bond of PdI<sub>2</sub> is particularly prone to undergo alkoxyacetylation in the presence of carbon monoxide and an alcohol, to give an

alkoxycarbonylpalladiumiodide intermediate with the elimination of HI. In the presence of an alkyne substrate, triple bond insertion may take place, usually in a regioselective manner, leading to an alkoxycarbonylvinylpalladium species, as depicted in Scheme 30. Under suitable conditions (depending on solvent, temperature, and CO pressure), this intermediate may further insert CO to give the corresponding acylpalladium complex, which undergoes nucleophilic displacement by the alcohol to give maleic diesters, HI, and Pd(0) (Section 2.1 and Scheme 30).



**Scheme 30.** Formation of alkoxycarbonylvinylpalladium iodide from PdI<sub>2</sub>, an alcohol, CO, and a terminal alkyne, and subsequent CO insertion and nucleophilic displacement leading to maleic diesters and Pd(0).

In particular, in highly polar MeOH as the solvent and nucleophile, maleic dimethyl esters are formed even at room temperature and under atmospheric pressure of CO, with reduction of PdI<sub>2</sub> to Pd(0), which readily decomposes to palladium black (stoichiometric oxidative dimethoxycarbonylation reaction) (Scheme 31a). As we have seen in Section 1, one way to achieve a catalytic process, affording maleic diesters, is to operate under oxidative conditions, that means, in the presence of an external oxidant (such as oxygen), able to reconvert Pd(0)+2HI into PdI<sub>2</sub> (Schemes 3 and 31b). In principle, however, a catalytic process can also take place under non-oxidative conditions, if the dialkoxycarbonylation process leading to maleates takes place under conditions able to stabilize the H-Pd-I species that can be in equilibrium with Pd(0)+HI. In this manner, in fact, palladium does not decompose to palladium black, while the H-Pd-I species may promote a parallel reduction process, leading to a second product corresponding to reductive carbonylation (a 2(5H)furanone, for example), with the regeneration of PdI<sub>2</sub> (Scheme 31c).




**Scheme 31.** (a) PdI<sub>2</sub>-promoted stoichiometric oxidative dimethoxycarbonylation of terminal alkynes, leading to maleic diesters and Pd(0); (b) possible reconversion of Pd(0) to PdI<sub>2</sub> under oxidative conditions (with oxygen as external oxidant); (c) possible reconversion of Pd(0) to PdI<sub>2</sub> under non-oxidative conditions, with concomitant formation of a reductive carbonylation product.

Many years ago we found that, in MeOH as the solvent and under particularly mild conditions (25 °C and 1 atm of CO), an appropriate ligand able to stabilize the H-Pd-I species deriving from the oxidative dimethoxycarbonylation of terminal alkynes could be thiourea (tu) [96]. In fact, using PdI<sub>2</sub> in conjunction with 3–3.5 equiv tu to PdI<sub>2</sub>, a catalytic process under non-oxidative conditions took place, corresponding to the combination between oxidative dimethoxycarbonylation (leading to maleic diesters) and reductive dicarbonylation (leading to 2(5H)-furanones). Accordingly, the maleic diesters and furanones were obtained in a ca. 1:1 molar ratio (Scheme 32) [96].




**Scheme 32.** PdI<sub>2</sub>-catalyzed carbonylation of terminal alkynes under non-oxidative conditions: combined oxidative dimethoxycarbonylation and reductive dicarbonylation of terminal alkynes leading to maleic diesters and 2(5H)-furanones, respectively [96].

In a simplified version of the mechanism, the tu-stabilized H-Pd-I species ensuing from the dimethoxycarbonylation of the triple bond may interact with the acylpalladium intermediate **I**, leading, after hydride exchange, to acylpalladiumhydride species **II**. The final furanone product would then be formed after the intramolecular nucleophilic attack of the hydride to the acyl moiety with simultaneous cyclization to give  $\pi$ -allylcomplex **III** followed by protonolysis, with the regeneration of PdI<sub>2</sub> (Scheme 33) [96].




**Scheme 33.** Simplified mechanistic proposal for the PdI<sub>2</sub>/tu-catalyzed combined oxidative dimethoxycarbonylation and reductive dicarbonylation of terminal alkynes leading to maleic diesters and 2(5H)-furanones, respectively [96].

Interestingly the same kind of reactivity was also observed in the butoxycarbonylation of terminal alkynes, when it was carried out with BuOH as a nucleophile (10 equiv with respect to the alkyne) in a polar aprotic solvent, such as MeCN [97]. In this case, the addition of tu as a ligand was not necessary, as the combination between the oxidative carbonylation and reductive carbonylation took place with PdI<sub>2</sub> in conjunction with KI as the catalyst (Scheme 34) [97]. This is likely due to the fact that, in the aprotic solvent, the equilibrium between H-Pd-I and Pd(0)+HI is much less shifted to the right with respect to MeOH, where HI is much more stabilized by hydrogen bonding.




**Scheme 34.** PdI<sub>2</sub>/KI-catalyzed combined oxidative dibutoxycarbonylation and reductive dicarbonylation of phenylacetylene leading to dibutyl 2-phenylmaleate and 3-phenylfuran-2(5H)-one, respectively [97].

Even more interesting was the observation that, shifting from polar MeCN ( $\epsilon = 37.5$ ) to apolar toluene ( $\epsilon = 2.4$ ), the reaction course changed dramatically, with the predominant formation of the hydrobutoxycarbonylation product (Scheme 35a) [97]. This product clearly derives from an additive carbonylation process, ensuing from protonolysis of the butoxycarbonylpalladium intermediate IV, with direct regeneration of PdI<sub>2</sub> (Scheme 36). The process worked even better in the absence of KI, as shown in Scheme 35b, and could also be generalized to other arylacetylenes (Scheme 35c) [97]. Apparently, in low-polar solvents, protonolysis by HI is much more favored with respect to further alkoxycarbonylation of IV (Scheme 36) [97].




**Scheme 35.** PdI<sub>2</sub>-catalyzed additive hydrobutoxycarbonylation reactions in toluene as the solvent: (a) of phenylacetylene with PdI<sub>2</sub>/KI as the catalytic system; (b) of phenylacetylene with PdI<sub>2</sub> alone as catalyst; (c) of arylacetylenes with PdI<sub>2</sub> as catalyst [97].



**Scheme 36.** Mechanism of the PdI<sub>2</sub>-catalyzed additive hydrobutoxycarbonylation of terminal alkynes in toluene as the solvent [97].


### 3.2. Combination Between Oxidative and Reductive Carbonylation within the Same Molecule and Sequential Homobimetallic Catalysis

Of particular interest is the possible combination between an oxidative carbonylation process and a reductive process within the same molecule, because this may result in the selective formation of a single molecule, under PdI<sub>2</sub>-catalyzed non-oxidative conditions. This kind of concept was successfully verified in the case of 3,3-diethoxyprop-1-yne, containing, besides the terminal triple bond, an acetalic moiety potentially reducible. This substrate was selectively converted into diethyl (*E*)-2-(ethoxymethylene)succinate in 65% yield in the presence of PdI<sub>2</sub>+3tu as the catalytic system under mild conditions (rt and 7 atm of CO), in EtOH as the solvent (Scheme 37). The initially formed maleate derivative reacted with H-Pd-I to give a  $\pi$ -allyl complex with the elimination of EtOH from the acetal function, and then protonolysis by HI afforded the final product (Scheme 37) [98].



**Scheme 37.** Synthesis of diethyl (*E*)-2-(ethoxymethylene)succinate from 3,3-diethoxyprop-1-yne by the combination between PdI<sub>2</sub>-catalyzed oxidative dialkoxycarbonylation and reduction within the same molecule [98].

Similar to an acetalic function, an allyl alcoholic moiety can also serve as “internal” reducible function, with the elimination of water, thus allowing the occurrence of catalytic carbonylation under non-oxidative conditions leading to a single product. Thus, 1,1'-(1,2-phenylene)bis(prop-2-yn-1-ol) was converted into methyl 2-(2-oxo-2,3-dihydronaphtho[1,2-*b*]furan-4-yl)acetate in 58% yields, by reduction of both the allyl alcoholic groups formed in the initial dimethoxycarbonylation process, under the catalysis of the Pdtu<sub>4</sub>I<sub>2</sub> complex (Scheme 38) [99]. The X-ray structure of this catalyst confirmed that tu coordinates through the sulfur atom [100].




**Scheme 38.**  $\text{Pdtu}_4\text{I}_2$ -catalyzed synthesis of methyl 2-(2-oxo-2,3-dihydronaphtho[1,2-*b*]furan-4-yl)acetate from 1,1'-(1,2-phenylene)bis(prop-2-yn-1-ol) by the combination between oxidative dialkoxycarbonylation and reduction within the same molecule [99].

More recently, we applied this concept in the non-oxidative carbonylation of 2-(1-hydroxyprop-2-ynyl)phenols, which, under suitable conditions, afforded coumarinacetate derivatives (Scheme 39) [101], which have shown an interesting potential herbicidal activity [102]. In this case, the initial oxidative cyclocarbonylation-methoxycarbonylation process was followed by reduction of the HO-C≡C-CO<sub>2</sub>Me moiety to give the final coumarinic derivative with the regeneration of PdI<sub>2</sub>. Notably, in this process, iodide was the only ligand used for palladium (no tu was needed), owing to the high reactivity of the HO-C≡C-CO<sub>2</sub>Me toward H-Pd-I to give a particularly stable β-allyl complex (Scheme 39) [101].

The possibility to combine oxidative carbonylation with a reduction within the same molecule was also assessed with 2-(1-hydroxylalk-2-ynyl)phenols, bearing an internal rather than a terminal triple bond. These substrates, however, turned out to be unstable, so we decided to start with 1-(2-allyloxyphenyl)-2-yn-1-ols, with the phenolic oxygen protected with the allyl group. In order to achieve the allyl deprotection in situ, we carried out the carbonylation experiments in the presence of two different kinds of catalysts: Pd(PPh<sub>3</sub>)<sub>4</sub> (able to promote substrate deallylation) and PdI<sub>2</sub>/KI (able to promote the oxidative carbonylation-reduction sequence). The use of these two catalysts simultaneously caused the selective formation of the desired product deriving from an ordered deallylation-oxidative carbonylation-reduction sequence, as shown in Scheme 40 [103,104]. For a process like this, consisting of two cycles catalyzed by the same metal, but in two different oxidation states, we coined the term “sequential homobimetallic catalysis” [103–105]. In more detail, the first cycle, catalyzed by Pd(0), relates to substrate deallylation, while the second cycle, catalyzed by Pd(II), corresponds to 5-*exo-dig* heterocyclization followed by oxidative alkoxy carbonylation and reduction of the HO-C≡C-CO<sub>2</sub>Me moiety (Scheme 40) [103–105]. No cyclocarbonylation product was observed,


because the intramolecular triple bond insertion into a phenoxyacetyl palladium intermediate was much less favored with an internal triple bond with respect to a terminal one. The benzofuran derivatives thus obtained were of particular interest, as they showed important antitumor activity against breast cancer cell lines [106]. Moreover, very recently, we have shown that it is possible to improve the delivery of this bioactive scaffold in vitro by chemically anchoring it (through the formation of amide bond) to suitably functionalized graphene quantum dots [107].




**Scheme 39.** Synthesis of methyl 2-(2-oxo-2H-chromen-3-yl)acetates from 2-(1-hydroxyprop-2-ynyl)phenols by the combination between PdI<sub>2</sub>-catalyzed oxidative alkoxyacylation and reduction within the same molecule [101].

Interestingly, the PPh<sub>3</sub>-stabilized Pd(0) complex could also be formed starting from PdI<sub>2</sub>+PPh<sub>3</sub>+H<sub>2</sub>O (Scheme 41) [103–105], since it is known that PdI<sub>2</sub> may be reduced to Pd(0) by the action of water and CO, through decarboxylation of the I-Pd-CO<sub>2</sub>H intermediate formed by the reaction between PdI<sub>2</sub>, CO, and H<sub>2</sub>O (Scheme 42) [33,108,109].


Amines could also be used as nucleophiles in this reaction, leading to the corresponding 2-benzofuran-2-ylacetamides in good to high yields (Scheme 43) [110]. More recently, we have also found that it is possible to carry out the sequential homobimetallic catalytic process in an unconventional solvent, such as BmimBF<sub>4</sub>, with the possibility to recycle the solvent-catalyst system [111].




1st cycle: Pd(0)-catalyzed deallylation




2nd cycle: Pd(II)-catalyzed carbonylative heterocyclization




**Scheme 40.** Sequential homobimetallic catalysis: synthesis of methyl 2-(benzofuran-2-yl)hexanoate from 1-[2-(allyloxy)phenyl]hept-2-yn-1-ol by the concatenation between Pd(0)-catalyzed deallylation – PdI<sub>2</sub>-catalyzed carbonylative heterocyclization [103–105].



**Scheme 41.** Synthesis of methyl 2-benzofuranacetates from 1-(2-allyloxyphenyl)-2-yn-1-ols by sequential homobimetallic catalysis, using PdI<sub>2</sub> as both the catalytic precursor for Pd(0) and as Pd(II) catalyst [103–105].




**Scheme 42.** In situ formation of Pd(0) from PdI<sub>2</sub>, CO, and H<sub>2</sub>O [33,108,109].



**Scheme 43.** Synthesis of 2-benzofuran-2-ylacetamides from 1-(2-allyloxyphenyl)-2-yn-1-ols by sequential homobimetallic catalysis using amines as nucleophiles [110].

The mechanistic sequence heterocyclization-oxidative alkoxyacylation-reduction of an HO-C-C=C-CO<sub>2</sub>Me moiety was also observed in the formation of indol-2-acetic esters starting from 1-(2-aminoaryl)-2-yn-1-ols, as shown in Scheme 44 [58].




**Scheme 44.** Synthesis of indol-2-acetic acids from 1-(2-aminoaryl)-2-yn-1-ols [58].

### 3.3. Substitutive Carbonylation of Allylic Alcohols to $\beta,\gamma$ -Unsaturated Acids or Esters





We have seen in Section 3.2 that the H-Pd-I species obtained from PdI<sub>2</sub>, possibly stabilized by tu, can efficiently promote the reduction of allyl alcoholic moieties through the formation of a  $\pi$ -allylpalladium complex with the elimination of water (Schemes 37–40 and 44). Under suitable conditions, the  $\pi$ -allylpalladiumiodide complex obtained from H-Pd-I and an allylic alcohol may undergo, instead of protonolysis, carbon monoxide insertion followed by nucleophilic displacement by water or an alcohol. This leads to the formation of a  $\beta,\gamma$ -unsaturated acid or ester, respectively, together with regeneration of H-Pd-I (Scheme 45). As we have described in Section 3.2 (Scheme 42), the easiest way to generate H-Pd-I from PdI<sub>2</sub> is to work in the presence of small amounts of water (either present as an impurity in the reaction solvent or added in the reaction mixture). Moreover, we have seen in Section 3.1 that the H-Pd-I species can be stabilized by the ligand tu.

Based on these concepts, we were able to directly convert (*E*)-3-phenylprop-2-ene-1-ol into 4-phenylbut-3-enoic acid in 81% yield, working in *N,N*-dimethylacetamide (DMA) as the solvent in the presence of water, using PdI<sub>2</sub> in conjunction with 3 equiv of tu as the catalytic system (Scheme 46a) [112]. Using methanol as the nucleophile under similar conditions, different allylic alcohols were also efficiently converted into the corresponding  $\beta,\gamma$ -unsaturated esters (Scheme 46b) [112]. The method also worked

nicely for the preparation of high value-added dimethyl hex-3-ene-1,6-dioate from but-2-ene-1,4-diol (Scheme 46c) and trimethyl aconitate from dimethyl hydroxymethylmaleate (Scheme 46d) [31,112].



**Scheme 45.** Formation of  $\beta,\gamma$ -unsaturated acids or esters by the reaction between allylic alcohols and H-Pd-I followed by CO insertion and nucleophilic displacement by water or an alcohol, respectively.




**Scheme 46.** PdI<sub>2</sub>/tu-catalyzed substitutive carbonylation of (a) cinnamic alcohol to cinnamic acid; (b) allylic alcohols to  $\beta,\gamma$ -unsaturated esters; (c) but-2-ene-1,4-diol to dimethyl hex-3-ene-1,6-dioate; (d) hydroxymethylmaleate to trimethyl aconitate [31,112].

### 3.4. Reductive Carbonylation of Terminal Alkynes

We have seen in Section 3.1 that, in MeOH as the solvent, the H-Pd-I species, ensuing from the dimethoxycarbonylation of the triple bond, and stabilized by thiourea as a ligand, is able to reduce the acylpalladium intermediate I (Scheme 33). This probably occurs through the formal formation of an acylpalladiumhydride species II, eventually leading (via the formation of  $\pi$ -allylpalladiumiodide complex III) to a furanone product, corresponding to reductive carbonylation of the triple bond (Scheme 33). We then also noticed that an H-Pd-I species can be relatively stable, even in the absence of tu as a ligand, in an aprotic solvent, such as MeCN (Scheme 35 and related text). On the other hand, in Section 3.2, we have seen that an H-Pd-I species, besides ensuing from oxidative alkoxy carbonylation

of acetylenic substrates (Section 3.1, Scheme 33), can also be easily formed from PdI<sub>2</sub>, CO, and H<sub>2</sub>O (Section 3.2, Scheme 42). Taken these considerations in mind, the reactivity that we observed with terminal alkynes when we allowed them to react with CO in the presence of the PdI<sub>2</sub>/KI catalytic system and small amounts of water in aprotic dioxane as the solvent can be easily interpreted (Scheme 47) [108,109,113]. Under these conditions, in fact, 2(5*H*)-furanones, corresponding to reductive carbonylation of the triple bond, were selectively obtained in good yields, with simultaneous oxidation of one equiv of CO to CO<sub>2</sub> [113] (Scheme 47). This transformation may be interpreted as occurring through a sequence of mechanistic steps involving the following: (a) formation of H-Pd-I from PdI<sub>2</sub>, CO, and H<sub>2</sub>O, with elimination of one equiv of HI and CO<sub>2</sub>; (b) formation of acylpalladium intermediate **V** from the terminal alkyne, PdI<sub>2</sub>, CO, and H<sub>2</sub>O; (c) formal reduction of **V** by H-Pd-I with formation of acylpalladiumhydride species **VI**; (d) lactonization of **VI** by intramolecular hydride attack to the carbonyl to give **III**; and (e) protonolysis of **III** by HI with regeneration of PdI<sub>2</sub> (Scheme 47) [108,109].




**Scheme 47.** Synthesis of 2(5*H*)-furanones by PdI<sub>2</sub>-catalyzed reductive carbonylation of terminal alkynes [108,109,113].

It is worth noting that we were able to observe the Pd-H bond by NMR in dioxane-*d*<sub>8</sub> (broad signal at −4.3 ppm) [109]. Moreover, an interesting phenomenon was observed when the same process was carried out in the presence of added CO<sub>2</sub> in excess (40 atm); in this case, in fact, the H-Pd-I species was partly reconverted to PdI<sub>2</sub> (through CO<sub>2</sub> insertion into the Pd-H bond, followed by protonolysis), with formation of maleic anhydrides besides furanones [108,109].

### 3.5. Propargylic Rearrangement Followed by Heterocyclization

Recently, an interesting process was realized by carbonylating 2-(alkynylthio)benzimidazoles under non-oxidative conditions, with the selective formation of 2-methyl-1-thia-4a,9-diazafluoren-4-ones (Scheme 48) [114,115]. Product formation corresponded to some kind of structural rearrangement, most likely occurring through Pd(0)-promoted propargyl-allene rearrangement. Palladium(0), under the reaction conditions, could be easily formed by the reaction of PdI<sub>2</sub> with CO and traces of water, as we have already seen in Section 3.3 (Scheme 42). The propargyl-allene rearrangement of the substrate was favored by the possibility to eliminate a thiolate as a leaving group, as shown in Scheme 48. The thiolate group then attacked the allenylpalladium intermediate to give a palladacycle, from which the final product was formed by reductive elimination (Scheme 48) [115].



**Scheme 48.** Synthesis of 2-methyl-1-thia-4a,9-diazafluoren-4-ones by PdI<sub>2</sub>-catalyzed carbonylation of 2-(alkynylthio)benzimidazoles [114,115].

## 4. Conclusions

In conclusion, we have seen how PdI<sub>2</sub>-based catalysis is very effective and versatile for realizing a variety of important carbonylation reactions, including oxidative carbonylations (of simple and functionalized alkynes, amines, diamines, β-amino alcohols, 2-aminophenols, 1,2- and 1,3-diols), additive carbonylations (of simple and functionalized alkynes), substitutive carbonylation of allylic alcohols, and reductive carbonylation of terminal alkynes.

PdI<sub>2</sub>-based catalysis works very nicely in different classical organic solvents, either nucleophilic (such as alcohols) or non-nucleophilic (such as *N,N*-dimethylacetamide, 1,2-dimethoxyethane, and 1,4-dioxane). Moreover, it can also efficiently occur, under suitable circumstances, in water as the cosolvent and in nonconventional solvents, such as ionic liquids.

The results achieved so far make this kind of catalysis one of the most important and versatile methods in carbonylation chemistry, and its use in the future, possibly in combination with

other cocatalysts or promoters, will allow achieving novel and even more complex carbonylative organic transformations.

**Author Contributions:** conceptualization, B.G. and R.M.; resources, all authors; data curation, all authors; writing—original draft preparation, B.G.; writing—review and editing, B.G.; visualization, all authors; supervision, B.G.

**Funding:** This research received no external funding.

**Acknowledgments:** Thanks are given to all the coauthors that appear in the cited references on PdI<sub>2</sub>-catalyzed carbonylation reactions.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Peng, J.B.; Geng, H.Q.; Wu, X.F. The Chemistry of CO: Carbonylation. *Chem* **2019**, *5*, 526–552. [[CrossRef](#)]
2. Ma, K.; Martin, B.S.; Yin, X.; Dai, M. Natural product syntheses via carbonylative cyclizations. *Nat. Prod. Rep.* **2019**, *36*, 174–219. [[CrossRef](#)] [[PubMed](#)]
3. Nielsen, D.U.; Neumann, K.T.; Lindhardt, A.T.; Skrydstrup, T. Recent developments in carbonylation chemistry using [<sup>13</sup>C]CO, [<sup>11</sup>C]CO, and [<sup>14</sup>C]CO. *J. Label. Compd. Radiopharm.* **2019**, *61*, 949–987. [[CrossRef](#)] [[PubMed](#)]
4. Li, Y.; Hu, Y.; Wu, X.F. Non-noble metal-catalysed carbonylative transformations. *Chem. Soc. Rev.* **2018**, *47*, 172–194. [[CrossRef](#)] [[PubMed](#)]
5. Peng, J.B.; Qi, X.; Wu, X.F. Recent Achievements in Carbonylation Reactions: A Personal Account. *Synlett* **2017**, *28*, 175–194.
6. Shen, C.; Wu, X.-F. Palladium-Catalyzed Carbonylative Multicomponent Reactions. *Chem. Eur. J.* **2017**, *23*, 2973–2987. [[CrossRef](#)]
7. Pineiro, M.; Dias, L.D.; Damas, L.; Aquino, G.L.; Calvete, M.J.; Pereira, M.M. Microwave irradiation as a sustainable tool for catalytic carbonylation reactions. *Inorganica Chim. Acta* **2017**, *455*, 364–377. [[CrossRef](#)]
8. Song, H.; Kang, M.; Jin, R.; Jin, F.; Chen, J. Application of Ionic Liquids to the Carbonylation Reactions. *Progr. Chem.* **2016**, *28*, 1313–1327.
9. Gehrtz, P.H.; Hirschbeck, V.; Ciszek, B.; Fleischer, I. Carbonylations of Alkenes in the Total Synthesis of Natural Compounds. *Synthesis* **2016**, *48*, 2679.
10. Wu, X.F. Palladium-catalyzed carbonylative transformation of aryl chlorides and aryl tosylates. *RSC Adv.* **2016**, *6*, 83831–83837. [[CrossRef](#)]
11. Wu, X.F.; Beller, M. Transition Metal Catalyzed Carbonylative Synthesis of Heterocycles. In *Topics in Heterocyclic Chemistry*; Heterocycles, H., Ed.; Springer: Berlin, Germany, 2016; Volume 42.
12. Kalck, P.; Urrutigoity, M. Recent improvements in the alkoxy carbonylation reaction catalyzed by transition metal complexes. *Inorganica Chim. Acta* **2015**, *431*, 110–121. [[CrossRef](#)]
13. Wu, L.; Fang, X.; Liu, Q.; Jackstell, R.; Beller, M.; Wu, X.F. Palladium-Catalyzed Carbonylative Transformation of C(sp<sup>3</sup>)-X Bonds. *ACS Catal.* **2014**, *4*, 2977–2989. [[CrossRef](#)]
14. Gadge, S.T.; Bhanage, B.M. Recent developments in palladium catalysed carbonylation reactions. *RSC Adv.* **2014**, *4*, 10367. [[CrossRef](#)]
15. Wu, X.F.; Neumann, H.; Beller, M. Synthesis of Heterocycles via Palladium-Catalyzed Carbonylations. *Chem. Rev.* **2013**, *113*, 1–35. [[CrossRef](#)]
16. Wu, X.F.; Neumann, H.; Beller, M. Palladium-Catalyzed Oxidative Carbonylation Reactions. *ChemSusChem* **2013**, *6*, 229–241. [[CrossRef](#)]
17. Wu, X.F.; Neumann, H. Ruthenium and Rhodium-Catalyzed Carbonylation Reactions. *ChemCatChem* **2012**, *4*, 447–458. [[CrossRef](#)]
18. Liu, Q.; Zhang, H.; Lei, A. Oxidative Carbonylation Reactions: Organometallic Compounds (R—M) or Hydrocarbons (R—H) as Nucleophiles. *Angew. Chem. Int. Ed.* **2011**, *50*, 10788–10799. [[CrossRef](#)]
19. Omae, I. Transition metal-catalyzed cyclocarbonylation in organic synthesis. *Coord. Chem. Rev.* **2011**, *255*, 139–160. [[CrossRef](#)]
20. Barnard, C.F.J. Palladium-Catalyzed Carbonylation—A Reaction Come of Age. *Organometallics* **2008**, *27*, 5402–5422. [[CrossRef](#)]

21. Kollár, L. *Modern Carbonylation Methods*; Wiley-VCH: Weinheim, Germany, 2008.
22. Beller, M. *Catalytic Carbonylation Reaction*; Springer: Berlin, Germany, 2006.
23. Gabriele, B. Recent advances in the PdI<sub>2</sub>-catalyzed carbonylative synthesis of heterocycles from acetylenic substrates: A personal account. *Targets Heterocycl Syst.* **2018**, *22*, 41–55.
24. Gabriele, B. Synthesis of Heterocycles by Palladium-Catalyzed Carbonylative Reactions. In *Advances in Transition-Metal Mediated Heterocyclic Synthesis*; Solé, D., Fernández, I., Eds.; Academic Press-Elsevier: London, UK, 2018; Volume 3, pp. 55–127.
25. Gabriele, B.; Mancuso, R.; Salerno, G. Oxidative Carbonylation as a Powerful Tool for the Direct Synthesis of Carbonylated Heterocycles. *Eur. J. Org. Chem.* **2012**, *2012*, 6825–6839. [[CrossRef](#)]
26. Gabriele, B.; Salerno, G.; Costa, M. Oxidative Carbonylations. *Top. Organomet. Chem.* **2006**, *18*, 239–272.
27. Gabriele, B.; Salerno, G. PdI<sub>2</sub>. In *e-EROS (Electronic Encyclopedia of Reagents for Organic Synthesis)*; Crich, D., Ed.; Wiley-Interscience: New York, NY, USA, 2006.
28. Gabriele, B.; Salerno, G.; Costa, M.; Chiusoli, G.P. Recent Advances in the Synthesis of Carbonyl Compounds by Palladium-Catalyzed Oxidative Carbonylation Reactions of Unsaturated Substrates. *Curr. Org. Chem.* **2004**, *35*, 919–946. [[CrossRef](#)]
29. Gabriele, B.; Salerno, G.; Costa, M. PdI<sub>2</sub>-Catalyzed Synthesis of Heterocycles. *Synlett* **2004**, *2004*, 2468–2483. [[CrossRef](#)]
30. Gabriele, B.; Salerno, G.; Costa, M.; Chiusoli, G.P. Recent developments in the synthesis of heterocyclic derivatives by PdI<sub>2</sub>-catalyzed oxidative carbonylation reactions. *J. Organomet. Chem.* **2003**, *687*, 219–228. [[CrossRef](#)]
31. Gabriele, B.; Costa, M.; Salerno, G.; Chiusoli, G.P. A new synthesis of trimethyl aconitate by palladium-catalysed triple carbonylation of propynyl alcohol. *J. Chem. Soc. Chem. Commun.* **1992**, *1992*, 1007–1008. [[CrossRef](#)]
32. Gabriele, B.; Costa, M.; Salerno, G.; Chiusoli, G.P. An efficient and selective palladium-catalysed oxidative dicarbonylation of alkynes to alkyl- or aryl-maleic esters. *J. Chem. Soc. Perkin Trans. 1* **1994**, *1994*, 83–87. [[CrossRef](#)]
33. Gabriele, B.; Veltri, L.; Salerno, G.; Costa, M.; Chiusoli, G.P. Synthesis of Maleic Anhydrides and Maleic Acids by Palladium Catalyzed Oxidative Dicarbonylation of Alk-1-yne. *Eur. J. Org. Chem.* **2003**, *2003*, 1722–1728. [[CrossRef](#)]
34. Zhu, C.; Yang, B.; Bäckvall, J.-E. Highly Selective Cascade C–C Bond Formation via Palladium-Catalyzed Oxidative Carbonylation–Carbocyclization–Carbonylation–Alkynylation of Enallenes. *J. Am. Chem. Soc.* **2015**, *137*, 11868–11871. [[CrossRef](#)]
35. Yang, B.; Qiu, Y.; Jiang, T.; Wulff, W.D.; Yin, X.; Zhu, C.; Bäckvall, J.E.; Bäckvall, J.E.; Bäckvall, J. Enantioselective Palladium-Catalyzed Carbonylative Carbocyclization of Enallenes via Cross-Dehydrogenative Coupling with Terminal Alkynes: Efficient Construction of  $\alpha$ -Chirality of Ketones. *Angew. Chem.* **2017**, *129*, 4606–4610. [[CrossRef](#)]
36. Zhu, C.; Yang, B.; Qiu, Y.; Bäckvall, J.E. Highly Selective Construction of Seven-Membered Carbocycles by Olefin-Assisted Palladium-Catalyzed Oxidative Carbocyclization-Alkoxy-carbonylation of Bisallenes. *Angew. Chem.* **2016**, *128*, 14617–14620. [[CrossRef](#)]
37. Gabriele, B.; Costa, M.; Salerno, G.; Chiusoli, G.P. Stereoselective synthesis of  $\beta$ -lactones containing  $\alpha$ -Z-alkoxy-carbonylmethylene chains by palladium-catalysed oxidative carbonylation of tertiary  $\alpha$ -hydroxyalkynes. *J. Chem. Soc. Chem. Commun.* **1994**, *1994*, 1429–1430. [[CrossRef](#)]
38. Gabriele, B.; Salerno, G.; De Pascali, F.; Costa, M.; Chiusoli, G.P. Stereoselective synthesis of (Z)- $\alpha$ -(alkoxy-carbonyl)methylene  $\beta$ - and  $\gamma$ -lactones by palladium-catalysed oxidative carbonylation of alkynols. *J. Chem. Soc. Perkin Trans. 1* **1997**, *1997*, 147–154. [[CrossRef](#)]
39. Bonardi, A.; Costa, M.; Gabriele, B.; Salerno, G.; Chiusoli, G.P. Versatile Synthesis of  $\beta$ -Lactams,  $\gamma$ -Lactams or Oxazolines by Palladium-catalysed Oxidative Carbonylation of 1-Substituted Prop-2-ynylamines. *Tetrahedron Lett.* **1995**, *36*, 7495–7498. [[CrossRef](#)]
40. Gabriele, B.; Salerno, G.; Veltri, L.; Costa, M.; Massera, C. Stereoselective Synthesis of (E)-3-(Methoxycarbonyl)methylene-1,3-dihydroindol-2-ones by Palladium-Catalyzed Oxidative Carbonylation of 2-Ethynylanilines. *Eur. J. Org. Chem.* **2001**, *2001*, 4607–4613. [[CrossRef](#)]

41. Gabriele, B.; Salerno, G.; Fazio, A.; Veltri, L. Versatile Synthesis of Pyrrole-2-acetic Esters and (Pyridine-2-one)-3-acetic amides by Palladium-Catalyzed, Carbon Dioxide-Promoted Oxidative Carbonylation of (Z)-(2-En-4-ynyl)amines. *Adv. Synth. Catal.* **2006**, *348*, 2212–2222. [[CrossRef](#)]
42. Mancuso, R.; Raut, D.S.; Marino, N.; De Luca, G.; Giordano, C.; Catalano, S.; Barone, I.; Andò, S.; Gabriele, B. A Palladium-Catalyzed Carbonylation Approach to Eight-Membered Lactam Derivatives with Antitumor Activity. *Chem. A Eur. J.* **2016**, *22*, 3053–3064. [[CrossRef](#)]
43. Mancuso, R.; Veltri, L.; Russo, P.; Grasso, G.; Cuocci, C.; Romeo, R.; Gabriele, B. Palladium-Catalyzed Carbonylative Synthesis of Functionalized Benzimidazopyrimidinones. *Synthesis* **2018**, *50*, 267–277.
44. Gabriele, B.; Salerno, G.; De Pascali, F.; Tomasi Scianò, G.; Costa, M.; Chiusoli, G.P. Novel synthesis of furan-2-acetic esters by palladium-catalysed oxidative cyclization-alkoxycarbonylation of (Z)-2-En-4-yn-1-ols. *Tetrahedron Lett.* **1997**, *38*, 6877–6880. [[CrossRef](#)]
45. Gabriele, B.; Salerno, G.; De Pascali, F.; Costa, M.; Chiusoli, G.P. An Efficient and General Synthesis of Furan-2-acetic Esters by Palladium-Catalyzed Oxidative Carbonylation of (Z)-2-En-4-yn-1-ols. *J. Org. Chem.* **1999**, *64*, 7693–7699. [[CrossRef](#)]
46. Gabriele, B.; Salerno, G.; De Pascali, F.; Costa, M.; Chiusoli, G.P. Palladium-catalysed synthesis of 2E-[(methoxycarbonyl)methylene]tetrahydrofurans: Oxidative cyclization-methoxycarbonylation of 4-yn-1-ols versus cycloisomerization-hydromethoxylation. *J. Organomet. Chem.* **2000**, *593*, 409–415. [[CrossRef](#)]
47. Bacchi, A.; Costa, M.; Della Ca', N.; Gabriele, B.; Salerno, G.; Cassoni, S. Heterocyclic Derivative Synthesis by Palladium-Catalyzed Oxidative Cyclization-Alkoxycarbonylation of Substituted  $\gamma$ -Oxoalkynes. *J. Org. Chem.* **2005**, *70*, 4971–4979. [[CrossRef](#)]
48. Della Ca', N.; Campanini, F.; Gabriele, B.; Salerno, G.; Massera, C.; Costa, M. Cascade Reactions: Catalytic Synthesis of Functionalized 1,3-Dihydroisobenzofuran and Tetrahydrofuran Derivatives by Sequential Nucleophilic Ring Opening-Heterocyclization- Oxidative Carbonylation of Alkynylloxiranes. *Adv. Synth. Catal.* **2009**, *351*, 2423–2432. [[CrossRef](#)]
49. Bacchi, A.; Chiusoli, G.P.; Costa, M.; Gabriele, B.; Righi, C.; Salerno, G. Palladium-catalysed sequential carboxylation-alkoxycarbonylation of acetylenic amines. *Chem. Commun.* **1997**, *1997*, 1209–1210. [[CrossRef](#)]
50. Chiusoli, G.; Costa, M.; Gabriele, B.; Salerno, G. Sequential reaction of carbon dioxide and carbon monoxide with acetylenic amines in the presence of a palladium catalyst. *J. Mol. Catal. A Chem.* **1999**, *143*, 297–310. [[CrossRef](#)]
51. Bacchi, A.; Chiusoli, G.P.; Costa, M.; Sani, C.; Gabriele, B.; Salerno, G. Nitrogen heterocycles by palladium-catalysed oxidative cyclization-alkoxycarbonylation of acetylenic ureas. *J. Organomet. Chem.* **1998**, *562*, 35–43. [[CrossRef](#)]
52. Bacchi, A.; Costa, M.; Gabriele, B.; Pelizzi, G.; Salerno, G. Efficient and General Synthesis of 5-(Alkoxycarbonyl)methylene-3-oxazolines by Palladium-Catalyzed Oxidative Carbonylation of Prop-2-ynylamides. *J. Org. Chem.* **2002**, *67*, 4450–4457. [[CrossRef](#)]
53. Gabriele, B.; Salerno, G.; Fazio, A.; Campana, F.B. Unprecedented carbon dioxide effect on a Pd-catalysed oxidative carbonylation reaction: A new synthesis of pyrrole-2-acetic esters. *Chem. Commun.* **2002**, *2002*, 1408–1409. [[CrossRef](#)]
54. Bacchi, A.; Costa, M.; Della Ca', N.; Fabbriatore, M.; Fazio, A.; Gabriele, B.; Nasi, C.; Salerno, G. Synthesis of 1-(Alkoxycarbonyl)methylene-1,3-dihydroisobenzofurans and 4-(Alkoxycarbonyl)benzo[c]pyrans by Palladium-Catalysed Oxidative Carbonylation of 2-Alkynylbenzyl Alcohols, 2-Alkynylbenzaldehydes and 2-Alkynylphenyl Ketones. *Eur. J. Org. Chem.* **2004**, *2004*, 574–585. [[CrossRef](#)]
55. Gabriele, B.; Veltri, L.; Maltese, V.; Spina, R.; Mancuso, R.; Salerno, G. Versatile Synthesis of Isoquinolines and Isochromenes by Pd-Catalyzed Oxidative Carbonylation of (2-Alkynyl)benzylideneamine Derivatives. *Eur. J. Org. Chem.* **2011**, *2011*, 5626–5635. [[CrossRef](#)]
56. Costa, M.; Della Ca', N.; Gabriele, B.; Massera, C.; Salerno, G.; Soliani, M. Synthesis of 4H-3,1-Benzoxazines, Quinazolin-2-ones, and Quinoline-4-ones by Palladium-Catalyzed Oxidative Carbonylation of 2-Ethynylaniline Derivatives. *J. Org. Chem.* **2004**, *69*, 2469–2477. [[CrossRef](#)]
57. Pancrazzi, F.; Motti, E.; Costa, M.; Mancuso, R.; Gabriele, B.; Della Ca', N. (Z)-4-(Carbomethoxymethylene)-2-(4-fluorophenyl)-4H-benzo[d][1,3] oxazine. *Molbank* **2017**, *2017*, M927. [[CrossRef](#)]
58. Gabriele, B.; Mancuso, R.; Salerno, G.; Lupinacci, E.; Ruffolo, G.; Costa, M. Versatile Synthesis of Quinoline-3-carboxylic Esters and Indol-2-Acetic Esters by Palladium-Catalyzed Carbonylation of 1-(2-Aminoaryl)-2-yn-1-ols. *J. Org. Chem.* **2008**, *73*, 4971–4977. [[CrossRef](#)]

59. Gabriele, B.; Veltri, L.; Salerno, G.; Mancuso, R.; Costa, M. Multicomponent Cascade Reactions: A Novel and Expedient Approach to Functionalized Indoles by an Unprecedented Nucleophilic Addition-Heterocyclization-Oxidative Alkoxyacylation Sequence. *Adv. Synth. Catal.* **2010**, *352*, 3355–3363. [[CrossRef](#)]
60. Gabriele, B.; Veltri, L.; Mancuso, R.; Salerno, G.; Costa, M. A General Synthesis of Indole-3-carboxylic Esters by Palladium-Catalyzed Direct Oxidative Carbonylation of 2-Alkynylaniline Derivatives. *Eur. J. Org. Chem.* **2012**, *2012*, 2549–2559. [[CrossRef](#)]
61. Mancuso, R.; Ziccarelli, I.; Armentano, D.; Marino, N.; Giofrè, S.V.; Gabriele, B. Divergent Palladium Iodide Catalyzed Multicomponent Carbonylative Approaches to Functionalized Isoindolinone and Isobenzofuranimine Derivatives. *J. Org. Chem.* **2014**, *79*, 3506–3518. [[CrossRef](#)]
62. Araniti, F.; Mancuso, R.; Ziccarelli, I.; Sunseri, F.; Abenavoli, M.R.; Gabriele, B. 3-[(Methoxycarbonyl)methylene]isobenzofuran-1-imines as a New Class of Potential Herbicides. *Molecules* **2014**, *19*, 8261–8275. [[CrossRef](#)]
63. Gabriele, B.; Veltri, L.; Mancuso, R.; Salerno, G.; Maggi, S.; Aresta, B.M. A Palladium Iodide-Catalyzed Carbonylative Approach to Functionalized Pyrrole Derivatives. *J. Org. Chem.* **2012**, *77*, 4005–4016. [[CrossRef](#)]
64. Gabriele, B.; Veltri, L.; Mancuso, R.; Plastina, P.; Salerno, G.; Costa, M. Palladium-Catalyzed Oxidative Heterocyclodehydration-Alkoxyacylation of 3-Yne-1,2-diols: A Novel and Expedient Approach to Furan-3-carboxylic Esters. *Tetrahedron Lett.* **2010**, *51*, 1663–1665. [[CrossRef](#)]
65. Gabriele, B.; Mancuso, R.; Maltese, V.; Veltri, L.; Salerno, G. Synthesis of Furan-3-carboxylic and 4-Methylene-4,5-dihydrofuran-3-carboxylic Esters by Direct Palladium Iodide Catalyzed Oxidative Carbonylation of 3-Yne-1, 2-diol Derivatives. *J. Org. Chem.* **2012**, *77*, 8657–8668. [[CrossRef](#)]
66. Mancuso, R.; Ziccarelli, I.; Fini, F.; Della Ca', N.; Marino, N.; Carfagna, C.; Gabriele, B. A Regio- and Stereoselective Carbonylative Approach to Alkyl (Z)-2-[3-Oxoisobenzofuran-1-(3H)-ylidene] acetates. *Adv. Synth. Catal.* **2019**, *361*, 690–695. [[CrossRef](#)]
67. Acerbi, A.; Carfagna, C.; Costa, M.; Mancuso, R.; Gabriele, B.; Della Ca', N. An Unprecedented Pd-Catalyzed Carbonylative Route to Fused Furo[3,4-b]indol-1-ones. *Chem. A Eur. J.* **2018**, *24*, 4835–4840. [[CrossRef](#)]
68. Mancuso, R.; Miliè, R.; Palumbo Piccionello, A.; Olivieri, D.; Della Ca', N.; Carfagna, C.; Gabriele, B. Catalytic Carbonylative Double Cyclization of 2-(3-Hydroxy-1-yn-1-yl) phenols in Ionic Liquids Leading to Furobenzofuranone Derivatives. *J. Org. Chem.* **2019**, *84*, 7303–7311. [[CrossRef](#)]
69. Mancuso, R.; Ziccarelli, I.; Chimento, A.; Marino, N.; Della Ca', N.; Sirianni, R.; Pezzi, V.; Gabriele, B. Catalytic Double Cyclization Process for Antitumor Agents against Breast Cancer Cell Lines. *iScience* **2018**, *3*, 279–288. [[CrossRef](#)]
70. Gabriele, B.; Salerno, G.; Veltri, L.; Costa, M. Synthesis of 2-ynamides by direct palladium-catalyzed oxidative aminocarbonylation of alk-1-yne. *J. Organomet. Chem.* **2001**, *622*, 84–88. [[CrossRef](#)]
71. Gabriele, B.; Salerno, G.; Plastina, P.; Costa, M.; Crispini, A. Expedient Synthesis of 4-Dialkylamino-5H-furan-2-ones by One-Pot Sequential Pd-Catalyzed Oxidative Carbonylation of 2-Yn-1-ols—Conjugate Addition-Lactonization. *Adv. Synth. Catal.* **2004**, *35*, 351–358. [[CrossRef](#)]
72. Plastina, P.; Gabriele, B.; Salerno, G. Palladium-Catalyzed Oxidative Aminocarbonylation of Alkynols. *Synthesis* **2007**, *2007*, 3083–3087. [[CrossRef](#)]
73. Gabriele, B.; Plastina, P.; Salerno, G.; Costa, M. A New Synthesis of 4-Dialkylamino-1, 5-dihydropyrrol-2-ones by Pd-Catalyzed Oxidative Aminocarbonylation of 2-Ynylamines. *Synlett* **2005**, *2005*, 935–938. [[CrossRef](#)]
74. Gabriele, B.; Salerno, G.; Plastina, P. A New Synthesis of 2-[(Dialkylcarbonyl)methylene]Tetrahydrofuran Derivatives by Pd-Catalyzed Oxidative Aminocarbonylation of 4-yn-1-ols. *Lett. Org. Chem.* **2004**, *1*, 134–136. [[CrossRef](#)]
75. Gabriele, B.; Salerno, G.; Veltri, L.; Mancuso, R.; Li, Z.; Crispini, A.; Bellusci, A. A New Synthesis of 2,3-Dihydrobenzo[1,4]dioxine and 3,4-Dihydro-2H-benzo[1,4]oxazine Derivatives by Tandem Palladium-Catalyzed Oxidative Aminocarbonylation—Cyclization of 2-Prop-2-ynyloxyphenols and 2-Prop-2-ynyloxyanilines. *J. Org. Chem.* **2006**, *71*, 7895–7898. [[CrossRef](#)]
76. Gabriele, B.; Plastina, P.; Salerno, G.; Mancuso, R. An Easy and Convenient Synthesis of 2-Furan-2-ylacetamides by Sequential Palladium-Catalyzed Oxidative Aminocarbonylation of (Z)-2-En-4-yn-1-ols/Conjugate Addition/Aromatization. *Synthesis* **2006**, *2006*, 4247–4251. [[CrossRef](#)]

77. Gabriele, B.; Mancuso, R.; Zicarelli, I.; Salerno, G. A new approach to isoindolinone derivatives by sequential palladium iodide-catalyzed oxidative aminocarbonylation–heterocyclization of 2-ethynylbenzamides. *Tetrahedron Lett.* **2012**, *53*, 6694–6696. [[CrossRef](#)]
78. Giofrè, S.V.; Cirimi, S.; Mancuso, R.; Nicolò, F.; Lanza, G.; Legnani, L.; Campisi, A.; Chiacchio, M.A.; Navarra, M.; Gabriele, B.; et al. Synthesis of spiro[isoindole-1,5'-isoxazolidin]-3(2H)-ones as potential inhibitors of the MDM2-p53 interaction. *Beilstein J. Org. Chem.* **2016**, *12*, 2793–2807. [[CrossRef](#)]
79. Veltri, L.; Giofrè, S.V.; Devo, P.; Romeo, R.; Dobbs, A.P.; Gabriele, B. A Palladium Iodide-Catalyzed Oxidative Aminocarbonylation–Heterocyclization Approach to Functionalized Benzimidazoimidazoles. *J. Org. Chem.* **2018**, *83*, 1680–1685. [[CrossRef](#)]
80. Veltri, L.; Grasso, G.; Rizzi, R.; Mancuso, R.; Gabriele, B. Palladium-Catalyzed Carbonylative Multicomponent Synthesis of Functionalized Benzimidazothiazoles. *Asian J. Org. Chem.* **2016**, *5*, 560–567. [[CrossRef](#)]
81. Gabriele, B.; Veltri, L.; Mancuso, R.; Carfagna, C. Cascade Reactions: A Multicomponent Approach to Functionalized Indane Derivatives by a Tandem Palladium-Catalyzed Carbamoylation/Carbocyclization Process. *Adv. Synth. Catal.* **2014**, *356*, 2547–2558. [[CrossRef](#)]
82. Gabriele, B.; Plastina, P.; Salerno, G.; Mancuso, R.; Costa, M. An Unprecedented Pd-Catalyzed, Water-Promoted Sequential Oxidative Aminocarbonylation-Cyclocarbonylation Process Leading to 2-Oxazolidinones. *Org. Lett.* **2007**, *9*, 3319–3322. [[CrossRef](#)]
83. Mancuso, R.; Maner, A.; Zicarelli, I.; Pomelli, C.; Chiappe, C.; Della Ca', N.; Veltri, L.; Gabriele, B. Auto-Tandem Catalysis in Ionic Liquids: Synthesis of 2-Oxazolidinones by Palladium-Catalyzed Oxidative Carbonylation of Propargylic Amines in EmimEtSO<sub>4</sub>. *Molecules* **2016**, *21*, 897. [[CrossRef](#)]
84. Veltri, L.; Mancuso, R.; Altomare, A.; Gabriele, B. Divergent Multicomponent Tandem Palladium-Catalyzed Aminocarbonylation-Cyclization Approaches to Functionalized Imidazothiazinones and Imidazothiazoles. *ChemCatChem* **2015**, *7*, 2206–2213. [[CrossRef](#)]
85. Maffei, M.; Giacoia, G.; Mancuso, R.; Gabriele, B.; Motti, E.; Costa, M.; Della Ca', N. A highly efficient Pd/CuI-catalyzed oxidative alkoxy carbonylation of  $\alpha$ -olefins to unsaturated esters. *J. Mol. Catal. A Chem.* **2017**, *426*, 435–443. [[CrossRef](#)]
86. Gabriele, B.; Mancuso, R.; Salerno, G.; Costa, M. A novel and efficient method for the Pd-catalysed oxidative carbonylation of amines to symmetrically and unsymmetrically substituted ureas. *Chem. Commun.* **2003**, *2003*, 486–487. [[CrossRef](#)]
87. Gabriele, B.; Salerno, G.; Mancuso, R.; Costa, M. Efficient Synthesis of Ureas by Direct Palladium-Catalyzed Oxidative Carbonylation of Amines. *J. Org. Chem.* **2004**, *69*, 4741–4750. [[CrossRef](#)]
88. Della Ca', N.; Bottarelli, P.; Dibenedetto, A.; Aresta, M.; Gabriele, B.; Salerno, G.; Costa, M. Palladium-catalyzed synthesis of symmetrical urea derivatives by oxidative carbonylation of primary amines in carbon dioxide medium. *J. Catal.* **2011**, *282*, 120–127.
89. Gabriele, B.; Mancuso, R.; Veltri, L.; Della Ca', N. Polemic against conclusions drawn in “Palladium/iodide catalyzed oxidative carbonylation of aniline to diphenylurea: Effect of ppm amounts of iron salts” (*J. Catal.* **2019**, *369* (2019) 257–266). *J. Catal.* **2019**, in press. [[CrossRef](#)]
90. Mancuso, R.; Raut, D.S.; Della Ca', N.; Fini, F.; Carfagna, C.; Gabriele, B. Catalytic Oxidative Carbonylation of Amino Moieties to Ureas, Oxamides, 2-Oxazolidinones, and Benzoxazolones. *ChemSusChem* **2015**, *8*, 2204–2211. [[CrossRef](#)]
91. Gabriele, B.; Salerno, G.; Brindisi, D.; Costa, M.; Chiusoli, G.P. Synthesis of 2-Oxazolidinones by Direct Palladium-Catalyzed Oxidative Carbonylation of 2-Amino-1-alkanols. *Org. Lett.* **2000**, *2*, 625–627. [[CrossRef](#)]
92. Gabriele, B.; Mancuso, R.; Salerno, G.; Costa, M. An Improved Procedure for the Palladium-Catalyzed Oxidative Carbonylation of  $\beta$ -Amino Alcohols to Oxazolidin-2-ones. *J. Org. Chem.* **2003**, *68*, 601–604. [[CrossRef](#)]
93. Mancuso, R.; Miliè, R.; Zicarelli, I.; Novello, M.; Della Ca', N.; Gabriele, B. (S)-4-Isopropyl-5,5-diphenyloxazolidin-2-one. *Molbank* **2018**, *2018*, M1017. [[CrossRef](#)]
94. Gabriele, B.; Mancuso, R.; Salerno, G.; Ruffolo, G.; Costa, M.; Dibenedetto, A. A novel and efficient method for the catalytic direct oxidative carbonylation of 1,2- and 1,3-diols to 5-membered and 6-membered cyclic carbonates. *Tetrahedron Lett.* **2009**, *50*, 7330–7332. [[CrossRef](#)]
95. Gabriele, B.; Mancuso, R.; Salerno, G.; Veltri, L.; Costa, M.; Dibenedetto, A. A General and Expedient Synthesis of 5- and 6-Membered Cyclic Carbonates by Palladium-Catalyzed Oxidative Carbonylation of 1,2- and 1,3-Diols. *ChemSusChem* **2011**, *4*, 1778–1786. [[CrossRef](#)]

96. Gabriele, B.; Salerno, G.; Costa, M.; Chiusoli, G.P. Combined oxidative and reductive carbonylation of terminal alkynes with palladium iodide-thiourea catalysts. *J. Organomet. Chem.* **1995**, *503*, 21–28. [[CrossRef](#)]
97. Queirolo, M.; Vezzani, A.; Mancuso, R.; Gabriele, B.; Costa, M.; Della Ca', N. Neutral vs anionic palladium iodide-catalyzed carbonylation of terminal arylacetylenes. *J. Mol. Catal. A Chem.* **2015**, *398*, 115–126. [[CrossRef](#)]
98. Bonardi, A.; Costa, M.; Gabriele, B.; Salerno, G.; Chiusoli, G.P. Coupling of oxidative and reductive processes: Catalytic carbonylation of acetals of prop-2-ynal. *J. Chem. Soc. Chem. Commun.* **1994**, *1994*, 2429–2430. [[CrossRef](#)]
99. Costa, M.; De Souza Santos, L.; Chiusoli, G.P.; Gabriele, B.; Salerno, G. Palladium-catalyzed ring closure to a naphthofuranoneacetic ester by selective carbonylation of diacetylenic precursors. *J. Mol. Catal.* **1993**, *78*, 151–158. [[CrossRef](#)]
100. De Munno, G.; Gabriele, B.; Salerno, G. X-ray structure of palladium (II) tetrakis-thiourea iodide, a catalyst for carbonylation reactions. *Inorg. Chim. Acta* **1995**, *234*, 181–183. [[CrossRef](#)]
101. Gabriele, B.; Mancuso, R.; Salerno, G.; Plastina, P. A Novel Palladium-Catalyzed Dicarboxylation Process Leading to Coumarins. *J. Org. Chem.* **2008**, *73*, 756–759. [[CrossRef](#)]
102. Araniti, F.; Mancuso, R.; Lupini, A.; Giofrè, S.V.; Sunseri, F.; Gabriele, B.; Abenavoli, M.R. Phytotoxic Potential and Biological Activity of Three Synthetic Coumarin Derivatives as New Natural-Like Herbicides. *Molecules* **2015**, *20*, 17883–17902. [[CrossRef](#)]
103. Gabriele, B.; Mancuso, R.; Salerno, G.; Veltri, L. Sequential homobimetallic catalysis: An unprecedented tandem Pd(0)-catalysed deprotection—Pd(II)-catalysed heterocyclization reaction leading to benzofurans. *Chem. Commun.* **2005**, *2005*, 271–273. [[CrossRef](#)]
104. Gabriele, B.; Mancuso, R.; Salerno, G.; Costa, M. Cascade Reactions: Sequential Homobimetallic Catalysis Leading to Benzofurans and  $\beta$ ,  $\gamma$ -Unsaturated Esters. *Adv. Synth. Catal.* **2006**, *37*, 1101–1109. [[CrossRef](#)]
105. Mancuso, R.; Gabriele, B. The Sequential Homobimetallic Catalysis Concept Applied to the Synthesis of Benzofuran Derivatives (Highlight). *Chem. Heterocycl. Compd.* **2014**, *50*, 160–170. [[CrossRef](#)]
106. Giordano, C.; Rovito, D.; Barone, I.; Mancuso, R.; Bonofiglio, D.; Giordano, F.; Catalano, S.; Gabriele, B.; Andò, S. Benzofuran-2-acetic ester derivatives induce apoptosis in breast cancer cells by upregulating p21 Cip/WAF1 gene expression in p53-independent manner. *DNA Repair* **2017**, *51*, 20–30. [[CrossRef](#)]
107. Iannazzo, D.; Pistone, A.; Celesti, C.; Triolo, C.; Patanè, S.; Giofrè, S.V.; Romeo, R.; Ziccarelli, I.; Mancuso, R.; Gabriele, B.; et al. A Smart Nanovector for Cancer Targeted Drug Delivery Based on Graphene Quantum Dots. *Nanomaterials* **2019**, *9*, 282. [[CrossRef](#)]
108. Gabriele, B.; Salerno, G.; Costa, M.; Chiusoli, G.P. Palladium-catalysed formation of maleic anhydrides from CO, CO<sub>2</sub> and alk-1-yne. *Chem. Commun.* **1999**, *1999*, 1381–1382. [[CrossRef](#)]
109. Chiusoli, G.P.; Costa, M.; Cucchia, L.; Gabriele, B.; Salerno, G.; Veltri, L. Carbon dioxide effect on palladium-catalyzed sequential reactions with carbon monoxide, acetylenic compounds and water. *J. Mol. Catal. A Chem.* **2003**, *204*, 133–142. [[CrossRef](#)]
110. Gabriele, B.; Mancuso, R.; Salerno, G.; Costa, M. Cascade Reactions: A New Synthesis of 2-Benzofuran-2-ylacetamides by Sequential Pd(0)-Catalyzed Deallylation—Pd(II)-Catalyzed Aminocarbonylative Heterocyclization of 1-(2-Allyloxyaryl)-2-yn-1-ols. *J. Org. Chem.* **2007**, *72*, 9278–9282. [[CrossRef](#)]
111. Gabriele, B.; Mancuso, R.; Lupinacci, E.; Salerno, G.; Veltri, L. Tandem catalysis in ionic liquids: A recyclable catalytic synthesis of benzofuran derivatives. *Tetrahedron* **2010**, *66*, 6156–6161. [[CrossRef](#)]
112. Gabriele, B.; Salerno, G.; Costa, M.; Chiusoli, G.P. A simple catalytic system for the substitutive carbonylation of allyl alcohols to  $\beta$ ,  $\gamma$ -unsaturated acids or esters. *J. Mol. Catal. A Chem.* **1996**, *111*, 43–48. [[CrossRef](#)]
113. Gabriele, B.; Salerno, G.; Costa, M.; Chiusoli, G.P. A new regioselective synthesis of 3-substituted furan-2(5H)-ones by palladium-catalysed reductive carbonylation of alk-1-yne. *Tetrahedron Lett.* **1999**, *40*, 989–990. [[CrossRef](#)]
114. Veltri, L.; Paladino, V.; Plastina, P.; Gabriele, B. A Palladium Iodide-Catalyzed Cyclocarbonylation Approach to Thiadiazafuorenones. *J. Org. Chem.* **2016**, *81*, 6106–6111. [[CrossRef](#)]
115. Veltri, L.; Amuso, R.; Cuocci, C.; Vitale, P.; Gabriele, B. Palladium-Catalyzed Cyclocarbonylation Approach to Thiadiazafuorenones: A Correction. *J. Org. Chem.* **2019**, *84*, 8743–8749. [[CrossRef](#)]

