



# UNIVERSITÀ DI PARMA

## ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Rational Design and Synthesis of Thioridazine Analogues as Enhancers of the Antituberculosis Therapy

This is the peer reviewed version of the following article:

*Original*

Rational Design and Synthesis of Thioridazine Analogues as Enhancers of the Antituberculosis Therapy / Pieroni, Marco; Machado, Diana; Azzali, Elisa; Santos Costa, Sofia; Couto, Isabel; Costantino, Gabriele; Viveiros, Miguel. - In: JOURNAL OF MEDICINAL CHEMISTRY. - ISSN 0022-2623. - 58:15(2015), pp. 5842-5853. [10.1021/acs.jmedchem.5b00428]

*Availability:*

This version is available at: 11381/2822802 since: 2021-10-14T10:43:34Z

*Publisher:*

American Chemical Society

*Published*

DOI:10.1021/acs.jmedchem.5b00428

*Terms of use:*

Anyone can freely access the full text of works made available as "Open Access". Works made available

*Publisher copyright*

note finali coverpage

(Article begins on next page)

## Rational Design and Synthesis of Thioridazine Analogues as Enhancers of the Antituberculosis Therapy

MARCO PIERONI, Diana Machado, Elisa Azzali, Sofia Santos Costa, Isabel Couto, Gabriele Costantino, and Miguel Viveiros

*J. Med. Chem.*, **Just Accepted Manuscript** • DOI: 10.1021/acs.jmedchem.5b00428 • Publication Date (Web): 21 Jul 2015

Downloaded from <http://pubs.acs.org> on July 27, 2015

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



# Rational Design and Synthesis of Thioridazine Analogues as Enhancers of the Anti-TB Therapy

*Marco Pieroni,<sup>†,#,\*</sup> Diana Machado,<sup>§,#</sup> Elisa Azzali,<sup>†</sup> Sofia Santos Costa,<sup>§</sup> Isabel Couto,<sup>§</sup>  
Gabriele Costantino<sup>†</sup> and Miguel Viveiros<sup>§,\*</sup>*

<sup>†</sup>Dipartimento di Farmacia, P4T group, University of Parma, Parco Area delle Scienze 27/A,  
Parma, 43124

<sup>§</sup>Grupo de Micobactérias, Unidade de Microbiologia Médica, Global Health and Tropical  
Medicine (GHTM), Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa  
(IHMT, UNL), Rua da Junqueira, 100, 1349-008 Lisbon, Portugal

KEYWORDS. Efflux pumps, Thioridazine, Tuberculosis, Antimicrobial Resistance, Medicinal  
Chemistry.

## Abstract

Tuberculosis, caused by *Mycobacterium tuberculosis*, is still one of the leading infectious diseases globally. Therefore, novel approaches are needed to face this disease. Efflux pumps are known to contribute to the emergence of *M. tuberculosis* drug resistance. Thioridazine has shown good anti-TB properties both *in vitro* and *in vivo*, likely due to its capacity to inhibit efflux mechanisms. Here we report the design and synthesis of a number of putative efflux inhibitors inspired by the structure of thioridazine. Compounds were evaluated for their *in vitro* and *ex vivo* activity against *M. tuberculosis* H37Rv. Compared to the parent molecule, some of the compounds synthesized showed higher efflux inhibitory capacity, less cytotoxicity, and a remarkable synergistic effect with anti-TB drugs both *in vitro* and in human macrophages, demonstrating their potential to be used as co-adjuvants for the treatment of tuberculosis.

## Introduction

Tuberculosis (TB) is a highly contagious airborne disease resulting from infection with *Mycobacterium tuberculosis*. The World Health Organization (WHO) currently estimates that one-third of the world's population is infected with *M. tuberculosis*, and in 2013, approximately 210 000 human lives were lost to TB.<sup>1</sup> The global health impact of *M. tuberculosis* is exacerbated by HIV co-infection, making TB one of the major AIDS-related opportunistic diseases.<sup>2</sup> This shocking failure to control TB is due to the combination of many factors, among which: the non-compliance of the patients to the required six months of multidrug therapy; the difficulty to provide it in a sustained and properly-dosed regimen, particularly in those parts of the world that are the least equipped to deal with the disease.<sup>3</sup> These issues have altogether led to the emergence of multidrug resistant (MDR) TB and extensively drug-resistant (XDR) TB. As

1  
2  
3 such, to address this global health crisis, the process of drug discovery must rely both on novel  
4 chemical anti-TB entities<sup>4,5</sup> and on an innovative approach that rationally prevents the raise of  
5 drug resistance.  
6  
7  
8

9  
10 After many years without delivering any new anti-TB drug, the discovery of bedaquiline for  
11 the treatment of MDR-TB, and a nourished pipeline of compounds already in clinical trials  
12 (<http://www.newtbdrugs.org/pipeline.php>) have renewed hope for the treatment of TB and  
13 especially MDR-TB. However, the lack of innovation in the overall therapeutic approach makes  
14 quite predictable that mutation-based resistant strains will inexorably emerge also for these novel  
15 drugs. The discovery of bedaquiline<sup>6</sup> and pretomanid<sup>7</sup> has showed that the whole cell phenotypic  
16 assay of large libraries of compounds, and the wise medicinal chemistry efforts toward the  
17 improvement of these molecules, are reliable methodological approaches to drive the research  
18 towards novel anti-TB agents.<sup>8</sup> Indeed, most of the efforts of some of the authors of this work  
19 has moved through this path.<sup>9-19</sup> While this approach has proved to be correct, it is nevertheless  
20 incomplete. It does not consider that, besides mutations, other mechanisms such as efflux play an  
21 important role in the development of resistance in *M. tuberculosis*.<sup>20-22</sup>  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

38 Efflux pumps are involved in several physiological processes such as detoxification of  
39 intracellular metabolites and cellular homeostasis.<sup>23</sup> Simultaneously, they contribute to the  
40 development of intrinsic and acquired drug resistance in many bacterial pathogens. Usually they  
41 confer low-to-intermediate level of resistance, however, the constant pressure of subinhibitory  
42 concentrations of the antibiotic promotes the selection of spontaneous mutants.<sup>20</sup> Therefore: *a)*  
43 efflux pumps are effectors of the innate drug-resistance machinery; *b)* they are crucial in  
44 conferring a low-level of drug resistance; *c)* they contribute to lower the concentration of the  
45 drug inside the mycobacterial cell, enabling the emergence of a high-level of resistance; *d)* it has  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

been demonstrated that efflux inhibitors (EIs) are able to improve or restore the activity of old drugs towards which the bacteria had become resistant.<sup>24</sup>

In spite of this knowledge, the extent to which an antimicrobial compound is a substrate of efflux is seldom measured, and it is not considered an important parameter in the hit-to-lead process for the optimization of novel anti-TB drugs. The recent discovery by Lee and colleagues of spectinamides as anti-TB drug candidates,<sup>25</sup> also by virtue of their scarce tendency to be extruded by efflux pumps, confirms the importance of taking into account efflux when the design of new drugs is planned.<sup>26</sup> In addition to these findings, although the mechanism of adaptation of *M. tuberculosis* to the hostile environment is still a matter of debate, Adams and colleagues have recently reported that bacterial efflux pumps mediate multi-drug tolerance and the bacterial survival in macrophages, leading to a long-lasting therapy.<sup>27,28</sup>

It is therefore possible to claim that inhibition of efflux would lead to a number of beneficial effects: *a)* increasing the activity of the anti-TB drugs subject to efflux; *b)* keeping the concentration of a given drug at the therapeutic dose, minimizing the possibility to select mutants; *c)* drastically shorten the duration of treatment by reducing multi-drug tolerance. This could be priceless in order to repurpose old drugs and preserve the efficacy of those novel compounds, such as bedaquiline, that are either next to be introduced in the anti-TB arena or that already are.

Thioridazine (TZ) is a neuroleptic with efflux inhibitory properties,<sup>29</sup> that has attracted the attention of many researchers due to its ability to hit drug resistant strains.<sup>30</sup> In a clinical trial in Argentina, TZ was used for the treatment of patients with XDR-TB in combination with anti-TB drugs, with good results.<sup>31</sup> It has been demonstrated that TZ has a dual effect against *M. tuberculosis*: it is an inhibitor of bacterial efflux pumps, and, also, it induces the acidification of

1  
2  
3 the phagosome, contributing to restriction of intracellular *M. tuberculosis* growth.<sup>30,32–34</sup> TZ is  
4  
5 not the only inhibitor of efflux investigated for the treatment of TB. Recently chlorpromazine,  
6  
7 another neuroleptic with efflux inhibitory properties, was tested along with its metabolites  
8  
9 against *M. smegmatis* and exhibited synergistic activity when administered in combination with  
10  
11 known anti-TB drugs.<sup>35,36</sup> Also the antiarrhythmic verapamil is an efflux pump inhibitor that has  
12  
13 been shown to have an antimicrobial-potentiating effect in the treatment of *M. tuberculosis* both  
14  
15 *in vitro* and *in vivo*.<sup>37</sup> It was shown that inhibition of the efflux pumps of *M. tuberculosis* by  
16  
17 verapamil reduces the bacterial drug tolerance induced in the intracellular compartment inside  
18  
19 the macrophages and in zebrafish granuloma-like lesions. Unfortunately, the use of TZ in the  
20  
21 therapy of TB is severely limited by its neuroleptic properties and a number of severe toxicity  
22  
23 issues. Inspired by these considerations, we herein report the design and synthesis of a number of  
24  
25 TZ derivatives, and their biological evaluation as efflux inhibitors and synergistic effect with  
26  
27 known anti-TB drugs both *in vitro* and on infected human monocyte-derived macrophages.  
28  
29 Among the novel compounds, when compared to the parent molecule TZ, compound **10** showed  
30  
31 higher efflux inhibitory ability, less toxicity, and a remarkable synergistic effect with first- and  
32  
33 second-line drugs, especially rifampicin. As such, it may represent an interesting starting point  
34  
35 for the development of more potent and less toxic efflux inhibitors to be used in combination  
36  
37 with the current first- and second-line anti-TB drugs.  
38  
39  
40  
41  
42  
43  
44

### 45 46 **Rational design**

47  
48 Since the lack of detailed information about the structure of the efflux systems, the new  
49  
50 compounds were prepared following a Ligand-Based Drug Design approach. TZ was used as  
51  
52 template, given its well-known and deeply investigated anti-TB properties when administered in  
53  
54 combination with first-line drugs, as mentioned above.<sup>29,32–34,38,39</sup> The aim of this work was  
55  
56  
57  
58  
59  
60

1  
2  
3 therefore to chemically modify TZ in order to obtain a compound with improved adjuvant  
4 properties while limiting the onset of side effects, especially at the CNS level. We reasoned that,  
5  
6 in first place, it was necessary to escape from the phenothiazine core because, according to the  
7  
8 SAR of antipsychotic drugs,<sup>40</sup> it is crucial in conferring the desired neuroleptic activity.  
9  
10 Moreover, although with a few exceptions,<sup>41</sup> a series of TZ analogues in which the  
11  
12 phenothiazine core was kept intact has already been reported as inhibitors of efflux in other  
13  
14 bacteria, but with scarce results.<sup>38,42</sup> In addition to this, efflux systems are known to be affine to  
15  
16 charged quaternary ammonium salts such as ethidium bromide (EtBr) and berberine, and to  
17  
18 molecules bearing basic amino groups (TZ, VP, reserpine), that likely occur to be protonated at  
19  
20 the acidic pH of the sites infected by *M. tuberculosis*. Given these premises, chemical  
21  
22 manipulation was carried out keeping intact the N-methylpiperidine moiety, whereas modifying  
23  
24 the nature of the heterocyclic attached to it and the length of the spacer.  
25  
26  
27  
28  
29  
30

31  
32 Several heterocycles, embodying a variable number of rings, were attached by the nitrogen  
33  
34 atom to an N-methyl-2-piperidinyl appendage through an ethylene linker (compounds **9-15**, chart  
35  
36 1).

37  
38 [Insert chart 1 here]

39  
40  
41 The carbazole (**9**) and the acridone (**12**), that are tricyclic flat structures, were chosen for their  
42  
43 close resemblance with the phenothiazine core. The indole (**10**), and the benzimidazole (**11**) are  
44  
45 among the most privileged scaffolds used in medicinal chemistry, and could afford extended  
46  
47 chemical manipulation of the series. The diphenylamines **13** and **14** were synthesized to check  
48  
49 whether the planar structure of the phenothiazine could be disrupted without loss of the  
50  
51 inhibitory activity. The pyrrolidine **15** was prepared to check whether a small ring could still be a  
52  
53 suitable substrate of efflux. Once the preliminary assays had corroborated the design of  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 compound **9**, other appendages were attached to the carbazole core in place of the N-methyl-2-  
4 piperidinyl one. To start, a less polar and flatter substrate such as the 2-ethylpyridine (**18**) was  
5 attached to the carbazole nitrogen; then, the distance between the nitrogen of carbazole and that  
6 of the appendage was shortened by one methylene unit in compound **19**.  
7  
8  
9  
10  
11

12 The synthesis of compounds **5** and **6** was accomplished in the attempt to couple a moiety that  
13 is thought to be responsible for the EI activity to a potent anti-TB agent;<sup>11,16</sup> this would have led  
14 to the synthesis of the first rationally designed TB-GEI (Growth and Efflux Inhibitor), which  
15 could represent a superior therapeutic option for the treatment of TB. Indeed, the recent finding  
16 that bedaquiline, along with its antibacterial activity, is also a weak inhibitor of efflux  
17 mechanisms, accounts for this reasoning.<sup>43</sup> Finally, since the recent interest in the use of VP and  
18 its derivatives in the treatment of TB,<sup>44,45</sup> we have tried to merge the two structures of TZ and  
19 VP in one hybrid molecule (Figure 1), and therefore compounds **7** and **8** (chart 1) were  
20 synthesized and tested.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32

33  
34 [Insert Figure 1 here]  
35

### 36 Chemistry

37  
38 The synthesis of derivatives **5–15** was achieved through reaction of the suitable heterocyclic  
39 precursor and 2-(2-chloroethyl)-N-methylpiperidine hydrochloride, in dry DMF using NaH in  
40 excess, either at room temperature or heating for several hours, when necessary (Scheme 1).  
41  
42 When not commercially available, the synthesis of the heterocyclic precursors was achieved in  
43 good yields according to a well-established procedure, as in the case of intermediates **1** and  
44 **2**.<sup>13,16</sup> Compound **3**<sup>46</sup> was prepared reacting 3,4-dimethoxyaniline with triethylamine and 2-  
45 iodopropane in refluxing methanol, differently from what already reported. For the synthesis of  
46 compound **18**, first 2-(hydroxymethyl)-pyridine was treated with *p*-toluensulfonyl chloride and  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 triethylamine in dichloromethane to give the tosyl derivative **16**, and then the ameliorated  
4  
5 leaving group is displaced by carbazole according to the conditions reported by Cheng (Scheme  
6  
7 2).<sup>47</sup> In the case of compound **19**, carbazole was first treated with dichloroethane to give N-  
8  
9 (chloroethyl)carbazole **17**, that was reacted with piperidine in DMF at room temperature using  
10  
11 K<sub>2</sub>CO<sub>3</sub> as the base to give **19** (Scheme 2).<sup>48</sup> For the synthesis of compound **8**, 3,4-  
12  
13 dimethoxyphenyl acetonitrile was first treated with sodium hydride and 2-bromopropane to give  
14  
15 intermediate **4**;<sup>49</sup> several attempts to couple this derivative with 2-(2-chloroethyl)-N-  
16  
17 methylpiperidine hydrochloride with the use of various bases, performed according to different  
18  
19 literature protocols,<sup>49-51</sup> failed to yield the desired product. The synthesis was finally  
20  
21 accomplished by heating the two reagents in DMF in a microwave oven, using an excess of  
22  
23 sodium amide as the base. Interestingly, the same synthetic conditions did not give any product  
24  
25 with conventional heating also after many days at 120 °C. Compound **8** was tested preliminarily  
26  
27 as a racemic mixture.  
28  
29  
30  
31  
32

33  
34 [Insert Scheme 1 here]

35  
36 [Insert Scheme 2 here]

### 37 38 **Results and Discussion.**

39  
40 A total of 13 compounds were synthesized in this study (**5-15**, **18**, **19**, chart 1), that underwent  
41  
42 biological assays. In order to set up a faster and cheaper model for screening the activity of the  
43  
44 compounds, they were first evaluated for their antimycobacterial and efflux inhibitory activity  
45  
46 against *M. smegmatis* mc<sup>2</sup>155 reference strain. For those compounds showing inhibitory activity  
47  
48 higher than TZ, cytotoxicity on human-monocyte derived macrophages was assessed.  
49  
50 Compounds showing less cytotoxicity and higher efflux inhibitory activity than that of TZ, were  
51  
52  
53  
54  
55 further evaluated on *M. tuberculosis* H37rv, as described herein.  
56  
57  
58  
59  
60

### Antimycobacterial and efflux inhibitory activity against *M. smegmatis* mc<sup>2</sup>155

The compounds were first tested for their capability to inhibit the growth of *M. smegmatis* strain mc<sup>2</sup>155 (Table 1). The compounds demonstrated extremely poor or absent antimycobacterial activity against *M. smegmatis*, especially if compared with TZ, which was used as positive control. This was considered a good result, as high bactericidal/bacteriostatic activity could bias the following experiments of synergy with first- and second-line anti-TB agents. Due to the reduced solubility and the low antimycobacterial activity, it was not possible to determine the exact MIC values for some compounds (*i.e.* **5-8**, **11**, **13**, **15**, **18**, **19**). As such, when bacterial growth is still noticed at the maximum concentration tested, the MIC was indicated by “>MIC”. For instance, >256 µg/mL means that the MIC of this compound is higher than 256 µg/mL, as it was technically impossible to determine it. After determining their MICs, the compounds were tested for their capability to inhibit efflux of EtBr from *M. smegmatis* mc<sup>2</sup>155 cells by real-time fluorometry, using TZ and VP as controls (Table 1, relative final fluorescence (RFF) values). According to a widely used protocol for this kind of assay,<sup>52</sup> in order not to compromise the cellular viability, all of the compounds were tested at ½ of their MIC. For those compounds for which the exact MIC could not be determined (above reported), the last concentration value that could be technically determined was considered ½ of the MIC. Despite the small set of compounds prepared for this preliminary study, a few SAR considerations can be made. In general, the majority of the compounds bearing the N-methyl-2-piperidinyl appendage had inhibitory effect on the efflux of EtBr. We were pleased to notice that compound **7**, designed as a hybrid molecule resembling both VP and TZ, had a very good inhibitory activity of EtBr efflux. However, the same does not apply to the other hybrid compound **8**. It might be then speculated that the N-methyl-2-piperidinyl appendage must be attached, through an ethylene

1  
2  
3 linker, to another nitrogen atom of the heterocyclic, yielding a more polar structure. When the N-  
4 methyl-2-piperidinyl appendage is attached through an ethylene linker either to tricyclic  
5  
6 (compounds **9**, **12**), bicyclic (compounds **10**, **11**) or more flexible ring structures (compounds **13**,  
7  
8 **14**), a good inhibitory activity, sometimes higher than that of the parent compound TZ, is  
9  
10 noticed. However, bulkier heterocyclic (compounds **5** and **6**) or small rings (compound **15**), have  
11  
12 a detrimental impact on the efflux inhibitory properties. Surprisingly, the benzimidazole  
13  
14 derivative **11** showed an exceptionally high activity, resulting by far more potent than VP. The  
15  
16 distance between the nitrogen of the appendage and that of the heterocyclic seems to have  
17  
18 pharmacological significance, since the activity of compound **19** is considerably low. Finally, the  
19  
20 presence of a tertiary aliphatic amino moiety seems to be determinant, being compound **18** the  
21  
22 least active of the series.  
23  
24  
25  
26  
27

### 28 **Cytotoxicity evaluation of compounds 7, 9, 10, 11 and 12**

29  
30  
31 Compounds **7**, **9**, **10**, **11** and **12**, that resulted more active than TZ in inhibiting the efflux of  
32  
33 EtBr out of *M. smegmatis* cells, were evaluated against human-monocyte derived macrophages  
34  
35 to assess their *ex vivo* cytotoxicity toward eukaryotic cells (Table 1). We were pleased to notice  
36  
37 that, with the exception of compound **12** ( $IC_{50} = 0.762 \mu\text{g/mL}$ ) all of the tested compounds were  
38  
39 found to be less cytotoxic than TZ in this assay. Compounds **9** and **10**, ( $IC_{50} = 8.8 \mu\text{g/mL}$  and  
40  
41  $10.5 \mu\text{g/mL}$ , respectively) are two-fold less cytotoxic than TZ ( $IC_{50} = 5.6 \mu\text{g/mL}$ ), whereas  
42  
43 compounds **7** and **11** ( $IC_{50} = 84.8 \mu\text{g/mL}$  and  $170.8 \mu\text{g/mL}$ ) were found to be safe towards human  
44  
45 macrophages. Overall, these preliminary data clearly suggest that the replacement of the  
46  
47 phenothiazine core with other ring systems allows maintaining the efflux inhibitory activity with  
48  
49 the possibility to decrease the cytotoxicity.  
50  
51  
52  
53  
54

55 [Insert Table 1 here]  
56  
57  
58  
59  
60

1  
2  
3 [Insert Figure 2 here]  
4

5 Compounds **7**, **9**, **10** and **11**, that resulted more active than TZ in inhibiting the efflux of EtBr  
6 from *M. smegmatis*, and less cytotoxic against human-monocyte derived macrophages, were  
7 selected for additional biological studies against *M. tuberculosis* H37Rv strain. Compound **12**,  
8 although presenting higher RFF values than that of TZ, was excluded based on its higher  
9 cytotoxicity.  
10  
11

12  
13  
14  
15  
16  
17  
18 **Evaluation of compounds 7, 9, 10 and 11 antimycobacterial and efflux inhibitory activity**  
19 **against *M. tuberculosis* H37Rv**  
20

21  
22 Compounds **7**, **9**, **10** and **11** were tested against the pan-susceptible reference strain H37Rv,  
23 following the same rationale used for *M. smegmatis*, *i.e.* the determination of antimycobacterial  
24 activity and the evaluation of their effect on EtBr efflux (Table 2). As observed for *M.*  
25 *smegmatis*, the selected derivatives showed no antimycobacterial activity against *M.*  
26 *tuberculosis*, therefore excluding the possibility of bias during the following assays. With regard  
27 to their activity as inhibitors of EtBr efflux, the compounds were tested with the same procedure  
28 used as in the case of *M. smegmatis*. Compound **9** showed similar activity to that of TZ, whereas  
29 compounds **7**, **10** and **11** were found to be 2-4 fold more potent than TZ, although less active  
30 than VP (Table 2, Figure 2). Although quite active, the benzimidazole derivative **11** failed to  
31 reproduce the extraordinary efflux inhibitory effect observed against *M. smegmatis*. This is not  
32 surprising and can be somehow explained on the basis of the faster metabolism of *M. smegmatis*  
33 compared to that of *M. tuberculosis*, that accounts for the higher susceptibility to the efflux  
34 inhibitory effect. Our own experience shows that RFF values are always scaled down in *M.*  
35 *tuberculosis* in comparison to those observed for the same compounds in *M. smegmatis*, although  
36 maintaining the inhibitory relation.<sup>53,54</sup> Regardless, we can claim that, in the conditions reported,  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 all of the tested compounds have maintained their superior capability of inhibiting efflux  
4  
5 compare to TZ against *M. tuberculosis*.  
6  
7

8 [Insert Table 2 here]  
9

10 **Evaluation of synergistic effect of the selected TZ analogues in combination with first-**  
11 **and second-line anti-TB drugs and determination of the fractional inhibitory concentration**  
12 **(FIC).**  
13  
14  
15

16  
17 Once that the inhibition of efflux was confirmed, the study went ahead evaluating compounds  
18 **7, 9, 10** and **11** for their effect in combination with the current first-line (isoniazid, INH;  
19 rifampicin, RIF) and second-line (amikacin, AMK; ofloxacin, OFX) drugs used for the treatment  
20 of TB. Their effect was also tested in combination with EtBr, to correlate its efflux with the  
21 efflux of the anti-TB drugs.<sup>55</sup> The MICs of INH, RIF, AMK and OFX were calculated in the  
22 absence and in the presence of scalar sub-inhibitory concentrations of each compound against *M.*  
23 *tuberculosis* H37Rv, that is known to have intrinsic and readable efflux activity of these anti-TB  
24 drugs.<sup>22,54</sup> (Table 3).  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 [Insert Table 3 here]  
37  
38

39 Initially, compounds **7, 9, 10** and **11** were tested at  $\frac{1}{2}$  of their MIC (see Table 3 for details),  
40 and then, according to the protocol widely used for this sort of investigations,<sup>52</sup> they were tested  
41 by two-dimensional broth microdilution checkerboard assay, to evaluate the extent of the  
42 synergism. Also in this case, as already explained in the case of the efflux inhibition assays, for  
43 those compounds for which the exact MIC could not be determined,  $\frac{1}{2}$  of the MIC was  
44 considered the concentration that was equal to the highest concentration where it was obtained  
45 growth. When tested at these concentrations, all of the newly synthesized compounds and the  
46 controls demonstrated very high synergistic effect with both the antibiotics and EtBr. In this  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 case, the MIC values of INH, RIF, AMK, OFX and EtBr could be reduced more than 64-fold  
4  
5 (Table 3). Compound **10**, for which the inhibitory effect on efflux was already assessed, had in  
6  
7 general a very high synergistic effect with the tested drugs and, in particular, with RIF. In fact,  
8  
9 compound **10** was able to improve the activity of RIF by 4-fold at concentrations as low as 8  
10  
11  $\mu\text{g/mL}$  (approx. 1/64 of its MIC), demonstrating a potent synergistic effect. Moreover, at a  
12  
13 concentration of 64  $\mu\text{g/mL}$  (approx. 1/8 of its MIC), compound **10** was also able to improve the  
14  
15 activity of AMK and OFX by more than 128-fold, and EtBr by 8-fold (Table 3). Importantly, we  
16  
17 have noticed that the synergistic activity demonstrated by the compounds in combination with  
18  
19 EtBr correlates well with the results obtained with the inhibition of EtBr efflux by RFF.  
20  
21  
22  
23  
24

25 One may argue that TZ, considering the absolute concentrations at which the synergistic  
26  
27 activity is measured, is slightly more active than compound **10**. However, seen under a more  
28  
29 widespread perspective, compound **10** is less toxic, more active with regard the inhibition of  
30  
31 efflux, and unlikely to possess CNS-related properties. This can be considered a good  
32  
33 achievement, in particular considering that the aim of this paper was the selection of compounds  
34  
35 devoid of antimycobacterial activity, with reduced toxicity to the human macrophage and with  
36  
37 higher efflux inhibitory capability, so as to promote the antimycobacterial activity of the anti-TB  
38  
39 drugs and be considered as a future adjuvant in the fighting against TB.  
40  
41  
42  
43

44 Surprisingly, despite the high activity in inhibiting the efflux, compound **11** was found to be  
45  
46 not active in the combination assay. Compounds **7** and **9** were found to be slightly less active  
47  
48 than **10** in the combination assay; however, in the case of compound **7**, it must be noticed that it  
49  
50 is able to improve the activity of RIF by 4-fold also at very low concentrations. It is intriguing to  
51  
52 notice that compounds **7** and **9**, as compound **10**, demonstrated to have a significant synergistic  
53  
54 effect in particular with RIF. The reason of this preferred synergism is worth of further  
55  
56  
57  
58  
59  
60

1  
2  
3 investigation but, at this moment, it can only be speculated that **10**, as well as the other  
4 derivatives belonging to this series, although to a lesser extent, have high affinity for interfering  
5 with efflux systems that extrude RIF. However, it cannot be ruled out that the two derivatives  
6 interfere with other mechanism of resistance used by the cell in the presence of RIF. Finally, it is  
7 worth of consideration the fact that compounds **7**, **9** and **10** were more active in the combination  
8 assay that VP, another drug often investigated for its potential role as adjuvant in the TB  
9 treatment.  
10

11  
12 The synergistic activity between the compounds was also represented as the fractional  
13 inhibitory concentration (FIC), that is a value expressing the synergistic activity of the  
14 compounds in combination with the antibiotics and EtBr (Figure 3). FIC was determined for  
15 each anti-TB drug by dividing the MIC of each drug when used in combination, by the MIC of  
16 each drug used alone. Since a drug cannot interact with itself, the effect of a self-drug  
17 combination will always be additive, with an FIC index of 1. As such, when the FIC value is  
18 lower than 0.25, a synergistic effect is noticed between the modulator and the antimicrobial.  
19 When above 2, there is antagonism between the compounds tested, whereas within these two FIC  
20 values, indifference is noticed.<sup>24</sup>  
21

22 [Insert figure 3 here]  
23

24 Isobolograms were constructed, by plotting changes in the MIC of antibiotics as a function of  
25 the tested compounds concentration (Figure 3 and S1 – example for compound **10**). As it can be  
26 seen, for the majority of the combinations of compound **10** with the antibiotics synergy was  
27 observed, with FIC values ranging from 1 to not possible to determine (ND) due to the very high  
28 synergistic effect, depending on the antibiotic combination (see legend of Figure 3 supporting  
29 information for details).  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



### Evaluation of the intracellular anti-TB activity of the selected TZ analogues.

Finally, we evaluated the intracellular antimycobacterial activity of compounds **7**, **9**, **10** and **11** alone and in combination with INH or RIF against *M. tuberculosis*-infected human monocyte-derived macrophages (Figure 4). This assay is a fair reproduction of what actual happens during the TB infection, and therefore it can help in obtaining a clearer idea of the potential of our compounds. INH and RIF were used at  $\frac{1}{2}$  of its MIC (0.05 and 0.5  $\mu\text{g/mL}$ , respectively), therefore having no direct inhibitory effect on the bacteria at these concentrations. The compounds **7**, **9**, **10** and **11** were tested at nontoxic concentrations (16  $\mu\text{g/mL}$ , 0.078  $\mu\text{g/mL}$ ; 1.25  $\mu\text{g/mL}$ , and 40  $\mu\text{g/mL}$ , respectively) determined against human-monocyte derived macrophages (Figure S2). Above these concentrations, macrophage viability is reduced below 90%, nullifying the assay. At these concentrations, the compounds tested showed to be devoid of any antimycobacterial activity against intracellular *M. tuberculosis*. Consistent with the findings above reported, we were pleased to notice that the co-administration of the compounds with INH or RIF (at a non-antibiotic inhibitory concentration) led to a strong enhancement of the killing activity of the human macrophages against the H37Rv strain. This is comparable with the results previously obtained with TZ and other efflux inhibitors that has been shown to have a dual bacterial-host boosting effect against *M. tuberculosis*.<sup>56</sup>

### Conclusion

The contribution of efflux systems to *M. tuberculosis* drug resistance and tolerance has been increasingly recognized, although only a few medicinal chemistry efforts have been made towards the synthesis of *M. tuberculosis* efflux inhibitors. Despite the amount of literature establishing TZ as a valuable agent against TB, the numerous toxicity issues have always hampered its safe administration as an adjuvant of TB therapy. To our knowledge, this is the first

1  
2  
3 medicinal chemistry campaign aimed at modifying TZ with regard to its potential as adjuvant of  
4 TB therapy, with specific assays such as the determination of the efflux inhibition and the  
5 synergistic activity in combination with known anti-TB drugs both *in vitro* and *ex vivo*.  
6  
7 Compared to TZ, the representative derivatives tested were found to be less toxic toward human  
8 macrophages, ranging from two-fold (compound **9** and **10**) to more than 50-fold (compound **11**).  
9  
10 Compound **7**, intended as a hybrid of TZ and VP, showed a higher inhibitory activity than that of  
11 TZ itself and much less toxicity, therefore confirming the rationale of the design. More  
12 importantly, we have escaped from the phenothiazine scaffold, thus assuming a reduction of the  
13 onset of side effects at the CNS level. The newly synthesized compounds showed high  
14 synergistic activity when tested in combination with some of the most important first-line (INH,  
15 RIF) and second-line (AMK, OFX) drugs used for the treatment of TB. In particular compound  
16 **10** was able to improve the activity of RIF at concentration as low as approximately 1/64 of its  
17 MIC. Additionally, these compounds were able to improve the activity of INH and RIF against  
18 intracellular *M. tuberculosis*. Considering the data on inhibition of EtBr efflux, coupled to the  
19 low antimycobacterial activity, it might be claimed that the synergetic activity is due to their  
20 ability to hamper the intrinsic mycobacterial drug efflux.  
21  
22

23  
24  
25 Finally, we confirmed *M. smegmatis* mc<sup>2</sup>155 as a suitable surrogate of *M. tuberculosis* for the  
26 preliminary assessment the efflux inhibitory activity, at a lower cost and time consuming rate.  
27  
28 Altogether, these findings provide a solid base to further investigate compound **10** as a booster of  
29 the antimycobacterial chemotherapy when associated with first- and second-line drugs, by virtue  
30 of its capability to block the intrinsic efflux activity of mycobacteria. This might represent a  
31 lateral approach toward the cure of TB. Studies expanding this versatile series and its mechanism  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 of action and activity toward *M. tuberculosis* drug resistant strains, especially resistant to  
4  
5  
6 isoniazid and rifampicin (MDR-TB), are currently underway in our research groups.  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## EXPERIMENTAL PART

### Chemistry

**General information.** All the reagents were purchased from Sigma-Aldrich, Alfa-Aesar and Enamine at reagent purity and, unless otherwise noted, were used without any further purification. Dry solvents used in the reactions were obtained by distillation of technical grade materials over appropriate dehydrating agents. MCRs were performed using CEM Microwave Synthesizer-Discover model. Reactions were monitored by thin layer chromatography on silica gel-coated aluminium foils (silica gel on Al foils, SUPELCO Analytical, Sigma-Aldrich) at both 254 and 365 nm wavelengths. When indicated, intermediates and final products were purified through silica gel flash chromatography (silica gel, 0.040-0.063 mm), using appropriate solvent mixtures.

$^1\text{H-NMR}$  and  $^{13}\text{C-NMR}$  spectra were recorded on a BRUKER AVANCE spectrometer at 300, 400 and 100 MHz respectively, with TMS as internal standard.  $^1\text{H-NMR}$  spectra are reported in this order: multiplicity and number of protons. Standard abbreviation indicating the multiplicity was used as follows: s = singlet, d = doublet, dd = doublet of doublets, t = triplet, q = quadruplet, m = multiplet and br = broad signal. HPLC/MS experiments were performed with HPLC: Agilent 1100 series, equipped with a Waters Symmetry C18, 3.5  $\mu\text{m}$ , 4.6 mm x 75 mm column and MS: Applied Biosystem/MDS SCIEX, with API 150EX ion source. HRMS experiments were performed with LTQ ORBITRAP XL THERMO.

All compounds were tested as 95% purity samples or higher (by HPLC/MS).

The synthesis and chemical characterizations of intermediates 1-4, 16, 17 is reported in the Supporting Information.

1  
2  
3       **General procedure for the synthesis of compounds 5, 6, 9-15, 18.** To a suspension of  
4 sodium hydride (60% suspension in mineral oil, 3 eq) in DMF (5 mL/mmol) was added the  
5 suitable heterocyclic compound (1 eq) at 0 °C. Reaction mixture was stirred for 15 minutes at the  
6 same temperature and then 2-(2-chloroethyl)-piperidine hydrochloride (1.3 eq) was added  
7 portion wise. The mixture was allowed to react at room temperature until consumption of the  
8 starting material. In some cases (**5, 6, 13**) reflux heating was necessary. After quenching with  
9 water, the mixture was extracted with ethyl acetate (3 × 10 mL), and the organic layers were  
10 washed with water and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced  
11 pressure. The crude material was purified through flash chromatography. Reaction times and  
12 conditions, yields, purification methods and other analytical data are reported in the supporting  
13 information.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29       **N-isopropyl-3,4-dimethoxy-N-(2-(1-methylpiperidin-2-yl)ethyl)aniline (7).** To a solution of  
30 **3** (60 mg; 0.38 mmol) in dry toluene (2 mL) was added sodium amide (120 mg; 3.07 mmol) and  
31 2-(2-chloroethyl)piperidine hydrochloride (244 mg; 1.23 mmol). The reaction mixture was  
32 stirred and heated in a microwave oven at 140 °C for 30 minutes. After quenching with brine (25  
33 mL), the mixture was extracted with ethyl acetate (3 × 10 mL), and the combined organic layers  
34 were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure. The crude material  
35 was then purified through flash chromatography eluting with dichloromethane/methanol 9:1.  
36 Yield: 16%. <sup>1</sup>H-NMR (400 MHz-CDCl<sub>3</sub>): □ 1.10-1.15 (m, 6 H); 1.20-1.87 (m; 8H), 2.30-2.35  
37 (bs; 5H); 2.95-3.10 (m; 2H), 3.15-3.25 (m, 1H); 3.65-3.75 (m, 1H); 3.84 (s, 3H); 3.87, (s, 3H);  
38 6.41 (d, J = 9 Hz, 1H); 6.49 (s, 1H); 6.79 (d, J = 9 Hz, 1H). <sup>13</sup>C-NMR (100.6 MHz-CDCl<sub>3</sub>): □  
39 19.4, 20.2, 23.3, 28.3, 41.3, 53.8, 55.9, 56.3, 105.6, 111.9, 142.6, 143.7, 149.5. HRMS (ESI)  
40 calculated for C<sub>19</sub>H<sub>32</sub>N<sub>2</sub>O<sub>2</sub> [M + H]<sup>+</sup> 321,2464, found 321.25365.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**2-(3,4-Dimethoxyphenyl)-2-isopropyl-4-(1-methylpiperidin-2-yl)butanenitrile (8).** To a solution of **4** (70 mg; 0.31 mmol) in dry toluene (2 mL) was added sodium amide (124 mg; 3.19 mmol) and 2-(2-chloroethyl)piperidine hydrochloride (252 mg; 1.25 mmol). The reaction mixture was stirred and heated in a microwave oven at 140 °C for 30 minutes. After quenching with brine (25 mL), the mixture was extracted with ethyl acetate (3 × 10 mL), and the combined organic layers were dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. The crude material was then purified through flash column chromatography eluting with dichloromethane/methanol 9:1. Yield: 50%. <sup>1</sup>H-NMR (300 MHz-CDCl<sub>3</sub>): □ 0.8 (dd, J<sub>1</sub> = 4 Hz, J<sub>2</sub> = 7 Hz, 3 H); 1.23-1.80 (m; 11H), 1.85-2.40 (m; 8H); 2.80-2.95 (m; 1H), 3.88 (s, 3H); 3.90 (s, 3H); 6.80-7.00 (m, 3H). <sup>13</sup>C-NMR (100.6 MHz-CDCl<sub>3</sub>): □ 18.7, 19.0, 24.0, 28.2, 29.9, 32.1, 32.5, 37.9, 53.4, 55.9, 56.0, 63.0, 109.1, 109.4, 111.0, 118.8, 121.1, 130.5, 148.3, 149.0. HRMS (ESI) calculated for C<sub>21</sub>H<sub>32</sub>N<sub>2</sub>O<sub>2</sub> [M + H]<sup>+</sup> 345,2464, found 345.25365.

**9-(2-(Piperidin-1-yl)ethyl)-9H-carbazole (19).** To a solution of piperidine (33 μl; 0.33 mmol) in DMF (15 mL/mmol) was added potassium carbonate (138 mg, 1 mmol), this mixture was stirred at room temperature for 20 minutes and then the intermediate **17** (10 mg, 0.43 mmol) was added. Reaction mixture was stirred to 60 °C overnight. After cooling, brine (25 mL) was added and the mixture was extracted with ethyl acetate (3 × 10 mL). The combined organic layers were washed with water and brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure. The crude material was purified through flash chromatography eluting with petroleum ether/ethyl acetate 95/5, yielding the title compound as a white solid. Yield: 19%. <sup>1</sup>H-NMR (400 MHz-CDCl<sub>3</sub>): □ 1.24-1.60 (m, 6H); 3.07 (bs, 2H); 3.39 (bs, 2H); 4.49 (t, J = 8 Hz, 2H); 4.62 (t, J = 8 Hz, 2H); 7.23-7.28 (m, 2H); 7.46-7.79 (m, 4H), 8.12 (d, J = 8 Hz, 2H). <sup>13</sup>C-NMR (100.6

1  
2  
3 MHz-CDCl<sub>3</sub>): □ 24.2, 25.4, 42.1, 44.7, 62.8, 108.7, 119.1, 120.3, 123.0, 125.7, 140.5, 155.0.

4  
5  
6 HRMS (ESI) calculated for C<sub>19</sub>H<sub>22</sub>N<sub>2</sub> [M + H]<sup>+</sup> 279,1783, found 279.39258

## 7 8 **Biology**

### 9 10 **Evaluation of the novel synthesized compounds against mycobacteria**

#### 11 12 **Reagents and mycobacterial strains**

13  
14  
15 Thioridazine (TZ), verapamil (VP), isoniazid (INH), rifampicin (RIF), amikacin (AMK),  
16  
17 ofloxacin (OFX), EtBr, phosphate-buffered saline (PBS), and glucose were purchased from  
18  
19 Sigma-Aldrich (St. Louis, MO, USA). All solutions were prepared in deionized water, except  
20  
21 rifampicin and the TZ analogues **5-15**, **18** and **19**, which were prepared in DMSO. All solutions  
22  
23 were prepared on the day of the experiment. Middlebrook 7H9 (MB7H9) and the OADC  
24  
25 supplement (oleic acid/albumin/dextrose/catalase) were purchased from Difco (Madrid, Spain)  
26  
27 and Becton and Dickinson (Sparks, MD, USA) respectively. The mycobacterial reference strains  
28  
29 *M. smegmatis* mc<sup>2</sup>155 ATCC700084 and *M. tuberculosis* H37Rv ATCC27294<sup>T</sup> were used to  
30  
31 evaluate the biologic activity of TZ analogues.  
32  
33  
34  
35

36  
37 The determination of the minimum inhibitory concentration (MIC), the evaluation of the  
38  
39 synergistic effect of TZ analogues **7**, **9**, **10** and **11** with selected antibiotics and EtBr by broth  
40  
41 microdilution checkerboard assay, the evaluation of efflux inhibitory activity of the compounds  
42  
43 by real-time fluorometry and the cytotoxicity evaluation are reported in detail in the Supporting  
44  
45 Information.  
46  
47

## 48 **References**

49  
50 (1) WHO | Global tuberculosis report 2014  
51  
52 [http://www.who.int/tb/publications/global\\_report/en/](http://www.who.int/tb/publications/global_report/en/) (accessed Feb 5, 2015).  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 (2) CDC - Reported Tuberculosis in the United States, 2011 - TB  
4  
5 <http://www.cdc.gov/tb/statistics/reports/2011/default.htm> (accessed Feb 5, 2015).  
6  
7

8  
9 (3) WHO | Multidrug-resistant tuberculosis (MDR-TB)  
10  
11 <http://www.who.int/tb/challenges/mdr/en/> (accessed Feb 5, 2015).  
12  
13

14 (4) Duncan, K. Progress in TB Drug Development and What Is Still Needed. *Tuberc. Edinb. Scotl.* **2003**, *83* (1-3), 201–207.  
15  
16  
17

18  
19 (5) PLOS guest blogger, Urgency Needed in the Global Response to Drug-resistant  
20 Tuberculosis [http://blogs.plos.org/speakingofmedicine/2013/11/01/urgency-needed-in-the-](http://blogs.plos.org/speakingofmedicine/2013/11/01/urgency-needed-in-the-global-response-to-drug-resistant-tuberculosis/)  
21  
22 [global-response-to-drug-resistant-tuberculosis/](http://blogs.plos.org/speakingofmedicine/2013/11/01/urgency-needed-in-the-global-response-to-drug-resistant-tuberculosis/) (accessed Mar 20, 2014).  
23  
24  
25  
26

27  
28 (6) Andries, K.; Verhasselt, P.; Guillemont, J.; Göhlmann, H. W. H.; Neefs, J.-M.; Winkler,  
29 H.; Gestel, J. V.; Timmerman, P.; Zhu, M.; Lee, E.; Williams, P.; Chaffoy, D. de; Huitric, E.;  
30 Hoffner, S.; Cambau, E.; Truffot-Pernot, C.; Lounis, N.; Jarlier, V. A Diarylquinoline Drug  
31 Active on the ATP Synthase of Mycobacterium Tuberculosis. *Science* **2005**, *307* (5707), 223–  
32 227.  
33  
34  
35  
36  
37  
38

39  
40 (7) Stover, C. K.; Warrener, P.; VanDevanter, D. R.; Sherman, D. R.; Arain, T. M.;  
41 Langhorne, M. H.; Anderson, S. W.; Towell, J. A.; Yuan, Y.; McMurray, D. N.; Kreiswirth, B.  
42 N.; Barry, C. E.; Baker, W. R. A Small-Molecule Nitroimidazopyran Drug Candidate for the  
43 Treatment of Tuberculosis. *Nature* **2000**, *405* (6789), 962–966.  
44  
45  
46  
47  
48  
49

50  
51 (8) Payne, D. J.; Gwynn, M. N.; Holmes, D. J.; Pompliano, D. L. Drugs for Bad Bugs:  
52 Confronting the Challenges of Antibacterial Discovery. *Nat. Rev. Drug Discov.* **2007**, *6* (1), 29–  
53 40.  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 (9) Lilienkampf, A.; Pieroni, M.; G. Franzblau, S.; R. Bishai, W.; P. Kozikowski, A.  
4  
5 Derivatives of 3-Isoxazolecarboxylic Acid Esters - A Potent and Selective Compound Class  
6  
7 against Replicating and Nonreplicating Mycobacterium Tuberculosis. *Curr. Top. Med. Chem.*  
8  
9 **2012**, *12* (7), 729–734.  
10  
11

12  
13 (10) Lilienkampf, A.; Pieroni, M.; Wan, B.; Wang, Y.; Franzblau, S. G.; Kozikowski, A. P.  
14  
15 Rational Design of 5-Phenyl-3-Isoxazolecarboxylic Acid Ethyl Esters as Growth Inhibitors of  
16  
17 *Mycobacterium Tuberculosis* . A Potent and Selective Series for Further Drug Development. *J.*  
18  
19 *Med. Chem.* **2010**, *53* (2), 678–688.  
20  
21  
22

23  
24 (11) Lun, S.; Guo, H.; Onajole, O. K.; Pieroni, M.; Gunosewoyo, H.; Chen, G.; Tipparaju, S.  
25  
26 K.; Ammerman, N. C.; Kozikowski, A. P.; Bishai, W. R. Indoleamides Are Active against Drug-  
27  
28 Resistant Mycobacterium Tuberculosis. *Nat. Commun.* **2013**, *4*, 2907.  
29  
30  
31

32 (12) Mao, J.; Yuan, H.; Wang, Y.; Wan, B.; Pieroni, M.; Huang, Q.; van Breemen, R. B.;  
33  
34 Kozikowski, A. P.; Franzblau, S. G. From Serendipity to Rational Antituberculosis Drug  
35  
36 Discovery of Mefloquine-Isoxazole Carboxylic Acid Esters. *J. Med. Chem.* **2009**, *52* (22), 6966–  
37  
38 6978.  
39  
40  
41

42 (13) Onajole, O. K.; Pieroni, M.; Tipparaju, S. K.; Lun, S.; Stec, J.; Chen, G.; Gunosewoyo,  
43  
44 H.; Guo, H.; Ammerman, N. C.; Bishai, W. R.; Kozikowski, A. P. Preliminary Structure–  
45  
46 Activity Relationships and Biological Evaluation of Novel Antitubercular Indolecarboxamide  
47  
48 Derivatives Against Drug-Susceptible and Drug-Resistant Mycobacterium Tuberculosis Strains.  
49  
50 *J. Med. Chem.* **2013**, *56* (10), 4093–4103.  
51  
52  
53

54  
55 (14) Pieroni, M.; Lilienkampf, A.; Wan, B.; Wang, Y.; Franzblau, S. G.; Kozikowski, A. P.  
56  
57 Synthesis, Biological Evaluation, and Structure–Activity Relationships for 5-[(*E*)-2-  
58  
59  
60

1  
2  
3 Arylethenyl]-3-Isoxazolecarboxylic Acid Alkyl Ester Derivatives as Valuable Antitubercular  
4  
5 Chemotypes. *J. Med. Chem.* **2009**, *52* (20), 6287–6296.

6  
7  
8  
9 (15) Pieroni, M.; Lilienkamp, A.; Wang, Y.; Wan, B.; Cho, S.; Franzblau, S. G.; Kozikowski,  
10  
11 A. P. NOC Chemistry for Tuberculosis-Further Investigations on the Structure-Activity  
12  
13 Relationships of Antitubercular Isoxazole-3-Carboxylic Acid Ester Derivatives. *ChemMedChem*  
14  
15 **2010**, *5* (10), 1667–1672.

16  
17  
18  
19 (16) Pieroni, M.; Tipparaju, S. K.; Lun, S.; Song, Y.; Sturm, A. W.; Bishai, W. R.;  
20  
21 Kozikowski, A. P. Pyrido[1,2-*a*]benzimidazole-Based Agents Active Against Tuberculosis (TB),  
22  
23 Multidrug-Resistant (MDR) TB and Extensively Drug-Resistant (XDR) TB. *ChemMedChem*  
24  
25 **2011**, *6* (2), 334–342.

26  
27  
28  
29 (17) Stec, J.; Huang, Q.; Pieroni, M.; Kaiser, M.; Fomovska, A.; Mui, E.; Witola, W. H.;  
30  
31 Bettis, S.; McLeod, R.; Brun, R.; Kozikowski, A. P. Synthesis, Biological Evaluation, and  
32  
33 Structure–Activity Relationships of *N*-Benzoyl-2-Hydroxybenzamides as Agents Active against  
34  
35 *P. Falciparum* (K1 Strain), Trypanosomes, and Leishmania. *J. Med. Chem.* **2012**, *55* (7), 3088–  
36  
37 3100.

38  
39  
40  
41  
42 (18) Tabarrini, O.; Sabatini, S.; Massari, S.; Pieroni, M.; Franzblau, S. G.; Cecchetti, V. 6-  
43  
44 Hydrogen-8-Methylquinolones Active Against Replicating and Non-Replicating *Mycobacterium*  
45  
46 *Tuberculosis*: Antimycobacterial 6-Desfluoroquinolones. *Chem. Biol. Drug Des.* **2012**, *80* (5),  
47  
48 781–786.

49  
50  
51  
52 (19) Tipparaju, S. K.; Joyasawal, S.; Pieroni, M.; Kaiser, M.; Brun, R.; Kozikowski, A. P. In  
53  
54 Pursuit of Natural Product Leads: Synthesis and Biological Evaluation of 2-[3-Hydroxy-2-[(3-  
55  
56 Hydroxypyridine-2-Carbonyl)amino]phenyl]benzoxazole-4-Carboxylic Acid (A-33853) and Its  
57  
58  
59  
60

1  
2  
3 Analogues: Discovery of *N*-(2-Benzoxazol-2-Ylphenyl)benzamides as Novel Antileishmanial  
4 Chemotypes. *J. Med. Chem.* **2008**, *51* (23), 7344–7347.

7  
8  
9 (20) Machado, D.; Couto, I.; Perdigão, J.; Rodrigues, L.; Portugal, I.; Baptista, P.; Veigas, B.;  
10 Amaral, L.; Viveiros, M. Contribution of Efflux to the Emergence of Isoniazid and Multidrug  
11 Resistance in Mycobacterium Tuberculosis. *PloS One* **2012**, *7* (4), e34538.

16  
17 (21) Louw, G. E.; Warren, R. M.; Gey van Pittius, N. C.; McEvoy, C. R. E.; Van Helden, P.  
18 D.; Victor, T. C. A Balancing Act: Efflux/influx in Mycobacterial Drug Resistance. *Antimicrob.*  
19 *Agents Chemother.* **2009**, *53* (8), 3181–3189.

24  
25 (22) Silva, P. E. A. da; Groll, A. V.; Martin, A.; Palomino, J. C. Efflux as a Mechanism for  
26 Drug Resistance in Mycobacterium Tuberculosis. *FEMS Immunol. Med. Microbiol.* **2011**, *63* (1),  
27 1–9.

32  
33 (23) Webber, M. A.; Piddock, L. J. V. The Importance of Efflux Pumps in Bacterial Antibiotic  
34 Resistance. *J. Antimicrob. Chemother.* **2003**, *51* (1), 9–11.

37  
38 (24) Coelho, T.; Machado, D.; Couto, I.; Maschmann, R.; Ramos, D.; von Groll, A.; Rossetti,  
39 M. L.; Silva, P. A.; Viveiros, M. Enhancement of Antibiotic Activity by Efflux Inhibitors against  
40 Multidrug Resistant Mycobacterium Tuberculosis Clinical Isolates from Brazil. *Front.*  
41 *Microbiol.* **2015**, *6*, 330.

47  
48 (25) Lee, R. E.; Hurdle, J. G.; Liu, J.; Bruhn, D. F.; Matt, T.; Scherman, M. S.; Vaddady, P.  
49 K.; Zheng, Z.; Qi, J.; Akbergenov, R.; Das, S.; Madhura, D. B.; Rathi, C.; Trivedi, A.; Villellas,  
50 C.; Lee, R. B.; Rakesh; Waidyarachchi, S. L.; Sun, D.; McNeil, M. R.; Ainsa, J. A.; Boshoff, H.  
51 I.; Gonzalez-Juarrero, M.; Meibohm, B.; Böttger, E. C.; Lenaerts, A. J. Spectinamides: A New  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Class of Semisynthetic Antituberculosis Agents That Overcome Native Drug Efflux. *Nat. Med.*  
4  
5 **2014**, *20* (2), 152–158.

6  
7  
8  
9 (26) Viveiros, M.; Pieroni, M. Spectinamides: A Challenge, a Proof, and a Suggestion. *Trends*  
10  
11 *Microbiol.* **2014**, *22* (4), 170–171.

12  
13  
14 (27) Adams, K. N.; Takaki, K.; Connolly, L. E.; Wiedenhoft, H.; Winglee, K.; Humbert, O.;  
15  
16 Edelstein, P. H.; Cosma, C. L.; Ramakrishnan, L. Drug Tolerance in Replicating Mycobacteria  
17  
18 Mediated by a Macrophage-Induced Efflux Mechanism. *Cell*, **2011**, *145* (1), 39–53.

19  
20  
21  
22 (28) Adams, K. N.; Szumowski, J. D.; Ramakrishnan, L. Verapamil, and Its Metabolite  
23  
24 Norverapamil, Inhibit Macrophage-Induced, Bacterial Efflux Pump-Mediated Tolerance to  
25  
26 Multiple Anti-Tubercular Drugs. *J. Infect. Dis.* **2014**, *210*: 456-66.

27  
28  
29  
30 (29) Rodrigues, L.; Wagner, D.; Viveiros, M.; Sampaio, D.; Couto, I.; Vavra, M.; Kern, W.  
31  
32 V.; Amaral, L. Thioridazine and Chlorpromazine Inhibition of Ethidium Bromide Efflux in  
33  
34 Mycobacterium Avium and Mycobacterium Smegmatis. *J. Antimicrob. Chemother.* **2008**, *61* (5),  
35  
36 1076–1082.

37  
38  
39  
40 (30) Martins, M.; Viveiros, M.; Kristiansen, J. E.; Molnar, J.; Amaral, L. The Curative  
41  
42 Activity of Thioridazine on Mice Infected with Mycobacterium Tuberculosis. *Vivo Athens*  
43  
44 *Greece* **2007**, *21* (5), 771–775.

45  
46  
47  
48 (31) Abbate, E.; Vescovo, M.; Natiello, M.; Cufre, M.; García, A.; Montaner, P. G.;  
49  
50 Ambroggi, M.; Ritacco, V.; Soolingen, D. van. Successful Alternative Treatment of Extensively  
51  
52 Drug-Resistant Tuberculosis in Argentina with a Combination of Linezolid, Moxifloxacin and  
53  
54 Thioridazine. *J. Antimicrob. Chemother.* **2012**, *67* (2), 473–477.  
55  
56  
57  
58  
59  
60

1  
2  
3 (32) Amaral, L.; Martins, M.; Viveiros, M. Enhanced Killing of Intracellular Multidrug-  
4 Resistant Mycobacterium Tuberculosis by Compounds That Affect the Activity of Efflux Pumps.  
5  
6

7  
8 *J. Antimicrob. Chemother.* **2007**, *59* (6), 1237–1246.  
9

10  
11 (33) Amaral, L.; Viveiros, M. Why Thioridazine in Combination with Antibiotics Cures  
12 Extensively Drug-Resistant Mycobacterium Tuberculosis Infections. *Int. J. Antimicrob. Agents*  
13  
14 **2012**, *39* (5), 376–380.  
15  
16

17  
18 (34) Martins, M.; Viveiros, M.; Couto, I.; Amaral, L. Targeting Human Macrophages for  
19 Enhanced Killing of Intracellular XDR-TB and MDR-TB. *Int. J. Tuberc. Lung Dis. Off. J. Int.*  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

30  
31  
32 (35) Kigundu, E. M.; Wasuna, A.; Warner, D. F.; Chibale, K. Pharmacologically Active  
33 Metabolites, Combination Screening and Target Identification-Driven Drug Repositioning in  
34 Antituberculosis Drug Discovery. *Bioorg. Med. Chem.* **2014**, *22* (16), 4453–4461.  
35  
36

37 (36) Kigundu, E. M.; Njoroge, M.; Singh, K.; Njuguna, N.; Warner, D. F.; Chibale, K.  
38 Synthesis and Synergistic Antimycobacterial Screening of Chlorpromazine and Its Metabolites.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

43 (37) Singh, M.; Jadaun, G. P. S.; Ramdas; Srivastava, K.; Chauhan, V.; Mishra, R.; Gupta, K.;  
44 Nair, S.; Chauhan, D. S.; Sharma, V. D.; Venkatesan, K.; Katoch, V. M. Effect of Efflux Pump  
45 Inhibitors on Drug Susceptibility of Ofloxacin Resistant Mycobacterium Tuberculosis Isolates.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

*Indian J. Med. Res.* **2011**, *133* (5), 535–540.

1  
2  
3 (38) Martins, M.; Schelz, Z.; Martins, A.; Molnar, J.; Hajös, G.; Riedl, Z.; Viveiros, M.;  
4 Yalcin, I.; Aki-Sener, E.; Amaral, L. In Vitro and Ex Vivo Activity of Thioridazine Derivatives  
5 against Mycobacterium Tuberculosis. *Int. J. Antimicrob. Agents* **2007**, *29* (3), 338–340.  
6  
7

8  
9  
10  
11 (39) Martins, M.; Viveiros, M.; Amaral, L. Inhibitors of Ca<sup>2+</sup> and K<sup>+</sup> Transport Enhance  
12 Intracellular Killing of M. Tuberculosis by Non-Killing Macrophages. *Vivo Athens Greece* **2008**,  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

(40) Aftab, D. T.; Ballas, L. M.; Loomis, C. R.; Hait, W. N. Structure-Activity Relationships  
of Phenothiazines and Related Drugs for Inhibition of Protein Kinase C. *Mol. Pharmacol.* **1991**,  
*40* (5), 798–805.

(41) Salie, S.; Hsu, N.-J.; Semanya, D.; Jardine, A.; Jacobs, M. Novel Non-Neuroleptic  
Phenothiazines Inhibit Mycobacterium Tuberculosis Replication. *J. Antimicrob. Chemother.*  
**2014**, *69* (6), 1551–1558.

(42) Takács, D.; Cerca, P.; Martins, A.; Riedl, Z.; Hajós, G.; Molnár, J.; Viveiros, M.; Couto,  
I.; Amaral, L. Evaluation of Forty New Phenothiazine Derivatives for Activity Against Intrinsic  
Efflux Pump Systems of Reference Escherichia Coli, Salmonella Enteritidis, Enterococcus  
Faecalis and Staphylococcus Aureus Strains. *In Vivo* **2011**, *25* (5), 719–724.

(43) Lu, P.; Villellas, C.; Koul, A.; Andries, K.; Lill, H.; Bald, D. The ATP Synthase Inhibitor  
Bedaquiline Interferes with Small-Molecule Efflux in Mycobacterium Smegmatis. *J. Antibiot.*  
*(Tokyo)* **2014**, *67* (12), 835–837.

1  
2  
3 (44) Gupta, S.; Tyagi, S.; Almeida, D. V.; Maiga, M. C.; Ammerman, N. C.; Bishai, W. R.  
4  
5 Acceleration of Tuberculosis Treatment by Adjunctive Therapy with Verapamil as an Efflux  
6  
7 Inhibitor. *Am. J. Respir. Crit. Care Med.* **2013**, *188* (5), 600–607.  
8  
9

10  
11 (45) Gupta, S.; Tyagi, S.; Bishai, W. R. Verapamil Increases the Bactericidal Activity of  
12  
13 Bedaquiline against Mycobacterium Tuberculosis in a Mouse Model. *Antimicrob. Agents*  
14  
15 *Chemother.* **2015**, *59* (1), 673–676.  
16  
17

18  
19 (46) Kumar, H. M. S.; Reddy, B. V. S.; Anjaneyulu, S.; Yadav, J. S. A Novel and Efficient  
20  
21 Approach to Mono-N-Alkyl Anilines via Addition of Grignard Reagents to Aryl Azides.  
22  
23 *Tetrahedron Lett.* **1999**, *40* (47), 8305–8306.  
24  
25

26  
27 (47) Cheng, K.; Kim, I. J.; Lee, M.-J.; Adah, S. A.; Raymond, T. J.; Bilsky, E. J.; Aceto, M.  
28  
29 D.; May, E. L.; Harris, L. S.; Coop, A.; Dersch, C. M.; Rothman, R. B.; Jacobson, A. E.; Rice, K.  
30  
31 C. Opioid Ligands with Mixed Properties from Substituted Enantiomeric N-Phenethyl-5-  
32  
33 Phenylmorphans. Synthesis of a M-Agonist  $\Delta$ -Antagonist and  $\Delta$ -Inverse Agonists. *Org. Biomol.*  
34  
35 *Chem.* **2007**, *5* (8), 1177–1190.  
36  
37

38  
39 (48) Bogdal, D.; Matras, K.; Sanetra, J.; Pokladko, M. Synthesis of Polymethacrylates with  
40  
41 Carbazole and Benzofuran Pendant Groups for Photovoltaic Applications. *Macromol. Symp.*  
42  
43 **2008**, *268* (1), 48–52.  
44  
45

46  
47 (49) Verbeek, J.; Syvänen, S.; Schuit, R. C.; Eriksson, J.; de Lange, E. C.; Windhorst, A. D.;  
48  
49 Luurtsema, G.; Lammertsma, A. A. Synthesis and Preclinical Evaluation of [11C]D617, a  
50  
51 Metabolite of (R)-[11C]verapamil. *Nucl. Med. Biol.* **2012**, *39* (4), 530–539.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 (50) Behrens, C. H.; Chatterjee, D.; Hajra, A. K.; He, Y.; Li, H. Y.; Priestley, T.; Rudra, S.;  
4 Saha, A. K.; Smith, R. A.; Thompson, S. K. Aminoindane Compounds and Use Thereof in  
5 Treating Pain, WO2012112969 A1 August 23, 2012.  
6  
7

8  
9  
10  
11 (51) Albani, D.; Arosio, R.; Merli, V. Process for Preparing N-Methyl-3, 4-  
12 Dimethoxyphenylethylamine, US patent 20090171110 A1, July 2, 2009.  
13  
14

15  
16  
17 (52) Ramón-García, S.; Ng, C.; Anderson, H.; Chao, J. D.; Zheng, X.; Pfeifer, T.; Av-Gay, Y.;  
18 Roberge, M.; Thompson, C. J. Synergistic Drug Combinations for Tuberculosis Therapy  
19 Identified by a Novel High-Throughput Screen. *Antimicrob. Agents Chemother.* **2011**, *55* (8),  
20 3861–3869.  
21  
22  
23

24  
25  
26  
27 (53) Rodrigues, L.; Machado, D.; Couto, I.; Amaral, L.; Viveiros, M. Contribution of Efflux  
28 Activity to Isoniazid Resistance in the Mycobacterium Tuberculosis Complex. *Infect. Genet.*  
29 *Evol. J. Mol. Epidemiol. Evol. Genet. Infect. Dis.* **2012**, *12* (4), 695–700.  
30  
31  
32

33  
34  
35 (54) Viveiros, M.; Rodrigues, L.; Martins, M.; Couto, I.; Spengler, G.; Martins, A.; Amaral, L.  
36 Evaluation of Efflux Activity of Bacteria by a Semi-Automated Fluorometric System. *Methods*  
37 *Mol. Biol. Clifton NJ* **2010**, *642*, 159–172.  
38  
39  
40

41  
42  
43 (55) Rodrigues, L.; Ramos, J.; Couto, I.; Amaral, L.; Viveiros, M. Ethidium Bromide  
44 Transport across Mycobacterium Smegmatis Cell-Wall: Correlation with Antibiotic Resistance.  
45 *BMC Microbiol.* **2011**, *11* (1), 35.  
46  
47  
48


49  
50  
51 (56) Viveiros M.; Martins, M.; Rodrigues, L.; Machado, D.; Couto, I.; Ainsa, J.; Amaral, L.  
52 Inhibitors of Mycobacterial Efflux Pumps as Potential Boosters for Anti-Tubercular Drugs.  
53 *Expert Rev Anti Infect Ther* **2012**, *10*, 983–998.  
54  
55  
56  
57  
58  
59  
60




1  
2  
3 (57) Moody, J. A. 1992. Synergy testing: broth microdilution checkerboard and broth  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

macrodilution methods, in *Clinical microbiology procedures handbook*, ed H. D. Eisenberg, American Society for Microbiology, Washington, DC, 1992; pp. 5.18.1-5.18.23.


**Figure 1.** Structure of thioridazine and verapamil. In red the backbone that the two molecules share is shown.




**Figure 2.** Effect of thioridazine (TZ), verapamil (VP) and compound 10 on the accumulation (left) and efflux (right) of ethidium bromide (EtBr) at 0.5  $\mu\text{g/mL}$  by *M. tuberculosis* H37Rv. **Left panel:** *M. tuberculosis* H37Rv was loaded with 0.5  $\mu\text{g/mL}$  of EtBr with and without compounds. Verapamil (VP) and thioridazine (TZ) were tested at  $\frac{1}{2}$  of their MIC whereas compound **10** was tested at 256  $\mu\text{g/mL}$ , (see text for details). The increase in the fluorescence is directly proportional to the amount of EtBr that is retained within the cells. **Right panel:** *M. tuberculosis* H37Rv was loaded with 0.5  $\mu\text{g/mL}$  of EtBr in the presence of VP at  $\frac{1}{2}$  of their MIC. Efflux of EtBr took place at 37  $^{\circ}\text{C}$  in the presence of glucose and was inhibited in the presence of the compounds at the same concentrations used on the accumulation assays (compare pink curve – no inhibitor – with the remaining curves – with inhibitor).




**Figure 3.** Fractional inhibitory concentration (FIC) determination for each combination against *M. tuberculosis* H37Rv. The effect of each compound on the activity of each antibiotic was represented by means of fractional inhibitory concentration (FIC) calculated as follows:  $FIC = MIC_{\text{antibacterial in combination}} / MIC_{\text{antibacterial alone}}$ . The FIC values were interpreted based on the criteria established as follows:  $FIC \leq 0.25$ , synergism;  $0.25 > FIC < 2$ , indifference and  $FIC \geq 2$ , antagonism.<sup>24</sup> The FIC value for the combinations was classified as ND (non-determinable) when the MICs of the compounds were greater than the highest, less than, or equal to the lowest concentration tested (Moody, 1992).<sup>57</sup> As such, FIC values were calculated at the following concentrations of the compounds: **7**, **10** and **11**, 128  $\mu\text{g/mL}$ ; **9**, 32  $\mu\text{g/mL}$ ; VP, 64  $\mu\text{g/mL}$ , and TZ, 3.75  $\mu\text{g/mL}$  (see Table 3 for further information).




**Figure 4.** Determination of the intracellular activity of compounds **7**, **9**, **10** and **11** against *M. tuberculosis*-infected human macrophages. For the evaluation of the intracellular synergistic activity against *M. tuberculosis*-infected human monocyte-derived macrophages, INH and RIF were used at  $\frac{1}{2}$  of its MIC (0.05 and 0.5  $\mu\text{g/mL}$ , respectively). Macrophage viability was determined after 3 days of treatment with the compounds and for the subsequent intracellular assays, these were used at concentrations that were shown to be non-toxic for the macrophages: compound **7**, 16  $\mu\text{g/mL}$ , **9**, 0.078  $\mu\text{g/mL}$ ; **10**, 1.25  $\mu\text{g/mL}$ , and **11**, 40  $\mu\text{g/mL}$ . Above these concentrations macrophage viability is reduced below 90%. The results are presented as a mean of the percentage of the survival  $\pm$  SD. (Figure S2)




Scheme 1.<sup>a,b</sup> Preparation of the thioridazine analogues.

<sup>a</sup>Reagents and conditions: a) Ammonium acetate, toluene,  $\alpha$ -benzyl-ethylacetoacetate, 150 °C, 1.5 h, 40%; b) 1-aminoadamantane, EDC·HCl, HOBT, triethylamine, DMF, rt, 24h, 49%; c) Isopropyl iodide, triethylamine, MeOH, reflux, 18h, 50%; d) Isopropyl bromide, NaH, DMF, rt, 18h, 47%; e) RNH, NaH, DMF, reflux, 18-72h, 9-70%; f) NaNH<sub>2</sub>, Toluene,  $\mu$ W, 110 °C, 30 min, 50%. <sup>b</sup>For complete structures see chart 1.

**Scheme 2.**<sup>a</sup> Preparation of the carbazole derivatives.

<sup>a</sup>Reagents and conditions: *a*) Dichloroethane, KOH, K<sub>2</sub>CO<sub>3</sub>, TBAF, 60 °C, 18h, 40%; *b*) piperidine, K<sub>2</sub>CO<sub>3</sub>, DMF, 60°C, 24h, 44%; *c*) *p*-Toluenesulfonyl chloride, triethylamine, DCM, rt, 18h, 75%; *d*) NaH, DMF, rt, 24h, 4%.

Chart 1. Structures of compounds 5-15<sup>a</sup>

<sup>a</sup>all compounds are mixture of enantiomers; <sup>b</sup>mixture of the diastereomers \*Indicates the point of attachment

**Table 1.** Screening of the antimycobacterial and efflux inhibitory activity of the compounds against *M. smegmatis* mc2 155 and their toxicity assessed by the index of cytotoxicity (IC) towards human monocyte derived macrophages.

comp	<i>M. smegmatis</i> mc <sup>2</sup> 155		IC <sub>50</sub> <sup>c</sup> μg/mL(μM)
	MIC (μg/mL) <sup>a</sup>	RFF ± SD <sup>b</sup>	
<b>5</b>	>64 <sup>d</sup>	0.66 ± 0.13*	nd <sup>e</sup>
<b>6</b>	>256 <sup>d</sup>	0.17 ± 0.14	nd
<b>7</b>	>256 <sup>d</sup>	<b>1.17 ± 0.46*</b>	84.8 (264.7)
<b>8</b>	>256 <sup>d</sup>	0.72 ± 0.19*	nd
<b>9</b>	<b>256</b>	<b>0.95 ± 0.15*</b>	8.8 (30)
<b>10</b>	<b>64</b>	<b>0.81 ± 0.27*</b>	10.5 (32.51)
<b>11</b>	>256 <sup>d</sup>	<b>15.34 ± 0.82***</b>	170.8 (701.8)
<b>12</b>	64	<b>0.94 ± 0.079**</b>	0.762 (2.37)
<b>13</b>	>64 <sup>d</sup>	0.61 ± 0.16*	nd
<b>14</b>	256	0.76 ± 0.04**	nd
<b>15</b>	>256 <sup>d</sup>	0.39 ± 0.16*	nd
<b>18</b>	>64	-0.17 ± 0.16	nd
<b>19</b>	>256 <sup>d</sup>	0.08 ± 0.18	nd
<b>TZ</b>	<b>30</b>	<b>0.82 ± 0.1*</b>	5.6 (13.78)
<b>VP</b>	<b>800</b>	<b>2.21 ± 0.14**</b>	50.5 (116.6)

<sup>a</sup> Determined by microdilution; <sup>b</sup> Relative final fluorescence based on accumulation of EtBr at 0.25 μg/mL; the results are presented as the average of three independent assays plus standard deviation (± SD). The results were considered significant when \*P<0.05 and highly significant when \*\*P<0.01 and \*\*\*P<0.001; <sup>c</sup> Index of Cytotoxicity (IC) determined in human monocyte derived macrophages; <sup>d</sup> Due to the reduced solubility of the compounds it was not possible to test at higher concentrations; <sup>e</sup> not determined.



**Table 2.** Evaluation of the antimycobacterial and efflux inhibitory activity of the selected compounds against *M. tuberculosis* H37Rv.

comp	<i>M. tuberculosis</i> H37Rv	
	MIC ( $\mu\text{g/mL}$ ) <sup>a</sup>	RFF $\pm$ SD <sup>b</sup>
<b>7</b>	>256 <sup>c</sup>	0.84 $\pm$ 0.02*
<b>9</b>	128	0.33 $\pm$ 0.11
<b>10</b>	>256 <sup>c</sup>	1.22 $\pm$ 0.09*
<b>11</b>	>256 <sup>c</sup>	1.00 $\pm$ 0.19
<b>TZ</b>	15	0.27 $\pm$ 0.02
<b>VP</b>	512	1.94 $\pm$ 0.005**

<sup>a</sup> Determined by microdilution; <sup>b</sup> Relative final fluorescence (RFF) based on accumulation of EtBr at 0.5  $\mu\text{g/mL}$ ; the results are presented as the average of three independent assays plus standard deviation ( $\pm$  SD). The results were considered significant when \*P<0.05 and highly significant when \*\*P<0.01; <sup>c</sup> Due to the reduced solubility of the compounds it was not possible to test at higher concentrations.

**Table 3.** Evaluation of the synergistic effect and determination of the modulation factor (MF) of the selected compounds with first- and second- line drugs against *M. tuberculosis* H37Rv.

comp	test conc <sup>a</sup>	MIC (μg/mL) against <i>M. tuberculosis</i> H37Rv (MF) <sup>g</sup>				
		INH <sup>b</sup>	RIF <sup>c</sup>	AMK <sup>d</sup>	OFX <sup>e</sup>	EtBr <sup>f</sup>
<b>no comp</b>		0.1	1	2	2	12.5
<b>7</b>	256	0.1	<0.0156(>↓64)	2	1	12.5
	128	0.1	0.25(↓4)	2	2	12.5
	64	0.1	0.25(↓4)	2	2	12.5
	32	0.1	0.25(↓4)	2	2	12.5
	16	0.1	0.25(↓4)	2	2	12.5
	8	0.1	0.25(↓4)	2	2	12.5
	4	0.1	0.25(↓4)	2	2	12.5
<b>9</b>	64	<0.00078 (↓>128)	<0.0156(>↓64)	<0.0156 (>↓128)	<0.0156(>↓128)	<0.195 (>↓64)
	32	0.05	0.0625 (↓16)	0.625 (↓4)	1	1.56 (↓4)
	16	0.1	0.5	1	1	6.25
	8	0.1	0.5	1	1	12.5
	4	0.1	1	1	1	12.5
	2	0.1	1	1	1	12.5
	1	0.1	1	1	1	12.5
<b>10</b>	256	<0.00078 (↓>128)	<0.0156(>↓64)	<0.0156 (>↓128)	<0.0156(>↓128)	<0.195 (>↓64)
	128	<0.0015 (>↓64)	<0.0156(>↓64)	<0.0156 (>↓128)	<0.0156(>↓128)	<0.195 (>↓64)
	64	0.05	<0.0156(>↓64)	>0.0156 (>↓128)	<0.0156(>↓128)	1.56 (↓8)
	32	0.1	0.0312(↓32)	1	1	6.25
	16	0.1	0.0625(↓16)	1	1	12.5
	8	0.1	0.25(↓4)	1	1	12.5
	4	0.1	0.5	1	1	12.5
<b>11</b>	256	<0.00078 (↓>128)	<0.0156(>↓64)	<0.0156 (>↓128)	<0.0156(>↓128)	<0.195 (>↓64)
	128	0.1	0.5	1	1	12.5
	64	0.1	0.5	1	1	12.5
	32	0.1	0.5	1	1	12.5
	16	0.1	0.5	1	1	12.5
	8	0.1	0.5	1	1	12.5
	4	0.1	0.5	1	1	12.5
<b>TZ</b>	7.5	<0.00078 (↓>128)	<0.0156(>↓64)	<0.0156 (>↓128)	<0.0156(>↓128)	<0.195 (>↓64)
	3.75	0.05	1	1	1	3.125 (↓4)
	1.88	0.1	1	1	1	6.25

	0.94	0.1	1	1	1	6.25
	0.47	0.1	1	1	1	6.25
	0.23	0.1	1	1	1	6.25
	0.12	0.1	1	1	1	6.25
	128	<0.00078 (↓>128)	<0.0156(>↓64)	<0.0156 (>↓128)	<0.0156(>↓128)	<0.195 (>↓64)
	64	0.05	0.25 (↓4)	1	1	3.125 (↓4)
	32	0.1	0.5	1	1	6.25
<b>VP</b>	16	0.1	0.5	1	1	6.25
	8	0.1	0.5	1	1	12.5
	4	0.1	1	1	1	12.5
	2	0.1	1	1	1	12.5

<sup>a</sup> Concentration at which the compound was tested (in µg/mL); <sup>b</sup> Isoniazid; <sup>c</sup> Rifampin; <sup>d</sup> Amikacin; <sup>e</sup> Ofloxacin; <sup>f</sup> Ethidium bromide; TZ, thioridazine; VP, verapamil. <sup>g</sup> modulation factor.

The modulation factor (MF) was calculated with the following formula:  $MF = \frac{MIC_{\text{antibiotic}}}{MIC_{\text{combination}}}$

The modulation factor reflects a reduction of the MIC values of a given antibiotic in the presence of an inhibitor and was considered significant when  $MF \geq 4$  (≥four-fold reduction). All experiments were repeated at least three times.

**Supporting Information Available:** <sup>1</sup>H NMR, <sup>13</sup>C NMR description of compounds **5**, **6**, **9-15**, **18**, the synthesis of compounds **1-4**, **16**, **17** and the biological methods used for the determination of the MIC, the RFF and the synergistic activity are reported in the supporting information. This material is available free of charge via the Internet at <http://pubs.acs.org>.

### Author information

#### Corresponding authors:

Marco Pieroni, [marco.pieroni@unipr.it](mailto:marco.pieroni@unipr.it), +39 0521905054

Miguel Viveiros, [mveiros@ihmt.unl.pt](mailto:mveiros@ihmt.unl.pt), +351 213652653

### Author Contributions

1  
2  
3 The manuscript was written through contributions of all authors. All authors have given approval  
4 to the final version of the manuscript. # These authors contributed equally.  
5  
6  
7


### 8 9 **Acknowledgments**

10  
11 Giovanni Guerreschi is kindly acknowledged for the help in the synthesis of derivatives. This  
12 work was partially supported by project PTDC/BIA-MIC/121859/2010 from Fundação para a  
13 Ciência e a Tecnologia (FCT), Portugal. Diana Machado and Sofia Santos Costa were supported  
14 by grants PTDC/BIA-MIC/121859/2010 (BI) and SFRH/BPD/97508/2013, respectively, from  
15 FCT, Portugal.  
16  
17  
18  
19  
20  
21  
22  
23

### 24 **Abbreviations**

25  
26 AMK, amikacin; DMF, *N,N*-dimethyl formamide; DOTS, directly observed therapy short-  
27 course; EI, efflux inhibitor; INH, isoniazid; MIC, minimum inhibitory concentration; MDR-TB,  
28 multidrug-resistant tuberculosis; *Mtb*, *Mycobacterium tuberculosis*; RIF, rifampicin; TB,  
29 tuberculosis; TEA, triethylamine ; XDR-TB, extensively drug-resistant tuberculosis.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60


## Table of Contents graphic



Thioridazine



Verapamil

Hybrid structure  
inhibitor of efflux  
7