UNIVERSITÀ DI PARMA
ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Current Trends in Ancient Grains-Based Foodstuffs: Insights into Nutritional Aspects and Technological Applications

This is the peer reviewd version of the followng article:

Original
Current Trends in Ancient Grains-Based Foodstuffs: Insights into Nutritional Aspects and Technological Applications / Boukid, F.; Folloni, S.; Sforza, S.; Vittadini, E.; Prandi, B.. - In: COMPREHENSIVE REVIEWS IN FOOD SCIENCE AND FOOD SAFETY. - ISSN 1541-4337. - 17:1(2018), pp. 123-136. [10.1111/1541-4337.12315]

Availability:
This version is available at: 11381/2838430 since: 2021-10-19T17:12:29Z

Publisher:
Blackwell Publishing Inc.

Published
DOI:10.1111/1541-4337.12315

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

25 April 2024

Current Trends in Ancient Grains-Based Foodstuffs: Insights into Nutritional Aspects and Technological Applications

Fatma Boukid[1*], Silvia Folloni[2], Stefano Sforza[1], Elena Vittadini[1], Barbara Prandi[1]

[1] Food and Drug Department, University of Parma, Parco Area delle Scienze 27/A, 43124 Parma, Italy

[2] Open Fields S.r.L., Collecchio, Parma, Italy

*Corresponding author:

Fatma Boukid, Food and Drug Department, University of Parma, Parco Area delle Scienze 27/A, 43124 Parma, Italy; e-mail: fatma.boukid@unipr.it; tel: +390521906079

Abstract

For centuries, ancient grains fed populations, but due to their low yield, they were abandoned and replaced by high-yielding species. However, currently, there is a renewed interest in ancient wheat and pseudocereal grains from consumers, farmers, and manufacturers. Ancient wheat such as einkorn, emmer, spelt, and Kamut®, are being reintegrated because of their low fertilizer input, high adaptability and important genetic diversity. New trends in pseudocereal products are also emerging, and they are mostly appreciated for their nutritional outcomes, particularly by the gluten-free market. Toward healthier lifestyle, ancient grains-based foodstuffs are a growing business and their industrialization is taking two pathways, either as a raw ingredient or a functional ingredient. This paper deals with these grain characteristics by focusing on the compositional profile and the technological potential.

Key words: Ancient Grains, Foodstuffs, Ancient Wheat, Pseudocereals, Gluten-Free, Quality.

1. Glossary

2. An overview of ancient grains

2.1. Fundamental classes and subclasses of ancient grains

2.2. The way forward: Reasons behind the shifting away from ancient grains

2.3. Back to basics: reasons behind the reintroduction of ancient grains

3. Composition of ancient grains versus modern grains

3.1. Ancient wheat

3.2. Pseudocereals

4. Ancient grains-based foodstuffs: nutritional added value and technological characteristics

4.1. Trends in ancient wheat-based foodstuffs

4.2. New advances in pseudocereals-based foodstuff formulations

4.2.1. Pseudocereals-based foodstuff

4.2.2. Pseudocereal incorporation in gluten-containing foodstuff

5. Landraces and old wheat varieties

6. Concluding remarks and future outlook

23

1. Glossary

Despite the many genetic and archeological data on the origins of agriculture, from gathering to cultivation to domestication to breeding, surprisingly there is no universal definition for modern or/and ancient grains. The case of wheat is complex because it includes several species with different degrees of ploidy, of which some evolved concurrently in different geographical areas. Herein, an attempt to define some debatable terms is performed, considering the degree of human intervention, the degree of breeding and the level of genetic evolution. The classifications, as well as the definition assigned to each category, are research-based, and scientifically sound (Table 1). It takes also into account several factors such as the history of breeding, the origin of selection, the crossing, and the pedigree.

In the last years, conserving the natural agrobiodiversity is challenging, and the risk of bottleneck situation in breeding keeps raising. Therefore, going back to ancient species might avoid or prevent this shifting. Furthermore, awareness was raised towards human health and nutrition. As a result, the consumers were interested in natural, unconventional and nutritional foods, which led to the development of new health-beneficial foods based on grain blends. The attention towards ancient species has also been renewed by the mounting demand for traditional products, the request for species suitable to be grown in marginal areas and the need to preserve genetic diversity. In this regard, this review aims to provide new insights into these grains' nutritional and technological characteristics as well as the current trends in their based foodstuffs, which might be extremely valuable for consumers, producers, processors and farmers.

2. An overview of ancient grains

2.1. Fundamental classes and subclasses of ancient grains

Ancient grains might be subdivided in several categories and subcategories (Figure 1A). Botanically, grains belong to monocotyledon "monocots" (one seed leaf) and dicotyledon "dicots" (two seed leaves) which are the two major subclasses of flowering plants "*angiosperms"*. Three main categories are: cereals, minor cereals, and pseudocereals. Cereals are made up of rice, wheat, and maize. Minor cereals

are mainly rye (*Secale cereale L.*), foxtail millet (*Setaria italica* L.), oat (*Avena sativa* L.), sorghum (*Sorghum bicolor* L.), barley (*Hordeum vulgare* L.), common millet (*Panicum miliaceum* L.) and teff (*Eragrostis tef* (*Zucc*.) Trotter) (Diao 2017). Cereals and minor cereals belong to *Poaceae* which is one of the most important families of the order *Poales* belonging to monocotyledon subclass (Figure 1B). For centuries, *Poacea*, formerly *Gramineae*, provided the most important share of human nutrition. *Poaceae*, also called grasses, contain over 600 genera and more than 10,000 species that dominate many ecological and agricultural systems (The International Brachypodium Initiative 2010). It was established also that the grasses family diverged into different subfamilies and tribes, with the sub-families being *Pooideae* (wheat, barley, and oats), *Ehrhartoideae* (rice), *Panicoideae* (maize, sorghum), and *Chloridoideae* (teff) (Charles and others 2009). Tribes are mainly *Triticeae* (wheat, rye, and barley), *Aveneae* (oat), *Paniceae* (millet), *Andropogonee* (sorghum and maize) and *Eragrosteae* (teff). On the other hand, pseudocereals are defined as non-grasses, dicots grains that diverge into several families such as *Polygonaceae*, *Amaranthaceae,* and *Lamiaceae* (Figure 1C). Amaranth (*Amaranthus caudatus L.; A. cruentus; A. hypochondriacus*), quinoa (*Chenopodium quinoa Willd.*), and buckwheat (*Fagopyrum esculentum Moench.*) are the best known pseudocereals (Fletcher 2016), while chia (*Salvia hispanica L.*) has been gaining interest recently due to its functional and nutritional properties.

Ancient grains might also be subdivided into gluten-containing and gluten-free grains (Figure 1A). Gluten containing grains are mainly ancient wheat, including einkorn, emmer, spelt, and Kamut® (Table 2), while the most popular gluten-free ancient grains are the pseudocereals buckwheat, quinoa, amaranth and chia (Table 3).

2.2. The way forward: Reasons behind the shifting away from ancient grains

The main reason behind the shifting away from ancient grains was their low yield. Plant improvement programs aimed at increasing grain yield to feed the growing populations (Okuno and others 2014) and, as a result, ancient grains were abandoned and replaced by high-yielding modern grains which contributed to the decrease of genetic diversity. Nowadays about 95% of the cultivated wheat worldwide

74 is *Triticum aestivum*, while most of the remaining 5% is *T. turgidum* susbp. *durum* (Brouns and others
75 2013).

76 Ancient wheat grew excessively tall, thereby becoming susceptible to lodging with consequent
77 significant yield loss (Okuno and others 2014). During evolution under domestication, spikelets have
78 undergone significant changes to keep the plant standing until harvest (Zhou and others 2015) to prevent
79 yield losses through seed shattering, to minimize seed dormancy, and to increase both seed size and
80 number (Peleg and others 2011; Sakuma and others 2011).

81 Later on, the green revolution, throughout the 1950s -1960s, led to the development of high-yielding,
82 disease-resistant wheat varieties with dwarfing genes (Lopes and others 2015). The new varieties were
83 selected through breeding protocols based on modern agronomic practices with high agricultural inputs
84 (Royo and others 2007; Longin and Würschum 2016).

85 ### 2.3. Back to basics: reasons behind the reintroduction of ancient grains

86 The increasing demand for traditional products, the request for high adaptability, and the need to
87 preserve genetic diversity are among the reasons behind the renewed attention towards ancient species
88 (Troccoli and Codianni 2005). The evolution of plant breeding resulted in genetic erosion and thereby
89 the loss of genetic diversity. Therefore, there is a serious need to go beyond the uniform model of today
90 agriculture and reintroduce germplasm characterized by high heterogeneity. Species diversity
91 contributes to the increase of crop productivity and stability (Hooper and others 2012; Khoury and others
92 2014). Indeed, ancient wheat is suitable for organic farming because of their adaptability to low
93 agronomic input; and where other wheat types would fail, they show high resistance to powdery mildew
94 and brown rust, and disadvantageous growing conditions (such as wet, cold soils, high altitudes, and
95 poor soils) (Konvalina and others 2010; Escarnot and others 2012). Pseudocereals are also known for
96 their high adaptability, low needs in terms of water, fertilizer, and energy as compared to traditional
97 cereals (Kang and others 2017; Santra and Schoenlechner 2017). As a result, the 2015 international

report from Health Focus International stated that the international awareness of ancient grains was up from 26% in 2012 to 28% in 2014, with 35% of the respondents expressing an interest in ancient grains.

Currently, ancient grains are gaining popularity as they often offer a better nutritional composition (Carnevali and others 2014). Ancient wheat have been rediscovered by consumers, bakers, millers, and farmers because they are good sources of proteins, lipids, fructans, trace elements, and several antioxidant compounds (Hidalgo and others 2014; Hidalgo and Brandolini 2014; Longin and others 2015). Pseudocereals have also a good protein quality content in terms of amino acid composition (Wang and Zhu 2015; Ngugi and others 2017). Thus, pseudocereals are integrated in gluten-free product formulations to improve their nutritional quality and, consequently, to avoid some complications such as nutrient deficiencies, bone disease, and lymphoma sensitivity (Alvarez-Jubete and others 2009a, b).

3. Composition of ancient grains versus modern grains

Wheat and cereals are largely consumed worldwide, supplying humans with energy and bioactive components (Lachman and others 2012). However, it should be kept it mind that the nutritional composition of cereal crops is closely associated with cultivation area, climatic conditions, agronomic practices, and genetic diversity (Miranda and others 2012; Hidalgo and Brandolini 2017).

3.1. Ancient wheat

The approximate chemical compositions of ancient wheat (einkorn, emmer, spelt, and Kamut), durum wheat and common wheat are displayed in Table 4. Table 4 also includes data relative to branded whole wheat flour products form the Food Composition Databases of the United States Department of Agriculture (USDA). The choice of branded products rather than research paper data was to give a concrete overview of what is available on the market. Modern wheat (durum and common) are also reported to underline their differences or/and similarities as compared to ancient wheat.

The main components of ancient wheat are carbohydrates, protein and fibers, similarly to modern wheat. Carbohydrate contents in durum and common wheat are slightly higher than in ancient wheat; indeed,

it was found that spelt (68%) provides the lowest carbohydrate contents together with einkorn (67%). Spelt contains the lowest fiber content (5.9%), while it was reported that it is a good source of fiber (11.4 %) by Ranhotra and others (1995), which implies large sample variability. Furthermore, einkorn flour has a lower content of total dietary fiber (6.7%), than common wheat (12.7%).

Important variability was observed between durum, common, and ancient wheat in terms of protein content. According to the intensive work of Hidalgo and Brandolini (2014), einkorn protein content is generally higher to that of common wheat as well as emmer and spelt. Kamut® has high protein, as reported by Sumczynski and others (2015), and it presents some prolamin alleles which are closely correlated to good pasta quality (Rodríguez-Quijano and others 2010).

Overall, the lipid content is a minor component in wheat ranging between 1.7 (einkorn) and 2.9% (spelt). Remarkably, ancient wheat, except for einkorn, has higher lipid contents than that of common wheat, which was consistent with the findings of Hidalgo and others (2009). Moreover. It was reported that einkorn has higher monounsaturated fatty acids, lower polyunsaturated fatty acids and lower saturated fatty acids than durum wheat, suggesting its beneficial effect on human health (Hidalgo and Brandolini 2014).

As for mineral content, Kamut® has comparable calcium, iron, magnesium, potassium, sodium, and zin contents to common wheat and durum wheat. Kamut® also showed higher levels for 8 out of 9 minerals compared to common wheat (Abdel-Aal and others 1998). Magnesium and zin contents of einkorn (200mg/100g and 15 mg/100g, respectively) also are higher than those of common wheat (90mg/100g and 3.5mg/100g, respectively) and durum wheat (144mg/100g and 4.2mg/100g, respectively), in concordance with previous studies (Suchowilska and others 2009; Erba and others 2011).

Regarding the vitamin content, Table 4 showed that Kamut® has higher vitamin A than common wheat and durum wheat. Despite the importance of micronutrients, wheat labels usually lack this indication, and more information in this regard are found in literature. Tocols and carotenoids were exclusively discussed in scientific works. Indeed, einkorn was found to have high tocol and carotenoid contents that

147 may make einkorn an interesting ingredient to be used in the development of new or special foods with
148 high nutritional quality (Hidalgo and others 2016). Einkorn tocols content was higher (42.7-70.2µg/g,
149 average =57) compared to durum wheat (40.1-62.7µg/g, average=48.1) (Lampi and others 2008). In
150 another study, it was reported that tocols in ancient wheat, einkorn (61.80–115.85 µg /g), emmer (62.7-
151 67.9µg/g), and spelt (62.7-67.5µg/g) were slightly higher than in durum (38.8-57.27µg/g) and common
152 (53.2-74.9µg/g) wheat (Hidalgo and others 2006). As for carotenoids, important variability was shown;
153 in common wheat, they are less abundant (1.4-3.05 mg/kg) than in durum wheat (1.9-5 mg/kg) and
154 Kamut® (4.4 mg/kg), while carotenoids in einkorn were the highest and varied between 5.3 and 13.6
155 mg/kg (Hidalgo and others 2006). The authors also highlighted the relevant impact of cultivation area
156 and genotype on the content of both tocols and carotenoids (Hidalgo and others 2006).

157 Overall, although only few available scientific data are found dealing with comparative studies in terms
158 of ancient wheat and modern wheat, compositional differences were observed between ancient wheat
159 and modern wheat, as well as among the different ancient grains.

160 ### 3.2. Pseudocereals

161 Table 5 presents the approximate chemical composition of pseudocereals and of the 2 major gluten-free
162 cereals used worldwide (maize and rice). As in Table 4, Table 5 summarized data relative to branded
163 whole flour products from the Food Composition Databases of the United States Department of
164 Agriculture (USDA). Rice and maize together with pseudocereals were also included to provide a full
165 comparative image of macro- and micronutrients.

166 Energy provided by chia is higher than by maize and rice flours, but it is mostly due to its higher lipid
167 content. According to the data reported in Table 5, maize and rice recorded the highest carbohydrate
168 contents (85.19 and 76.5 g/100g, respectively), while amaranth and buckwheat are better sources of
169 fiber (8.9 and 10 g/100, respectively). Quinoa had the lowest fiber content (7g/100g), but it remains
170 highly dependent of the type of variety. Indeed, quinoa's fiber content was reported to range from 7.7%
171 to 13.8% (Li and Zhu 2017) and 8.8-14.1% (Nowak and Charrondière 2016), while chia was found to
172 be the richest in fiber 34.4%, and rather consistent (Ixtaina and others 2008) (30-33g/100g).

Pseudocereal protein contents (13.25-16.5 g/100g) were significantly higher than those of rice (7.23 g/100g) and maize (7.4g/100g). Furthermore, the quality of the proteins of pseudocereals is correlated with their nutritional added-value compared to rice and maize. Quinoa proteins, containing essential amino acids lysine, threonine, and methionine are balanced a micronutrient composition (Wang and Zhu, 2015). Amaranth has also satisfactory lysine and tryptophan contents according to FAO/WHO standards, which make it a valuable fortification ingredient for meals with limited contents like maize or sorghum (Ngugi and others 2017). Chia seeds provide a high-quality protein (about 16.5%) with good amino acids balance, especially methionine and cysteine (Ayerza 2013).

As for lipid fraction, chia showed the highest value (30.7 g/100g and it is among the richest natural source of the essential fatty acid α-linolenic (Menga and others 2017). Amaranth and quinoa showed quite similar lipid contents (6.7 ad 6.1 g/100g, respectively), which were higher than those of rice and maize (3.7 and 2.78 g/100g). The oils obtained from amaranth contain linoleic and α-linolenic fatty acids. Quinoa was reported to be particularly rich in linoleate and linolenate (Chillo and others 2009), while chia seeds are rich in those essential fatty acids (omega 3 and omega 6) (Mohd Ali and others 2012; Costantini and others 2014).

Regarding micronutrients, chia had interesting minerals composition in particular about twice of the values of Ca and P as compared to rice and maize. Buckwheat, amaranth, and quinoa grains are considered good sources of minerals such as Zn, Cu, Mn, K, Na, Ca, and Mg (Chillo and others 2009; Singh and Singh, 2011). Chia was remarkably rich in vitamin A (16.2 µg/100g). Quinoa (2.44 µg/100g) showed high content in vitamin E as compared to rice (0.6 µg/100g). Buckwheat seeds also are rich in thiamine (vitamin B1), riboflavin (vitamin B2) and pyridoxine (vitamin B6) (Dziadek and others 2016; Guo and others 2017), and they also are abundant in several natural antioxidants, such as tocopherols, rutin, quercetin, flavonoids, and phenolic acids.

Taking in account all the above-mentioned, the good composition of pseudocereals is not only suitable to individuals with medical needs, but also to consumers seeking “healthy” foodstuffs (Pellegrini and Agostoni, 2015; Balestra and others 2015).

4. Ancient grains-based foodstuffs: nutritional added value and technological characteristics

Because ancient grains have a rich and balanced micronutrient composition, they might be suitable raw ingredients to enhance technological quality or/and health benefits (Arzani 2011; Randall and others 2012; Chandi and others 2015). From an economic point of view, ancient grains are low-inputs crops with lower needs in fertilizers and they might be a suitable corp in lower-income countries. Quinoa, for example, is known for its tolerance to soil salinity indicating its suitability to harsh conditions (Wu and others 2016).

Thus, ancient crops rediscovering to be a part of the daily human diet might be reinforced by a further understanding of their technological properties and applications.

4.1. Trends in ancient wheat-based foodstuffs

The increasing popularity of ancient wheat as environmentally friendly cereal crops is stimulating research into their utilization in both traditional and new foods (Messia and others 2012). Hulled wheat species (einkorn, emmer, and spelt) are often used as whole grains for salads or soups, while they are used differently for processing, with einkorn and emmer mainly used for pasta products and spelt mainly used for bakery products (Benincasa and others 2015).

Bread-making

Einkorn flour is generally reported as not suitable for bread-making because of its sticky dough and poor rheological properties, but the existence of accessions with high bread-making quality has been confirmed (Hidalgo and Brandolini 2011; Brandolini and Hidalgo 2011). It was reported that the screening of a wide collection of einkorns (>1000 accessions) allowed the identification of approximately 16% of the total accessions with sodium dodecyl sulfate (SDS) sedimentation values corresponding to the threshold value for bread-making potential (Borghi and others 1996). Later, Brandolini and others (2008) conducted a survey of 65 einkorn samples to study their pasting properties and concluded that einkorn had higher peak viscosity and final viscosity than modern wheat. The

differences are probably related to the smaller size and different grading of einkorn starch granules as well as to the lower amylose percentage of einkorn flour (Brandolini and Hidalgo 2011). Regarding bread color, einkorn has a color lighter than common wheat and durum wheat suggesting that einkorn undergoes lower heat damage than modern wheat during baking because low α- and β- amylases limit the degradation of starch (Brandolini and Hidalgo, 2011). As a result, the reduced generation of reducing sugars in the dough limited the Maillard reactions during food processing. Low lipoxygenase activities in einkorn dough also limits the degradation of carotenoids (Hidalgo and Brandolini 2014).

Kamut® bread showed good sensory properties and loaf volumes, highly resembling bread obtained from modern wheat (Pasqualone and others 2011). Indeed, it was found that it is more suitable than durum wheat for the fermentation processes at acidic conditions because an increase in the bread volume and the metabolic heat production by yeast were observed (Balestra and others 2015).

A comparative study with spelt varieties showed acceptable sensory scores with significant differences among the varieties (Korczyk Szabó and Lacko Bartošová 2013), leading to conclude that spelt might be a suitable raw material for bread making, but it remains closely related to the choice of spelt variety (Korczyk Szabó and Lacko Bartošová 2015). Compared to common bread, spelt genotypes had high crumb elasticity, but low crumb cell homogeneity, which are probably due to its special dough rheological attributes (Callejo and others 2015). Nutritionally, these breads had less total starch, more resistant starch, and less rapidly digested proteins in comparison to bread made with modern wheat flours (Bonafaccia and others 2000). Spelt and emmer sourdoughs had slightly higher pH values than wheat sourdough but titratable acidity, concentration of free amino acids, and phytase activity were higher than in common wheat sourdough (Coda and others 2010). Specific volume and crumb of spelt breads showed higher resemblance to those of wheat breads than emmer. Sensory analysis also revealed that spelt and emmer can be made into acceptable bread products (Coda and others 2010).

Pasta making

Little information was found in the scientific literature on 100% ancient wheat-based pasta due to its low pasting properties (Brandolini and others 2008). Indeed, 100% einkorn pasta showed less compact

structure than durum wheat, resulting in high cooking losses and low ability in binding water (Pasini and others 2015). However, pasta made by a mix of 50/50 semolina/einkorn showed a high aggregation of gluten (La Gatta and others 2017). Higher carotenoid levels also were found during kneading because of the low enzyme activities in einkorn (Hidalgo and others 2010). Marconi and others (2002) assessed gluten properties of 3 spelt genotypes (Rouquin, Redoute, and HGQ Rouquin (Rouquin improved for gluten quality)) using SDS sedimentation, gluten index values, and alveograph and farinograph parameters. Compared to durum wheat pasta, spelt pasta dried at high temperature had darker color which might be attributed to higher furosine, probably correlated with higher reducing sugars or damaged starch contents in the semolina (Marconi and others 1999; 2002). Although ancient crops are commonly reported to have a low technological quality of proteins (gluten strength) (Brandolini and others 2008), spelt (HGQ Rouquin) allowed the production of pasta with satisfactory cooking quality (Marconi and others 2002). As a matter of fact, 100% emmer pasta had improved organoleptic value and lower glycemic index than durum pasta (Fares and others 2008). Moreover, pasta cooking did not damage the polysaccharide composition (soluble and insoluble), but it induced a drastic loss in terms of tocopherol and carotenoid contents (Fares and others 2008).

Baked goods, snacks, breakfast cereals

Einkorn flour was reported for its excellent aptitude to produce many foodstuffs and it is currently trending as a material for manufacturing new special foods with high nutritional quality such as crackers and snacks (Hidalgo and Brandolini 2010; Brandolini and Hidalgo 2011; Hidalgo and others 2016, Hidalgo and Brandolini 2017). For instance, compared to einkorn, puffed modern kernels seem more appealing (Hidalgo and others 2016). However, from a nutritional point of view, puffed einkorn kernels had good composition, in terms of proteins and bioactive compounds (Hidalgo and others 2016). Compared to commercial products, flakes and muesli made by Kamut® and spelt had acceptable sensory features (appearance, consistency and flavor), and they showed the highest total phenolic, flavonoid, and crude fiber contents (Sumczynski and others 2015). Furthermore, tortillas partially substituted by whole Kamut® flour (60%) showed high resemblance with that standard (Carini and others 2010). Spelt flour addition (5, 10 and 15%) to corn grits decreased the expansion ratio and fracturability, whereas

bulk density and hardness of extrudates increased (Jozinović and others 2016). Moreover, color changed, the peaks of viscosity (hot and cold) decreased, and less retrogradation was observed.

Beverage making

A few studies have focused on ancient wheat-based alcoholic and/or nonalcoholic beverages. Einkorn wort exhibits standard properties, and the resulting beers showed excellent foam stability and distinct pleasing carbonation taste. As a result, it is suggested as a potentially new raw material to produce organic beer (Fogarasi and others 2015). Einkorn malts had high radical cation scavenging activities, as measured by DPPH and ABTS methods, but the phenolic content was lower when using wheat (Fogarasi and others 2015). As well, emmer malt is characterized by a very high extract yield and good saccharification time, and good foam stability, but weak final attenuation, low polyphenol content, and darker color than the barley malt beers (Esslinger 2009; Mayer and others 2011). Barley malt enriched by emmer showed different sensory profiles as well as different compositions in terms of concentrations of organic acids, carbohydrates, amino acids, dietary fibers, vitamins, and antioxidant and phytase activities (Coda and others 2011). Spelt malt had an appropriate extract yield and apparent attenuation limit in comparison with barley and wheat malt (Muñoz-Insa and others 2013). However, low spelt soluble nitrogen caused a low Kolbach index; high viscosity caused lautering and filtration problems. As a result, this cereal can be used in "normal" barley malt houses under commonly used malting conditions, but it remains interesting to optimize beer production from mixes of ancient wheat malt and barley malt. Furthermore, hulled malts exhibit lower soluble protein contents with higher free amino nitrogen and total free amino acid values than dehulled malts (Marconi and others 2013). More studies are required to determine the influence of the glume or husk on malt and technological aspects.

Blends of rice and emmer flours (ratio 94:6) also allowed the production of a good yogurt-like beverage in terms of textural, sensory, and nutritional properties (Coda and others 2012).

4.2. New advances in pseudocereals-based foodstuff formulations

The development of new products based on pseudocereals is, currently, more of necessity rather than a choice. The attention towards these ancient grains have been renewed by the increasing demand for natural and health-beneficial foods. They are also naturally gluten-free seed perfectly suitable as a reinforcement of the gluten-free market which is nowadays mainly based on maize and rice.

4.2.1. Pseudocereals-based foodstuff

Gluten-free foodstuff-making is a challenge for technologists and nutritionists due the absence of gluten, which provides the viscoelastic properties of dough (De la Barc and others 2010). Nutritionally, pseudocereals represent a healthy alternative to compensate for the deficiencies of a gluten-free diet (Saturni and others 2010), because most of the gluten-free products currently available in the market are made basically from refined flour or starches which are characterized by low contents of high-quality protein, fiber, calcium, and iron (Cabrera-Chávez and others 2012; Molina-Rosell and others 2015).

Bread-making

The fortification of breads using buckwheat (50%) or quinoa (50%) flours increased the volume and softened the crumb (Alvarez-Jubete and others 2009b, 2010). Buckwheat addition (from 10 to 30%) also decreased starch retrogradation which enhanced the anti-staling properties (Torbica and others 2010). Furthermore, 100% buckwheat bread had low specific volume due to its dense structure (Hager and others 2012). The enrichment of bread by amaranth (40%) enhanced the physical, and rheological gluten-free dough as well as bread final attributes compared to corn starch-based formulations (Mariotti and others 2009). Pseudocereals addition also improved bread nutritional profiles, fitting perfectly with the expert nutritional recommendations for a gluten-free diet and gluten-free foods (Alvarez-Jubete and others 2009b, 2010; Sakac and others 2011; Wronkowska and others 2010; Koppalu and others, 2016).

Amaranth and quinoa starches are used also for gluten-free food-making because they are characterized by a low tendency to retrogradation (Abugoch 2009; Singh and Singh, 2011). They showed also higher gelatinization temperature and peak viscosity but lower swelling capacities compared to those of fine flour and middling fractions (Kumar and others 2016; Sakhare and others 2017). More attention,

however, should be paid to amaranth technological processing because it requires innovative approaches due to its small seed size (Santra and Schoenlechner 2017).

The combination of sourdough lactic acid bacteria and pseudocereals was successfully employed in the formulation of new pseudocereals-based products (Dallagnol and others 2012; Gobbetti and others 2014). During sourdough fermentation, several reactions, such as acidification and proteolysis, greatly contribute to the increase in bread extensibility, softness, and volume (Nionelli and Rizzello, 2012). For instance, quinoa addition (3.75%) improved shelf-life of bread (Coda and others 2010; Dallagnol and others 2012). Likewise, 100% bread made from buckwheat sourdough improved bread quality, thereby reducing the need of additives (Moroni and others 2009, 2010, 2011). Sourdoughs obtained with teff (5, 10 and 20%) and buckwheat (15%) also enhanced bread aroma and increased fruity, cereal, and toasty notes as well as increased the perceived elasticity (Campo and others 2016). On the other hand, fermentation stimulated protein hydrolysis, which increased the concentrations of free amino acids (Dallagnol and others 2012), offering bioactive peptides and amino acids unavailable in gluten-free products.

Recently, steeping and germination were suggested to be able to reduce the levels of antinutrients, thereby enhancing the bioavailability of minerals in amaranth grains, resulting in a nutrient-dense complementary food (Ngugi and others 2017).

Pasta-making

Commonly, corn and rice flours are used as main ingredients for gluten-free pasta production (Cabrera-Chávez and others 2012), while recently there is increasing interest in new formulations based on gluten-free blended flours (cereals and pseudocereals) to improve the nutritional aspect as well as the technological features. For instance, pasta made from corn flour and low quinoa content (0.16%) had acceptable physical properties compared to common wheat pasta (Caperuto and others 2000). About 25% of amaranth enrichment gave good pasta-making results due to the good interaction between rice starch and amaranth proteins. Moreover, Fiorda and others (2013) suggested that gluten-free pasta fortified by 30% amaranth flour had higher quality and sensory scores to those made with regular and

whole wheat flour pasta. Besides improved technological quality, these blends had higher protein and fiber contents than gluten-free pasta. Noodles produced from amaranth flour (20%) had low firmness and high cooking losses (Schoenlechner and others 2011). However, its nutritional composition, in terms of mineral and fiber contents, and protein digestibility were improved when a novel and adequate extrusion-cooking process was used (Cabrera-Chávez and others 2012). Pasta fortified by chia (5 and 10%) showed that chia seeds and mucilage might be a good natural thickening agent (Menga and others 2017). Furthermore, this addition increased total phenolic acids and dietary fiber as compared to the pasta made from rice (Menga and others 2017).

Baked good, snacks, and breakfast cereals

New trends and formulations of pseudocereal foodstuffs are spreading to produce health-beneficial and tasty products, particularly baked good, snacks, and breakfast cereals. For example, quinoa was largely used as an ingredient for baked goods (Rizzello and others 2015). Muffins fortified with 25% quinoa flour are soft and have good overall consumer acceptability. However, 100% quinoa flour muffin had low overall consumer acceptability due to the bitter taste of quinoa flour (Bhaduri 2013). An increase in quinoa supplementation (30%) also increased the hardness of cookies, which was attributed to the high content of fiber and protein (Brito and others 2015). The specific volume also was reduced resulting in less bulky cookies, and also the final color was darker (Brito and others 2015).

On the other hand, Burgos and Armada (2015) demonstrated amaranth suitability to precooked products due its high expansion and yellow index. Chia flour was incorporated at different levels in cakes (25%, 50%, and 75%) (Borneo and others 2010) and corn tortillas (5%, 10%, 15%, and 20%) (Rendón-Villalobos and others 2012), and gluten-free bread (from 4% to 15%) (Moreira and others 2013; Steffolani and others 2014; Da Mota Huerta and others 2016). Chia flour played the role of hydrocolloid, indeed, because it significantly increased the water-holding capacity of the dough (Da Mota Huerta and others 2016; Olivos-Lugo and others 2010; Steffolani and others 2014).

Beverage-making

A fermented quinoa-based beverage was recently formulated using 2 varieties (Pasankalla and Rosada de Huancayo) (Ludena Urquizo and others 2017). The obtained drinks had viable and stable microbiota during storage (28 days) and the fermentation was mostly homolactic (Ludena Urquizo and others 2017). Pasankalla-derived drinks had higher protein contents, lower saponin concentration, and lower loss of viscosity during the fermentation process compared to Rosada de Huancayo drinks (Ludena Urquizo and others 2017). Buckwheat showed lower malt extracts, longer saccharification times, higher total protein and fermentable amino nitrogen content, and higher values of the iodine test and color as compared to barley malts (Deželak and others 2014). However, fermentability values, wort pH, soluble protein content, and volatile compounds were comparable and, consequently, the organoleptic perception of the buckwheat beverage was good (Deželak and others 2014).

The aqueous extract of pseudocereal flours was incorporated in fermented lactose-free dairy products. High incorporation of quinoa extracts (70 and 100%) increased the viscosity due to the high protein content (Bianchi and others 2015), while low addition (30%) was more appreciated by the tasters, as well as the obtained drink presented better nutritional features (protein, fat, ash, and total solid levels) (Bianchi and others 2015). Regarding lactose-containing products, El-Deeb and others (2014) suggested that 75 or 100% quinoa addition allowed the production of fermented milk beverages without altering bacterial growth. After 10 days, these beverages had the highest scores in terms of body, texture, color, and appearance. Moreover, the nutritional value was enhanced such as iron and some amino acids (phenylalanine, methionine, histidine and leucine).

Meat industry

The addition of chia was studied to develop lipid meat products (Herrero and others 2017). Because of uncertainties in relation with the potential allergenicity, the daily intake of chia seeds should not exceed 48 g/day, according to the 2000 US Dietary Guidelines), and 15 g/day according to the European commission. Chia-incorporated meat products showed relevant results. For example, frankfurters formulated with 1% chia flour reported significant improvements in water-binding properties (Pintado and others 2015). Furthermore, scanning electron microscopy showed that chia addition improved

emulsification and juiciness and, consequently, the overall acceptance was comparable to products with added fat (Ding and others 2017). The high content of fiber in chia is correlated with higher water-holding, absorption, and emulsifying activity and stability (Alfredo and others 2009). Chia oil addition to meat batter also resulted in products with good stability and homogeneous structure (Cofrades and others 2014). Furthermore, this fortification enhanced the nutritional aspect by increasing linolenic acid and reducing processing and purges (Pintado and others 2016; De Souza and others 2015). Thus, meat products containing chia might be suitable for special nutritional label (Pintado and others 2016).

4.2.2.Pseudocereal incorporation in gluten-containing foodstuff

Fortification of staple foods is an effective strategy to deliver and increase the intake of micronutrients in the diet and can reduce micronutrient deficiencies (Yusufali and others 2012). As amply mentioned above, pseudocereals have a good chemical composition in terms of bioactive components (Padalino and others 2016).

Common wheat bread fortification could enable the development of a range of new baking products with enhanced nutritional value (Stikic and others 2012). Indeed, bread supplemented with quinoa flour had high acceptance with high nutritional value depending on the substitution level (Calderelli and others 2010; Iglesias-Puig and others 2015). Indeed, bread made from blends containing 5% or 10% of quinoa whole flour showed good bread-making properties, while blends with 15% of quinoa flour were not acceptable (Enriquez and others 2003). Rodriguez-Sandoval and others (2012) also showed that partial substitution with quinoa whole flour (10 and 20 %) resulted in breads with decreased specific volume. However, with up to 20% addition of purified quinoa, the rheological characteristics of dough and sensory characteristics were improved (Stikic and others 2012). A 50% substitution induced weak final quality (Alvarez-Jubete and others 2010). Indeed, bread crumb hardness increased (Iglesias-Puig and others 2015), about 6 times higher than sourdough bread with 100% substitution (Wolter and others 2014), as well as density and chewiness (Wang and others 2014). Specific volume and yellowness also were reduced (Wang and others 2014). The addition of chia (5, 10, and 15%) improved gas retention in dough and cut the time required to reach maximum dough development (Verdú and others 2015). The

obtained bread had a reduced water activity and contained the same amount of moisture compared with the control (Verdú and others 2015). Indeed, the high content of fiber in chia might be the main cause of the high water-holding capacity, as it forms an active hydrocolloid which interacts better with gluten proteins (Verdú and others 2015). Therefore, the inclusion of chia increased the overall acceptability by consumers (Iglesias-Puig and Haros 2013).

Recently, Rizzello and others (2016) carried out a study to investigate wheat bread enrichment by quinoa flour sourdough. They concluded important improvements in chemical composition, namely free amino acids, soluble fibers, total phenols, phytase, and in antioxidant activities (Rizzello and others 2016). The sensory features of wheat bread made using 20% of quinoa sourdough were also improved with regards to the use of quinoa flour (Rizzello and others 2016). However, in buckwheat and wheat sourdough breads, acidification increased crumb porosity compared to control breads (Wolter and others 2013). Blends made by 30% of buckwheat and common wheat flours were more effective in enhancing antioxidant activity in comparison with amaranth and quinoa (Chlopicka and others 2012). Moreover, it might improve bread quality attributes such as taste, color, and odor (Chlopicka and others 2012). Fortified bread with 2.5% buckwheat showed acceptable crust and crumb color and taste, and also odor, elasticity and the appropriate bread volume (Gawlik-Dziki and others 2009). Likewise, the addition of chia increased the content of fiber, total antioxidant activity, and ω-3 fatty acid in the final products (Coelho and de las Mercedes Salas-Mellado 2015; Constantini and others 2014). Chia flour significantly increased water absorption and reduced the extensibility of dough (Steffolani and others 2014). It was reported also that chia had thickening potential (Vázquez-Ovando and 2009) and it might replace as much as 25% of oil or eggs in cakes, while yielding a more nutritious product with acceptable sensory characteristics (Borneo and others (2010)

Regarding the pasta industry, pseudocereals and durum wheat blends are still a challenge since the addition of alternative ingredients markedly affects technological and sensory properties (Rizzello and others 2017). Recently, Lorusso and others (2017) revealed that the substitution of 20% of semolina with quinoa flour improved the nutritional aspect of pasta, including free amino acids, total phenols, and

455 the antioxidant activity of pasta; while the resulting pasta tenacity increased. Despite the great nutritional
456 input, more work is required on the balance between substitution level and quality requirement.

457 5. Landraces and old wheat varieties

458 Landraces and old wheat varieties were the most cultivated until the late 19th century (Belderok
459 2000). Therefore, modern varieties were developed to create more productive plants with modified
460 chemical composition and others quality attributes. Comparative studies on old and modern varieties
461 have focused mainly on the physiological basis of yield (Giunta and others 2007). On the other hand, to
462 discriminate between wheat varieties, qualitative and quantitative gliadin and glutenin compositions
463 are the traits commonly adopted (Vita and others 2016). However, few studies were dedicated to
464 screen the compositional and technological aspects of landraces and old genotypes.

465 For instance, Dinelli and others (2013) studied 2 Italian durum wheat genotypes: Senatore Cappelli
466 (1915, selection from the exotic Tunisian landrace "Jneh Khottifa") and Urria 12 (1900) cultivated under
467 a low-input agricultural system. Senatore Cappelli showed the lowest starch content (50.5%), while it
468 had the highest protein content (16.38 %) as well as gluten content (12%) as compared to modern
469 genotypes. Likewise, protein concentration showed a decreasing trend over time of cultivar releasing,
470 dropping from about 18% in the old cultivars to about 16.5% in the modern durum wheat cultivars (De
471 Vita and others 2007), which was consistent with the findings of Fois and others (2011). The low protein
472 content in modern cultivars was not due to a reduced nitrogen uptake, but to the dilution effect caused
473 by the heavier grains of modern durum cultivars (Motzo and others 2004). Regarding fiber and lipid
474 contents, few differences were found between both groups (Dinelli and others 2013). Carotenoid and
475 total polyphenol contents showed a non-significant trend from old to modern common wheat varieties
476 (Dinelli and others 2013). However, phenolic profiles of landraces and old genotypes showed a number
477 of total compounds and isomer forms much higher than those in the modern cultivars, particularly the
478 landrace Gentil Rosso, whih had a much higher amount of total, free, and bound polyphenols (Dinelli
479 and others 2009, 2012). As for minerals, landraces and old cultivars had higher concentrations as
480 compared to more modern material (Hussain and others 2010, 2012), because genetic breeding did not

focus on the improvement of mineral content (Hussain and others 2012). Overall, Di Silvestro and others (2012) found that old common wheat varieties had better nutrient contents when cultivated under low-input conditions compared to the modern ones, which are strictly dependent on high levels of fertilization. Therefore, beside breeding history, the biosynthesis and accumulation of micronutrients are closely influenced by genotype, environmental conditions (Migliorini and others 2016), and farming systems (Rizzello and others 2015).

From a technological point of view, De Vita and others (2007) studied the rheological properties of some Italian landraces, including Timilia (released in 1900, indigenous landrace population from Sicily), Russello S.G.7 (1910, selection from landrace "Russie"), Senatore Cappelli and Aziziah (1925, selection from landraces). Old genotypes (Timilia, Russello S.G.7 and Senatore Cappelli) were characterized by the lowest baking strength (alveographic W value), and dough-gluten properties (P and L alveographic values) (De Vita and others 2007). Modern cultivars showed about twice and thrice the dough W values of the landraces, and about a 50% and 100% greater P, respectively (Sanchez-Garcia and others 2015). However, no significant differences were observed between 7 old (Sieve, Verna, Gentil Rosso, Andriolo, Gambo di ferro, Frassineto, and Abbondanza), 2 mixtures, and 4 modern (Bolero, Blasco, Arabia, and Bologna) varieties, in termS of P/L (Migliorini and others 2016). Gluten Index (GI) was higher for modern cultivars than for landraces and old durum wheat varieties (Motzo and others 2004, Fois and others 2011). Breeding induced a notable increase in GI, which was reflected by an improvement of grain protein quality (Motzo and others 2004). Nevertheless, protein content, rather than gluten quality, has a dominant role in determining pasta cooking quality of high-temperature-dried pasta (Dexter and others 1977; D'Egidio and other 1990), explaining why some durum wheat old varieties such as Senatore Cappelli and others give good pasta texture (Fois and others 2011). Regarding bread-making, breads from landraces were scored as acceptable by consumers (Migliorini and others 2016). Although modern bread wheat varieties showed great variability in bread-making quality attributes (Sanchez-Garcia and others 2015), the landrace Andriolo showed interesting sensory aspects (Migliorini and others 2016). In term of aroma profile, landraces and older cultivars of bread wheat had a soft aroma, while modern varieties had a much stronger aroma (Starr and others 2013).

508 Discussing the quality attributes of big wheat collections is expensive and time-consuming, and the
509 available comparative studies on wheat quality traits are usually restricted to small sets. Moreover, many
510 modern varieties were selected to fulfill specific technological transformations, as for example the
511 Italian durum wheat cultivar Svevo and, consequently, processing landraces and old varieties using the
512 same technologies might explain the difference in the end-product quality. Thus, drawing sound
513 conclusions about the poor technological quality of landraces lacks strong evidences and pleads to much
514 argumentation. Besides breeding new adapted lines for the low-input sector, landraces and old genotypes
515 are naturally suitable to produce organic food and support environmentally friendly practices (Di
516 Silvestro and others 2012).

517 6. Concluding remarks and future outlook

518 The rediscovery of ancient grains provides new alternatives to farmers, consumers, and the food
519 industry, such as seen in terms of gluten-free (non-wheat ancient grains) and gluten-containing
520 foodstuffs.

521 From a breeding point of view, shifting away from ancient species and the gradual shift towards a model
522 of agriculture based on uniformity, steered by the search for higher yields, has increased the risk of
523 genetic erosion (Dwivedi and others 2016). Diversity loss, here as in all the cultivated species, is indeed
524 critical and dangerous, also because ancient species might help us to face threats to food security
525 (Ceccarelli and Grando 2000; Dwivedi and others 2016). Indeed, ancient species, landraces, and old
526 genotypes, even though much less productive than the modern ones, are perfectly suitable for marginal
527 areas and low-input and high-stress conditions (Ceccarelli and Grando 1996) and, consequently, they
528 could represent a solution for local communities where the commonly grown varieties are not cultivable
529 (Migliorini and others 2016). From a nutritional point of view, reintroducing pseudocereals in the daily
530 diet as a fortifying agent with functional added-value features, might offer to consumers a richer variety
531 of beneficial compounds without altering the technological quality. Ancient wheat-based foodstuffs are
532 now an increasing trend in the food market, substituting durum or/and common wheat flour for creating
533 new lines of products. Their technological defects, as discussed, are not always associated with inner

characteristics of seeds, because traditionally made products have often good-quality features. Indeed, modern processing methods or/and industrial machinery, even if perfect for modern varieties, was not be suitable for older grains. Therefore, more work is needed in order to optimize technological processing and formulations to fit their compositional and morphological characteristics.

Thus, ancient grains might constitute an alternative, which can co-exist in the current market with the undoubtedly needed modern high-productive varieties. Certainly, methods and technologies to obtain novel products with high technological quality and health-beneficial and affordable price are needed. Also, blends of pseudocereals and ancient wheat might be considered for creating new health-beneficial food products.

References

Aaron GJ, Laillou A, Wolfson J, Moench-Pfanner R. 2012. Fortification of staple cereal flours with iron and other micronutrients: cost implications of following World Health Organization-endorsed recommendations. Food Nutr Bull 33:S336-43.

Abdel-Aal ESM, Sosulski FW, Hucl P. 1998. Origins, characteristics and potentials of ancient wheats. Cereal Foods World 43:708-15.

Abugoch JLE. 2009. Quinoa (Chenopodium quinoa Wild.): composition, chemistry, nutritional, and functional properties. Adv Food Nutr Res 58:1-3.

Calderelli VAS, de Toledo M, Visentainer JV, Matioli G. 2010. Quinoa and flaxseed: potential ingredients in the production of bread with functional quality. Braz Arch Biol Technol 53:981-6.

Alfredo VO, Gabriel RR, Luis CG, David BA. 2009. Physicochemical properties of a fibrous fraction from chia (Salvia hispanica L.). LWT-Food Sci Technol 42:168-73.

Alvarez-Jubete L, Arendt EK, Gallaghera E. 2009a. Nutritive value of pseudocereals and their increasing use as functional gluten-free ingredients. Trends Food Sci Technol 21:106-13.

Alvarez-Jubete L, Arendt EK, Gallagher E. 2009b. Nutritive value and chemical composition of pseudocereals as gluten-free ingredients. Int J Food Sci Nutr 60:240-57.

Alvarez-Jubete L, Auty M, Arendt EK, Gallagher E. 2010. Baking properties and microstructure of pseudocereal flours in gluten-free bread formulations. Eur Food Res Technol 230:437.

Arzani A. 2011. Emmer (Triticum turgidum ssp. dicoccum) flour and bread. In: Preedy VR, Watson RR, Patel VB, editors. Flour and fortification in healthand disease prevention. Amsterdam: Elsevier, Academic Press Imprint. p 69–78.

564 Ayerza R. 2013. Seed composition of two chia (Salvia hispanica L.) genotypes which differ in seed
565 color. Emir J Food Agric 25:495-500.

566 Balestra F, Laghi L, Saa DT, Gianotti A, Rocculi P, Pinnavaia G. 2015. Physico-chemical and
567 metabolomic characterization of KAMUT® Khorasan and durum wheat fermented dough. Food Chem
568 187:451-59

569 Belderok B. 2000. Developments in bread-making processes. Plant Foods Hum Nutr 55:1-86.

570 Benincasa P, Galieni A, Manetta AC, Pace R, Guiducci M, Pisante M, Stagnari F. 2015. Phenolic
571 compounds in grains, sprouts and wheatgrass of hulled and non-hulled wheat species. J Sci Food Agric
572 95:1795-803.

573 Bhaduri S. 2013. A comprehensive study on physical properties of two gluten-free flour fortified
574 muffins. J Food Process Technol 4:251.

575 Bianchi F, Rossi EA, Gomes RG, Sivieri K. 2015. Potentially synbiotic fermented beverage with
576 aqueous extracts of quinoa (Chenopodium quinoa Willd) and soy. Food Sci Technol Int 21:403-15.

577 Bonafaccia G, Galli V, Francisci R, Mair V, Skrabanja V, Kreft I. 2000. Characteristics of spelt wheat
578 products and nutritional value of spelt wheat-based bread. Food Chem 68:437–41.

579 Borghi B, Castagna R, Corbellini M, Heun M, Salamini F. 1996. Breadmaking quality of einkorn wheat
580 (Triticum monococcum ssp. monococcum). Cereal Chem 73:208–14

581 Borneo R, Aguirre A, León AE. 2010. Chia (Salvia hispanica L) gel can be used as egg or oil replacer
582 in cake formulations. J Am Diet Assoc110:946-9.

583 Brandolini A, Hidalgo A, Moscaritolo S. 2008. Chemical composition and pasting properties of einkorn
584 (Triticum monococcum L. subsp. monococcum) whole meal flour. J Cereal Sci 47:599–609.

585 Brito IL, de Souza EL, Felex SS, Madruga MS, Yamashita F, Magnani M. 2015. Nutritional and sensory
586 characteristics of gluten-free quinoa (Chenopodium quinoa Wild)-based cookies development using an
587 experimental mixture design. J Food Sci Technol 52: 5866-73.

588 Brouns F, van Buula VJ, Shewry, PR. 2013. Does wheat make us fat and sick? J Cereal Sci 58:209–15.

589 Burgos VE, Armada M. 2015. Characterization and nutritional value of precooked products of kiwicha
590 grains (Amaranthus caudatus). Food Sci. Technol (Campinas) 35: 531-8.

591 Cabrera-Chávez F, Iametti S, Miriani M, de la Barca AM, Mamone G, Bonomi F. 2012. Maize
592 prolamins resistant to peptic-tryptic digestion maintain immune-recognition by IgA from some celiac
593 disease patients. Plant Foods Hum Nut 67:24.

594 Callejo MJ, Vargas-Kostiuk ME, Rodríguez-Quijano M. 2015.Selection, training and validation process
595 of a sensory panel for bread analysis: Influence of cultivar on the quality of breads made from common
596 wheat and spelt wheat. J Cereal Sci 61:55-62.

Campo E, del Arco L, Urtasun L, Oria R, Ferrer-Mairal A. 2016. Impact of sourdough on sensory properties and consumers' preference of gluten-free breads enriched with teff flour. J Cereal Sci 67:75–82.

Caperuto L, Amaya-Farfan J, Camargo CR. 2000. Performance of quinoa (Chenopodium quinoa wild) flour in the manufacture of gluten-free spaghetti. J Sci Food Agric 81:95-101.

Carini E, Curti E, Vittadini E. 2010. Effect of long-term storage on water status and physicochemical properties of nutritionally enhanced tortillas. Food Biophys 5:300-8.

Carnevali A, Gianotti A, Benedetti S, Tagliamonte MC, Primiterra M, Laghi L, Danesi F, Valli V, Ndaghijimana M, Capozzi F. 2014. Role of Kamut VR brand khorasan wheat in the counteraction of non-celiac wheat sensitivity and oxidative damage. Food Res Int 63:218-26

Ceccarelli S, Grando S. 2000. Barley landraces from the Fertile Crescent: a lesson for plant breeders. In: Brush SB, editor. Genes in the field: On-farm conservation of crop diversity. Florida: CRC Press p 51.76,

Ceccarelli S. 1996. Adaptation to low /high-input cultivation. Euphytica 92:203-14.

Chandi GK, Lok CW, Jie NY, Seetharaman K. 2015. Functionality of Kamut and millet flours in macro-wire-cut cookie systems. J Food Sci Technol Mys 52:556-61.

Chatzav M, Peleg Z, Ozturk L, Yazici A, Fahima T, Cakmak I, Saranga Y. 2009. Genetic diversity for grain nutrients in wild emmer wheat: potential for wheat improvement. Ann Bot 105:1211-20.

Chillo S, Civica V, Iannetti M, Suriano N, Mastromatteo M, Del Nobile MA. 2009. Properties of quinoa and oat spaghetti loaded with carboxymethylcellulose sodium salt and pregelatinized starch as structuring agents. Carbohydr Polym 78:932-7.

Chlopicka J, Pasko P, Gorinstein A, Zagrodzki P. 2012. Total phenolic and total flavonoid content, antioxidant activity and sensory evaluation of pseudocereal breads. LWT–Food Scie Tech 46:548-55.

Coda R, Nionelli L, Rizzello CG, De Angelis M, Tossut P, Gobbetti M. 2010. Spelt and emmer flours: characterization of the lactic acid bacteria microbiota and selection of mixed starters for bread making. J Appl Microbiol 108:925-35.

Coda R, Rizzello CG, Gobbetti M. 2011. Use of sourdough fermentation and pseudo-cereals and leguminous flours for the making of a functional bread enriched of γ-aminobutyric acid (GABA). Int J Food Microbiol137:236-45.

Coda R, Rizzello CG, Pinto D, Gobbetti M. 2012. Selected lactic acid bacteria synthesize antioxidant peptides during sourdough fermentation of cereal flours. Appl Environ Microb. 78:1087-96.

Coelho MS, Salas-Mellado MM. 2015. Effects of substituting chia (Salvia hispanica L.) flour or seeds for wheat flour on the quality of the bread. Food Sci Technol 60:729-36.

630 Cofrades S, Santos-López JA, Freirea M, Benedí J, Sánchez-Munizc FJ, Jiménez-Colmeneroa F. 2014.
631 Oxidative stability of meat systems made with W1/O/W2 emulsions prepared with hydroxytyrosol and
632 chia oil as lipid phase. LWT-Food Sci Technol 59:941-7.

633 Costantini L, Lukšič L, Molinari R, Kreft I, Bonafaccia G, Manzi L, Merendino N. 2014. Development
634 of gluten-free bread using tartary buckwheat and chia flour rich in flavonoids and omega-3 fatty acids
635 as ingredients. Food Chem 165:232-40.

636 Dallagnol AM, Pescuma M, De Valdez GF, Rollán G. 2013. Fermentation of quinoa and wheat slurries
637 by Lactobacillus plantarum CRL 778: proteolytic activity. Appl Microbiol Biotechnol 97:3129-40.

638 De la Barc AMC, Rojas-Martínez ME, Islas-Rubio AR, Cabrera-Chávez F. 2010. Gluten-free breads
639 and cookies of raw and popped amaranth flours with attractive technological and nutritional qualities.
640 Plant Foods Hum Nutr 65:241-6.

641 D'Egidio MG, Mariani BM, Nardi S, Novaro P and Cubadda R. 1990. Chemical and technological
642 variables and their relationships: A predictive equation for pasta cooking quality. Cereal Chem 67:275-
643 81.

644 de Souza RJ, Mente A, Maroleanu A, Cozma AI, Ha V, Kishibe T, Uleryk E, Budylowski P,
645 Schünemann H, Beyene J, Anand SS. 2015. Intake of saturated and trans unsaturated fatty acids and risk
646 of allcause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis
647 of observational studies. BMJ. 351:h3978.

648 De Vita P, Li Destri Nicosia O, Nigro F, Platani C, Riefolo C, Di Fonzo N, Cattivelli L. 2007. Breeding
649 progress in morpho-physiological, agronomical and qualitative traits of durum wheat cultivars released
650 in Italy during the 20th century. Eur J Agron 26:39-53.

651 Dexter JE, Matsuo RR. 1977. Influence of protein content on some durum wheat quality parameters.
652 Can J Plant Sci 57:717-27.

653 Deželak M, Zarnkow M, Becker T, Košir IJ. 2014. Processing of bottom-fermented gluten-free beer-
654 like beverages based on buckwheat and quinoa malt with chemical and sensory characterization. J Inst
655 Brew 120:360-70.

656 Di Silvestro R, Marotti I, Bosi S, Bregola V, Segura-Carretero A, Sedej I, Mandic A, Sakac M,
657 Benedettelli S, Dinelli G. 2012. Health-promoting phytochemicals of Italian common wheat varieties
658 grown under low-input agricultural management. J Sci Food Agric 92:2800-10.

659 Diao XP. 2017. Production and genetic improvement of minor cereals in China. Crop J 5:103-14.

660 Dinelli G, Marotti I, Di Silvestro R, Bosi S, Bregola V, Accorsi M, Di Loreto A, Benedettelli S, Ghiselli
661 L, Catizone P. 2013. Agronomic, nutritional and nutraceutical aspects of durum wheat (Triticum durum
662 Desf.) cultivars under low-input agricultural management. Ital J Agron 8:85-93.

663 Dinelli G, Segura-Carretero A, Di Silvestro R, Marotti I, Arráez-Román D, Benedettelli S, Ghiselli L,
664 Fernandez-Gutierrez A. 2011. Profiles of phenolic compounds in modern and old common wheat

varieties determined by liquid chromatography coupled with time-of-flight mass spectrometry. J Chromatogr A 1218:7670-81.

Dinelli G, Segura-Carretero A, Di Silvestro R, Marotti I, Fu S, Benedettelli S, Ghiselli L. 2009. Determination of phenolic compounds in modern and old varieties of durum wheat using liquid chromatography coupled with time-of-flight mass spectrometry. J Chromatogr 1216:7229-40.

Dziadek K, Kopeć A, Pastucha E, Piątkowska E, Leszczyńska T, Pisulewska E, Witkowicz R, Francik R. 2016. Basic chemical composition and bioactive compounds content in selected cultivars of buckwheat whole seeds, dehulled seeds and hulls. J Cereal Sci 69:1-8.

El-Deeb AM, Hassan NSY, Hassanein AM. 2014. Preparation and properties of flavored fermented beverage based on partial or complete replacement of milk with quinoa seeds water extract (QSWE). Inter J Dairy Sci 9:96-1.

Enriquez N, Peltzer M, Raimundi A, Tosi V, Pollio ML. 2003. Characterization of the wheat and quinoa flour blends in relation to their bread making quality. J Arg Chem Soc 91: 47-54.

Erba D, Hidalgo A, Bresciani J and Brandolini A. 2011. Environmental and genotypic influences on trace element and mineral concentrationsin whole meal flour of einkorn (Triticum monococcum L. subsp.monococcum). J Cereal Sci 54:250-4.

Escarnot E, Jacquemin JM, Agneessens R, Paquot M. 2012. Comparative study of the content and profiles of macronutrients in spelt and wheat, a review. Biotechnol Agron Soc Environ 16:243-56.

Fares C, Codianni P, Nigro F, Platani C, Scazzina F, Pellegrini N. 2008. Processing and cooking effects on chemical, nutritional and functional properties of pasta obtained from selected emmer genotypes. J Sci Food Agric 88:2435-44.

Faris JD, Zhang Z, Chao S. 2014. Map-based analysis of the tenacious glume gene Tg-B1 of wild emmer and its role in wheat domestication. Gene 542:198-208.

Fiorda FA, Soares M, da Silva FA, Souto LRF, Grosmann MVE. 2013. Amaranth flour, cassava starch and cassava bagasse in the production of gluten-free pasta: Technological and sensory aspects International. Int J Food Sci Technol 48:1977-84.

Fogarasi AL, Kun S, Tankó G, Stefanovits-Bányai E, Hegyesné-Vecseri B. 2015. A comparative assessment of antioxidant properties, total phenolic content of einkorn, wheat, barley and their malts. Food Chem 167:1-6.

Fois S, Schlichting L, Marchylo B, Dexter J, Motzo R, Giunta F. 2011. Environmental conditions affect semolina quality in durum wheat (Triticum turgidum ssp. durum L.) cultivars with different gluten strength and gluten protein composition. J Sci Food Agric 91:2664-73.

Fuller DQ. 2007. Contrasting patterns in crop domestication and domestication rates: Recent archaeobotanical insights from the Old World. Ann Bot (Lond) 100:903-24.

699 Gawlik-Dziki U, Dziki D, Baraniak B, Lin R. 2009. The effect of simulated digestion in vitro on
700 bioactivity of wheat bread with Tartary buckwheat flavones addition. LWT - Food Sci Technol 42:137-
701 43.

702 Gegas VC, Nazari A, Griffiths S, Simmonds J, Fish L, Orford S, Sayers L, Doonan JH, Snapea JW.
703 2010. A Genetic framework for grain size and shape variation in wheat. Plant Cell 22:1046-56.

704 Giambanelli E, Ferioli F, Kocaoglu B, Jorjadze M, Alexieva I, Darbinyan N, Antuono F. 2013. A
705 comparative study of bioactive compounds in primitive wheat populations from Italy, Turkey, Georgia,
706 Bulgaria and Armenia. J Sci Food Agric 93:3490-501.

707 Giunta F, Motzo R, Pruneddu G. 2007. Trends since 1900 in the yield potential of Italian-bred durum
708 wheat cultivars. Europ J Agronomy 27:12–24.

709 Gobbetti M, Rizzello CG, Di Cagno R, De Angelis M. 2014. How the sourdough may affect the
710 functional features of leavened baked goods. Food Microbiol 37:30-40.

711 Graf BL, Rojas-Silva P, Rojo LE, Delatorre-Herrera J, Baldeón ME, Raskin I. 2015. Innovations in
712 health value and functional food development of quinoa (Chenopodium quinoa Wild.). Comp Rev Food
713 Sci Food 14:431-45.

714 Guo CJ, Chang FY, Wyche TP, Backus KM, Acker TM, Funabashi M, Taketani M, Donia MS, Nayfach
715 S, Pollard KS. 2017. Discovery of reactive microbiota-derived metabolites that inhibit host proteases.
716 Cell 168:517-26.

717 Guzmán C, Caballero L, Martín LM, Alvarez JB. 2013. Waxy genes from spelt wheat: new alleles for
718 modern wheat breeding and new phylogenetic inferences about the origin of this species. Ann Bot
719 110:1161-71.

720 Hager AS, Wolter A, Czerny M, Bez J, Zannini E, Arendt EK, Czerny M. 2012. Investigation of product
721 quality, sensory profile and ultrastructure of breads made from a range of commercial gluten-free flours
722 compared to their wheat counterparts. Eur Food Res Technol 235:333-44.

723 Hammer K. 1984. Das Domestikationssyndrom. Kulturpflanze 2:11–34.

724 Hebelstrup KH. 2017. Differences in nutritional quality between wild and domesticated forms of barley
725 and emmer wheat. Plant Sci 256:1-4.

726 Herrero AM, Ruiz-Capillas C, Pintado T, Carmona P, Jimenez-Colmenero F. 2017. Infrared
727 spectroscopy used to determine effects of chia and olive oil incorporation strategies on lipid structure of
728 reduced-fat frankfurters. Food Chem 221:1333-9.

729 Hidalgo A, Brandolini A, Pompeia C, Piscozzi R. 2006. Carotenoids and tocols of einkorn wheat
730 (Triticum monococcum ssp. monococcum L.) J Cereal Sci 44:182-93.

731 Hidalgo A, Brandolini A, Ratti S. 2009. Influence of genetic and environmental factors on selected
732 nutritional traits of Triticum monococcum. JAgric Food Chem 57:6342-8.

733 Hidalgo A, Brandolini A, Pompei C. 2010. Carotenoids evolution during pasta, bread and water biscuit
734 preparation from wheat flours. Food Chem 121:746-51.

735 Hidalgo A, Brandolini A. 2014. Nutritional properties of einkorn wheat (Triticum monococcum L.). J
736 Sci Food Agric 94:601-12.

737 Hidalgo A, Brandolini A. 2017. Nitrogen fertilisation effects on technological parameters and
738 carotenoid, tocol and phenolic acid content of einkorn (Triticum monococcum L. subsp. monococcum):
739 A two-year evaluation. J Cereal Sci 73:18-24

740 Hidalgo A, Brandolini A. 2011. Heat damage of water biscuits from einkorn, durum and bread wheat
741 flours. Food Chem 128:471-8.

742 Hidalgo A, Brusco M, Plizzari L, Brandolini A. 2014. Polyphenol oxidase, alpha-amylase and beta-
743 amylase activities of T. monococcum, T. turgidum and T. aestivum: a two-year study. J Cereal Sci
744 58:51-8.

745 Hidalgo A, Yilmaz VA, Brandolini A. 2016. Influence of water biscuit processing and kernel puffing
746 on the phenolic acid content and the antioxidant activity of einkorn and bread wheat. J Food Sci Technol
747 53:541-50.

748 Hooper DU, Chapin III F.S, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM, Loreau
749 M, Naeem S, Schmid B, Setälä H, Symstad AJ, Vandermeer J, Wardle DA. 2005. Effects of biodiversity
750 on ecosystem functioning: A consensus of current knowledge. Ecol Monogr 75:3-35.

751 Huerta KM, Alves JS, da Silva AFC, Kubota EH, da Rosa CS. 2016. Sensory response and physical
752 characteristics of gluten-free and gum-free bread with chia flour. Food Sci Technol (Campinas) 36:15-
753 8.

754 Hussain A, Larsson H, Kuktaite R, Johansson E. 2010. Mineral composition of organically grown wheat
755 genotypes: contribution to daily minerals intake. Int J Environ Res Public Health 7: 3442-56.

756 Hussain A, Larsson H, Kuktaite R, Johansson E. 2012. Concentration of some heavy metals in
757 organically grown primitive, old and modern wheat genotypes: implications for human health. J Environ
758 Sci Health B 47:751-8.

759 Iglesias-Puig E, Haros M. 2013. Evaluation of performance of dough and bread incorporating chia
760 (Salvia hispanica L.). Eur Food Res Technol 237:865–74.

761 Iglesias-Puig E, Monedero V, Haros M. 2015. Bread with whole quinoa flour and bifidobacterial
762 phytases increases dietary mineral intake and bioavailability. LWT-Food Sci Technol 60:71-7.

763 Ixtaina VY, Nolasco SM, Tomas MC. 2008. Physical properties of chia (Salvia hispanica L.) seeds. Ind
764 Crop Prod 28:286–93.

765 Jozinović A, Šubarić D, Ačkar Đ, Babić J, Miličević B. 2016. Influence of spelt flour addition on
766 properties of extruded products based on corn grits. J Food Eng 172:31-37

767 Kang M, Zhai FeH, Li XX, Cao JL, Han JR. 2017.Total phenolic contents and antioxidant properties of
768 buckwheat fermented by three strains of Agaricus. J Cereal Sci 73:138-142

769 Kohajdová Z, Karovičov J. 2008. Nutritional value and baking applications of spelt wheat. Acta Sci Pol
770 Technol Aliment 7:5-14.

771 Konvalina P, Capouchová I, Stehno Z, Moudrý J, Moudrý J. 2010. Weaknesses of emmer wheat genetic
772 resources and possibilities of its improvement for low-input and organic farming systems. Int j food
773 agric and environ 8:376-82.

774 Koppalu V, Kumara P, Dharmaraja U, Sakhareb SD, Inamdarb A A. 2016. Preparation of protein and
775 mineral rich fraction from grain amaranth and evaluation of its functional characteristics. J Cereal Sci
776 69:358-62.

777 Korczyk-Szabó J, Lacko-Bartošová M. 2013. Crumb texture of spelt bread. J Cent Eur Agr 14:1343-52.

778 Korczyk-Szabó J, Lacko-Bartošová M. 2015. Textural properties of spelt noodles. Acta fytotechnica et
779 zootechnica 18:43-4.

780 Kucek LK, Dyck E, Russell J, Clark L, Hamelman J, Leader SB, Senders S, Jones J, Benscher D, Davis
781 M, Roth G, Zwinger S, Sorrells ME, Dawson JC. 2017. Evaluation of wheat and emmer varieties for
782 artisanal baking, pasta making, and sensory quality. J Cereal Sci 74:19-27.

783 Kumar KV, Dharmaraj U, Sakhare SD, Inamdar AA. 2016. Effect of grain moisture content during
784 milling on pasting profile and functional properties of amaranth fractions. J Food Sci Technol 53:2434-
785 42.

786 la Gatta B, Rutigliano M, Rusco G, Petrella G, Di Luccia A. 2017. Evidence for different supramolecular
787 arrangements in pasta from durum wheat (Triticum durum) and einkorn (Triticum monococcum) flours.
788 J Cereal Sci 73:76-83.

789 Lachman J, Musilova J, Kotikova Z, Hejtmankova K, Orsak M, Pribyl J. 2012. Spring, einkorn and
790 emmer wheat species-potential rich sources offree ferulic acid and other phenolic compounds. Plant Soil
791 Environ 58:347-53.

792 Lampi AM, Nurmi T, Ollilainen V, Piironen V. 2008. Tocopherols and tocotrienols in wheat genotypes
793 in the HEALTHGRAIN diversity screen. J Agric Food Chem 56: 9716-21.

794 Li G, Zhu F. 2017. Physicochemical properties of quinoa flour as affected by starch interactions. Food
795 Chem 221: 560–8.

796 Longin CFH, Würschum T. 2016. Back to the future – tapping into ancient grains for food diversity.
797 Trends in Plant Science 21:731-7.

798 Lopes MS, El-Basyoni I, Baenziger PS, Singh S, Royo C, Ozbek K, Aktas H, Ozer E, Ozdemir F,
799 Manickavelu A, Ban T, Vikram P. 2015. Exploiting genetic diversity from landraces in wheat breeding
800 for adaptation to climate change. J Exp Bot 66:3477-86.

Lorusso A, Verni M, Montemurro M, Coda R, Gobbetti M, Rizzello CG. 2017. Use of fermented quinoa flour for pasta making and evaluation of the technological and nutritional features. LWT - Food Sci Technol 78:215-21.

Ludena Urquizo FE, García Torres SM, Tolonen T, Jaakkola M, Pena-Niebuhr MG, von Wright A, Repo-Carrasco-Valencia R, Korhonen H, Plumed-Ferrer C. 2017. Development of a fermented quinoa-based beverage. Food Sci Nutr 5:602-8.

Marconi E, Carcea M, Graziano M, Cubadda R. 1999. Kernel properties and pasta-making quality of five European spelt wheat (Triticum spelta L.) cultivars. Cereal Chem 76:25-9.

Marconi E, Carcea M, Schiavone M, Cubadda R. 2002. Spelt (Triticum spelta L.) Pasta quality: combined effect of flour properties and drying conditions. Cereal Chem 79:634-9.

Marconi O, Mayer H, Chiacchieroni F, Ricci E, Perretti G, Fantozzi P. 2013. The influence of glumes on malting and brewing of hulled wheats. J Am Soc Brew Chem 71:41-8.

Mariottia M, Lucisanoa M, Ambrogina Pagania M, Perry K.W. 2009. The role of corn starch, amaranth flour, pea isolate, and Psyllium flour on the rheological properties and the ultrastructure of gluten-free doughs. Food Res Int 42:963-75.

Mayer H, Marconi O, Perretti G, Sensidoni M, Fantozzi P. 2011. Investigation of the suitability of hulled wheats for malting and brewing. J Am Soc Brew Chem 69:116-20.

Menga V, Amato M, Phillips TD, Angelino D, Morreale F, Fares C. 2017. Gluten-free pasta incorporating chia (Salvia hispanica L.) as thickening agent: An approach to naturally improve the nutritional profile and the in vitro carbohydrate digestibility. Food Chem 221:1954-61.

Messia MC, Iafelice G, Marconi E. 2012. Effect of parboiling on physical and chemical characteristics and non-enzymatic browning of emmer (Triticum dicoccon Schrank). J Cereal Sci 56:147-52.

Michalcová V, Dušinský R, Sabo M, Al Beyroutiová. 2014. Taxonomical classification and origin of Kamut wheat. Plant Syst Evol 300:1749-57.

Miranda M, Vega-Gálvez A, Uribe E, Lópeza J, Martínez E, Rodrígueza MJ, Quispea I, Di Scalac K. 2012. Physico-chemical analysis, antioxidant capacity and vitamins of six ecotypes of Chilean quinoa (Chenopodium quinoa Willd). Procedia Food Science, 1, 1439–1446.

Moreira R, Chenlo F, Torres MD. 2013. Effect of chia (Sativa hispanica L.) and hydrocolloids on the rheology of gluten-free doughs based on chestnut flour. LWT Food Sci Technol 50: 160-6.

Mohammadi R, Sadeghzadeh B, Ahmadi H, Bahrami N, Amri A. 2015. Field evaluation of durum wheat landraces for prevailing abiotic and biotic stresses in highland rainfed regions of Iran. Crop J 3:423-33.

Mohan D, Singh AM, Ahlawat AK, Gupta RK. 2014. Analogy between agronomic and grain quality attributes of wheat for response to crop seasons, locations, site-year and genotype-environment interactions. J Wheat Res 6:27–32.

Mohd Ali N, Yeap SK, Ho WY, Beh BK, Tan SW, Tan SG. 2012. The Promising Future of Chia, Salvia hispanica L. J Biomed Biotechnol 2012:171956.

Moreno ML, Comino I, Sousa C. 2014. Alternative grains as potential raw material for gluten-free food development in the diet of celiac and gluten-sensitive patients. Austin J Nutri Food Sci 2:9.

Moroni AV, Arendt EK, Dal Bello F. 2011. Biodiversity of lactic acid bacteria and yeasts in spontaneously-fermented buckwheat and teff sourdoughs. Food Microbiol 28:497-502.

Moroni AV, Arendt EK, Morrissey JP, Dal Bello F. 2010. Development of buckwheat and teff sourdoughs with the use of commercial starters. Int J Food Microbiol142:142-8.

Moroni AV, Dal Bello F, Arendt EK. 2009. Sourdough in gluten-free bread-making: an ancient technology to solve a novel issue? Food Microbiol 26:676-84.

Motzo R, Fois S, Giunta F. 2004. Relationship between grain yield and quality of durum wheats from different eras of breeding. Euphytica 140:147-54.

Muñoz-Insa A, Selciano H, Zarnkow M, Becker T, Gastl M. 2013. Malting process optimization of spelt (Triticum spelta L.) for the brewing process. LWT - Food Sci Technol 50:99–109.

Nazco R, Villegas D, Ammar K, Peña RJ, Moragues M, Royo C. 2012. Can Mediterranean durum wheat landraces contribute to improved grain quality attributes in modern cultivars? Euphytica 185:1–17.

Ngugi CC, Oyoo-Okoth E, Manyala JO, Fitzsimmons K, Kimotho A. 2017. Characterization of the nutritional quality of amaranth leaf protein concentrates and suitability of fish meal replacement in Nile tilapia feeds. Aquaculture Reports 5:62-9

Nionelli L, Rizzello CG. 2016. Sourdough-based biotechnologies for the production of gluten-free foods. Foods 5:65.

Nowak V, Du J, Charrondière UR. 2016. Assessment of the nutritional composition of quinoa (Chenopodium quinoa Wild.). Food Chem 193:47-54.

Okuno A, Hirano K, Asano K, Takase W, Masuda R, Morinaka Y, Ueguchi-Tanaka M, Kitano H, Matsuoka M. 2014. Approach to increasing rice lodging resistance and biomass yield through the use of high gibberellin producing varieties. PLoS One 9:e86870.

Olivos-Lugo BL, Valdivia-López MÁ, Tecante A. 2010. Thermal and physicochemical properties and nutritional value of the protein fraction of mexican chia seed (Salvia hispanica L.). Food Sci Technol Int 16:89-96.

Ozkan H, Willcox G, Graner A, Salamini F, Kilian B. 2011. Geographic distribution and domestication of wild emmer wheat (Triticum dicoccoides). Genet Resour Crop Evol 58:11–53

Pasini G, Greco F, . Cremonini MA, Brandolini A, Consonni R, Gussoni M. 2015. The structural and nutritional properties of pasta from triticum monococcum and triticum durum species. A combined 1H NMR, MRI, and digestibility study. J Agric Food Chem 63:5072-82.

869 Pasqualone A, Piergiovanni AR, Caponio F, Paradiso VM, Summo C, Simeone R. 2011. Evaluation of
870 the technological characteristics and bread-making quality of alternative wheat cereals in comparison
871 with common and durum wheat. Food Sci Technol Int 17:135–42.

872 Peleg Z, Fahima T, Korol AB, Abbo S, Saranga Y. 2011. Genetic analysis of wheat domestication and
873 evolution under domestication. J Exp Bot 62:5051–61.

874 Pellegrini N, Agostoni C. 2015. Nutritional aspects of gluten-free products. J Sci Food Agric 95:2380-
875 5.

876 Pintado T, Herrero AM, Jiménez-Colmenero F, Ruiz-Capillas C. 2016. Strategies for incorporation of
877 chia (Salvia hispanica L.) in frankfurters as a health-promoting ingredient. Meat Sci 114:75-84.

878 Pintado T, Ruiz-Capillas C, Jiménez-Colmenero F, Carmona P, Herrero AM. 2015. Oil-in-water
879 emulsion gels stabilized with chia (Salvia hispanica L.) and cold gelling agents: Technological and
880 infrared spectroscopic characterization. Food Chem 185:470-8.

881 Randall P, Johnson Q, Verster A. 2012. Fortification of wheat flour and maize meal with different iron
882 compounds: results of a series of baking trials. Food Nutr Bull 33:S344-59.

883 Ranhotra GS, Gelroth JA, Glaser BK, Lorenz KJ. 1996. Nutrient composition of spelt wheat. J Food
884 Comp Anal 9:81-4.

885 Rendón-Villalobos R, Ortíz-Sánchez A, Solorza-Feria J, Trujillo-Hernández CA. 2012. Formulation,
886 physicochemical, nutritional and sensorial evaluation of corn tortillas supplemented with chía seed
887 (Salvia hispanica L.). Czech J Food Sci 30:118-25.

888 Rizzello CG, Cavoski I, Turk J, Ercolini D, Nionelli L, Pontonio E, De Angelis M, De Filippis F,
889 Gobbetti M, Di Cagno R. 2015. Organic cultivation of Triticum turgidum subsp. durum is reflected in
890 the flour-sourdough fermentation-bread axis. Appl Environ Microbiol 81:3192-204.

891 Rizzello CG, Lorusso A, Montemurro M, Gobbetti M. 2016. Use of sourdough made with quinoa
892 (Chenopodium quinoa) flour and autochthonous selected lactic acid bacteria for enhancing the
893 nutritional, textural and sensory features of white bread. Food Microbiol 56:1-13.

894 Rizzello CG, Lorusso A, Russo V, Pinto D, Marzani B, Gobbetti M. 2017. Improving the antioxidant
895 properties of quinoa flour through fermentation with selected autochthonous lactic acid bacteria. Int J
896 Food Microbiol 241:252-61.

897 Rodriguez-Sandoval E, Sandoval G, Cortes-Rodríguez M. 2012. Effect of quinoa and potato flours on
898 the thermomechanical and breadmaking properties of wheat flour. Braz Arch Biol Technol 29:503-10.

899 Royo C, Álvaro F, Martos V, Ramdani A, Isidro J, Villegas D, García del Moral LF. 2007. Genetic
900 changes in durum wheat yield components and associated traits in Italian and Spanish varieties during
901 the 20th century. Euphytica 155:259-70.

902 Sakac M, Torbica A, Sedej I, Hadnadev M. 2011. Influence of bread making on antioxidant capacity of
903 gluten free breads based on rice and buckwheat flours. Food Res Int 44:2806-13.

Sakuma S, Salomon B, Komatsuda T. 2011. The domestication syndrome genes responsible for the major changes in plant form in the triticeae. Crops Plant Cell Physiol 52:738-49.

Salamini F, Ozkan H, Brandolini A, Schafer-Pregl R, Marin W. 2002. Genetics and geography of wild cereal domestication in the Near East. Nat Rev Genet 3:420-41.

Salse J, Chagué V, Bolot S, Magdelenat G, Huneau C, Pont C, Belcram H, Couloux A, Gardais S, Evrard A, Segurens B, Charles M, Ravel C, Samain S, Charmet G, Boudet N, Chalhoub B. 2008. New insights into the origin of the B genome of hexaploid wheat: evolutionary relationships at the SPA genomic region with the S genome of the diploid relative Aegilops speltoides. BMC Genomics 9:555.

Sanchez-Garcia M, Álvaro F. Peremarti A, Martín-Sánchez JA, Royo C. 2015. Changes in bread-making quality attributes of bread wheat varieties cultivated in Spain during the 20th century. Europ J Agronomy 63:79-88

Santra DK, Schoenlechner R. 2017. Amaranth part 2 sustainability, processing, and applications of amaranth sustainable protein sources. In Nadathur S, Wanasundara JPD, Scanlin L, editors. Sustainable Protein Sources. Amsterdam: Elsevier Inc. p 257–64.

Saturni L, Ferretti G, Bacchetti T. 2010. The gluten-free diet: safety and nutritional quality. Nutrients 2:16–34.

Schoenlechner R, Drausinger J, Ottenschlaeger V, Jurackova K, Berghofer E. 2011. Functional properties of gluten-free pasta produced from amaranth, quinoa and buckwheat. Plant Foods Hum Nutr 65:339-49.

Singh N, Singh P. 2011. Amaranth: Potential source for flour enrichment. In Preedy V, Watson R, Patel V, editors. Flour and breads and their fortification in health and disease prevention. Amsterdam: Elsevier Inc. p 101-11.

Soriano JM, Villegas D, Aranzana MJ, García Del Moral LF, Royo C. 2016. Genetic structure of modern durum wheat cultivars and mediterranean landraces matches with their agronomic performance. PLoS One 11:e0160983.

Starr G, Bredie WLP, Hansen ÅS. 2013. Sensory profiles of cooked grains from wheat species and varieties. J Cereal Sci 57:295-303.

Steffolani ME, León AE, Pérez GT. 2014. Study of the physicochemical and functional characterization of quinoa and kañiwa starches. Starch-Stärke 65:976-83.

Stikic R, Glamoclija D, Demin M, Vucelic-Radovic B, Jovanovic Z, Milojkovic Opsenica D. 2012. Agronomical and nutritional evaluation of quinoa seeds (Chenopodium quinoa Wild.) as an ingredient in bread formulations. J Cereal Sci 55:132-8.

Suchowilska E, Wiwart M, Borejszo Z, Packa D, Kandler W, Krska R. 2009. Discriminant analysis of selected yield components and fatty acidcomposition of chosen Triticum monococcum, Triticum dicoccum and Triticum spelta accessions. J Cereal Sci 49:310–5.

939 Sumczynski D, Bubelova Z, Sneyd J, Erb-Weber S, Mlcek J. 2015. Total phenolics, flavonoids,
940 antioxidant activity, crude fiber and digestibility in non-traditional wheat flakes and muesli. Food Chem
941 174:319–25.

942 The International Brachypodium Initiative, 2010

943 The international report from Health Focus International 2015

944 Torbica A, Hadnađev M, Dapčević T. 2010. Rheological, textural and sensory properties of gluten-free
945 bread formulations based on rice and buckwheat flour. Food Hydrocoll 24:626–32.

946 Troccoli A, Codianni P. 2005. Appropriate seeding rate for einkorn, emmer, and spelt grown under
947 rainfed conditions in southern Italy. Eur J Agron 22: 293-300.

948 United States Department of Agriculture. USDA Food Composition Database. Available from:
949 http://ndb.nal.usda.gov/.

950 Vega-Gálvez A, Miranda M, Vergara J, Uribe E, Puente L, Martínez EA. 2010. Nutrition facts and
951 functional potential of quinoa (Chenopodium quinoa willd.), an ancient Andean grain: a review. J Sci
952 Food Agric 90, 2541-7.

953 Verdú S, Vásquez F, E. Ivorra E, Sánchez AJ, Barat JM, Grau R. 2015. Physicochemical effects of chia
954 (Salvia Hispanica) seed flour on each wheat bread-making process phase and product storage. J Cereal
955 Sci 65:67–73.

956 Vita F, Taiti C, Pompeiano A, Gu Z, Presti EL, Whitney L, Monti M, Di Miceli G, Giambalvo D, Ruisi
957 P, Guglielminetti L, Mancuso S. 2016. Aromatic and proteomic analyses corroborate the distinction
958 between Mediterranean landraces and modern varieties of durum wheat. Sci Rep 6:34619

959 Wang S, Opassathavorn A, Zhu F. 2014. Influence of quinoa flour on quality characteristics of cookies,
960 bread, and Chinese steamed bread. J Texture Stud 46:281-92.

961 Wang S, Zhu F. 2015. Formulation and quality attributes of quinoa food product. Food Bioproc Tech 9:
962 49-68.

963 Wolter A, Hager A, Zannini E, Czerny M, Arendt EK. 2014. Influence of dextran-producing Weissella
964 cibaria on baking properties and sensory profile of gluten-free and wheat breads. Inter J Food Microbiol
965 172:83-9.

966 Wolter A, Hager AS, Zannini E, Galleb S, Gänzleb MG, Watersa, DM Arendt EK. 2013. Evaluation of
967 exopolysaccharide producing Weissella cibaria MG1 strain for the production of sourdough from
968 various flours. Food Microbiol 37:44-50.

969 Wronkowska M, Zielinska D, Szawara-Nowak D, Troszynska A, Soral-Semietana M. 2010.
970 Antioxidative and reducing capacity, macroelements content and sensorial properties of buckwheat-
971 enhanced gluten-free bread. Inter J Food Sci Tech 5:1993–2000.

Wu G, Peterson AJ, Morris CF, Murphy KM. 2016. Quinoa seed quality response to sodium chloride and sodium sulfate salinity. Front Plant Sci 7:790.

Zhou Z, Jiang Y, Wang Z, Gou Z, Lyu J, Li W, Yu Y, Shu L, Zhao Y, Ma Y, Fang C, Shen Y, Liu T, Li C, Li Q, Wu M, Wang M, Wu Y, Dong Y, Wan W, Wang X, Ding Z, Gao Y, Xiang H, Zhu B, Lee S, Wang W, Tian Z. 2015. Resequencing 302 wild and cultivated accessions identifies genes related to domestication and improvement in soybean. Nature Biotech 33:408-14.

Zohary D, Hopf M. 2000. Cereals. In: Domestication of plants in the old world: the origin and spread of cultivated plants in West Asia, Europe, and the Nile Valley. New York: Oxford University Press. p 316.

Figure caption

Figure 1: Major classes and subclasses of ancient grains. A: Classification of ancient grains; B: Taxonomy of the most cultivated cereals; C: Taxonomy of the most known pseudocereals. This figure explains the classes and subclasses of ancient grains in terms of ancient wheat and pseudocereals.

Table caption

Table 1: Glossary. This table gives the definitions of the key terms used for wheat and pseudocereals.

Table 2: Gluten-containing grains. This table focuses on the main ancient wheat characteristics.

Table 3: Gluten-free grains. This table shows the main characteristics of pseudocereals.

Table 4: Approximate chemical composition of ancient wheat, durum wheat, and common wheat. This table contains data reported about the chemical composition of ancient wheat in terms of macro- and micronutrients in comparison with durum and common wheat.

Table 5: Approximate chemical composition of pseudocereals compared to maize and rice. This table contains data reporting on the chemical composition of pseudocereals in terms of macro- and micronutrients in comparison with maize and rice.

A



1019

B



1020

C



1021

Figure 1: Major classes and subclasses of ancient grains. A: Classification of ancient grains; B: Taxonomy of the most cultivated cereals; C: Taxonomy of the most known pseudocereals.

Table 1: Glossary.

Key terms	Definition
Wild species or wild ancestors	are the species naturally grown in the Old World, before any cultivation and domestication*. *Triticum urartu* (AA) and *Triticum boeoticum* (wild einkorn, AA), *Triticum turgidum* ssp. *dicoccoides* (wild emmer, AABB) are the wild progenitors of modern wheat.
Domesticated* wheat	is obtained by the selection of novel spontaneous mutations or recessive alleles in cultivated populations or among the wild populations (Hebelstrup 2017).
Ancient grains	are represented by populations of **primitive grains**, which were not subject to any modern breeding or selection, and which retained characters of wild ancestors, such as large individual variability, ear height, brittle rachis, and low harvest index (Giambanelli and others 2013).
Ancient wheat	refers to emmer, einkorn, Khorasan wheat (Oriental wheat) and spelt.
Pseudocereals	are mainly amaranth, quinoa, buckwheat, and chia.
Landraces and old varieties	are developed by natural and human selection, genetically heterogeneous, locally adapted, and they too were cultivated until the middle of the 20th century (Nazco and others 2012; Lopes and others 2015; Mohammadi and others 2015; Soriano and others 2016).
Modern varieties	are the result of the continued modern breeding progress, aiming to select homogeneous lines with stable and improved characters, mainly RHt dwarfing gene to avoid lodging. Currently, thanks to international breeding programs, these varieties are cultivated worldwide ensuring higher productivity than landraces.

**Note: The shift from wild to domesticated involved an evolutionary process of morphological, physiological, and genetic events, referred to as 'domestication syndrome' (Hammer 1984). The two most important events of the domestication syndrome are: a non- brittle rachis mutation that resulted in non-shattering domesticated wheat and a non-hulled mutation which resulted in free-threshing domesticated wheat, where the husk covering the seed comes off during threshing. Domestication involved other traits, such as increase in both seed size and number, loss of germination inhibition, lower grain protein and mineral concentrations, and increased grain carbohydrate content (Zohary and Hopf 2000; Fuller 2007; Gegas and others 2010; Sakuma and others 2011; Peleg and others 2011). Triticum aestivum L. and Triticum turgidum subsp. durum, which are free-threshing wheat, represent the final step of Triticum domestication (Salamini and others 2002).*

Table 2: Gluten-containing grains.

Common name	Binomial name	Origin/History	References
Einkorn wheat	*Triticum monococcum* L., subsp. *monococcum* (2n = 2x = 14, AA),	Domesticated diploid wheat from ssp. *aegilopoides* through the acquisition of a non-brittle rachis.	(Hidalgo and Brandolini 2017; Faris and others 2014).
Emmer wheat, also known as farro in Italy	*Triticum turgidum dicoccum* (2n = 4x =28, genome AABB)	The domesticated form of wild emmer, which derived from *T. urartu*, (2n = 2x = 14, AA), donor of the genome A, and another unknown species of the *Sitopsis* section, donor of the B genome, for which the closest known relative is goat grass (*Aegilops speltoides*, 2n = 2x = 14, SS).	(Salse and others 2008; Chatzav and others 2009; Ozkan and others 2010; Peng and others 2011).
Spelt wheat	*Triticum aestivum subsp. spelta* (2n = 6x= 42; genome AABBDD)	It is suggested to be the ancestral form of *T. aestivum*; however, it is also hypothesized to be probably derived from a secondary hybridization between emmer wheat and a hexaploid wheat (*T. aestivum* L. ssp. *compactum* Host em).	(Dovrak and others 2011; Guzmán and others 2012)
Khorasan wheat or Kamut®	*Triticum turgidum* subsp. *turanicum* (2n = 4x= 28, genome AABB)	This wheat originated in the Khorasan region and was always known in the Mediterranean basin, where many populations exist. Its origin is probably as far in time as durum wheat. Kamut® is the registered trademark for Khorasan wheat produced under the controlled value chain of Kamut International.	(Michalcová and others 2014)

1079 **Table 3:** Gluten-free grains.

Common name	Main species	Origin/History	References
Buckwheat	The two major species are common buckwheat (sweet) (*Fagopyrum esculentum* Moench) and tartary buckwheat (*Fagopyrum tataricum* Gaertn.)	One of the traditional crops cultivated in Asia and Central and Eastern Europe.	(Moreno and others 2014; Kang and others 2017)
Quinoa	The two commercial varieties are *Amarilla de Marangani* and *Blanca de Junın*	Among the most popular crops for the people of rural South America.	(Vega-Gálvez and others 2010; Graf and others 2015)
Amaranth	60 plant species, *Amaranthus cruentus*, *A. hypochondriacus*, and *A. Caudatus* are the main cultivated amaranth species for grain, whereas *A. cruentus, A. blitum, A. dubius*, and *A. tricolor* are used as leafy vegetables	It was once a staple food of the Aztecs.	(Singh and Singh, 2011).
Chia	Three main species: *Salvia columbariae* Benth., *Salvia polytachya* Cavan., *and Salvia hispanica L.*.	It was found growing wild in Mexico.	(Verdú and others 2015).

1080

1081

1082

Table 4: Approximate chemical composition of ancient wheat, durum wheat and common wheat.

	Einkorn	Emmer	Spelt	Kamut®	Durum wheat	Common wheat
Energy (kcal/100g)	333	362	324	337	339	340
Carbohydrate (g/100g)	67	72	68	71	71	75
Protein (g/100g)	13.3	12.8	14.7	14.5	13.7	10.7
Fiber (g/100g)	6.7	10.6	5.9	11.1	11.6	12.7
Lipid (g/100g)	1.7	2.1	2.9	2.1	2.5	2
Minerals						
Calcium (mg/100g)	Nr	Nr	17.6	22	34	34
Iron (mg/100g)	3.6	1.5	3.1	3.8	3.2	5.4
Magnesium (mg/100g)	200	128	Nr	130	144	90
Phosphorus (mg/100g)	Nr	Nr	Nr	364	508	402
Potassium (mg/100g)	Nr	Nr	Nr	403	431	435
Sodium (mg/100g)	Nr	Nr	Nr	5	2	2
Zinc (mg/100g)	15	4.8	Nr	3.7	4.2	3.5
Vitamins (µg/100g)						
Vitamin A	Nr	Nr	Nr	0.3	0	0
Vitamin B_6	0.4	Nr	Nr	0.26	0.42	0.38
Vitamin C	Nr	Nr	Nr	0	0	0
Vitamin E	Nr	Nr	Nr	0.61	0	1.01

Nr: Not reported

United States Department of Agriculture. USDA Food Composition Database. Available from: http://ndb.nal.usda.gov/.

Table 5: Approximate chemical composition of pseudocereals compared to maize and rice.

	Pseudocereals				Maize	Rice
	Buckwheat	Quinoa	Amaranth	Chia		
Energy (kcal/100g)	343	368	378	486	370	363
Carbohydrate (g/100g)	71.5	64.2	66.7	18.4	85.19	76.5
Protein (g/100g)	13.25	14.1	15.5	16.5	7.4	7.23
Fiber (g/100g)	10	7	8.9	34.4	7.4	4.6
Lipid (g/100g)	3.4	6.1	6.7	30.7	3.7	2.78
Minerals (g/100g)						
Calcium (mg/100g)	18	47	133	631	Nr	11
Iron (mg/100g)	2.8	4.57	8	7.72	2.67	1.98
Magnesium (mg/100g)	231	197	Nr	335	259	112
Phosphorus (mg/100g)	347	457	Nr	860	Nr	337
Potassium (mg/100g)	460	563	Nr	407	Nr	289
Sodium (mg/100g)	1	5	22	16	Nr	8
Zinc (mg/100g)	2.4	3.1	Nr	4.58	Nr	2.45
Vitamins (µg/100g)						
Vitamin A	Nr	4.2	Nr	16.2	Nr	Nr
Vitamin B_6	0.21	0.487	Nr	Nr	Nr	0.736
Vitamin C	Nr	Nr	5.3	1.6	Nr	Nr
Vitamin E	Nr	2.44	Nr	0.5	Nr	0.6

Nr. Not reported

United States Department of Agriculture. USDA Food Composition Database. Available from: http://ndb.nal.usda.gov/.