



UNIVERSITÀ DI PARMA

ARCHIVIO DELLA RICERCA

University of Parma Research Repository

Clinical efficacy of medical hydrology: an umbrella review

This is the peer reviewed version of the following article:

Original

Clinical efficacy of medical hydrology: an umbrella review / Antonelli, M.; Donelli, D.; Veronesi, L.; Vitale, M.; Pasquarella, C.. - In: INTERNATIONAL JOURNAL OF BIOMETEOROLOGY. - ISSN 0020-7128. - 65:10(2021), pp. 1597-1614. [10.1007/s00484-021-02133-w]

Availability:

This version is available at: 11381/2914091 since: 2022-01-21T16:46:25Z

Publisher:

Springer Science and Business Media Deutschland GmbH

Published

DOI:10.1007/s00484-021-02133-w

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

12 August 2025

International Journal of Biometeorology

Clinical efficacy of Medical Hydrology: an umbrella review

--Manuscript Draft--

Manuscript Number:	IJBM-D-20-00565R1
Full Title:	Clinical efficacy of Medical Hydrology: an umbrella review
Article Type:	Review Paper
Keywords:	Medical hydrology; Clinical efficacy; Public health; Integrative medicine; Umbrella review.
Corresponding Author:	Michele Antonelli, M.D. AUSL-IRCCS Reggio Emilia ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	AUSL-IRCCS Reggio Emilia
Corresponding Author's Secondary Institution:	
First Author:	Michele Antonelli, M.D.
First Author Secondary Information:	
Order of Authors:	Michele Antonelli, M.D.
	Davide Donelli, M.D.
	Licia Veronesi, M.D.
	Marco Vitale, M.D.
	Cesira Pasquarella, M.D.
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	<p>Introduction. The aim of this research was to summarize available scientific evidence on the efficacy of Medical Hydrology for the management of any health condition.</p> <p>Methods. The search was conducted on March 26th, 2021, in the following databases: Medline (via PubMed), EMBASE, Web of Science, Cochrane Library, and Google Scholar. All relevant literature reviews investigating the clinical efficacy of interventions characterized by the use of natural mineral waters and muds were included. The quality of studies was assessed with the "AMSTAR 2" tool.</p> <p>Results. After article screening, 49 reviews were included in this work. Overall, retrieved scientific evidence suggests that spa therapy is beneficial for patients affected by some specific musculoskeletal conditions, with improvements potentially lasting up to 9 months. Moreover, balneotherapy can be an integrative support for the management of chronic venous insufficiency and some inflammatory skin diseases like psoriasis. The role of spa therapy in rehabilitation appears relevant as well. More limited, although interesting evidence exists for inhalation and hydropinic therapies.</p> <p>Discussion. Globally, retrieved evidence suggests that, besides individual wellbeing, Medical Hydrology can be useful for public health. In particular, higher-quality studies seem to support the integrative use of spa-related interventions for conditions like osteoarthritis, fibromyalgia, low back pain of rheumatic origin, and chronic venous insufficiency. However, the body of evidence has some limitations and further clinical trials should be designed for each relevant application to consolidate and expand acquired knowledge.</p>

First of all, we would like to sincerely thank the editors and referees for providing very important feedback on our research work. We tried to address all raised issues point by point in a thorough way, as reported below. Our ultimate goal was to significantly improve the quality of an article that may help Medical Hydrology upgrade its framework of scientific evidence. All text changes were highlighted in the full-text version of our revised manuscript.

Reviewer #1: Dear authors this is a very well written review, but I have some points to criticize.

REPLY: Thanks for the appreciation of our article.

You judge the quality of the reviews, which have quite often a high quality, but on this way it gets lost, that in good reviews there can be quite a lot of trials with low quality, and this is very hard to see in your umbrella review as for example the review of Stier and Jarmer Stier-Jarmer M, Kus S, Frisch D, et al (2015) Health resort medicine in non-musculoskeletal 1019 disorders: is there evidence of its effectiveness? Int J Biometeorol 59:1523-1544. 1020 <https://doi.org/10.1007/s00484-015-0953-6>

So the evidence appears higher reading your review than it actually is.

REPLY: We have thoroughly re-assessed the methodological quality of included systematic reviews with another appraisal tool, the so called "AMSTAR 2", an internationally validated 16-item system that can assist in grading the review quality in much more detail (it is based on a 4-tier hierarchy: good, moderate, low and very low quality). Additional domains were taken into account to better evaluate the quality and risk of bias of any relevant systematic review. For transparency, full details of the review quality assessment were reported in the Electronic Supplementary Materials. Then, to specifically account for the actual risk of bias of clinical trials, we reported in Table 1 the overall quality of any set of primary studies included in each literature review, as per the review authors' global judgement. Thus, the reader can now see at a glance both the quality of included reviews and the quality of primary studies analyzed in each review. The Discussion section has been modified accordingly.

Then you cite sometimes only abstracts and I know well the reviews of Beer 2018 and Schuh 2009 and therefore know, that it is possible to get the whole article.

Beer A, Kleinschmidt J, Jagenburg L (2018) Zur Wirksamkeit der künftlichen Heilorttherapie: eine

860 Literaturübersicht aus neueren Veröffentlichungen. Phys Med Rehab Kuror 28:365-371.

861 <https://doi.org/10.1055/a-0713-0694>

Schuh A (2009) Die Evidenz der Klima- und Thalassotherapie. Ein Review. Schweizerische 1015 Zeitschrift für Ganzheitsmedizin / Swiss Journal of Integrative Medicine 21:96-104

REPLY: These studies were only assessed on the basis of information provided in their abstract because their full-text version was written in German and we were unable to directly translate it (languages that we could read on our own were listed in the Methods section and included English, French, Italian, Spanish and Portuguese). We have now sought help from a translator to extract relevant data from these studies too.

In many different reviews appear the same trials which can cause a bias, as it appears that there is more evidence, when more reviews with the same trials are published.

You should address this topic better.

REPLY: We decided to tackle this issue in a practical way, by providing the reader with an example. We selected a representative sub-topic (balneotherapy for fibromyalgia) and extracted the list of all clinical studies included in any relevant review analyzed in our research. Then, we put them in a double-entry “matching table” to highlight redundancy of clinical evidence across different literature reviews and overlaps in reporting trial results (see the Electronic Supplementary Materials for further details).

For knee osteoarthritis, another example can be found here: D’Angelo, D., Coclite, D., Napoletano, A., Fauci, A. J., Latina, R., Gianola, S., ... & Iannone, P. (2021). The efficacy of balneotherapy, mud therapy and spa therapy in patients with osteoarthritis: an overview of reviews. *International Journal of Biometeorology*, 1-17.

<https://doi.org/10.1007/s00484-021-02102-3>

The results of this analysis were briefly discussed in the Limitations section of the manuscript.

I your search in pubmed you use [Title/abstract]. Given the problems with the precise terminology (especially in English) in the field of Medical Hydrology as you correctly describe in limimtatipns, this may be a big bias missing relevant reviews.

you also use this for systematic review and meta-nalysis, but should better use the filters instead.

your search strategy in Google scholar is quite different from the other databases and this point can only be seen in the appendix.

You should use a similar strategy as in the other databases or at least describe how and why the search in Scholar was performed in this way.

This may also explain why you have only 20% duplicates by using 5 different databases with the "same" search strategy. This is nearly impossible and you should adapt the search strategies or address this fact at least in the discussion or limitations.

REPLY: The search strategy has now been homogenized across different scientific databases. We chose not to change the search strategy used for Google Scholar to narrow down our search and improve its precision (this detail was reported in the Methods section). Following the referee’s recommendation, we have run the search again applying proper filters and performed the selection of eligible articles another time from the beginning. The flowchart summarizing the article selection process was updated accordingly. For transparency, a complete list of articles excluded after their full-text assessment (with reasons) was provided in the Electronic Supplementary Materials. A few experts were contacted for some general feedback, and regulatory papers/literature overviews were searched as well through snowballing to avoid missing any relevant article. Finally, four new systematic reviews were included in our umbrella overview.

Many thanks again for your insightful and kind help.

Title

Clinical efficacy of Medical Hydrology: an umbrella review

Abstract

Introduction. The aim of this research was to summarize available scientific evidence on the efficacy of Medical Hydrology for the management of any health condition.

Methods. The search was conducted on March 26th, 2021, in the following databases: Medline (via PubMed), EMBASE, Web of Science, Cochrane Library, and Google Scholar. All relevant literature reviews investigating the clinical efficacy of interventions characterized by the use of natural mineral waters and muds were included. The quality of studies was assessed with the “AMSTAR 2” tool.

Results. After article screening, 49 reviews were included in this work. Overall, retrieved scientific evidence suggests that spa therapy is beneficial for patients affected by some specific musculoskeletal conditions, with improvements potentially lasting up to 9 months. Moreover, balneotherapy can be an integrative support for the management of chronic venous insufficiency and some inflammatory skin diseases like psoriasis. The role of spa therapy in rehabilitation appears relevant as well. More limited, although interesting evidence exists for inhalation and hydropinic therapies.

Discussion. Globally, retrieved evidence suggests that, besides individual wellbeing, Medical Hydrology can be useful for public health. In particular, higher-quality studies seem to support the integrative use of spa-related interventions for conditions like osteoarthritis, fibromyalgia, low back pain of rheumatic origin, and chronic venous insufficiency. However, the body of evidence has some limitations and further clinical trials should be designed for each relevant application to consolidate and expand acquired knowledge.

Keywords: Medical hydrology; Clinical efficacy; Public health; Integrative medicine; Umbrella review.

Authors

First author (corresponding author*): **Michele Antonelli**. Affiliation: AUSL-IRCCS Reggio Emilia, Reggio Emilia (Italy). *Address: Via Giovanni Amendola 2, 42122 Reggio Emilia (Italy). Email: michele.antonelli@ausl.re.it ORCID: 0000-0002-5941-6604

Second author: **Davide Donelli**. Affiliation: AUSL-IRCCS Reggio Emilia, Reggio Emilia (Italy). Email: donelli.davide@gmail.com ORCID: 0000-0002-7895-2840

Third author: **Licia Veronesi**. Affiliation: Department of Medicine and Surgery, University of Parma, Parma (Italy). Email: licia.veronesi@unipr.it

Fourth author: **Marco Vitale**. Affiliation (1): Department of Medicine and Surgery, University of Parma, Parma (Italy). Affiliation (2): FoRST Foundation, Rome (Italy). Email: marco.vitale@unipr.it

Last author: **Cesira Pasquarella**. Affiliation: Department of Medicine and Surgery, University of Parma, Parma (Italy). Email: cesiraisabellamaria.pasquarella@unipr.it ORCID: 0000-0001-7579-4827

Funding: this research work was not funded.

Conflicts of interest/Competing interests: no conflict of interest or competing interest has to be declared by the authors.

Ethics approval: not applicable.

Introduction

Background and definitions

Medical Hydrology (or Balneology) is a biomedical discipline with a long-standing tradition, which investigates clinical uses and health-related applications of natural mineral waters and muds for preventive, therapeutic, and rehabilitative purposes (Nappi 2001; Gutenbrunner et al. 2010; Maraver and Karagülle 2012). These treatments are usually administered in health facilities labelled as “spa centers”, where “spa” stands for “*salus per aquam*” or “*sanitas per aquam*”, an ancient Latin expression which literally refers to the pursuit of “health through water” (van Tubergen and van der Linden 2002).

According to the European Union (EU) law, as stated in the Council Directive 80/777/EEC, a natural mineral water is defined as a microbiologically wholesome water (originating in an underground water table or deposit and emerging from a spring), which can be distinguishable from ordinary drinking water by its nature and original purity (The European Parliament and the Council of the European Union 2009). In particular, natural mineral waters are characterized by a specific mineral content and biochemical composition, and by certain effects, including their potential pharmacological, physiological, and clinical action (The European Parliament and the Council of the European Union 2009). In several European countries, this regulatory definition is also accepted by the scientific community (Gutenbrunner et al. 2010). Therefore, in this article, we referred to “natural mineral waters” or, simply, “mineral waters” following the above mentioned description and indicating spring waters used for preventive, therapeutic or rehabilitative purposes in spa centers. With the aim of unequivocally classifying all spa-based interventions and considering a common set of definitions used in the scientific literature (Pittler et al. 2006; Gutenbrunner et al. 2010; Gomes et al. 2013; Fioravanti et al. 2017; Antonelli and Donelli 2018a), the following terms were used in this research work:

- “Balneotherapy”: any treatment which only involves the partial or total-body immersion in natural mineral waters (Antonelli and Donelli 2018a; Antonelli et al. 2018). When sea water was used, the term “Talassotherapy” was adopted (Maraver et al. 2011).
- “Mud therapy”: any treatment which only involves the application over skin of muds or other peloids, which were defined as natural products used as therapeutic agents and consisting of a mixture of minero-medicinal water with organic and inorganic material, like clay minerals (Carretero 2002). Although usually classified among balneo-therapeutic interventions (Fioravanti et al. 2017), mud therapy was distinguished from standard balneotherapy for its specific characteristics.
- “Spa therapy”: any spa-based multicomponent treatment which includes at least a treatment involving the therapeutic use of natural mineral waters (Pittler et al. 2006; Karagülle and

Karagülle 2015; Antonelli and Donelli 2018a). Due to its complexity and integrated therapeutic action, fango-balneotherapy, namely mud baths followed or preceded by standard balneotherapy with mineral waters, was classified under this category.

- “Inhalation therapy”: any treatment characterized by the inhalation of natural mineral waters and their gases in the form of vapors, nebulizations, aerosols, politzers, and humages (Costantino et al. 2006; Keller et al. 2014). Oral and nasal irrigations were also included in this category.
- “Hydropinic therapy”: any treatment which implies the oral ingestion of natural mineral waters with medicinal properties (Albertini et al. 2007).

Sometimes, especially in Italy or France, the old-style term “chreno-therapy” (from the Greek “krēnē”: spring, well, fountain) is adopted to indicate all interventions based on the external and/or internal administration of mineral waters and muds (Nappi 2001; Gutenbrunner et al. 2010; Maraver and Karagülle 2012). However, this term was not used in this article to avoid potential misunderstandings. Whenever hydrologic treatments involved the sole use of tap non-mineral water, for example in the control group, the generic term “hydrotherapy” was adopted.

Rationale

In Italy, costs of treatments based on natural mineral waters and therapeutic muds can be partially or fully covered by the National Healthcare System, and the same happens in other countries, both within and outside Europe (Gutenbrunner et al. 2010). In accordance with a regulatory document approved by the Italian Ministry of Health, conditions which can benefit from Medical Hydrology include osteoarthritis, extra-articular rheumatisms, chronic sinusitis and rhinosinusitis, vasomotor rhinopathy, chronic laryngopharyngitis, chronic catarrhal otitis, tubaric stenosis of flogistic origin, chronic bronchitis, psoriasis, eczema and atopic dermatitis, seborrheic dermatitis, chronic venous insufficiency, constipation due to irritable bowel syndrome, dyspepsia caused by gastroenteric or biliary dysfunction, recurrent kidney stones, and some forms of chronic vaginitis (Ministero della Sanità 1995).

Globally, these conditions are mostly caused by chronic diseases with an epidemiologically considerable impact on health-related quality of life (QoL). For example, osteoarthritis is a frequent degenerative disorder of the musculoskeletal system, especially among elderly subjects, with around 14 million patients only in the USA, and it is expected to become one of the leading causes of disability worldwide (Vina and Kwoh 2018). A relevant impact on public health is also shared by other health conditions which can benefit from spa-based interventions, such as fibromyalgia (Queiroz 2013), chronic venous insufficiency (Al Shammeri et al. 2014), chronic inflammatory respiratory diseases (Ferrante et al. 2017), and phlogistic skin disorders like psoriasis (Rachakonda et al. 2014). However, the above mentioned clinical indications were last updated and revised years ago (Ministero della Sanità 1995), and the most recent attempt to collect all available evidence on the topic with a

cross-cutting approach dates back to March 2014, when the “Hydroglobe” project was published by a panel of experts with the technical support of the World Health Organization (WHO) (Vv.Aa. 2014). In the light of what stated above, considering that Medical Hydrology has a long-standing tradition (a), that mineral waters and therapeutic muds are easily available as natural resources (b), that spa-based interventions are usually prescribed for epidemiologically relevant health conditions (c), and that scientific evidence rapidly evolves (d), it can be useful to synthesize and critically appraise available findings on the topic with a systematic approach, thus following the basic principles of Evidence-Based Medicine (EBM). This would help to outline an updated list of evidence-based indications for treatments with mineral waters and therapeutic muds, and to guide further research in this field, which is important for both individual wellbeing and public health.

Study objective

The aim of this research work was to summarize available scientific evidence on the efficacy of Medical Hydrology for the management of any health condition with a critical assessment of all relevant literature reviews.

Methods

Study design and protocol registration

A systematic review of literature reviews and meta-analyses (umbrella review) was conducted following the internationally-accepted PRISMA (*Preferred Reporting Items for Systematic Reviews and Meta-Analyses*) guidelines (Moher et al. 2009). Additional methodological recommendations for umbrella reviews were taken into account to improve the overall quality of this work (Fusar-Poli and Radua 2018). The review protocol was registered in OSF (*Open Science Framework*) under the following DOI: 10.17605/OSF.IO/2NJ5X (<https://osf.io/2nj5x>).

Eligibility criteria

The following PICOS criteria were applied for the inclusion and exclusion of studies in this review:

- Population: healthy subjects and/or individuals affected by any disease diagnosed in accordance with validated clinical criteria.
- Intervention: all spa interventions based on the use of natural mineral waters and therapeutic muds (balneotherapy, mud therapy, spa therapy, inhalation/irrigation therapy, hydropinic therapy). Reviews were excluded when they focused on studies with non-mineral tap water.
- Control: any type, including no control.

- Outcomes: any relevant clinical outcome (symptomatic and functional improvements, variations in laboratory parameters, reduction of drug consumption, effects on health-related quality of life).
- Study design: systematic reviews and meta-analyses of clinical studies. Reviews were excluded when they had no description of research methods and when they included only pre-clinical laboratory studies. Relevant high-quality narrative reviews (reporting a full description of their research methods and based on a quite extensive literature search) were also included along with systematic reviews, but they were synthesized in a separated section of the manuscript. Following the Cochrane recommendations, systematic reviews were defined as any “review [that] attempts to identify, appraise and synthesize all the empirical evidence that meets pre-specified eligibility criteria to answer a specific research question” (The Cochrane Group). Reviews had to be published in a refereed journal to be eligible for inclusion.

Information sources

Following shared recommendations for optimal database combinations (Bramer et al. 2017), the literature search was conducted in Medline (via PubMed), EMBASE, Web of Science, Cochrane Library, and Google Scholar from inception to March 2021.

Search

The search was conducted on March 26th, 2021. A first tentative pilot search was performed by one author only (M.A.) on January 7th, 2020, then, after refinements, the entire search was updated and all articles were screened again by two authors independently (M.A., D.D.).

The following keywords were used: “balneotherapy”, “hydrotherapy”, “thalassotherapy”, “spa therapy”, “water therapy”, “aquatic therapy”, “mud therapy”, “peloid therapy”, psammotherapy, “inhalation therapy”, “endotympanic insufflation”, “politzer”, “cave therapy”, “hydropinotherapy”, “mineral water*”, “thermal water*”, “hot spring water*”. Specific search strategies adopted for all screened databases were disclosed in the Electronic Supplementary Materials.

In order to narrow down the search and increase its precision, keywords used in Google Scholar were markedly simplified. References of important regulatory papers and overviews of reviews were screened with a “snowballing technique” for an additional check. An author (M.V.), with his long-standing experience in Medical Hydrology research, performed a supplementary search to make sure not to have missed any highly relevant articles in this field of study.

Study selection

The article screening process was conducted with the help of EndNote® software (version X4) by two authors independently (M.A., D.D.). In cases of discrepancies, another author (C.P.) was consulted

and disagreements were discussed until consensus was reached. Only systematic reviews and meta-analyses (along with the most relevant and extensive narrative reviews) matching the above-mentioned PICOS criteria were included in this umbrella review. All studies written in English, French, Italian, Spanish, and Portuguese were considered eligible for inclusion. Studies matching inclusion criteria but written in other languages (German, Japanese) were consulted with the support of a translator. In order to maximize retrievable evidence on the topic and to reduce the risk of publication bias, even articles with only an English abstract available for consultation were included in this umbrella review. This detail was reported in the table describing the main characteristics of included systematic reviews, although it was not possible to assess the quality of these research works due to the lack of detailed methodological information.

Data collection process

Data extraction was conducted manually with a predefined Excel® table designed in accordance with the PICOS criteria. Data extraction was performed by one author (M.A.) with a second check by another author (C.P.). In any case of missing data, authors were contacted via email or through ResearchGate®. The full-text version of a review article was retrieved in this way (Raza et al. 2020).

Data items

Data items extracted from included studies were the following ones: the number and main characteristics of study populations, the type of intervention and control, all relevant clinical outcomes, and the study design (namely whether each review was systematic or narrative, and if it was coupled or not with a meta-analysis).

To properly account for the reliability and consistency of clinical evidence on the topic, it was decided to also report the overall quality of primary studies analyzed in all systematic reviews eligible for inclusion. In this regard, the set of trials included in each review was globally judged as characterized by a good (1), fair (2) or poor (3) quality, depending on the risk-of-bias assessment performed by the review authors. This system is based on the three-tier quality rating of scientific studies recommended by the American National Institutes of Health (National Institutes of Health).

Risk of bias and quality of studies

The quality of included systematic reviews was independently evaluated by two authors (M.A., D.D.) with a dedicated appraisal tool called “AMSTAR 2”, specifically developed and validated by a team of expert methodologists for this purpose (Shea et al. 2017). In cases of disagreement, items were discussed with another author (C.P.) until consensus was reached. The “AMSTAR 2” tool provides a 16-item checklist aimed to explore different methodological domains, including the appropriateness of inclusion/exclusion criteria, search strategies, article selection, data extraction, risk-of-bias assessment, heterogeneity evaluation, quantitative synthesis and critical discussion. Each item

corresponds to a question, which can be answered with “yes” or “no” (sometimes “yes, but partially” is available as a middle response). Included reviews were evaluated one by one and their overall quality was rated as:

- High quality (A) if zero or one non-critical weaknesses were found.
- Moderate quality (B) if two or more non-critical weaknesses were found.
- Low quality (C) if one critical flaw (with or without non-critical weaknesses) was found.
- Very low quality (D) if two or more critical flaws (with or without non-critical weaknesses) were found.

As reported in our study protocol, it was originally planned to use the appraisal tool developed by the American NIH (National Institutes of Health) to assess the quality of included systematic reviews. However, in consideration of the need for a deeper methodological analysis of retrieved reviews, it was eventually decided to resort to the more specific and widely used “AMSTAR 2” tool.

The methodological quality of narrative reviews was assessed with the SANRA scale (Baethge et al. 2019) during the article selection process and only higher-quality narrative reviews were included in this work. The SANRA is a 6-item scale which evaluates the relevance/importance of a narrative review (1), whether its aim is sufficiently focused (2), if the literature search is broad enough (3), whether referencing (4), scientific reasoning (5), and presentation of data (6) are appropriate. Each item score can vary from 0 to 2, and the overall review quality score can range from 0 to 12 (high-quality narrative reviews usually score 9 or more SANRA points).

Publication bias and potential biases across studies were only qualitatively assessed because no quantitative synthesis was feasible due to the detection of a high level of heterogeneity across included studies.

Synthesis of results

The main characteristics of included reviews were reported in two tables, then retrieved evidence was qualitatively synthesized and critically discussed. Results of the study quality assessment were used for a critical discussion. Included reviews were also grouped on the basis of intervention type, health condition of interest, study design (systematic/narrative) and methodological quality. Systematic reviews specifically providing regional data, namely reviews appraising evidence from clinical studies only conducted in a given country of the world, were summarized in another table and mentioned in the “Discussion” section for better comprehensiveness.

Results

Study selection

Overall, the literature search yielded 803 results and, after screening and selection of potentially eligible articles, 49 reviews (41 systematic and 8 narrative reviews) were eventually included in this research work. Details of the article selection process, along with the main reasons for exclusion of non-eligible studies, were summarized in a dedicated flowchart (Figure 1). The list of all articles eligible for a full-text assessment and then excluded after a thorough evaluation was provided in the Electronic Supplementary Materials. The quality of a review available as a conference abstract in a refereed journal (Cao et al. 2020) was assessed on the basis of information retrieved from the corresponding preprint (<https://www.researchsquare.com/article/rs-16293/v1>). An article was kindly provided by the authors after our direct inquiry (Raza et al. 2020). It was not possible to consult the full-text version of a review but this study was included all the same for better comprehensiveness (essential data were extracted from the abstract) (Zhen-han et al. 2014). A meta-analysis found through snowballing and published as a Ph.D. thesis was excluded from the main search but mentioned in the Discussion section to better analyze the mechanisms of action of mud therapy for osteoarthritis (Crespin 2017).

Characteristics of included studies

The main characteristics of all included studies, collected and described in accordance with the PICOS criteria, were reported in Table 1 (systematic reviews) and Table 2 (narrative reviews), along with a brief summary of the authors' conclusions and, for systematic reviews, with their overall methodological quality evaluated in accordance with the AMSTAR-2 recommendations (Bohmer et al. 2000; Brosseau et al. 2002; 泮青 et al. 2006; Pittler et al. 2006; Verhagen et al. 2007, 2015; Forestier and Françon 2008; Schuh 2009; Harzy et al. 2009; Falagas et al. 2009; Langhorst et al. 2009; Guidelli et al. 2012; Fraioli et al. 2013, 2018; Espejo-Antúnez et al. 2013; Liu et al. 2013; Zhen-han et al. 2014; Roques 2014; Keller et al. 2014; Naumann and Sadaghiani 2014; Karagülle and Karagülle 2015; Françon et al. 2015; Tenti et al. 2015; Stier-Jarmer et al. 2015; Fortunati et al. 2016; Xiang et al. 2016; Forestier et al. 2016, 2017; Bender 2016; Santos et al. 2016; Matsumoto et al. 2017; Naumann et al. 2017; Passali et al. 2017; Morer et al. 2017; Casale et al. 2018; Antonelli and Donelli 2018a; Antonelli et al. 2018; Beer et al. 2018; An et al. 2019; Corvillo et al. 2019; de Moraes Silva et al. 2019; Bai et al. 2019; Yuan et al. 2019; Raza et al. 2020; Sulaiman et al. 2020; Hou et al. 2020; Cao et al. 2020; Cacciapuoti et al. 2020; Gravelier et al. 2020).

Population

The number of study participants whose data were analyzed within included systematic reviews varied from a minimum of 54 (Bohmer et al. 2000) to a maximum of 13782 (Stier-Jarmer et al. 2015), with a median value of 731. In the majority of included reviews, regardless of their design (systematic or narrative), study participants were patients with chronic conditions, such as rheumatic (osteoarthritis, fibromyalgia, back pain of rheumatic origin, rheumatoid arthritis), cardiovascular

(chronic venous insufficiency, hypertension), dermatologic, respiratory, otorhinolaryngological, neurologic, digestive, and urologic diseases (Figure 2). In a review, the effects of hydropinic therapy on healthy subjects were studied (Bohmer et al. 2000), whereas studies with both healthy and diseased individuals were analyzed in three research works (Antonelli and Donelli 2018a; An et al. 2019; Sulaiman et al. 2020).

Intervention

Most included reviews analyzed the efficacy of balneotherapy (n=10) and mud therapy (n=6) alone or in combination with other non-spa-related treatments (n=26) (Figure 3). In some cases, spa therapy also involved physical rehabilitation, relaxing massage, diet prescriptions for weight loss or for preventive purposes, pharmacological treatments, and psychological support. Only in a few reviews the efficacy of hydropinic therapy (n=3) and inhalation therapy or nasal irrigations (n=3) were investigated. In one (n=1) research work, the efficacy of any type of spa-related therapy was studied (Stier-Jarmer et al. 2015).

Control

Most analyzed reviews did not have specific restrictions in terms of control type for primary study inclusion. In a systematic review, subjects with cervical pain undergoing balneotherapy were compared to individuals sharing the same health condition who did not receive any treatment or who were administered a cycle of standard rehabilitation (Corvillo et al. 2019). In two research works, the therapeutic efficacy of spa therapy was compared to hydrotherapy with tap non-mineral water (Morer et al. 2017; Sulaiman et al. 2020). Finally, in other systematic reviews, the health effect of orally taking two different mineral waters was compared with modifications induced by consuming other foods and nutritional products (Bohmer et al. 2000; Naumann et al. 2017).

Outcomes

Main clinical outcomes of analyzed reviews included the following ones: symptoms (mostly pain, evaluated with a Visual Analogue Scale), functionality and disability (sometimes assessed in combination with pain using algo-functional scales), drug consumption (especially with regard to the long-term intake of painkillers), quality of life (measured with specific questionnaires), results of various diagnostic investigations and laboratory parameter assessment (including biomarkers of inflammation, metabolic indices, circulating levels of hormones and other biochemical substances).

Study design

Thirtyseven (41) included research works were systematic literature reviews, and seventeen (17) of them were also coupled with a meta-analysis (Pittler et al. 2006; Verhagen et al. 2007, 2015; Forestier and Françon 2008; Langhorst et al. 2009; Liu et al. 2013; Zhen-han et al. 2014; Keller et al. 2014;

Xiang et al. 2016; Matsumoto et al. 2017; Naumann et al. 2017; Antonelli et al. 2018; de Moraes Silva et al. 2019; Bai et al. 2019; Raza et al. 2020; Hou et al. 2020; Cao et al. 2020). Eight (8) included reviews had a narrative design, but they still provided an extensive, quasi-systematic and valuable overview of the scientific literature (Guidelli et al. 2012; Roques 2014; Françon et al. 2015; Tenti et al. 2015; Fortunati et al. 2016; Bender 2016; Passali et al. 2017; Cacciapuoti et al. 2020), thus outlining a state-of-the-art description of specific subtopics, such as the efficacy of spa therapy for respiratory illnesses (Passali et al. 2017), skin diseases (Cacciapuoti et al. 2020), or hand osteoarthritis (Fortunati et al. 2016). The number of clinical studies included in the systematic reviews varied from a minimum of 3 (Brosseau et al. 2002; Fortunati et al. 2016) to a maximum of 41 (Stier-Jarmer et al. 2015), with a median of 12. Such primary studies were mostly Randomized Controlled Trials, although some of them were characterized by a different design, like nonrandomized trials, uncontrolled studies, and observational investigations. The quality of clinical studies tended to range from fair to poor, with a potential risk of bias mostly arising from small sample size, lack of adequate control, no randomization of trial participants, and poor information about blinding of intervention.

Quality of included studies

After the methodological assessment of included systematic reviews with the “AMSTAR 2” tool, the quality of analyzed research works was judged as good (A) to moderate (B) in 11 cases (Pittler et al. 2006; Verhagen et al. 2007, 2015; Harzy et al. 2009; Langhorst et al. 2009; Naumann and Sadaghiani 2014; Stier-Jarmer et al. 2015; Santos et al. 2016; Matsumoto et al. 2017; de Moraes Silva et al. 2019; Hou et al. 2020) and low (C) to very low (D) in the remaining cases, as reported in Table 1 (see the Supplementary Materials for further details). It was not possible to assess the quality of a systematic review because its full-text version was unavailable (Zhen-han et al. 2014). In general, the most frequent methodological weaknesses found within included reviews were excessively narrow search strategies, a poor description of study selection and evaluation processes, and limited consideration of the trial risk-of-bias assessment for informing a critical discussion. Major strengths were usually a clear definition of the research question and, where applicable, a good methodological level of meta-analyses.

Discussion

Efficacy of interventions

Scientific evidence from included reviews indicates that balneotherapy, mud therapy, and spa therapy can significantly improve clinical parameters like pain, joint functionality, mobility, and quality of life of patients with chronic musculoskeletal conditions, mostly osteoarthritis, fibromyalgia, and other

pain-related rheumatic diseases. For these subjects, beneficial effects may last up to 9 months after intervention (on average, 3 to 6 months) (Forestier et al. 2016, 2017), and clinical improvements are associated with a reduced intake of analgesic drugs (Françon et al. 2015; Tenti et al. 2015; Forestier et al. 2017; Fraioli et al. 2018; Antonelli et al. 2018). Furthermore, in two included research works, it was demonstrated that balneotherapy is superior to the same treatment administered in pools with tap water (“sham balneotherapy”) in terms of clinical benefits and improved quality of life among patients with osteoarthritis and fibromyalgia, thus underscoring the contribution of the specific biochemical composition of natural mineral waters to the overall therapeutic effect (Morer et al. 2017; Antonelli et al. 2018).

With regard to vascular health, balneotherapy can be useful for the amelioration of pain, quality of life, and skin pigmentation due to chronic venous insufficiency in lower limbs (Stier-Jarmer et al. 2015; de Moraes Silva et al. 2019), and it appears not to have negative effects on blood pressure levels (Yuan et al. 2019). More limited evidence suggests that balneotherapy, including talassotherapy (sea water baths), can be beneficial for patients with skin diseases like psoriasis and atopic dermatitis (Schuh 2009; Falagas et al. 2009; Stier-Jarmer et al. 2015; Cacciapuoti et al. 2020), and possibly for the integrative treatment of burn scars (Gravelier et al. 2020).

Balneotherapy, mud therapy, and spa therapy can also improve mental wellbeing and psychophysical stress, and this is demonstrated by studies in which stress hormone levels were measured (Roques 2014; Stier-Jarmer et al. 2015; Antonelli and Donelli 2018a). An interesting role for orthopedic or neurological patients of aquatic rehabilitation in pools with natural mineral waters has been underscored by several authors, with beneficial effects on the most important clinical outcomes (Falagas et al. 2009; Stier-Jarmer et al. 2015).

With regard to inhalation therapy, clinical improvements were observed in patients with diseases of the upper and lower respiratory tract, mostly chronic rhinosinusitis and bronchitis (Schuh 2009; Keller et al. 2014; Stier-Jarmer et al. 2015; Casale et al. 2018), but even in subjects with chattral otitis (Stier-Jarmer et al. 2015). Scientific evidence suggests an interesting role of highly mineralized or sulphur waters for these illnesses (Schuh 2009; Keller et al. 2014; Casale et al. 2018).

Considering hydropinic therapy, study results indicate that the bioavailability of calcium from calcium-rich waters is comparable to that one derived from dairy products (Bohmer et al. 2000), and that the consumption of bicarbonate- and magnesium-rich waters may be beneficial for kidney stone prevention and for an improvement of glycemic control (Naumann et al. 2017; Sulaiman et al. 2020). Moreover, hydropinic therapy with highly mineralized waters can be useful for constipation due to irritable bowel syndrome (Stier-Jarmer et al. 2015).

Systematic reviews including only clinical studies conducted in a specific country or region of the world were collected in Table 3 (Karagülle and Karagülle 2004; Roques et al. 2012; Katz et al. 2012; Bender et al. 2014; Stanhope et al. 2018; Khalilzadeh et al. 2019; Drobnik and Stebel 2020). These research works were mostly carried out in regions where Medical Hydrology is widely spread and

well known, such as Europe or the Middle East, and they outlined an efficacy profile of balneotherapy, mud therapy, and spa therapy which is similar to that one already described in literature reviews collected in Table 1 and Table 2. Additionally, some interesting findings suggested that balneotherapy with specific waters characterized by a high mineral content, like Dead Sea water, can be useful for the treatment of psoriasis (Katz et al. 2012; Khalilzadeh et al. 2019). Some authors also underscored the lack of relevant studies in continents like Oceania, thus urging the need for clinical investigations in Australia or New Zealand, where mineral water springs exist, but they are currently underused for medicinal purposes (Stanhope et al. 2018).

Globally, systematic reviews characterized by a higher overall quality of their methodological design tended to support the efficacy of balneotherapy, mud therapy, and spa therapy for the integrative management of osteoarthritis (especially knee osteoarthritis), fibromyalgia, low back pain of rheumatic origin, and chronic venous insufficiency (Pittler et al. 2006; Verhagen et al. 2007, 2015; Harzy et al. 2009; Langhorst et al. 2009; Naumann and Sadaghiani 2014; Stier-Jarmer et al. 2015; Santos et al. 2016; Matsumoto et al. 2017; de Moraes Silva et al. 2019; Hou et al. 2020). However, the authors underscored that clinical evidence needs to grow before firm conclusions can be drawn, and this is even more relevant for non-rheumatic conditions (for example, respiratory or skin illnesses). All the same, in general, considering both trial results and empirical observations, spa-based treatments appear useful for rehabilitation and chronic disease management, because they seem capable of exerting a beneficial action on symptom control and psychophysical wellbeing.

Safety and tolerability of interventions

Globally, evidence from included studies suggests that balneotherapy, mud therapy, spa therapy, inhalation therapy, and hydropinic therapy are quite safe and well tolerated by patients, provided that all necessary medical and hygienic precautions are taken in advance.

The most important contraindications to treatments based on mineral waters and therapeutic muds are mainly derived from tradition and they can be grouped into three categories (Nappi 2001; Vv.Aa. 2014):

- Contraindications related to the disease stage: spa-based interventions are not to be administered when a patient is affected by an acute disease or during symptomatic relapses of chronic conditions.
- Contraindications related to the patient's illnesses and comorbidities: they include infectious conditions, cancer, and unstable or poorly controlled diseases (severe heart failure, advanced kidney insufficiency, uncontrolled hypertension, cirrosis, medically-unresponsive epilepsy...).
- Contraindications related to the type of intervention: they depend on specific characteristics of single mineral waters and therapeutic muds.

Possible side effects of spa-based interventions are usually mild and often resolve spontaneously after treatment discontinuation: they are mostly due to an individual response, thus being highly “patient-specific”, and their occurrence is worsened by an improper or unsupervised administration of intervention (Nappi 2001; Vv.Aa. 2014). Side effects generally include symptoms like headache, dizziness and nausea, mild relapses of local chronic pain, sleep disturbances, heart palpitations, or a general sensation of irritability and fatigue (rarely coupled with a short-lasting low-grade fever) (Vv.Aa. 2014). Additionally, intervention-specific side effects can occur, such as diarrhea and hydroelectrolytic imbalances due to an overconsumption of highly mineralized waters, a temporary increase of fluid secretions in the airways (with runny nose and cough) after some inhalation therapies, or cutaneous irritation caused by hot mud application on irritable skin. Extreme caution is also advised in more fragile individuals like pediatric patients, pregnant women, and very elderly subjects, whose clinical response to spa-based interventions can be less predictable with a potential higher incidence of more severe side effects. For all these reasons, a medical check and supervision are strongly advised for an appropriate prescription of spa-based interventions, not only to make the most of them on the basis of the patient’s characteristics and disease, but even to avoid the onset of adverse events, thus optimizing the safety and tolerability of such treatments.

Mechanisms of action: hypotheses and evidence-based explanations

In light of scientific evidence described by expert authors of the “*HydroGlobe*” project, effects on health of Medical Hydrology-related treatments have been reported to be the following ones (Vv.Aa. 2014):

- Antalgic effect,
- Myorelaxant action,
- Activation of microcirculation,
- Immunomodulation,
- Neuro-hormonal stimulation,
- Improvement of fat and carbohydrate metabolism.

In the same project, traditional uses of different types of natural mineral waters (each of them characterized by specific biochemical component/s) were collected, with general clinical indications formulated for various health conditions, as reported in Table 4 (Nappi 2001; Vv.Aa. 2014; Quattrini et al. 2016).

In general, treatments used in Medical Hydrology can be classified into three main categories on the basis of their route of administration (external or internal) and the state of matter of the therapeutic medium (liquid or gaseous):

1. Balneo-therapeutic treatments, such as baths with natural mineral waters and mud therapy.

2. Inhalation-based treatments, such as vapours or aerosols derived from natural mineral waters inhaled for medicinal purposes.

3. Hydropinic treatments, when natural mineral waters are taken orally as therapeutic drinks.

Balneo-therapeutic treatments are believed to exert their global therapeutic effect on the body thanks to a synergistic combination of mechanical (hydrostatic pressure), thermal (high temperature), and biochemical actions, the latter due to both the mineral (osmotic pressure and direct activity) and the organic (with anti-inflammatory and immunomodulatory properties) components of waters and muds (Vv.Aa. 2014; Fioravanti et al. 2017; Antonelli and Donelli 2018b). From a physiological point of view, balneo-therapeutic treatments can increase serum β -endorphins and can modulate cortisol levels in such a way as to improve individual stress resilience without disrupting circadian rhythms of this hormone (Fioravanti et al. 2011; Antonelli and Donelli 2018a). The long-term increase of cortisol awakening response due to a cycle of balneo-therapeutic treatments may be the reason why, in some clinical studies, the effects on health of balneotherapy, mud therapy, and spa therapy were observed to last for a few months after intervention (Forestier et al. 2016; Antonelli and Donelli 2018a). If we consider inflammatory mediators, mud applications followed by baths with natural mineral waters can reduce the production of prostaglandin E2 (PGE2), leukotriene B4 (LTB4), interleukin-1 β (IL-1 β), and tumor necrosis factor alpha (TNF- α) (Fioravanti et al. 2011). Quantitative data from the “gray” literature also show that, at least for osteoarthritis, therapeutic mud applications are superior to simple hot packs to improve joint functionality and pain, thus underscoring the therapeutic importance of the organic component (Crespin 2017). It is possible that the effect of balneo-therapeutic treatments on the inflammatory response and interleukin production is due to modifications of microRNA expressions induced by thermal and mechanical stimuli, as observed in a cohort of patients with osteoarthritis (Giannitti et al. 2017). At a joint level, balneo-therapeutic treatments may stimulate cartilage metabolism through mediators like the insulin-like growth factor-1 (IGF1), they may exert an antioxidant effect by reducing the release of Reactive Oxygen and Nitrogen Species (Fioravanti et al. 2011; Masselli et al. 2020) and also modulate intracellular mediators like protein kinases involved in cartilage growth and cell proliferation (Queirolo et al. 2016; Martini et al. 2018). Mud baths are also associated with a decrease in serum levels of adiponectin and resistin, hormonal substances probably implied in the progression of chronic degenerative diseases like osteoarthritis (Fioravanti et al. 2015). Furthermore, transcutaneous absorption of antiphlogistic substances released by water and mud microflora may contribute to the overall pharmacological effect of balneo-therapeutic treatments (Vv.Aa. 2014; Antonelli and Donelli 2018b). More details about possible mechanism of action of balneotherapy, as hypothesized on the basis of in-vitro laboratory studies, have recently been collected in a comprehensive literature review, demonstrating the anti-inflammatory, antioxidant, chondroprotective, and immunosuppressive role of this type of intervention at a cellular level (Cheleschi et al. 2020). Regarding balneotherapy and peripheral venous circulation, it is believed that the main therapeutic action of baths is determined by the hydrostatic and osmotic pressure of natural

mineral waters, which seems capable of reducing pain and oedema (de Moraes Silva et al. 2019). The combination of baths with hydrojet massage, Kneipp therapy (in which hot and cold baths are alternated), and physical exercise can determine a useful compression of peripheral veins, thanks to an external (water pressure) and internal (muscle contraction) synergistic action (de Moraes Silva et al. 2019). In particular, beneficial effects for the cardiovascular system seem to be more frequently associated with balneo-therapeutic sessions in carbon dioxide-rich water, which may be responsible for lowering peripheral vascular resistance and increasing blood flow in a more pronounced way if compared with other water types (Pagourelas et al. 2011). Additionally, balneo-therapeutic treatments, especially those ones based on sulphur-rich waters, seem to have anti-inflammatory, keratolytic, and regenerative effects on skin due to a direct pharmacological action of the mineral component and to interactions between thermal and cutaneous microflora with a potential modulation of local immune functions (Gobbi et al. 2009; Mirandola et al. 2011; Katz et al. 2012; Antonelli and Donelli 2018b; Eliasse et al. 2020).

Inhalation treatments with vaporized natural mineral waters can have anti-inflammatory, mucolytic, and antimicrobial properties, whereas irrigations with liquid-phase waters also have an action of mechanical washing on the upper respiratory tract (Vv.Aa. 2014; Keller et al. 2014; Casale et al. 2018). Among others, waters with sulphur seem to promote mucociliary clearance, regulate local immunity, inflammation, and have antiallergic effects (Rinaldi et al. 2006; Mirandola et al. 2007, 2013; Keller et al. 2014; Viegas et al. 2019; Carubbi et al. 2019). For all these reasons, sulphur-rich water inhalations have been proposed as an integrative treatment for patients with chronic obstructive pulmonary disease (Khaltaev et al. 2020).

Health effects of hydropinic treatments are mainly mediated by the intake of water minerals (osmotic and prokinetic action on the intestine when the mineral content is high, diuretic effect when the mineral content is low) and by the interaction between water and gut microflora (Vv.Aa. 2014). Globally, the mechanisms of action of spa-related treatments, although not fully understood, seem to be determined by a synergistic action of all water and mud components, capable of eliciting beneficial effects both locally and at a systemic level.

Limitations of this study and new perspectives for future research

First, a major limitation of both primary (clinical) and secondary (review) studies on the topic, which is responsible for hindering optimal retrieval and dissemination of scientific information, is the lack of a widely accepted consensus on a precise English terminology in the field of Medical Hydrology. In other words, terms like balneotherapy, hydrotherapy, and spa therapy are given diverse (and sometimes misleading or confusing) meanings when used by different authors, and this aspect has already been underscored even by several experts (Gutenbrunner et al. 2010; Fioravanti et al. 2017). For this reason, a list of specific definitions was provided in the Introduction section to avoid possible misunderstandings and to address this issue as best as possible.

Even if the number of reviews included in this research work is considerable, the actual basis of clinical evidence in support of spa-related treatments is quite limited. In fact, there is a demonstrated substantial degree of clinical evidence overlap among analyzed reviews (see the Electronic Supplementary Materials for an example about balneotherapy for fibromyalgia or refer to a recently published overview of literature reviews for another example about spa therapy for knee osteoarthritis (D'Angelo et al. 2021)). For this reason, it is important to keep on studying Medical Hydrology with further clinical research projects to expand the existing evidence basis.

With regard to the quality of clinical trials analyzed in included reviews, the most frequent limitations were reported to be the low number of study participants, poor information about proper randomization, and the lack of adequate control. Instead, if we consider the quality of included reviews, major limitations involved methodological issues like excessively narrow search strategies (with the literature search sometimes only run in PubMed), poor description of the details about article selection and quality assessment, and an often inadequate evaluation of the risk of publication bias. The trial quality assessment was not adequately used by some review authors for informing a critical discussion, with potentially inflated results and biased conclusions. The high level of heterogeneity across primary and secondary studies on the topic makes it difficult to synthesize available data, especially from a quantitative point of view, thus urging the necessity to conduct further investigations adopting a more homogeneous design.

As a proposal, it would be useful to promote a wider and better application of the PRISMA guidelines among researchers who want to conduct systematic reviews about Medical Hydrology, since, if we consider included research works, only some authors actually followed these internationally-accepted methodological recommendations in a thorough way (Moher et al. 2009). Poor compliance with the PRISMA guidelines can result in omitting essential information which is useful for clinicians and policymakers to translate research findings into practice (external validity), or, even worse, this lack of transparency can hide substantial flaws in the conduction of the review, thus undermining the conclusion reliability (internal validity). For this reason, our umbrella review was conducted in accordance with the PRISMA statement, and methodological quality of included systematic reviews was weighted as best as possible.

Moreover, authors of some included reviews underscored the difficulty to find a specifically tailored assessment tool to evaluate the quality of primary studies on the topic, which regard atypical, non-pharmacological and hard-to-blind interventions like balneotherapy. Different solutions were adopted by review authors to find an adequate solutions, ranging from the use of a modified version of the standard Cochrane tool (Higgins et al. 2011) to other tools often employed for the evaluation of public health interventions like the Canadian "Quality Assessment Tool for Quantitative Studies" (Armijo-Olivo et al. 2012). To properly address this issue, world experts should agree on a set of essential domains that any assessment tool should include to be adequately usable for evaluating the quality and risk of bias of Medical Hydrology-related trials. In fact, some efforts have already been made

towards this direction, for example by developing the “SPAC” checklist (Kamioka et al. 2013) or by evaluating the biasing impact of unblinded balneotherapy (Verhagen et al. 1998). Finally, in order to optimize the collection of available evidence on the topic, any effort was made to retrieve all relevant data reported both in the scientific and in the so called “gray” literature (Petticrew et al. 2008), including findings only displayed in conference proceedings and lectures published in peer-reviewed journals (Roques 2014; Bender 2016; Cao et al. 2020). However, the risk of publication bias, although minimized, could not be fully excluded, and it was not possible to statistically assess it because no quantitative synthesis was feasible.

Conclusions

Globally, retrieved evidence suggests that Medical Hydrology can be useful for individual wellbeing and public health. In particular, higher-quality studies support the use of spa-related interventions for conditions like osteoarthritis, fibromyalgia, and chronic back pain, but also for chronic venous insufficiency. However, as discussed above, the existing body of evidence has some relevant limitations, especially with regard to non-rheumatic diseases. For this reason, further high-quality clinical trials and observational studies should be designed to confirm the beneficial effects of Medical Hydrology and to thoroughly estimate its effectiveness outside experimental settings in real-life clinical practice.

References

- Albertini MC, Dacha M, Teodori L, Conti ME (2007) Drinking mineral waters: biochemical effects and health implications the state-of-the-art. *International Journal of Environment and Health* 1:153
- Al Shammeri O, AlHamdan N, Al-Hothaly B, et al (2014) Chronic Venous Insufficiency: prevalence and effect of compression stockings. *Int J Health Sci* 8:231–236. <https://doi.org/10.12816/0023975>
- An J, Lee I, Yi Y (2019) The Thermal Effects of Water Immersion on Health Outcomes: An Integrative Review. *Int J Environ Res Public Health* 16.: <https://doi.org/10.3390/ijerph16071280>
- Antonelli M, Donelli D (2018a) Effects of balneotherapy and spa therapy on levels of cortisol as a stress biomarker: a systematic review. *Int J Biometeorol* 62:913–924. <https://doi.org/10.1007/s00484-018-1504-8>
- Antonelli M, Donelli D (2018b) Mud therapy and skin microbiome: a review. *Int J Biometeorol* 62:2037–2044. <https://doi.org/10.1007/s00484-018-1599-y>
- Antonelli M, Donelli D, Fioravanti A (2018) Effects of balneotherapy and spa therapy on quality of life of patients with knee osteoarthritis: a systematic review and meta-analysis. *Rheumatol Int* 38:1807–1824. <https://doi.org/10.1007/s00296-018-4081-6>
- Armijo-Olivo S, Stiles CR, Hagen NA, et al (2012) Assessment of study quality for systematic

659 reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public
660 Health Practice Project Quality Assessment Tool: methodological research. *J Eval Clin Pract*
661 18:12–18. <https://doi.org/10.1111/j.1365-2753.2010.01516.x>

662 Baethge C, Goldbeck-Wood S, Mertens S (2019) SANRA—a scale for the quality assessment of
663 narrative review articles. *Research Integrity and Peer Review* 4

664 Bai R, Li C, Xiao Y, et al (2019) Effectiveness of spa therapy for patients with chronic low back pain:
665 An updated systematic review and meta-analysis. *Medicine* 98:e17092.
666 <https://doi.org/10.1097/MD.00000000000017092>

667 Beer A, Kleinschmidt J, Jagenburg L (2018) Zur Wirksamkeit der kurörtlichen Heiltorftherapie: eine
668 Literaturübersicht aus neueren Veröffentlichungen. *Phys Med Rehab Kuror* 28:365–371.
669 <https://doi.org/10.1055/a-0713-0694>

670 Bender T (2016) SP0165 Balneotherapy:Efficacy and Safety in Osteoarthritis. *Ann Rheum Dis*
671 75:40.4–40. <https://doi.org/10.1136/annrheumdis-2016-eular.6267>

672 Bender T, Bálint G, Prohászka Z, et al (2014) Evidence-based hydro- and balneotherapy in
673 Hungary—a systematic review and meta-analysis. *International Journal of Biometeorology*
674 58:311–323

675 Bohmer H, Müller H, Resch KL (2000) Calcium supplementation with calcium-rich mineral waters: a
676 systematic review and meta-analysis of its bioavailability. *Osteoporos Int* 11:938–943.
677 <https://doi.org/10.1007/s001980070032>

678 Bramer WM, Rethlefsen ML, Kleijnen J, Franco OH (2017) Optimal database combinations for
679 literature searches in systematic reviews: a prospective exploratory study. *Syst Rev* 6:245.
680 <https://doi.org/10.1186/s13643-017-0644-y>

681 Brosseau L, MacLeay L, Robinson V, et al (2002) Efficacy of balneotherapy for osteoarthritis of the
682 knee: A systematic review. *Phys Ther Rev* 7:209–222.
683 <https://doi.org/10.1179/108331902235001976>

684 Cacciapuoti S, Luciano MA, Megna M, et al (2020) The Role of Thermal Water in Chronic Skin
685 Diseases Management: A Review of the Literature. *J Clin Med Res* 9.:
686 <https://doi.org/10.3390/jcm9093047>

687 Cao C-F, Ma K-L, Li Q-L, et al (2020) Benefits of balneotherapy in the management of fibromyalgia
688 syndrome: a systematic review and meta-analysis of randomized controlled trials. *Pain Practice*

689 Carretero MI (2002) Clay minerals and their beneficial effects upon human health. A review. *Appl*
690 *Clay Sci* 21:155–163. [https://doi.org/10.1016/S0169-1317\(01\)00085-0](https://doi.org/10.1016/S0169-1317(01)00085-0)

691 Carubbi C, Masselli E, Calabrò E, et al (2019) Sulphurous thermal water inhalation impacts
692 respiratory metabolic parameters in heavy smokers. *Int J Biometeorol* 63:1209–1216.
693 <https://doi.org/10.1007/s00484-019-01737-7>

694 Casale M, Moffa A, Cassano M, et al (2018) Saline nasal irrigations for chronic rhinosinusitis: From
695 everyday practice to evidence-based medicine. An update. *Int J Immunopathol Pharmacol*
696 32:2058738418802676. <https://doi.org/10.1177/2058738418802676>

697 Cheleschi S, Gallo I, Tenti S (2020) A comprehensive analysis to understand the mechanism of action
698 of balneotherapy: why, how, and where they can be used? Evidence from in vitro studies
699 performed on human and animal samples. *Int J Biometeorol* 64:1247–1261.
700 <https://doi.org/10.1007/s00484-020-01890-4>

701 Corvillo I, Armijo F, Álvarez-Badillo A, et al (2019) Efficacy of aquatic therapy for neck pain: a
702 systematic review. *Int J Biometeorol*. <https://doi.org/10.1007/s00484-019-01738-6>

703 Costantino M, Lampa E, Nappi G (2006) Effectiveness of sulphur spa therapy with politzer in the
704 treatment of rhinogenic deafness. *Acta Otorhinolaryngol Ital* 26:7–13

705 Crespin M (2017) The effects of mud pack therapy versus hot pack treatment to improve functionality
706 and decrease pain in older adults suffering with knee osteoarthritis. PhD, College of Health and
707 Human Services California State University

708 D'Angelo D, Coclite D, Napoletano A, et al (2021) The efficacy of balneotherapy, mud therapy and
709 spa therapy in patients with osteoarthritis: an overview of reviews. *Int J Biometeorol*.
710 <https://doi.org/10.1007/s00484-021-02102-3>

711 de Moraes Silva MA, Nakano LC, Cisneros LL, Miranda F Jr (2019) Balneotherapy for chronic
712 venous insufficiency. *Cochrane Database Syst Rev* 8:CD013085.
713 <https://doi.org/10.1002/14651858.CD013085.pub2>

714 Drobnik J, Stebel A (2020) Central European ethnomedical and officinal uses of peat, with special
715 emphasis on the Tolpa peat preparation (TPP): An historical review. *J Ethnopharmacol*
716 246:112248. <https://doi.org/10.1016/j.jep.2019.112248>

717 Eliasse Y, Redoules D, Espinosa E (2020) Impact of Avène Thermal Spring Water on immune cells. *J*
718 *Eur Acad Dermatol Venereol* 34 Suppl 5:21–26. <https://doi.org/10.1111/jdv.16335>

719 Espejo-Antúnez L, Cardero-Durán MA, Garrido-Ardila EM, et al (2013) Clinical effectiveness of
720 mud pack therapy in knee osteoarthritis. *Rheumatology* 52:659–668.
721 <https://doi.org/10.1093/rheumatology/kes322>

722 Falagas ME, Zarkadoulia E, Rafailidis PI (2009) The therapeutic effect of balneotherapy: evaluation
723 of the evidence from randomised controlled trials. *Int J Clin Pract* 63:1068–1084.
724 <https://doi.org/10.1111/j.1742-1241.2009.02062.x>

725 Ferrante G, Baldissera S, Campostrini S (2017) Epidemiology of chronic respiratory diseases and
726 associated factors in the adult Italian population. *Eur J Public Health* 27:1110–1116.
727 <https://doi.org/10.1093/eurpub/ckx109>

728 Fioravanti A, Cantarini L, Guidelli GM, Galeazzi M (2011) Mechanisms of action of spa therapies in
729 rheumatic diseases: what scientific evidence is there? *Rheumatology International* 31:1–8

730 Fioravanti A, Giannitti C, Cheleschi S, et al (2015) Circulating levels of adiponectin, resistin, and
731 visfatin after mud-bath therapy in patients with bilateral knee osteoarthritis. *International Journal*
732 *of Biometeorology* 59:1691–1700

733 Fioravanti A, Karagülle M, Bender T, Karagülle MZ (2017) Balneotherapy in osteoarthritis: Facts,
734 fiction and gaps in knowledge. *Eur J Integr Med* 9:148–150.
735 <https://doi.org/10.1016/j.eujim.2017.01.001>

736 Forestier R, Erol Forestier FB, Francon A (2016) Spa therapy and knee osteoarthritis: A systematic
737 review. *Ann Phys Rehabil Med* 59:216–226. <https://doi.org/10.1016/j.rehab.2016.01.010>

738 Forestier R, Françon A (2008) Crenobalneotherapy for limb osteoarthritis: systematic literature
739 review and methodological analysis. *Joint Bone Spine* 75:138–148.
740 <https://doi.org/10.1016/j.jbspin.2007.06.009>

741 Forestier RJ, Erol Forestier FB, Francon A (2017) Crénothérapie (crénobalnéothérapie) dans la
742 lombalgie chronique : une revue critique. *Revue du Rhumatisme Monographies* 84:91–98.

743 <https://doi.org/10.1016/j.monrhu.2016.11.008>

744 Fortunati NA, Fioravanti A, Seri G, et al (2016) May spa therapy be a valid opportunity to treat hand
 745 osteoarthritis? A review of clinical trials and mechanisms of action. *Int J Biometeorol* 60:1–8.
 746 <https://doi.org/10.1007/s00484-015-1030-x>

747 Fraioli A, Grassi M, Mennuni G, et al (2013) Clinical researches on the efficacy of spa therapy in
 748 fibromyalgia. A systematic review. *Ann Ist Super Sanita* 49:219–229.
 749 https://doi.org/10.4415/ANN_13_02_13

750 Fraioli A, Mennuni G, Fontana M, et al (2018) Efficacy of Spa Therapy, Mud-Pack Therapy,
 751 Balneotherapy, and Mud-Bath Therapy in the Management of Knee Osteoarthritis. A Systematic
 752 Review. *Biomed Res Int* 2018:1042576. <https://doi.org/10.1155/2018/1042576>

753 Françon A, Duplan B, Forestier FBE, Forestier R (2015) Thermalisme et douleur chronique en
 754 rhumatologie : revue des essais cliniques randomisés (ECR) et des méta-analyses. *Douleur analg*
 755 28:47–53. <https://doi.org/10.1007/s11724-015-0410-7>

756 Fusar-Poli P, Radua J (2018) Ten simple rules for conducting umbrella reviews. *Evid Based Ment*
 757 *Health* 21:95–100. <https://doi.org/10.1136/ebmental-2018-300014>

758 Giannitti C, De Palma A, Pascarelli NA, et al (2017) Can balneotherapy modify microRNA
 759 expression levels in osteoarthritis? A comparative study in patients with knee osteoarthritis. *Int J*
 760 *Biometeorol* 61:2153–2158. <https://doi.org/10.1007/s00484-017-1420-3>

761 Gobbi G, Ricci F, Malinverno C, et al (2009) Hydrogen sulfide impairs keratinocyte cell growth and
 762 adhesion inhibiting mitogen-activated protein kinase signaling. *Lab Invest* 89:994–1006.
 763 <https://doi.org/10.1038/labinvest.2009.61>

764 Gomes C, Carretero MI, Pozo M, et al (2013) Peloids and pelotherapy: Historical evolution,
 765 classification and glossary. *Appl Clay Sci* 75-76:28–38.
 766 <https://doi.org/10.1016/j.clay.2013.02.008>

767 Gravelier C, Kanny G, Adetu S, Goffinet L (2020) Spa therapy and burn scar treatment: a systematic
 768 review of the literature. *Int J Biometeorol* 64:2195–2203. [https://doi.org/10.1007/s00484-020-](https://doi.org/10.1007/s00484-020-01988-9)
 769 [01988-9](https://doi.org/10.1007/s00484-020-01988-9)

770 Guidelli GM, Tenti S, De Nobili E, Fioravanti A (2012) Fibromyalgia syndrome and spa therapy:
 771 myth or reality? *Clin Med Insights Arthritis Musculoskelet Disord* 5:19–26.
 772 <https://doi.org/10.4137/CMAMD.S8797>

773 Gutenbrunner C, Bender T, Cantista P, Karagülle Z (2010) A proposal for a worldwide definition of
 774 health resort medicine, balneology, medical hydrology and climatology. *Int J Biometeorol*
 775 54:495–507. <https://doi.org/10.1007/s00484-010-0321-5>

776 Harzy T, Ghani N, Akasbi N, et al (2009) Short- and long-term therapeutic effects of thermal mineral
 777 waters in knee osteoarthritis: a systematic review of randomized controlled trials. *Clin*
 778 *Rheumatol* 28:501–507. <https://doi.org/10.1007/s10067-009-1114-2>

779 Higgins JPT, Altman DG, Gøtzsche PC, et al (2011) The Cochrane Collaboration’s tool for assessing
 780 risk of bias in randomised trials. *BMJ* 343:d5928. <https://doi.org/10.1136/bmj.d5928>

781 Hou C, Liang L, Chu X, et al (2020) The short-term efficacy of mud therapy for knee osteoarthritis: A
 782 meta-analysis. *Medicine* 99:e19761. <https://doi.org/10.1097/MD.00000000000019761>

783 Kamioka H, Kawamura Y, Tsutani K, et al (2013) A checklist to assess the quality of reports on spa
 784 therapy and balneotherapy trials was developed using the Delphi consensus method: the SPAC

785 checklist. *Complement Ther Med* 21:324–332. <https://doi.org/10.1016/j.ctim.2013.05.002>

786 Karagülle M, Karagülle MZ (2015) Effectiveness of balneotherapy and spa therapy for the treatment
787 of chronic low back pain: a review on latest evidence. *Clin Rheumatol* 34:207–214.
788 <https://doi.org/10.1007/s10067-014-2845-2>

789 Karagülle MZ, Karagülle M (2004) [Balneotherapy and spa therapy of rheumatic diseases in Turkey:
790 a systematic review]. *Forsch Komplementarmed Klass Naturheilkd* 11:33–41.
791 <https://doi.org/10.1159/000077194>

792 Katz U, Shoenfeld Y, Zakin V, et al (2012) Scientific evidence of the therapeutic effects of dead sea
793 treatments: a systematic review. *Semin Arthritis Rheum* 42:186–200.
794 <https://doi.org/10.1016/j.semarthrit.2012.02.006>

795 Keller S, König V, Mösges R (2014) Thermal water applications in the treatment of upper respiratory
796 tract diseases: a systematic review and meta-analysis. *J Allergy* 2014:943824.
797 <https://doi.org/10.1155/2014/943824>

798 Khalilzadeh S, Shirbeigi L, Naghizadeh A, et al (2019) Use of mineral waters in the treatment of
799 psoriasis: Perspectives of Persian and conventional medicine. *Dermatol Ther* 32:e12969.
800 <https://doi.org/10.1111/dth.12969>

801 Khaltayev N, Solimene U, Vitale F, Zanasi A (2020) Balneotherapy and hydrotherapy in chronic
802 respiratory disease. *J Thorac Dis* 12:4459–4468. <https://doi.org/10.21037/jtd-gard-2019-009>

803 Langhorst J, Musial F, Klose P, Häuser W (2009) Efficacy of hydrotherapy in fibromyalgia
804 syndrome--a meta-analysis of randomized controlled clinical trials. *Rheumatology* 48:1155–
805 1159. <https://doi.org/10.1093/rheumatology/kep182>

806 Liu H, Zeng C, Gao S-G, et al (2013) The effect of mud therapy on pain relief in patients with knee
807 osteoarthritis: a meta-analysis of randomized controlled trials. *J Int Med Res* 41:1418–1425.
808 <https://doi.org/10.1177/0300060513488509>

809 Maraver F, Karagülle MZ (2012) *Medical Hydrology and Balneology: Environmental Aspects*.
810 *Publicaciones Universidad Complutense de Madrid*

811 Maraver F, Michán A, Morer C, Aguilera L (2011) Is thalassotherapy simply a type of
812 climatotherapy? *Int. J. Biometeorol.* 55:107–108

813 Martini S, Soliman T, Gobbi G, et al (2018) PKCε Controls Mitotic Progression by Regulating
814 Centrosome Migration and Mitotic Spindle Assembly. *Mol Cancer Res* 16:3–15.
815 <https://doi.org/10.1158/1541-7786.MCR-17-0244>

816 Masselli E, Pozzi G, Vaccarezza M, et al (2020) ROS in Platelet Biology: Functional Aspects and
817 Methodological Insights. *Int J Mol Sci* 21.: <https://doi.org/10.3390/ijms21144866>

818 Matsumoto H, Hagino H, Hayashi K, et al (2017) The effect of balneotherapy on pain relief, stiffness,
819 and physical function in patients with osteoarthritis of the knee: a meta-analysis. *Clin Rheumatol*
820 36:1839–1847. <https://doi.org/10.1007/s10067-017-3592-y>

821 Ministero della Sanità (1995) Modificazioni all’elenco delle patologie che possono trovare reale
822 beneficio dalle cure termali e proroga della sua validità. *Gazzetta Ufficiale*

823 Mirandola P, Gobbi G, Malinverno C, et al (2013) Impact of sulphurous water politzer inhalation on
824 audiometric parameters in children with otitis media with effusion. *Clin Exp Otorhinolaryngol*
825 6:7–11. <https://doi.org/10.3342/ceo.2013.6.1.7>

826 Mirandola P, Gobbi G, Micheloni C, et al (2011) Hydrogen sulfide inhibits IL-8 expression in human
827 keratinocytes via MAP kinase signaling. *Lab Invest* 91:1188–1194.
828 <https://doi.org/10.1038/labinvest.2011.76>

829 Mirandola P, Gobbi G, Sponzilli I, et al (2007) Exogenous hydrogen sulfide induces functional
830 inhibition and cell death of cytotoxic lymphocytes subsets. *J Cell Physiol* 213:826–833.
831 <https://doi.org/10.1002/jcp.21151>

832 Moher D, Liberati A, Tetzlaff J, et al (2009) Preferred reporting items for systematic reviews and
833 meta-analyses: the PRISMA statement. *PLoS Med* 6:e1000097.
834 <https://doi.org/10.1371/journal.pmed.1000097>

835 Morer C, Roques C-F, Françon A, et al (2017) The role of mineral elements and other chemical
836 compounds used in balneology: data from double-blind randomized clinical trials. *Int J*
837 *Biometeorol* 61:2159–2173. <https://doi.org/10.1007/s00484-017-1421-2>

838 Nappi G (2001) *Medicina e clinica termale*

839 National Institutes of Health Study Quality Assessment Tools. [https://www.nhlbi.nih.gov/health-](https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools)
840 [topics/study-quality-assessment-tools](https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools). Accessed 2 Apr 2021

841 Naumann J, Biehler D, Lüty T, Sadaghiani C (2017) Prevention and Therapy of Type 2 Diabetes-
842 What Is the Potential of Daily Water Intake and Its Mineral Nutrients? *Nutrients* 9.:
843 <https://doi.org/10.3390/nu9080914>

844 Naumann J, Sadaghiani C (2014) Therapeutic benefit of balneotherapy and hydrotherapy in the
845 management of fibromyalgia syndrome: a qualitative systematic review and meta-analysis of
846 randomized controlled trials. *Arthritis Res Ther* 16:R141. <https://doi.org/10.1186/ar4603>

847 Pagourelas ED, Zorou PG, Tsaligopoulos M, et al (2011) Carbon dioxide balneotherapy and
848 cardiovascular disease. *Int J Biometeorol* 55:657–663. [https://doi.org/10.1007/s00484-010-0380-](https://doi.org/10.1007/s00484-010-0380-7)
849 7

850 Passali D, Gabelli G, Passali GC, et al (2017) Radon-enriched hot spring water therapy for upper and
851 lower respiratory tract inflammation. *Otolaryngol Pol* 71:8–13.
852 <https://doi.org/10.5604/01.3001.0010.2242>

853 Petticrew M, Egan M, Thomson H, et al (2008) Publication bias in qualitative research: what becomes
854 of qualitative research presented at conferences? *J Epidemiol Community Health* 62:552–554.
855 <https://doi.org/10.1136/jech.2006.059394>

856 Pittler MH, Karagülle MZ, Karagülle M, Ernst E (2006) Spa therapy and balneotherapy for treating
857 low back pain: meta-analysis of randomized trials. *Rheumatology* 45:880–884.
858 <https://doi.org/10.1093/rheumatology/kei018>

859 Quattrini S, Pampaloni B, Brandi ML (2016) Natural mineral waters: chemical characteristics and
860 health effects. *Clin Cases Miner Bone Metab* 13:173–180.
861 <https://doi.org/10.11138/ccmbm/2016.13.3.173>

862 Queirolo V, Galli D, Masselli E, et al (2016) PKC ϵ is a regulator of hypertrophic differentiation of
863 chondrocytes in osteoarthritis. *Osteoarthritis Cartilage* 24:1451–1460.
864 <https://doi.org/10.1016/j.joca.2016.04.003>

865 Queiroz LP (2013) Worldwide epidemiology of fibromyalgia. *Curr Pain Headache Rep* 17:356.
866 <https://doi.org/10.1007/s11916-013-0356-5>

867 Rachakonda TD, Schupp CW, Armstrong AW (2014) Psoriasis prevalence among adults in the United

- 868 States. *J Am Acad Dermatol* 70:512–516. <https://doi.org/10.1016/j.jaad.2013.11.013>
- 869 Raza HMA, Krutulyte G, Rimdeikiene I, Savickas R (2020) Efficacy of Balneotherapy and Mud
870 Therapy in Patients with Knee Osteoarthritis: A Systematic Literature Review. *Aktuelle*
871 *Rheumatologie*
- 872 Rinaldi L, Gobbi G, Pambianco M, et al (2006) Hydrogen sulfide prevents apoptosis of human PMN
873 via inhibition of p38 and caspase 3. *Lab Invest* 86:391–397.
874 <https://doi.org/10.1038/labinvest.3700391>
- 875 Roques C, -C. Mancret R, Tabone W (2012) Beneficial effect of balneotherapy: Achievements of the
876 French association of balneotherapy research and methodological difficulties. *Annals of Physical*
877 *and Rehabilitation Medicine* 55:e351
- 878 Roques CF (2014) Balneotherapy actual medical benefit. Data of evidence for the last twenty years.
879 *Ann Phys Rehabil Med* 57:e159. <https://doi.org/10.1016/j.rehab.2014.03.576>
- 880 Santos I, Cantista P, Vasconcelos C (2016) Balneotherapy in rheumatoid arthritis-a systematic review.
881 *Int J Biometeorol* 60:1287–1301. <https://doi.org/10.1007/s00484-015-1108-5>
- 882 Schuh A (2009) Die Evidenz der Klima- und Thalassotherapie. Ein Review. *Schweizerische*
883 *Zeitschrift für Ganzheitsmedizin / Swiss Journal of Integrative Medicine* 21:96–104
- 884 Shea BJ, Reeves BC, Wells G, et al (2017) AMSTAR 2: a critical appraisal tool for systematic
885 reviews that include randomised or non-randomised studies of healthcare interventions, or both.
886 *BMJ* 358:j4008. <https://doi.org/10.1136/bmj.j4008>
- 887 Stanhope J, Weinstein P, Cook A (2018) Do natural spring waters in Australia and New Zealand
888 affect health? A systematic review. *Journal of Water and Health* 16:1–13
- 889 Stier-Jarmer M, Kus S, Frisch D, et al (2015) Health resort medicine in non-musculoskeletal
890 disorders: is there evidence of its effectiveness? *Int J Biometeorol* 59:1523–1544.
891 <https://doi.org/10.1007/s00484-015-0953-6>
- 892 Sulaiman SK, Enakshee J, Traxer O, Somani BK (2020) Which Type of Water Is Recommended for
893 Patients with Stone Disease (Hard or Soft Water, Tap or Bottled Water): Evidence from a
894 Systematic Review over the Last 3 Decades. *Curr Urol Rep* 21:6. <https://doi.org/10.1007/s11934-020-0968-3>
- 896 Tenti S, Cheleschi S, Galeazzi M, Fioravanti A (2015) Spa therapy: can be a valid option for treating
897 knee osteoarthritis? *Int J Biometeorol* 59:1133–1143. <https://doi.org/10.1007/s00484-014-0913-6>
- 898 The Cochrane Group About Cochrane Reviews. <https://www.cochranelibrary.com/about/about-cochrane-reviews>. Accessed 25 Dec 2020
- 900 The European Parliament and the Council of the European Union (2009) Directive 2009/54/EC of the
901 European Parliament and of the Council of 18 June 2009 on the exploitation and marketing of
902 natural mineral waters. European Union law
- 903 van Tubergen A, van der Linden S (2002) A brief history of spa therapy. *Ann Rheum Dis* 61:273–
904 275. <https://doi.org/10.1136/ard.61.3.273>
- 905 Verhagen AP, Bierma-Zeinstra SMA, Boers M, et al (2007) Balneotherapy for osteoarthritis.
906 *Cochrane Database Syst Rev* CD006864. <https://doi.org/10.1002/14651858.CD006864>
- 907 Verhagen AP, Bierma-Zeinstra SMA, Boers M, et al (2015) Balneotherapy (or spa therapy) for
908 rheumatoid arthritis. *Cochrane Database Syst Rev* CD000518.

909 <https://doi.org/10.1002/14651858.CD000518.pub2>

910 Verhagen AP, de Vet HCW, de Bie RA, et al (1998) Balneotherapy and Quality Assessment:
 911 Interobserver Reliability of the Maastricht Criteria List and the Need for Blinded Quality
 912 Assessment. *J Clin Epidemiol* 51:335–341. [https://doi.org/10.1016/S0895-4356\(97\)00297-7](https://doi.org/10.1016/S0895-4356(97)00297-7)

913 Viegas J, Esteves AF, Cardoso EM, et al (2019) Biological Effects of Thermal Water-Associated
 914 Hydrogen Sulfide on Human Airways and Associated Immune Cells: Implications for
 915 Respiratory Diseases. *Front Public Health* 7:128. <https://doi.org/10.3389/fpubh.2019.00128>

916 Vina ER, Kwoh CK (2018) Epidemiology of osteoarthritis: literature update. *Curr Opin Rheumatol*
 917 30:160–167. <https://doi.org/10.1097/BOR.0000000000000479>

918 Vv.Aa. (2014) Hydroglobe. Definizione di un quadro di riferimento complessivo mondiale per
 919 l'idroterapia. Narcissus.me

920 Xiang J, Wu D, Li J 'an (2016) Clinical Efficacy of Mudpack Therapy in Treating Knee
 921 Osteoarthritis: A Meta-Analysis of Randomized Controlled Studies. *Am J Phys Med Rehabil*
 922 95:121–131. <https://doi.org/10.1097/PHM.0000000000000354>

923 Yuan D, Yu Z-X, Wang W, Chen Y (2019) Head-out immersion in natural thermal mineral water for
 924 the management of hypertension: a review of randomized controlled trials. *Int J Biometeorol*
 925 63:1707–1718. <https://doi.org/10.1007/s00484-019-01780-4>

926 Zhen-han D, Tuo Y, Hui L, et al (2014) A meta-analysis of mud therapy on knee osteoarthritis pain.
 927 *Journal of Clinical Rehabilitative Tissue Engineering Research* 2455–2460

928 洋晴上岡, 律雄黒柳, 泰喜小松, et al (2006) 温泉の治療と健康増進の効果に関する. *日温気物医誌* 69:155–166.
 929 <https://doi.org/10.11390/onki1962.69.155>




Figure 1. Flow-chart describing the study selection process.

The flow-chart was adapted from the model recommended by the PRISMA guidelines (Moher et al. 2009).




Figure 2. Number of included reviews for each health condition.

● Balneotherapy ● Mud therapy ● Spa therapy ● Hydroponic therapy
● Inhalation/irrigation therapy ● Any mineral water/mud-based therapy




Figure 3. Number of included reviews for each type of therapy.

TABLE 1. Main characteristics (PICOS, methodological quality and study authors' conclusions) of included systematic reviews.

Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
An et al. 2019 (An et al. 2019)	Both healthy and diseased subjects (212)	BT	Any type	Various (physiological and clinical)	SR (13 clinical studies of any type)	D	?	Demonstration of various cardiovascular and neuromuscular effects, but uncertain conclusions.
Brosseau et al. 2013 (Brosseau et al. 2002)	Patients with knee OA (172)	BT	Any type	Pain and functionality	SR (3 RCTs)	C	2	Only short-term effects were demonstrated, with significant benefits in terms of pain relief exclusively shown for baths in Dead Sea or hot sulphur water.
Cao et al. 2020 (Cao et al. 2020)	Patients with fibromyalgia (611)	BT	Any type	Pain, QoL and mood	SR + MA (10 RCTs) - CP	C	3	Pooled evidence from analyzed RCTs (follow-up: from 12 to 48 weeks) indicates that BT may reduce pain and improve QoL of patients with fibromyalgia.
Corvillo et al. 2019 (Corvillo et al. 2019)	Patients with neck pain of different origins (658)	BT	No treatment or standard rehabilitation	Pain, functionality, QoL and mood	SR (13 clinical studies of any type)	D	2	Beneficial effects for all studied algofunctional outcomes, and for psychophysical wellbeing.
Falagas et al. 2009 (Falagas et al. 2009)	Patients with any disease (1720 subjects with rheumatic conditions)	BT	Any type	Any relevant clinical outcome	SR (29 RCTs)	C	2	Clinical improvement of several rheumatic diseases (OA, fibromyalgia, ankylosing spondylitis, rheumatoid arthritis). Preliminary evidence of benefits for psoriasis and Parkinson's disease.
Harzy et al. 2009 (Harzy et al. 2009)	Patients with knee OA (493)	BT	Any type	Pain, functionality, drugs	SR (9 RCTs)	B	2	Beneficial effects on pain and joint functionality lasting up to 24 weeks from intervention.
Moraes Silva et al. 2019 (de Moraes Silva et al. 2019)	Patients with chronic venous insufficiency in leg veins (891)	BT	Any type	Any relevant clinical and QoL outcome	SR+MA (7 RCTs)	A	3	Low-to-moderate evidence indicates beneficial effects in terms of pain, QoL, and skin pigmentation due to chronic venous insufficiency.
Schuh 2009 (Schuh 2009)	Patients with any disease (?)	BT (TT)	Any type	Any relevant clinical outcome	SR (12 studies related to marine climatotherapy and TT)	D	?	A combination of thalassotherapy with marine climatotherapy can exert positive effects on psoriasis, atopic dermatitis, and bronchial asthma.
Yuan et al. (2019) (Yuan et al. 2019)	Patients with arterial hypertension (1122)	BT	Any type	Variations of blood pressure	SR (12 RCTs)	C	3	No worsening of blood pressure parameters.
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Beer et al. 2018 (Beer et al. 2018)	Patients with any disease (728)	MT	Any type	Any relevant clinical outcome	SR (35 clinical studies)	D	2	High-quality evidence indicates a beneficial effect for the symptomatic treatment of OA and fibromyalgia.
Espejo-Antunez et al. 2013 (Espejo-Antunez et al. 2013)	Patients with knee OA (2102)	MT	Any type	Symptoms and QoL	SR (20 studies of any type)	D	2	Significant improvement of pain, general symptoms and QoL.
Hou et al. 2020 (Hou et al. 2020)	Patients with knee OA (1106)	MT	Any type	Pain and functionality	SR + MA (11 RCTs)	B	3	Significant amelioration of knee pain and function.
Liu et al. 2013 (Liu et al. 2013)	Patients with knee OA (410)	MT	Any type	Pain and functionality	SR + MA (7 RCTs)	D	2	Significant improvement of OA-related pain.
Xiang et al. 2016 (Xiang et al. 2016)	Patients with knee OA (1010)	MT	Any type	Functionality	SR + MA (10 RCTs)	C	2	No significant improvement in joint functionality.
Zhen-han et al. 2014	Patients with knee OA (410)	MT	Any type	Pain	SR + MA (7 RCTs) - abstract	?	?	Mud therapy can significantly attenuate knee osteoarthritis pain.

(Zhen-han et al. 2014)					only			
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Antonelli & Donelli 2018 (Antonelli and Donelli 2018a)	Both healthy and diseased subjects (684)	SPA-T	Any type	Variations of salivary or serum cortisol levels	SR (15 clinical studies of any type)	C	2	The effect on cortisol levels suggests that intervention can have an anti-stress action and improve stress resilience.
Antonelli et al. 2018 (Antonelli et al. 2018)	Patients with knee OA (1599)	SPA-T	Any type	QoL, algofunctional indices, drugs	SR+MA (17 RCTs)	C	3	Significant improvement of QoL. Beneficial effects on algofunctional indices and painkiller intake.
Bai et al. 2019 (Bai et al. 2019)	Patients with chronic LBP (1038)	SPA-T	Any type	Back pain and mobility	SR+MA (12 RCTs)	C	2	Significant improvement for back pain and functionality.
Forestier & Francon 2008 (Forestier and Francon 2008)	Patients with OA of the limbs (1658)	SPA-T	Any type	Pain, functionality, QoL	SR+MA (19 RCTs)	D	2	Suggestive evidence of a possible beneficial effect, but analyzed studies have some limitations.
Forestier et al. 2016 (Forestier et al. 2016)	Patients with knee OA (2917)	SPA-T	Any type	Any relevant clinical outcome	SR (30 RCTs)	C	2	Evidence of relevant clinical improvements lasting, on average, from 3 to 6 months (and up to 9 months) after intervention.
Forestier et al. 2017 (Forestier et al. 2017)	Patients with chronic LBP (2146)	SPA-T	Any type	Pain, functionality, QoL, drugs	SR (18 RCTs)	C	2	Evidence of improvements in pain, disability, and QoL, lasting 3-6 months after intervention. Possible reduction of painkiller intake.
Fraioli et al. 2013 (Fraioli et al. 2013)	Patients with fibromyalgia (271)	SPA-T	Any type	Pain, symptoms and mood	SR (7 studies)	D	?	Evidence of symptomatic improvement, including pain and mood (depression).
Fraioli et al. 2018 (Fraioli et al. 2018)	Patients with knee OA (1649)	SPA-T	Any type	Pain, functionality, QoL, drugs	SR (12 clinical studies)	D	?	Improvement of pain, joint mobility, NSAID intake, and QoL.
Gravelier et al. 2020 (Gravelier et al. 2020)	Patients with burn scars (115)	SPA-T	Any type	Pain, skin elasticity, QoL	SR (2 RCTs)	C	3	Potential beneficial effects for burn scar recovery, but further scientific evidence is needed.
Kamioka et al. 2006 (菅井 et al. 2006)	Patients with any disease (1425)	SPA-T	Any type	Pain, QoL, drugs, occupational functionality and healthcare costs	SR (18 RCTs)	D	2	Amelioration of all analyzed outcomes with high-quality studies mostly supporting the efficacy of intervention for rheumatic disorders.
Karagülle M. & Karagülle M.Z. 2015 (Karagülle and Karagülle 2015)	Patients with chronic LBP (769)	SPA-T	Any type	Any relevant clinical outcome	SR (8 RCTs)	D	?	Evidence of an improvement in symptomatic management, but further studies are needed to confirm these results.
Langhorst et al. 2009 (Langhorst et al. 2009)	Patients with fibromyalgia (446)	SPA-T	Any type	Any clinical and QoL-related outcome	SR+MA (10 RCTs)	A	2	Moderate evidence of beneficial effects on health-related QoL.
Matsumoto et al. 2017 (Matsumoto et al. 2017)	Patients with knee OA (734)	SPA-T	Any type	Pain and functionality	SR+MA (8 RCTs)	B	3	Possible beneficial effects, but high heterogeneity is found across studies and evidence is not of sufficiently high quality.
Morer et al. 2017 (Morer et al. 2017)	Patients with any rheumatic condition (1118)	SPA-T	HT	Pain, functionality, drugs, QoL, laboratory parameters	SR (27 RCTs)	D	3	Possible beneficial effects, but high heterogeneity and potential risk of bias is reported within and across available studies.
Naumann &	Patients with	SPA-T	Any type	Pain, functionality,	SR+MA (12	B	2	Significant improvements of symptoms and

Sadaghiani 2014 (Naumann and Sadaghiani 2014)	fibromyalgia (553)			QoL, mood	RCTs about SPA-T and 12, excluded, about HT)			QoL with potentially long-lasting effects on pain. No significant effect was observed on depressive symptoms
Pittler et al. 2006 (Pittler et al. 2006)	Patients with chronic LBP (674)	SPA-T	Any type	Pain, functionality, QoL, drugs	SR+MA (5 RCTs)	B	2	Scant but promising evidence which suggests a possible clinical benefit.
Raza et al. 2020 (Raza et al. 2020)	Patients with knee OA (831)	SPA-T	Any type	Pain and functionality	SR+MA (10 RCTs)	C	2	Significant improvement in pain and functionality, as measured with the WOMAC scale.
Santos et al. 2016 (Santos et al. 2016)	Patients with rheumatoid arthritis (496)	SPA-T	Any type	Pain, functionality, QoL, drugs, laboratory parameters	SR (8 RCTs)	B	2	Evidence of a beneficial effect on studied outcomes even up to 3 months after intervention.
Verhagen et al. 2007 (Verhagen et al. 2007)	Patients with OA (498)	SPA-T	Any type	Pain, functionality, QoL	SR+ MA (7 RCTs)	A	3	A positive effect is found when intervention is compared to no treatment, but the quality of evidence is low.
Verhagen et al. 2015 (Verhagen et al. 2015)	Patients with rheumatoid arthritis (579)	SPA-T	Any type	Pain, functionality, QoL	SR+MA (9 RCTs)	A	3	Insufficient evidence to draw conclusions on the topic.
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Casale et al. 2018 (Casale et al. 2018)	Patients with rhinosinusitis (663)	IT/IR	Any type	Functionality of upper airways	SR (11 RCTs)	D	2	Nasal irrigations with mineral waters can be clinically beneficial for patients with rhinosinusitis in terms of endoscopic scores and mucociliary clearance if compared to isotonic solution.
Keller et al. 2014 (Keller et al. 2014)	Patients with rhinosinusitis (840)	IT/IR	Any type	Functionality of upper airways	SR + MA (13 clinical studies of any type)	D	2	Mineral waters (for example, those ones rich in sulphur) can have an integrative role in the management of chronic inflammatory diseases of the upper respiratory tract.
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Böhmer et al. 2000 (Bohmer et al. 2000)	Healthy subjects (54)	HPT (mineral waters with high calcium content)	Dairy products	Bioavailability of calcium	SR (4,controlled studies)	D	?	Bioavailability of calcium from calcium-rich waters is comparable to that one from dairy product consumption.
Naumann et al. 2017 (Naumann et al. 2017)	Patients with cardiovascular risk factors (1089)	HPT (various mineral waters)	Other drinks	Glycemic control	SR (15 RCTs)	C	3	Bicarbonate- and magnesium-rich waters may have a positive impact on glycemic control.
Sulaiman et al. 2020 (Sulaiman et al. 2020)	Both healthy and diseased subjects at risk of kidney stone formation (470 involved in interventional studies)	HPT (mineral waters with high calcium, bicarbonate or magnesium content)	Tap water	Kidney stone prevention, especially calcium stones	SR (10 clinical interventional studies and 5 observational studies)	C	2	Bicarbonate- and magnesium-rich waters may be useful against kidney stone formation. Consumption of waters with a high calcium content can lead to hypercalciuria.
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Stier-Jarmer et al. 2015 (Stier-Jarmer et al. 2015)	Patients with any non-musculoskeletal disease (13782)	Any Hydrologic Therapy	Any type	Any improvement assessed clinically or with diagnostic devices	SR (41 clinical studies of any type)	B	3	Clinical benefits for various skin diseases (atopic dermatitis, psoriasis, ictiosis), respiratory and ENT illnesses (rhinosinusitis, COPD, catarrhal otitis), vascular problems (hypertension, chronic venous insufficiency), digestive complaints (dyspepsia, irritable bowel syndrome), and neuropsychiatric

								conditions (peripheral neuropathy, Parkinson's' disease, psychophysical stress).
--	--	--	--	--	--	--	--	--

Legends:

BT=Balneotherapy (only baths with mineral waters)

COPD=Chronic Obstructive Pulmonary Disease

CP=Conference Proceedings (review presented at a conference/lecture/symposium and then published in a refereed journal as an abstract)

HPT=Hydropinic therapy (drinking mineral waters)

HT=Hydrotherapy (use of tap non-mineral water)

IT/IR=Inhalation therapy/irrigations (aerosols, vapors, nebulizations, humages, and oral/nasal irrigations)

LBP=Low Back Pain

MA=Meta-analysis

MT=Mud therapy (use of therapeutic muds)

NSAID=NonSteroidal Anti-Inflammatory Drug

OA=Osteoarthritis

QoL=Quality of Life

RCT=Randomized Controlled Trial

RQ=Review Quality or overall quality assessed on the basis of review methods (A=high quality, B=moderate quality, C=low quality,

D=very low quality, ?=the review quality was not assessable because the full-text version was irretrievable)

SPA-T=Spa therapy (multicomponent spa-based treatments)

SR=Systematic Review

TQ=Trial Quality or average quality of clinical studies included in each analyzed review (1=high quality, 2=fair quality, 3=poor quality,

?=trial quality was not assessed by the authors of included reviews)

TT=Thalassotherapy (baths with sea water)

Caption:

Included reviews are grouped on the basis of analyzed intervention and alphabetically sorted according to the first author's surname.

TABLE 2. Main characteristics (PICOS and study authors' conclusions) of included high-quality narrative reviews.

Reference	Population (n)	Intervention	Control	Outcomes	Study design	Conclusions
Cacciapuoti et al. 2020 (Cacciapuoti et al. 2020)	Patients with chronic inflammatory skin diseases (?)	BT	Any type	Various health-related outcomes	NR (?)	Benefits for psoriasis and atopic dermatitis, but also for pruritus, prurigo, lichen ruber planus, acne vulgaris, and seborrheic dermatitis.
Bender 2016 (Bender 2016)	Patients with OA (?)	SPA-T	Any type	Pain and QoL	NR (50+ RCTs) - CP	Positive effect on pain and QoL of patients with OA of various joints.
Fortunati et al. 2016 (Fortunati et al. 2016)	Patients with hand OA (168)	SPA-T	Any type	Pain, functionality, QoL	NR (3 RCTs)	Suggestive evidence of a potential benefit, but further investigation is advised to draw conclusions.
Francon et al. 2015 (Françon et al. 2015)	Patients with chronic pain of rheumatic origin (2905)	SPA-T	Any type	Pain, functionality, QoL, drugs	NR (28 RCTs)	Improvement of pain, joint mobility, painkiller intake, and QoL.
Guidelli et al. 2012 (Guidelli et al. 2012)	Patients with fibromyalgia (314)	SPA-T	Any type	Pain, functionality, QoL, mood	NR (8 RCTs)	Evidence of benefits in terms of pain, functionality, QoL, and mood, lasting 3 to 9 months after intervention.
Roques 2014 (Roques 2014)	Patients with any disease (?)	SPA-T	Any type	Any relevant clinical outcome	NR (90 RCTs) - CP	Beneficial effects mostly for arthro-rheumatic diseases, but even for psoriasis, atopic dermatitis, vessel disorders of lower limbs, gynecological conditions, and psychosomatic diseases.
Tenti et al. 2015 (Tenti et al. 2015)	Patients with knee OA (1198)	SPA-T	Any type	Pain, functionality, QoL, drugs	NR (14 RCTs)	Evidence of a positive effect on pain, disability, and QoL, which may last up to 6-9 months.
Passali et al. 2017 (Passali et al. 2017)	Patients with chronic rhinosinusitis, allergic rhinitis or asthmatic bronchitis (110)	IT	Any type	Functionality of upper and lower airways	NR (4 clinical studies of any type)	A 2-week inhalation therapy with radon-enriched water may improve nasal function and respiratory obstruction in patients with allergic respiratory diseases.

Legends:

BT=Balneotherapy (only baths with mineral waters)

COPD=Chronic Obstructive Pulmonary Disease

CP=Conference Proceedings (review presented at a conference/lecture/symposium and then published in a refereed journal as an abstract)

HT=Hydrotherapy (use of tap non-mineral water)

IT/IR=Inhalation therapy/irrigations (aerosols, vapors, nebulizations, humages, and oral/nasal irrigations)

LBP=Low Back Pain

MT=Mud therapy (use of therapeutic muds)

NR=Narrative Review

NSAID=NonSteroidal Anti-Inflammatory Drug

OA=Osteoarthritis

QoL=Quality of Life

RCT=Randomized Controlled Trial

SPA-T=Spa therapy (multicomponent spa-based treatments)

Caption:

Included reviews are grouped on the basis of analyzed intervention and alphabetically sorted according to the first author's surname.

1128 **TABLE 3. Systematic reviews including only clinical studies conducted in a specific country or region of the world.**

Reference	Country	Conclusions
Bender et al. 2014 (Bender et al. 2014)	Hungary	Some evidence of efficacy of Hungarian mud baths for several rheumatic conditions, mostly osteoarthritis.
Drobnik and Stebel 2020 (Drobnik and Stebel 2020)	Poland and Austria	Scientific evidence on the “Tolpa” peloid is not strong enough to formulate specific clinical indications.
Karagulle and Karagulle 2004 (Karagülle and Karagülle 2004)	Turkey	Efficacy of spa therapy with Turkish mineral waters and muds for various rheumatic conditions, including osteoarthritis, fibromyalgia, and rheumatoid arthritis.
Katz et al. 2012 (Katz et al. 2012)	Israel	Baths in Dead Sea water and mud can be useful for the treatment of rheumatic diseases and psoriasis.
Khalilzadeh et al. 2019 (Khalilzadeh et al. 2019)	Iran	Possible efficacy of balneotherapy with Persian mineral waters for the management of psoriasis.
Roques et al. 2012 (Roques et al. 2012)	France	Spa therapy with French mineral waters and muds can be effective for the complementary treatment of osteoarthritis, tendinopathies, benign chronic low back pain, and leg chronic venous insufficiency. Beneficial effects were found for anxiety. Preliminary studies also reported some improvements in patients with neurologic and metabolic conditions.
Stanhope et al. 2018 (Stanhope et al. 2018)	Australia and New Zealand	No evidence of efficacy due to complete lack of clinical studies on the topic.

1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152

1153

TABLE 4. Water types and clinical indications according to the “HydroGlobe” study (2014).

Affected apparatus	Example	Administration	Recommended water types
Ear-Nose-Throat and Respiratory tract	Chronic rhinosinusitis or bronchitis	Inhalations, irrigations	Sulphurous; Salt, bromine and iodine; Bicarbonate; Arsenical-ferruginous.
Cardiovascular system	Chronic venous insufficiency	Baths	Carbonic.
Gynecological apparatus	Chronic vaginitis	Irrigations	Sulphurous; Salt, bromine; Bicarbonate; Sulphate.
Urinary tract	Recurrent kidney stones	Oral intake	Oligomineral (low mineral content); Bicarbonate.
Gastrointestinal system	Irritable bowel disease with constipation	Oral intake	Bicarbonate; Sulphate; Salt.
Skin	Psoriasis, atopic dermatitis	Baths	Salt, bromine and iodine; Radioactive; Bicarbonate; Sulphurous.
Musculoskeletal system	Osteoarthritis, fibromyalgia	Baths	Sulphurous; Salt, bromine and iodine; Radioactive.

Legends:

According to relevant directives of the European Union and to traditional definitions, mineral waters are labelled as follows (Nappi 2001; Vv.Aa. 2014; Quattrini et al. 2016):

- “Arsenical-ferruginous” waters: when they contain both arsenic and iron, either as a ferrous or ferric ion. Waters are considered “ferrous” or “ferruginous” when iron content is > 1 mg/L.
- “Bicarbonate-rich” waters: when bicarbonate content is > 600 mg/L. If the calcium content of these waters is > 150 mg/L and magnesium content is > 50 mg/L, they are also defined as “calcium- and magnesium-rich” waters.
- “Carbonic” waters: they spring up naturally with a detectable content of free carbon dioxide. When CO₂ content is > 250 mg/L, they are defined as “acid waters” due to their low pH.
- “Oligomineral” waters: characterized by a low mineral content, with a fixed residue at 180°C inferior to 500 mg/L. On the contrary, when the fixed residue exceeds 1500 mg/L, waters are defined as “highly mineralized”.
- “Radioactive” waters: they have a radioactivity of at least 1 nCi/L, mostly due to their content of Radon.
- “Sulphate-rich” waters: when sulphate content is > 200 mg/L. If the calcium content of these waters is > 150 mg/L and magnesium content is > 50 mg/L, they are also defined as “calcium- and magnesium-rich” waters.
- “Sulphurous” waters: they have a high content of bivalent sulphur.
- “Waters rich in salt, bromine and iodine”: they have a high mineral content and, like seawater, they are rich in sodium (in the form of NaCl) and other minerals. If sodium content is > 200 mg/L, they are labelled as “sodium-rich” waters. They can also contain elements like bromine and iodine.

Title

Clinical efficacy of Medical Hydrology: an umbrella review

Abstract

Introduction. The aim of this research was to summarize available scientific evidence on the efficacy of Medical Hydrology for the management of any health condition.

Methods. The search was conducted on March 26th, 2021, in the following databases: Medline (via PubMed), EMBASE, Web of Science, Cochrane Library, and Google Scholar. All relevant literature reviews investigating the clinical efficacy of interventions characterized by the use of natural mineral waters and muds were included. The quality of studies was assessed with the “AMSTAR 2” tool.

Results. After article screening, 49 reviews were included in this work. Overall, retrieved scientific evidence suggests that spa therapy is beneficial for patients affected by some specific musculoskeletal conditions, with improvements potentially lasting up to 9 months. Moreover, balneotherapy can be an integrative support for the management of chronic venous insufficiency and some inflammatory skin diseases like psoriasis. The role of spa therapy in rehabilitation appears relevant as well. More limited, although interesting evidence exists for inhalation and hydropinic therapies.

Discussion. Globally, retrieved evidence suggests that, besides individual wellbeing, Medical Hydrology can be useful for public health. In particular, higher-quality studies seem to support the integrative use of spa-related interventions for conditions like osteoarthritis, fibromyalgia, low back pain of rheumatic origin, and chronic venous insufficiency. However, the body of evidence has some limitations and further clinical trials should be designed for each relevant application to consolidate and expand acquired knowledge.

Keywords: Medical hydrology; Clinical efficacy; Public health; Integrative medicine; Umbrella review.

Authors

First author (corresponding author*): **Michele Antonelli**. Affiliation: AUSL-IRCCS Reggio Emilia, Reggio Emilia (Italy). *Address: Via Giovanni Amendola 2, 42122 Reggio Emilia (Italy). Email: michele.antonelli@ausl.re.it ORCID: 0000-0002-5941-6604

Second author: **Davide Donelli**. Affiliation: AUSL-IRCCS Reggio Emilia, Reggio Emilia (Italy). Email: donelli.davide@gmail.com ORCID: 0000-0002-7895-2840

Third author: **Licia Veronesi**. Affiliation: Department of Medicine and Surgery, University of Parma, Parma (Italy). Email: licia.veronesi@unipr.it

Fourth author: **Marco Vitale**. Affiliation (1): Department of Medicine and Surgery, University of Parma, Parma (Italy). Affiliation (2): FoRST Foundation, Rome (Italy). Email: marco.vitale@unipr.it

Last author: **Cesira Pasquarella**. Affiliation: Department of Medicine and Surgery, University of Parma, Parma (Italy). Email: cesiraisabellamaria.pasquarella@unipr.it ORCID: 0000-0001-7579-4827

Funding: this research work was not funded.

Conflicts of interest/Competing interests: no conflict of interest or competing interest has to be declared by the authors.

Ethics approval: not applicable.

Introduction

Background and definitions

Medical Hydrology (or Balneology) is a biomedical discipline with a long-standing tradition, which investigates clinical uses and health-related applications of natural mineral waters and muds for preventive, therapeutic, and rehabilitative purposes (Nappi 2001; Gutenbrunner et al. 2010; Maraver and Karagülle 2012). These treatments are usually administered in health facilities labelled as “spa centers”, where “spa” stands for “*salus per aquam*” or “*sanitas per aquam*”, an ancient Latin expression which literally refers to the pursuit of “health through water” (van Tubergen and van der Linden 2002).

According to the European Union (EU) law, as stated in the Council Directive 80/777/EEC, a natural mineral water is defined as a microbiologically wholesome water (originating in an underground water table or deposit and emerging from a spring), which can be distinguishable from ordinary drinking water by its nature and original purity (The European Parliament and the Council of the European Union 2009). In particular, natural mineral waters are characterized by a specific mineral content and biochemical composition, and by certain effects, including their potential pharmacological, physiological, and clinical action (The European Parliament and the Council of the European Union 2009). In several European countries, this regulatory definition is also accepted by the scientific community (Gutenbrunner et al. 2010). Therefore, in this article, we referred to “natural mineral waters” or, simply, “mineral waters” following the above mentioned description and indicating spring waters used for preventive, therapeutic or rehabilitative purposes in spa centers. With the aim of unequivocally classifying all spa-based interventions and considering a common set of definitions used in the scientific literature (Pittler et al. 2006; Gutenbrunner et al. 2010; Gomes et al. 2013; Fioravanti et al. 2017; Antonelli and Donelli 2018a), the following terms were used in this research work:

- “Balneotherapy”: any treatment which only involves the partial or total-body immersion in natural mineral waters (Antonelli and Donelli 2018a; Antonelli et al. 2018). When sea water was used, the term “Talassotherapy” was adopted (Maraver et al. 2011).
- “Mud therapy”: any treatment which only involves the application over skin of muds or other peloids, which were defined as natural products used as therapeutic agents and consisting of a mixture of minero-medicinal water with organic and inorganic material, like clay minerals (Carretero 2002). Although usually classified among balneo-therapeutic interventions (Fioravanti et al. 2017), mud therapy was distinguished from standard balneotherapy for its specific characteristics.
- “Spa therapy”: any spa-based multicomponent treatment which includes at least a treatment involving the therapeutic use of natural mineral waters (Pittler et al. 2006; Karagülle and

Karagülle 2015; Antonelli and Donelli 2018a). Due to its complexity and integrated therapeutic action, fango-balneotherapy, namely mud baths followed or preceded by standard balneotherapy with mineral waters, was classified under this category.

- “Inhalation therapy”: any treatment characterized by the inhalation of natural mineral waters and their gases in the form of vapors, nebulizations, aerosols, politzers, and humages (Costantino et al. 2006; Keller et al. 2014). Oral and nasal irrigations were also included in this category.
- “Hydropinic therapy”: any treatment which implies the oral ingestion of natural mineral waters with medicinal properties (Albertini et al. 2007).

Sometimes, especially in Italy or France, the old-style term “chreno-therapy” (from the Greek “krēnē”: spring, well, fountain) is adopted to indicate all interventions based on the external and/or internal administration of mineral waters and muds (Nappi 2001; Gutenbrunner et al. 2010; Maraver and Karagülle 2012). However, this term was not used in this article to avoid potential misunderstandings. Whenever hydrologic treatments involved the sole use of tap non-mineral water, for example in the control group, the generic term “hydrotherapy” was adopted.

Rationale

In Italy, costs of treatments based on natural mineral waters and therapeutic muds can be partially or fully covered by the National Healthcare System, and the same happens in other countries, both within and outside Europe (Gutenbrunner et al. 2010). In accordance with a regulatory document approved by the Italian Ministry of Health, conditions which can benefit from Medical Hydrology include osteoarthritis, extra-articular rheumatisms, chronic sinusitis and rhinosinusitis, vasomotor rhinopathy, chronic laryngopharyngitis, chronic catarrhal otitis, tubaric stenosis of flogistic origin, chronic bronchitis, psoriasis, eczema and atopic dermatitis, seborrheic dermatitis, chronic venous insufficiency, constipation due to irritable bowel syndrome, dyspepsia caused by gastroenteric or biliary dysfunction, recurrent kidney stones, and some forms of chronic vaginitis (Ministero della Sanità 1995).

Globally, these conditions are mostly caused by chronic diseases with an epidemiologically considerable impact on health-related quality of life (QoL). For example, osteoarthritis is a frequent degenerative disorder of the musculoskeletal system, especially among elderly subjects, with around 14 million patients only in the USA, and it is expected to become one of the leading causes of disability worldwide (Vina and Kwoh 2018). A relevant impact on public health is also shared by other health conditions which can benefit from spa-based interventions, such as fibromyalgia (Queiroz 2013), chronic venous insufficiency (Al Shammeri et al. 2014), chronic inflammatory respiratory diseases (Ferrante et al. 2017), and phlogistic skin disorders like psoriasis (Rachakonda et al. 2014). However, the above mentioned clinical indications were last updated and revised years ago (Ministero della Sanità 1995), and the most recent attempt to collect all available evidence on the topic with a

cross-cutting approach dates back to March 2014, when the “Hydroglobe” project was published by a panel of experts with the technical support of the World Health Organization (WHO) (Vv.Aa. 2014). In the light of what stated above, considering that Medical Hydrology has a long-standing tradition (a), that mineral waters and therapeutic muds are easily available as natural resources (b), that spa-based interventions are usually prescribed for epidemiologically relevant health conditions (c), and that scientific evidence rapidly evolves (d), it can be useful to synthesize and critically appraise available findings on the topic with a systematic approach, thus following the basic principles of Evidence-Based Medicine (EBM). This would help to outline an updated list of evidence-based indications for treatments with mineral waters and therapeutic muds, and to guide further research in this field, which is important for both individual wellbeing and public health.

Study objective

The aim of this research work was to summarize available scientific evidence on the efficacy of Medical Hydrology for the management of any health condition with a critical assessment of all relevant literature reviews.

Methods

Study design and protocol registration

A systematic review of literature reviews and meta-analyses (umbrella review) was conducted following the internationally-accepted PRISMA (*Preferred Reporting Items for Systematic Reviews and Meta-Analyses*) guidelines (Moher et al. 2009). Additional methodological recommendations for umbrella reviews were taken into account to improve the overall quality of this work (Fusar-Poli and Radua 2018). The review protocol was registered in OSF (*Open Science Framework*) under the following DOI: 10.17605/OSF.IO/2NJ5X (<https://osf.io/2nj5x>).

Eligibility criteria

The following PICOS criteria were applied for the inclusion and exclusion of studies in this review:

- Population: healthy subjects and/or individuals affected by any disease diagnosed in accordance with validated clinical criteria.
- Intervention: all spa interventions based on the use of natural mineral waters and therapeutic muds (balneotherapy, mud therapy, spa therapy, inhalation/irrigation therapy, hydropinic therapy). Reviews were excluded when they focused on studies with non-mineral tap water.
- Control: any type, including no control.

- Outcomes: any relevant clinical outcome (symptomatic and functional improvements, variations in laboratory parameters, reduction of drug consumption, effects on health-related quality of life).
- Study design: systematic reviews and meta-analyses of clinical studies. Reviews were excluded when they had no description of research methods and when they included only pre-clinical laboratory studies. Relevant high-quality narrative reviews (reporting a full description of their research methods and based on a quite extensive literature search) were also included along with systematic reviews, but they were synthesized in a separated section of the manuscript. Following the Cochrane recommendations, systematic reviews were defined as any “review [that] attempts to identify, appraise and synthesize all the empirical evidence that meets pre-specified eligibility criteria to answer a specific research question” (The Cochrane Group). Reviews had to be published in a refereed journal to be eligible for inclusion.

Information sources

Following shared recommendations for optimal database combinations (Bramer et al. 2017), the literature search was conducted in Medline (via PubMed), EMBASE, Web of Science, Cochrane Library, and Google Scholar from inception to March 2021.

Search

The search was conducted on March 26th, 2021. A first tentative pilot search was performed by one author only (M.A.) on January 7th, 2020, then, after refinements, the entire search was updated and all articles were screened again by two authors independently (M.A., D.D.).

The following keywords were used: “balneotherapy”, “hydrotherapy”, “thalassotherapy”, “spa therapy”, “water therapy”, “aquatic therapy”, “mud therapy”, “peloid therapy”, psammotherapy, “inhalation therapy”, “endotympanic insufflation”, “politzer”, “cave therapy”, “hydropinotherapy”, “mineral water*”, “thermal water*”, “hot spring water*”. Specific search strategies adopted for all screened databases were disclosed in the Electronic Supplementary Materials.

In order to narrow down the search and increase its precision, keywords used in Google Scholar were markedly simplified. References of important regulatory papers and overviews of reviews were screened with a “snowballing technique” for an additional check. An author (M.V.), with his long-standing experience in Medical Hydrology research, performed a supplementary search to make sure not to have missed any highly relevant articles in this field of study.

Study selection

The article screening process was conducted with the help of EndNote® software (version X4) by two authors independently (M.A., D.D.). In cases of discrepancies, another author (C.P.) was consulted

and disagreements were discussed until consensus was reached. Only systematic reviews and meta-analyses (along with the most relevant and extensive narrative reviews) matching the above-mentioned PICOS criteria were included in this umbrella review. All studies written in English, French, Italian, Spanish, and Portuguese were considered eligible for inclusion. Studies matching inclusion criteria but written in other languages (German, Japanese) were consulted with the support of a translator. In order to maximize retrievable evidence on the topic and to reduce the risk of publication bias, even articles with only an English abstract available for consultation were included in this umbrella review. This detail was reported in the table describing the main characteristics of included systematic reviews, although it was not possible to assess the quality of these research works due to the lack of detailed methodological information.

Data collection process

Data extraction was conducted manually with a predefined Excel® table designed in accordance with the PICOS criteria. Data extraction was performed by one author (M.A.) with a second check by another author (C.P.). In any case of missing data, authors were contacted via email or through ResearchGate®. The full-text version of a review article was retrieved in this way (Raza et al. 2020).

Data items

Data items extracted from included studies were the following ones: the number and main characteristics of study populations, the type of intervention and control, all relevant clinical outcomes, and the study design (namely whether each review was systematic or narrative, and if it was coupled or not with a meta-analysis).

To properly account for the reliability and consistency of clinical evidence on the topic, it was decided to also report the overall quality of primary studies analyzed in all systematic reviews eligible for inclusion. In this regard, the set of trials included in each review was globally judged as characterized by a good (1), fair (2) or poor (3) quality, depending on the risk-of-bias assessment performed by the review authors. This system is based on the three-tier quality rating of scientific studies recommended by the American National Institutes of Health (National Institutes of Health).

Risk of bias and quality of studies

The quality of included systematic reviews was independently evaluated by two authors (M.A., D.D.) with a dedicated appraisal tool called “AMSTAR 2”, specifically developed and validated by a team of expert methodologists for this purpose (Shea et al. 2017). In cases of disagreement, items were discussed with another author (C.P.) until consensus was reached. The “AMSTAR 2” tool provides a 16-item checklist aimed to explore different methodological domains, including the appropriateness of inclusion/exclusion criteria, search strategies, article selection, data extraction, risk-of-bias assessment, heterogeneity evaluation, quantitative synthesis and critical discussion. Each item

corresponds to a question, which can be answered with “yes” or “no” (sometimes “yes, but partially” is available as a middle response). Included reviews were evaluated one by one and their overall quality was rated as:

- High quality (A) if zero or one non-critical weaknesses were found.
- Moderate quality (B) if two or more non-critical weaknesses were found.
- Low quality (C) if one critical flaw (with or without non-critical weaknesses) was found.
- Very low quality (D) if two or more critical flaws (with or without non-critical weaknesses) were found.

As reported in our study protocol, it was originally planned to use the appraisal tool developed by the American NIH (National Institutes of Health) to assess the quality of included systematic reviews. However, in consideration of the need for a deeper methodological analysis of retrieved reviews, it was eventually decided to resort to the more specific and widely used “AMSTAR 2” tool.

The methodological quality of narrative reviews was assessed with the SANRA scale (Baethge et al. 2019) during the article selection process and only higher-quality narrative reviews were included in this work. The SANRA is a 6-item scale which evaluates the relevance/importance of a narrative review (1), whether its aim is sufficiently focused (2), if the literature search is broad enough (3), whether referencing (4), scientific reasoning (5), and presentation of data (6) are appropriate. Each item score can vary from 0 to 2, and the overall review quality score can range from 0 to 12 (high-quality narrative reviews usually score 9 or more SANRA points).

Publication bias and potential biases across studies were only qualitatively assessed because no quantitative synthesis was feasible due to the detection of a high level of heterogeneity across included studies.

Synthesis of results

The main characteristics of included reviews were reported in two tables, then retrieved evidence was qualitatively synthesized and critically discussed. Results of the study quality assessment were used for a critical discussion. Included reviews were also grouped on the basis of intervention type, health condition of interest, study design (systematic/narrative) and methodological quality. Systematic reviews specifically providing regional data, namely reviews appraising evidence from clinical studies only conducted in a given country of the world, were summarized in another table and mentioned in the “Discussion” section for better comprehensiveness.

Results

Study selection

Overall, the literature search yielded 803 results and, after screening and selection of potentially eligible articles, 49 reviews (41 systematic and 8 narrative reviews) were eventually included in this research work. Details of the article selection process, along with the main reasons for exclusion of non-eligible studies, were summarized in a dedicated flowchart (Figure 1). The list of all articles eligible for a full-text assessment and then excluded after a thorough evaluation was provided in the Electronic Supplementary Materials. The quality of a review available as a conference abstract in a refereed journal (Cao et al. 2020) was assessed on the basis of information retrieved from the corresponding preprint (<https://www.researchsquare.com/article/rs-16293/v1>). An article was kindly provided by the authors after our direct inquiry (Raza et al. 2020). It was not possible to consult the full-text version of a review but this study was included all the same for better comprehensiveness (essential data were extracted from the abstract) (Zhen-han et al. 2014). A meta-analysis found through snowballing and published as a Ph.D. thesis was excluded from the main search but mentioned in the Discussion section to better analyze the mechanisms of action of mud therapy for osteoarthritis (Crespin 2017).

Characteristics of included studies

The main characteristics of all included studies, collected and described in accordance with the PICOS criteria, were reported in Table 1 (systematic reviews) and Table 2 (narrative reviews), along with a brief summary of the authors' conclusions and, for systematic reviews, with their overall methodological quality evaluated in accordance with the AMSTAR-2 recommendations (Bohmer et al. 2000; Brosseau et al. 2002; 泮青 et al. 2006; Pittler et al. 2006; Verhagen et al. 2007, 2015; Forestier and Françon 2008; Schuh 2009; Harzy et al. 2009; Falagas et al. 2009; Langhorst et al. 2009; Guidelli et al. 2012; Fraioli et al. 2013, 2018; Espejo-Antúnez et al. 2013; Liu et al. 2013; Zhen-han et al. 2014; Roques 2014; Keller et al. 2014; Naumann and Sadaghiani 2014; Karagülle and Karagülle 2015; Françon et al. 2015; Tenti et al. 2015; Stier-Jarmer et al. 2015; Fortunati et al. 2016; Xiang et al. 2016; Forestier et al. 2016, 2017; Bender 2016; Santos et al. 2016; Matsumoto et al. 2017; Naumann et al. 2017; Passali et al. 2017; Morer et al. 2017; Casale et al. 2018; Antonelli and Donelli 2018a; Antonelli et al. 2018; Beer et al. 2018; An et al. 2019; Corvillo et al. 2019; de Moraes Silva et al. 2019; Bai et al. 2019; Yuan et al. 2019; Raza et al. 2020; Sulaiman et al. 2020; Hou et al. 2020; Cao et al. 2020; Cacciapuoti et al. 2020; Gravelier et al. 2020).

Population

The number of study participants whose data were analyzed within included systematic reviews varied from a minimum of 54 (Bohmer et al. 2000) to a maximum of 13782 (Stier-Jarmer et al. 2015), with a median value of 731. In the majority of included reviews, regardless of their design (systematic or narrative), study participants were patients with chronic conditions, such as rheumatic (osteoarthritis, fibromyalgia, back pain of rheumatic origin, rheumatoid arthritis), cardiovascular

(chronic venous insufficiency, hypertension), dermatologic, respiratory, otorhinolaryngological, neurologic, digestive, and urologic diseases (Figure 2). In a review, the effects of hydropinic therapy on healthy subjects were studied (Bohmer et al. 2000), whereas studies with both healthy and diseased individuals were analyzed in three research works (Antonelli and Donelli 2018a; An et al. 2019; Sulaiman et al. 2020).

Intervention

Most included reviews analyzed the efficacy of balneotherapy (n=10) and mud therapy (n=6) alone or in combination with other non-spa-related treatments (n=26) (Figure 3). In some cases, spa therapy also involved physical rehabilitation, relaxing massage, diet prescriptions for weight loss or for preventive purposes, pharmacological treatments, and psychological support. Only in a few reviews the efficacy of hydropinic therapy (n=3) and inhalation therapy or nasal irrigations (n=3) were investigated. In one (n=1) research work, the efficacy of any type of spa-related therapy was studied (Stier-Jarmer et al. 2015).

Control

Most analyzed reviews did not have specific restrictions in terms of control type for primary study inclusion. In a systematic review, subjects with cervical pain undergoing balneotherapy were compared to individuals sharing the same health condition who did not receive any treatment or who were administered a cycle of standard rehabilitation (Corvillo et al. 2019). In two research works, the therapeutic efficacy of spa therapy was compared to hydrotherapy with tap non-mineral water (Morer et al. 2017; Sulaiman et al. 2020). Finally, in other systematic reviews, the health effect of orally taking two different mineral waters was compared with modifications induced by consuming other foods and nutritional products (Bohmer et al. 2000; Naumann et al. 2017).

Outcomes

Main clinical outcomes of analyzed reviews included the following ones: symptoms (mostly pain, evaluated with a Visual Analogue Scale), functionality and disability (sometimes assessed in combination with pain using algo-functional scales), drug consumption (especially with regard to the long-term intake of painkillers), quality of life (measured with specific questionnaires), results of various diagnostic investigations and laboratory parameter assessment (including biomarkers of inflammation, metabolic indices, circulating levels of hormones and other biochemical substances).

Study design

Thirtyseven (41) included research works were systematic literature reviews, and seventeen (17) of them were also coupled with a meta-analysis (Pittler et al. 2006; Verhagen et al. 2007, 2015; Forestier and Françon 2008; Langhorst et al. 2009; Liu et al. 2013; Zhen-han et al. 2014; Keller et al. 2014;

Xiang et al. 2016; Matsumoto et al. 2017; Naumann et al. 2017; Antonelli et al. 2018; de Moraes Silva et al. 2019; Bai et al. 2019; Raza et al. 2020; Hou et al. 2020; Cao et al. 2020). Eight (8) included reviews had a narrative design, but they still provided an extensive, quasi-systematic and valuable overview of the scientific literature (Guidelli et al. 2012; Roques 2014; Françon et al. 2015; Tenti et al. 2015; Fortunati et al. 2016; Bender 2016; Passali et al. 2017; Cacciapuoti et al. 2020), thus outlining a state-of-the-art description of specific subtopics, such as the efficacy of spa therapy for respiratory illnesses (Passali et al. 2017), skin diseases (Cacciapuoti et al. 2020), or hand osteoarthritis (Fortunati et al. 2016). The number of clinical studies included in the systematic reviews varied from a minimum of 3 (Brosseau et al. 2002; Fortunati et al. 2016) to a maximum of 41 (Stier-Jarmer et al. 2015), with a median of 12. Such primary studies were mostly Randomized Controlled Trials, although some of them were characterized by a different design, like nonrandomized trials, uncontrolled studies, and observational investigations. The quality of clinical studies tended to range from fair to poor, with a potential risk of bias mostly arising from small sample size, lack of adequate control, no randomization of trial participants, and poor information about blinding of intervention.

Quality of included studies

After the methodological assessment of included systematic reviews with the “AMSTAR 2” tool, the quality of analyzed research works was judged as good (A) to moderate (B) in 11 cases (Pittler et al. 2006; Verhagen et al. 2007, 2015; Harzy et al. 2009; Langhorst et al. 2009; Naumann and Sadaghiani 2014; Stier-Jarmer et al. 2015; Santos et al. 2016; Matsumoto et al. 2017; de Moraes Silva et al. 2019; Hou et al. 2020) and low (C) to very low (D) in the remaining cases, as reported in Table 1 (see the Supplementary Materials for further details). It was not possible to assess the quality of a systematic review because its full-text version was unavailable (Zhen-han et al. 2014). In general, the most frequent methodological weaknesses found within included reviews were excessively narrow search strategies, a poor description of study selection and evaluation processes, and limited consideration of the trial risk-of-bias assessment for informing a critical discussion. Major strengths were usually a clear definition of the research question and, where applicable, a good methodological level of meta-analyses.

Discussion

Efficacy of interventions

Scientific evidence from included reviews indicates that balneotherapy, mud therapy, and spa therapy can significantly improve clinical parameters like pain, joint functionality, mobility, and quality of life of patients with chronic musculoskeletal conditions, mostly osteoarthritis, fibromyalgia, and other

pain-related rheumatic diseases. For these subjects, beneficial effects may last up to 9 months after intervention (on average, 3 to 6 months) (Forestier et al. 2016, 2017), and clinical improvements are associated with a reduced intake of analgesic drugs (Françon et al. 2015; Tenti et al. 2015; Forestier et al. 2017; Fraioli et al. 2018; Antonelli et al. 2018). Furthermore, in two included research works, it was demonstrated that balneotherapy is superior to the same treatment administered in pools with tap water (“sham balneotherapy”) in terms of clinical benefits and improved quality of life among patients with osteoarthritis and fibromyalgia, thus underscoring the contribution of the specific biochemical composition of natural mineral waters to the overall therapeutic effect (Morer et al. 2017; Antonelli et al. 2018).

With regard to vascular health, balneotherapy can be useful for the amelioration of pain, quality of life, and skin pigmentation due to chronic venous insufficiency in lower limbs (Stier-Jarmer et al. 2015; de Moraes Silva et al. 2019), and it appears not to have negative effects on blood pressure levels (Yuan et al. 2019). More limited evidence suggests that balneotherapy, including talassotherapy (sea water baths), can be beneficial for patients with skin diseases like psoriasis and atopic dermatitis (Schuh 2009; Falagas et al. 2009; Stier-Jarmer et al. 2015; Cacciapuoti et al. 2020), and possibly for the integrative treatment of burn scars (Gravelier et al. 2020).

Balneotherapy, mud therapy, and spa therapy can also improve mental wellbeing and psychophysical stress, and this is demonstrated by studies in which stress hormone levels were measured (Roques 2014; Stier-Jarmer et al. 2015; Antonelli and Donelli 2018a). An interesting role for orthopedic or neurological patients of aquatic rehabilitation in pools with natural mineral waters has been underscored by several authors, with beneficial effects on the most important clinical outcomes (Falagas et al. 2009; Stier-Jarmer et al. 2015).

With regard to inhalation therapy, clinical improvements were observed in patients with diseases of the upper and lower respiratory tract, mostly chronic rhinosinusitis and bronchitis (Schuh 2009; Keller et al. 2014; Stier-Jarmer et al. 2015; Casale et al. 2018), but even in subjects with chatarral otitis (Stier-Jarmer et al. 2015). Scientific evidence suggests an interesting role of highly mineralized or sulphur waters for these illnesses (Schuh 2009; Keller et al. 2014; Casale et al. 2018).

Considering hydropinic therapy, study results indicate that the bioavailability of calcium from calcium-rich waters is comparable to that one derived from dairy products (Bohmer et al. 2000), and that the consumption of bicarbonate- and magnesium-rich waters may be beneficial for kidney stone prevention and for an improvement of glycemic control (Naumann et al. 2017; Sulaiman et al. 2020). Moreover, hydropinic therapy with highly mineralized waters can be useful for constipation due to irritable bowel syndrome (Stier-Jarmer et al. 2015).

Systematic reviews including only clinical studies conducted in a specific country or region of the world were collected in Table 3 (Karagülle and Karagülle 2004; Roques et al. 2012; Katz et al. 2012; Bender et al. 2014; Stanhope et al. 2018; Khalilzadeh et al. 2019; Drobnik and Stebel 2020). These research works were mostly carried out in regions where Medical Hydrology is widely spread and

well known, such as Europe or the Middle East, and they outlined an efficacy profile of balneotherapy, mud therapy, and spa therapy which is similar to that one already described in literature reviews collected in Table 1 and Table 2. Additionally, some interesting findings suggested that balneotherapy with specific waters characterized by a high mineral content, like Dead Sea water, can be useful for the treatment of psoriasis (Katz et al. 2012; Khalilzadeh et al. 2019). Some authors also underscored the lack of relevant studies in continents like Oceania, thus urging the need for clinical investigations in Australia or New Zealand, where mineral water springs exist, but they are currently underused for medicinal purposes (Stanhope et al. 2018).

Globally, systematic reviews characterized by a higher overall quality of their methodological design tended to support the efficacy of balneotherapy, mud therapy, and spa therapy for the integrative management of osteoarthritis (especially knee osteoarthritis), fibromyalgia, low back pain of rheumatic origin, and chronic venous insufficiency (Pittler et al. 2006; Verhagen et al. 2007, 2015; Harzy et al. 2009; Langhorst et al. 2009; Naumann and Sadaghiani 2014; Stier-Jarmer et al. 2015; Santos et al. 2016; Matsumoto et al. 2017; de Moraes Silva et al. 2019; Hou et al. 2020). However, the authors underscored that clinical evidence needs to grow before firm conclusions can be drawn, and this is even more relevant for non-rheumatic conditions (for example, respiratory or skin illnesses). All the same, in general, considering both trial results and empirical observations, spa-based treatments appear useful for rehabilitation and chronic disease management, because they seem capable of exerting a beneficial action on symptom control and psychophysical wellbeing.

Safety and tolerability of interventions

Globally, evidence from included studies suggests that balneotherapy, mud therapy, spa therapy, inhalation therapy, and hydropinic therapy are quite safe and well tolerated by patients, provided that all necessary medical and hygienic precautions are taken in advance.

The most important contraindications to treatments based on mineral waters and therapeutic muds are mainly derived from tradition and they can be grouped into three categories (Nappi 2001; Vv.Aa. 2014):

- Contraindications related to the disease stage: spa-based interventions are not to be administered when a patient is affected by an acute disease or during symptomatic relapses of chronic conditions.
- Contraindications related to the patient's illnesses and comorbidities: they include infectious conditions, cancer, and unstable or poorly controlled diseases (severe heart failure, advanced kidney insufficiency, uncontrolled hypertension, cirrosis, medically-unresponsive epilepsy...).
- Contraindications related to the type of intervention: they depend on specific characteristics of single mineral waters and therapeutic muds.

Possible side effects of spa-based interventions are usually mild and often resolve spontaneously after treatment discontinuation: they are mostly due to an individual response, thus being highly “patient-specific”, and their occurrence is worsened by an improper or unsupervised administration of intervention (Nappi 2001; Vv.Aa. 2014). Side effects generally include symptoms like headache, dizziness and nausea, mild relapses of local chronic pain, sleep disturbances, heart palpitations, or a general sensation of irritability and fatigue (rarely coupled with a short-lasting low-grade fever) (Vv.Aa. 2014). Additionally, intervention-specific side effects can occur, such as diarrhea and hydroelectrolytic imbalances due to an overconsumption of highly mineralized waters, a temporary increase of fluid secretions in the airways (with runny nose and cough) after some inhalation therapies, or cutaneous irritation caused by hot mud application on irritable skin. Extreme caution is also advised in more fragile individuals like pediatric patients, pregnant women, and very elderly subjects, whose clinical response to spa-based interventions can be less predictable with a potential higher incidence of more severe side effects. For all these reasons, a medical check and supervision are strongly advised for an appropriate prescription of spa-based interventions, not only to make the most of them on the basis of the patient’s characteristics and disease, but even to avoid the onset of adverse events, thus optimizing the safety and tolerability of such treatments.

Mechanisms of action: hypotheses and evidence-based explanations

In light of scientific evidence described by expert authors of the “*HydroGlobe*” project, effects on health of Medical Hydrology-related treatments have been reported to be the following ones (Vv.Aa. 2014):

- Antalgic effect,
- Myorelaxant action,
- Activation of microcirculation,
- Immunomodulation,
- Neuro-hormonal stimulation,
- Improvement of fat and carbohydrate metabolism.

In the same project, traditional uses of different types of natural mineral waters (each of them characterized by specific biochemical component/s) were collected, with general clinical indications formulated for various health conditions, as reported in Table 4 (Nappi 2001; Vv.Aa. 2014; Quattrini et al. 2016).

In general, treatments used in Medical Hydrology can be classified into three main categories on the basis of their route of administration (external or internal) and the state of matter of the therapeutic medium (liquid or gaseous):

1. Balneo-therapeutic treatments, such as baths with natural mineral waters and mud therapy.

2. Inhalation-based treatments, such as vapours or aerosols derived from natural mineral waters inhaled for medicinal purposes.

3. Hydropinic treatments, when natural mineral waters are taken orally as therapeutic drinks.

Balneo-therapeutic treatments are believed to exert their global therapeutic effect on the body thanks to a synergistic combination of mechanical (hydrostatic pressure), thermal (high temperature), and biochemical actions, the latter due to both the mineral (osmotic pressure and direct activity) and the organic (with anti-inflammatory and immunomodulatory properties) components of waters and muds (Vv.Aa. 2014; Fioravanti et al. 2017; Antonelli and Donelli 2018b). From a physiological point of view, balneo-therapeutic treatments can increase serum β -endorphins and can modulate cortisol levels in such a way as to improve individual stress resilience without disrupting circadian rhythms of this hormone (Fioravanti et al. 2011; Antonelli and Donelli 2018a). The long-term increase of cortisol awakening response due to a cycle of balneo-therapeutic treatments may be the reason why, in some clinical studies, the effects on health of balneotherapy, mud therapy, and spa therapy were observed to last for a few months after intervention (Forestier et al. 2016; Antonelli and Donelli 2018a). If we consider inflammatory mediators, mud applications followed by baths with natural mineral waters can reduce the production of prostaglandin E2 (PGE2), leukotriene B4 (LTB4), interleukin-1 β (IL-1 β), and tumor necrosis factor alpha (TNF- α) (Fioravanti et al. 2011). Quantitative data from the “gray” literature also show that, at least for osteoarthritis, therapeutic mud applications are superior to simple hot packs to improve joint functionality and pain, thus underscoring the therapeutic importance of the organic component (Crespin 2017). It is possible that the effect of balneo-therapeutic treatments on the inflammatory response and interleukin production is due to modifications of microRNA expressions induced by thermal and mechanical stimuli, as observed in a cohort of patients with osteoarthritis (Giannitti et al. 2017). At a joint level, balneo-therapeutic treatments may stimulate cartilage metabolism through mediators like the insulin-like growth factor-1 (IGF1), they may exert an antioxidant effect by reducing the release of Reactive Oxygen and Nitrogen Species (Fioravanti et al. 2011; Masselli et al. 2020) and also modulate intracellular mediators like protein kinases involved in cartilage growth and cell proliferation (Queirolo et al. 2016; Martini et al. 2018). Mud baths are also associated with a decrease in serum levels of adiponectin and resistin, hormonal substances probably implied in the progression of chronic degenerative diseases like osteoarthritis (Fioravanti et al. 2015). Furthermore, transcutaneous absorption of antiphlogistic substances released by water and mud microflora may contribute to the overall pharmacological effect of balneo-therapeutic treatments (Vv.Aa. 2014; Antonelli and Donelli 2018b). More details about possible mechanism of action of balneotherapy, as hypothesized on the basis of in-vitro laboratory studies, have recently been collected in a comprehensive literature review, demonstrating the anti-inflammatory, antioxidant, chondroprotective, and immunosuppressive role of this type of intervention at a cellular level (Cheleschi et al. 2020). Regarding balneotherapy and peripheral venous circulation, it is believed that the main therapeutic action of baths is determined by the hydrostatic and osmotic pressure of natural

mineral waters, which seems capable of reducing pain and oedema (de Moraes Silva et al. 2019). The combination of baths with hydrojet massage, Kneipp therapy (in which hot and cold baths are alternated), and physical exercise can determine a useful compression of peripheral veins, thanks to an external (water pressure) and internal (muscle contraction) synergistic action (de Moraes Silva et al. 2019). In particular, beneficial effects for the cardiovascular system seem to be more frequently associated with balneo-therapeutic sessions in carbon dioxide-rich water, which may be responsible for lowering peripheral vascular resistance and increasing blood flow in a more pronounced way if compared with other water types (Pagourelas et al. 2011). Additionally, balneo-therapeutic treatments, especially those ones based on sulphur-rich waters, seem to have anti-inflammatory, keratolytic, and regenerative effects on skin due to a direct pharmacological action of the mineral component and to interactions between thermal and cutaneous microflora with a potential modulation of local immune functions (Gobbi et al. 2009; Mirandola et al. 2011; Katz et al. 2012; Antonelli and Donelli 2018b; Eliasse et al. 2020).

Inhalation treatments with vaporized natural mineral waters can have anti-inflammatory, mucolytic, and antimicrobial properties, whereas irrigations with liquid-phase waters also have an action of mechanical washing on the upper respiratory tract (Vv.Aa. 2014; Keller et al. 2014; Casale et al. 2018). Among others, waters with sulphur seem to promote mucociliary clearance, regulate local immunity, inflammation, and have antiallergic effects (Rinaldi et al. 2006; Mirandola et al. 2007, 2013; Keller et al. 2014; Viegas et al. 2019; Carubbi et al. 2019). For all these reasons, sulphur-rich water inhalations have been proposed as an integrative treatment for patients with chronic obstructive pulmonary disease (Khaltaev et al. 2020).

Health effects of hydropinic treatments are mainly mediated by the intake of water minerals (osmotic and prokinetic action on the intestine when the mineral content is high, diuretic effect when the mineral content is low) and by the interaction between water and gut microflora (Vv.Aa. 2014). Globally, the mechanisms of action of spa-related treatments, although not fully understood, seem to be determined by a synergistic action of all water and mud components, capable of eliciting beneficial effects both locally and at a systemic level.

Limitations of this study and new perspectives for future research

First, a major limitation of both primary (clinical) and secondary (review) studies on the topic, which is responsible for hindering optimal retrieval and dissemination of scientific information, is the lack of a widely accepted consensus on a precise English terminology in the field of Medical Hydrology. In other words, terms like balneotherapy, hydrotherapy, and spa therapy are given diverse (and sometimes misleading or confusing) meanings when used by different authors, and this aspect has already been underscored even by several experts (Gutenbrunner et al. 2010; Fioravanti et al. 2017). For this reason, a list of specific definitions was provided in the Introduction section to avoid possible misunderstandings and to address this issue as best as possible.

Even if the number of reviews included in this research work is considerable, the actual basis of clinical evidence in support of spa-related treatments is quite limited. In fact, there is a demonstrated substantial degree of clinical evidence overlap among analyzed reviews (see the Electronic Supplementary Materials for an example about balneotherapy for fibromyalgia or refer to a recently published overview of literature reviews for another example about spa therapy for knee osteoarthritis (D'Angelo et al. 2021)). For this reason, it is important to keep on studying Medical Hydrology with further clinical research projects to expand the existing evidence basis.

With regard to the quality of clinical trials analyzed in included reviews, the most frequent limitations were reported to be the low number of study participants, poor information about proper randomization, and the lack of adequate control. Instead, if we consider the quality of included reviews, major limitations involved methodological issues like excessively narrow search strategies (with the literature search sometimes only run in PubMed), poor description of the details about article selection and quality assessment, and an often inadequate evaluation of the risk of publication bias. The trial quality assessment was not adequately used by some review authors for informing a critical discussion, with potentially inflated results and biased conclusions. The high level of heterogeneity across primary and secondary studies on the topic makes it difficult to synthesize available data, especially from a quantitative point of view, thus urging the necessity to conduct further investigations adopting a more homogeneous design.

As a proposal, it would be useful to promote a wider and better application of the PRISMA guidelines among researchers who want to conduct systematic reviews about Medical Hydrology, since, if we consider included research works, only some authors actually followed these internationally-accepted methodological recommendations in a thorough way (Moher et al. 2009). Poor compliance with the PRISMA guidelines can result in omitting essential information which is useful for clinicians and policymakers to translate research findings into practice (external validity), or, even worse, this lack of transparency can hide substantial flaws in the conduction of the review, thus undermining the conclusion reliability (internal validity). For this reason, our umbrella review was conducted in accordance with the PRISMA statement, and methodological quality of included systematic reviews was weighted as best as possible.

Moreover, authors of some included reviews underscored the difficulty to find a specifically tailored assessment tool to evaluate the quality of primary studies on the topic, which regard atypical, non-pharmacological and hard-to-blind interventions like balneotherapy. Different solutions were adopted by review authors to find an adequate solutions, ranging from the use of a modified version of the standard Cochrane tool (Higgins et al. 2011) to other tools often employed for the evaluation of public health interventions like the Canadian “Quality Assessment Tool for Quantitative Studies” (Armijo-Olivo et al. 2012). To properly address this issue, world experts should agree on a set of essential domains that any assessment tool should include to be adequately usable for evaluating the quality and risk of bias of Medical Hydrology-related trials. In fact, some efforts have already been made

towards this direction, for example by developing the “SPAC” checklist (Kamioka et al. 2013) or by evaluating the biasing impact of unblinded balneotherapy (Verhagen et al. 1998). Finally, in order to optimize the collection of available evidence on the topic, any effort was made to retrieve all relevant data reported both in the scientific and in the so called “gray” literature (Petticrew et al. 2008), including findings only displayed in conference proceedings and lectures published in peer-reviewed journals (Roques 2014; Bender 2016; Cao et al. 2020). However, the risk of publication bias, although minimized, could not be fully excluded, and it was not possible to statistically assess it because no quantitative synthesis was feasible.

Conclusions

Globally, retrieved evidence suggests that Medical Hydrology can be useful for individual wellbeing and public health. In particular, higher-quality studies support the use of spa-related interventions for conditions like osteoarthritis, fibromyalgia, and chronic back pain, but also for chronic venous insufficiency. However, as discussed above, the existing body of evidence has some relevant limitations, especially with regard to non-rheumatic diseases. For this reason, further high-quality clinical trials and observational studies should be designed to confirm the beneficial effects of Medical Hydrology and to thoroughly estimate its effectiveness outside experimental settings in real-life clinical practice.

References

- Albertini MC, Dacha M, Teodori L, Conti ME (2007) Drinking mineral waters: biochemical effects and health implications the state-of-the-art. *International Journal of Environment and Health* 1:153
- Al Shammeri O, AlHamdan N, Al-Hothaly B, et al (2014) Chronic Venous Insufficiency: prevalence and effect of compression stockings. *Int J Health Sci* 8:231–236. <https://doi.org/10.12816/0023975>
- An J, Lee I, Yi Y (2019) The Thermal Effects of Water Immersion on Health Outcomes: An Integrative Review. *Int J Environ Res Public Health* 16.: <https://doi.org/10.3390/ijerph16071280>
- Antonelli M, Donelli D (2018a) Effects of balneotherapy and spa therapy on levels of cortisol as a stress biomarker: a systematic review. *Int J Biometeorol* 62:913–924. <https://doi.org/10.1007/s00484-018-1504-8>
- Antonelli M, Donelli D (2018b) Mud therapy and skin microbiome: a review. *Int J Biometeorol* 62:2037–2044. <https://doi.org/10.1007/s00484-018-1599-y>
- Antonelli M, Donelli D, Fioravanti A (2018) Effects of balneotherapy and spa therapy on quality of life of patients with knee osteoarthritis: a systematic review and meta-analysis. *Rheumatol Int* 38:1807–1824. <https://doi.org/10.1007/s00296-018-4081-6>
- Armijo-Olivo S, Stiles CR, Hagen NA, et al (2012) Assessment of study quality for systematic

- 659 reviews: a comparison of the Cochrane Collaboration Risk of Bias Tool and the Effective Public
660 Health Practice Project Quality Assessment Tool: methodological research. *J Eval Clin Pract*
661 18:12–18. <https://doi.org/10.1111/j.1365-2753.2010.01516.x>
- 662 Baethge C, Goldbeck-Wood S, Mertens S (2019) SANRA—a scale for the quality assessment of
663 narrative review articles. *Research Integrity and Peer Review* 4
- 664 Bai R, Li C, Xiao Y, et al (2019) Effectiveness of spa therapy for patients with chronic low back pain:
665 An updated systematic review and meta-analysis. *Medicine* 98:e17092.
666 <https://doi.org/10.1097/MD.00000000000017092>
- 667 Beer A, Kleinschmidt J, Jagenburg L (2018) Zur Wirksamkeit der kurörtlichen Heiltorftherapie: eine
668 Literaturübersicht aus neueren Veröffentlichungen. *Phys Med Rehab Kuror* 28:365–371.
669 <https://doi.org/10.1055/a-0713-0694>
- 670 Bender T (2016) SP0165 Balneotherapy: Efficacy and Safety in Osteoarthritis. *Ann Rheum Dis*
671 75:40.4–40. <https://doi.org/10.1136/annrheumdis-2016-eular.6267>
- 672 Bender T, Bálint G, Prohászka Z, et al (2014) Evidence-based hydro- and balneotherapy in
673 Hungary—a systematic review and meta-analysis. *International Journal of Biometeorology*
674 58:311–323
- 675 Bohmer H, Müller H, Resch KL (2000) Calcium supplementation with calcium-rich mineral waters: a
676 systematic review and meta-analysis of its bioavailability. *Osteoporos Int* 11:938–943.
677 <https://doi.org/10.1007/s001980070032>
- 678 Bramer WM, Rethlefsen ML, Kleijnen J, Franco OH (2017) Optimal database combinations for
679 literature searches in systematic reviews: a prospective exploratory study. *Syst Rev* 6:245.
680 <https://doi.org/10.1186/s13643-017-0644-y>
- 681 Brosseau L, MacLeay L, Robinson V, et al (2002) Efficacy of balneotherapy for osteoarthritis of the
682 knee: A systematic review. *Phys Ther Rev* 7:209–222.
683 <https://doi.org/10.1179/108331902235001976>
- 684 Cacciapuoti S, Luciano MA, Megna M, et al (2020) The Role of Thermal Water in Chronic Skin
685 Diseases Management: A Review of the Literature. *J Clin Med Res* 9.:
686 <https://doi.org/10.3390/jcm9093047>
- 687 Cao C-F, Ma K-L, Li Q-L, et al (2020) Benefits of balneotherapy in the management of fibromyalgia
688 syndrome: a systematic review and meta-analysis of randomized controlled trials. *Pain Practice*
- 689 Carretero MI (2002) Clay minerals and their beneficial effects upon human health. A review. *Appl*
690 *Clay Sci* 21:155–163. [https://doi.org/10.1016/S0169-1317\(01\)00085-0](https://doi.org/10.1016/S0169-1317(01)00085-0)
- 691 Carubbi C, Masselli E, Calabrò E, et al (2019) Sulphurous thermal water inhalation impacts
692 respiratory metabolic parameters in heavy smokers. *Int J Biometeorol* 63:1209–1216.
693 <https://doi.org/10.1007/s00484-019-01737-7>
- 694 Casale M, Moffa A, Cassano M, et al (2018) Saline nasal irrigations for chronic rhinosinusitis: From
695 everyday practice to evidence-based medicine. An update. *Int J Immunopathol Pharmacol*
696 32:2058738418802676. <https://doi.org/10.1177/2058738418802676>
- 697 Cheleschi S, Gallo I, Tenti S (2020) A comprehensive analysis to understand the mechanism of action
698 of balneotherapy: why, how, and where they can be used? Evidence from in vitro studies
699 performed on human and animal samples. *Int J Biometeorol* 64:1247–1261.
700 <https://doi.org/10.1007/s00484-020-01890-4>

701 Corvillo I, Armijo F, Álvarez-Badillo A, et al (2019) Efficacy of aquatic therapy for neck pain: a
702 systematic review. *Int J Biometeorol*. <https://doi.org/10.1007/s00484-019-01738-6>

703 Costantino M, Lampa E, Nappi G (2006) Effectiveness of sulphur spa therapy with politzer in the
704 treatment of rhinogenic deafness. *Acta Otorhinolaryngol Ital* 26:7–13

705 Crespin M (2017) The effects of mud pack therapy versus hot pack treatment to improve functionality
706 and decrease pain in older adults suffering with knee osteoarthritis. PhD, College of Health and
707 Human Services California State University

708 D'Angelo D, Coclite D, Napoletano A, et al (2021) The efficacy of balneotherapy, mud therapy and
709 spa therapy in patients with osteoarthritis: an overview of reviews. *Int J Biometeorol*.
710 <https://doi.org/10.1007/s00484-021-02102-3>

711 de Moraes Silva MA, Nakano LC, Cisneros LL, Miranda F Jr (2019) Balneotherapy for chronic
712 venous insufficiency. *Cochrane Database Syst Rev* 8:CD013085.
713 <https://doi.org/10.1002/14651858.CD013085.pub2>

714 Drobnik J, Stebel A (2020) Central European ethnomedical and officinal uses of peat, with special
715 emphasis on the Tolpa peat preparation (TPP): An historical review. *J Ethnopharmacol*
716 246:112248. <https://doi.org/10.1016/j.jep.2019.112248>

717 Eliasse Y, Redoules D, Espinosa E (2020) Impact of Avène Thermal Spring Water on immune cells. *J*
718 *Eur Acad Dermatol Venereol* 34 Suppl 5:21–26. <https://doi.org/10.1111/jdv.16335>

719 Espejo-Antúnez L, Cardero-Durán MA, Garrido-Ardila EM, et al (2013) Clinical effectiveness of
720 mud pack therapy in knee osteoarthritis. *Rheumatology* 52:659–668.
721 <https://doi.org/10.1093/rheumatology/kes322>

722 Falagas ME, Zarkadoulia E, Rafailidis PI (2009) The therapeutic effect of balneotherapy: evaluation
723 of the evidence from randomised controlled trials. *Int J Clin Pract* 63:1068–1084.
724 <https://doi.org/10.1111/j.1742-1241.2009.02062.x>

725 Ferrante G, Baldissera S, Campostrini S (2017) Epidemiology of chronic respiratory diseases and
726 associated factors in the adult Italian population. *Eur J Public Health* 27:1110–1116.
727 <https://doi.org/10.1093/eurpub/ckx109>

728 Fioravanti A, Cantarini L, Guidelli GM, Galeazzi M (2011) Mechanisms of action of spa therapies in
729 rheumatic diseases: what scientific evidence is there? *Rheumatology International* 31:1–8

730 Fioravanti A, Giannitti C, Cheleschi S, et al (2015) Circulating levels of adiponectin, resistin, and
731 visfatin after mud-bath therapy in patients with bilateral knee osteoarthritis. *International Journal*
732 *of Biometeorology* 59:1691–1700

733 Fioravanti A, Karagülle M, Bender T, Karagülle MZ (2017) Balneotherapy in osteoarthritis: Facts,
734 fiction and gaps in knowledge. *Eur J Integr Med* 9:148–150.
735 <https://doi.org/10.1016/j.eujim.2017.01.001>

736 Forestier R, Erol Forestier FB, Francon A (2016) Spa therapy and knee osteoarthritis: A systematic
737 review. *Ann Phys Rehabil Med* 59:216–226. <https://doi.org/10.1016/j.rehab.2016.01.010>

738 Forestier R, Françon A (2008) Crenobalneotherapy for limb osteoarthritis: systematic literature
739 review and methodological analysis. *Joint Bone Spine* 75:138–148.
740 <https://doi.org/10.1016/j.jbspin.2007.06.009>

741 Forestier RJ, Erol Forestier FB, Francon A (2017) Crénothérapie (crénobalnéothérapie) dans la
742 lombalgie chronique : une revue critique. *Revue du Rhumatisme Monographies* 84:91–98.

743 <https://doi.org/10.1016/j.monrhu.2016.11.008>

744 Fortunati NA, Fioravanti A, Seri G, et al (2016) May spa therapy be a valid opportunity to treat hand
 745 osteoarthritis? A review of clinical trials and mechanisms of action. *Int J Biometeorol* 60:1–8.
 746 <https://doi.org/10.1007/s00484-015-1030-x>

747 Fraioli A, Grassi M, Mennuni G, et al (2013) Clinical researches on the efficacy of spa therapy in
 748 fibromyalgia. A systematic review. *Ann Ist Super Sanita* 49:219–229.
 749 https://doi.org/10.4415/ANN_13_02_13

750 Fraioli A, Mennuni G, Fontana M, et al (2018) Efficacy of Spa Therapy, Mud-Pack Therapy,
 751 Balneotherapy, and Mud-Bath Therapy in the Management of Knee Osteoarthritis. A Systematic
 752 Review. *Biomed Res Int* 2018:1042576. <https://doi.org/10.1155/2018/1042576>

753 Françon A, Duplan B, Forestier FBE, Forestier R (2015) Thermalisme et douleur chronique en
 754 rhumatologie : revue des essais cliniques randomisés (ECR) et des méta-analyses. *Douleur analg*
 755 28:47–53. <https://doi.org/10.1007/s11724-015-0410-7>

756 Fusar-Poli P, Radua J (2018) Ten simple rules for conducting umbrella reviews. *Evid Based Ment*
 757 *Health* 21:95–100. <https://doi.org/10.1136/ebmental-2018-300014>

758 Giannitti C, De Palma A, Pascarelli NA, et al (2017) Can balneotherapy modify microRNA
 759 expression levels in osteoarthritis? A comparative study in patients with knee osteoarthritis. *Int J*
 760 *Biometeorol* 61:2153–2158. <https://doi.org/10.1007/s00484-017-1420-3>

761 Gobbi G, Ricci F, Malinverno C, et al (2009) Hydrogen sulfide impairs keratinocyte cell growth and
 762 adhesion inhibiting mitogen-activated protein kinase signaling. *Lab Invest* 89:994–1006.
 763 <https://doi.org/10.1038/labinvest.2009.61>

764 Gomes C, Carretero MI, Pozo M, et al (2013) Peloids and pelotherapy: Historical evolution,
 765 classification and glossary. *Appl Clay Sci* 75-76:28–38.
 766 <https://doi.org/10.1016/j.clay.2013.02.008>

767 Gravelier C, Kanny G, Adetu S, Goffinet L (2020) Spa therapy and burn scar treatment: a systematic
 768 review of the literature. *Int J Biometeorol* 64:2195–2203. [https://doi.org/10.1007/s00484-020-](https://doi.org/10.1007/s00484-020-01988-9)
 769 [01988-9](https://doi.org/10.1007/s00484-020-01988-9)

770 Guidelli GM, Tenti S, De Nobili E, Fioravanti A (2012) Fibromyalgia syndrome and spa therapy:
 771 myth or reality? *Clin Med Insights Arthritis Musculoskelet Disord* 5:19–26.
 772 <https://doi.org/10.4137/CMAMD.S8797>

773 Gutenbrunner C, Bender T, Cantista P, Karagülle Z (2010) A proposal for a worldwide definition of
 774 health resort medicine, balneology, medical hydrology and climatology. *Int J Biometeorol*
 775 54:495–507. <https://doi.org/10.1007/s00484-010-0321-5>

776 Harzy T, Ghani N, Akasbi N, et al (2009) Short- and long-term therapeutic effects of thermal mineral
 777 waters in knee osteoarthritis: a systematic review of randomized controlled trials. *Clin*
 778 *Rheumatol* 28:501–507. <https://doi.org/10.1007/s10067-009-1114-2>

779 Higgins JPT, Altman DG, Gøtzsche PC, et al (2011) The Cochrane Collaboration’s tool for assessing
 780 risk of bias in randomised trials. *BMJ* 343:d5928. <https://doi.org/10.1136/bmj.d5928>

781 Hou C, Liang L, Chu X, et al (2020) The short-term efficacy of mud therapy for knee osteoarthritis: A
 782 meta-analysis. *Medicine* 99:e19761. <https://doi.org/10.1097/MD.00000000000019761>

783 Kamioka H, Kawamura Y, Tsutani K, et al (2013) A checklist to assess the quality of reports on spa
 784 therapy and balneotherapy trials was developed using the Delphi consensus method: the SPAC

785 checklist. *Complement Ther Med* 21:324–332. <https://doi.org/10.1016/j.ctim.2013.05.002>

786 Karagülle M, Karagülle MZ (2015) Effectiveness of balneotherapy and spa therapy for the treatment
787 of chronic low back pain: a review on latest evidence. *Clin Rheumatol* 34:207–214.
788 <https://doi.org/10.1007/s10067-014-2845-2>

789 Karagülle MZ, Karagülle M (2004) [Balneotherapy and spa therapy of rheumatic diseases in Turkey:
790 a systematic review]. *Forsch Komplementarmed Klass Naturheilkd* 11:33–41.
791 <https://doi.org/10.1159/000077194>

792 Katz U, Shoenfeld Y, Zakin V, et al (2012) Scientific evidence of the therapeutic effects of dead sea
793 treatments: a systematic review. *Semin Arthritis Rheum* 42:186–200.
794 <https://doi.org/10.1016/j.semarthrit.2012.02.006>

795 Keller S, König V, Mösges R (2014) Thermal water applications in the treatment of upper respiratory
796 tract diseases: a systematic review and meta-analysis. *J Allergy* 2014:943824.
797 <https://doi.org/10.1155/2014/943824>

798 Khalilzadeh S, Shirbeigi L, Naghizadeh A, et al (2019) Use of mineral waters in the treatment of
799 psoriasis: Perspectives of Persian and conventional medicine. *Dermatol Ther* 32:e12969.
800 <https://doi.org/10.1111/dth.12969>

801 Khaltayev N, Solimene U, Vitale F, Zanasi A (2020) Balneotherapy and hydrotherapy in chronic
802 respiratory disease. *J Thorac Dis* 12:4459–4468. <https://doi.org/10.21037/jtd-gard-2019-009>

803 Langhorst J, Musial F, Klose P, Häuser W (2009) Efficacy of hydrotherapy in fibromyalgia
804 syndrome--a meta-analysis of randomized controlled clinical trials. *Rheumatology* 48:1155–
805 1159. <https://doi.org/10.1093/rheumatology/kep182>

806 Liu H, Zeng C, Gao S-G, et al (2013) The effect of mud therapy on pain relief in patients with knee
807 osteoarthritis: a meta-analysis of randomized controlled trials. *J Int Med Res* 41:1418–1425.
808 <https://doi.org/10.1177/0300060513488509>

809 Maraver F, Karagülle MZ (2012) *Medical Hydrology and Balneology: Environmental Aspects*.
810 *Publicaciones Universidad Complutense de Madrid*

811 Maraver F, Michán A, Morer C, Aguilera L (2011) Is thalassotherapy simply a type of
812 climatotherapy? *Int. J. Biometeorol.* 55:107–108

813 Martini S, Soliman T, Gobbi G, et al (2018) PKCε Controls Mitotic Progression by Regulating
814 Centrosome Migration and Mitotic Spindle Assembly. *Mol Cancer Res* 16:3–15.
815 <https://doi.org/10.1158/1541-7786.MCR-17-0244>

816 Masselli E, Pozzi G, Vaccarezza M, et al (2020) ROS in Platelet Biology: Functional Aspects and
817 Methodological Insights. *Int J Mol Sci* 21.: <https://doi.org/10.3390/ijms21144866>

818 Matsumoto H, Hagino H, Hayashi K, et al (2017) The effect of balneotherapy on pain relief, stiffness,
819 and physical function in patients with osteoarthritis of the knee: a meta-analysis. *Clin Rheumatol*
820 36:1839–1847. <https://doi.org/10.1007/s10067-017-3592-y>

821 Ministero della Sanità (1995) Modificazioni all’elenco delle patologie che possono trovare reale
822 beneficio dalle cure termali e proroga della sua validità. *Gazzetta Ufficiale*

823 Mirandola P, Gobbi G, Malinverno C, et al (2013) Impact of sulphurous water politzer inhalation on
824 audiometric parameters in children with otitis media with effusion. *Clin Exp Otorhinolaryngol*
825 6:7–11. <https://doi.org/10.3342/ceo.2013.6.1.7>

826 Mirandola P, Gobbi G, Micheloni C, et al (2011) Hydrogen sulfide inhibits IL-8 expression in human
827 keratinocytes via MAP kinase signaling. *Lab Invest* 91:1188–1194.
828 <https://doi.org/10.1038/labinvest.2011.76>

829 Mirandola P, Gobbi G, Sponzilli I, et al (2007) Exogenous hydrogen sulfide induces functional
830 inhibition and cell death of cytotoxic lymphocytes subsets. *J Cell Physiol* 213:826–833.
831 <https://doi.org/10.1002/jcp.21151>

832 Moher D, Liberati A, Tetzlaff J, et al (2009) Preferred reporting items for systematic reviews and
833 meta-analyses: the PRISMA statement. *PLoS Med* 6:e1000097.
834 <https://doi.org/10.1371/journal.pmed.1000097>

835 Morer C, Roques C-F, Françon A, et al (2017) The role of mineral elements and other chemical
836 compounds used in balneology: data from double-blind randomized clinical trials. *Int J*
837 *Biometeorol* 61:2159–2173. <https://doi.org/10.1007/s00484-017-1421-2>

838 Nappi G (2001) *Medicina e clinica termale*

839 National Institutes of Health Study Quality Assessment Tools. [https://www.nhlbi.nih.gov/health-](https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools)
840 [topics/study-quality-assessment-tools](https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools). Accessed 2 Apr 2021

841 Naumann J, Biehler D, Lüty T, Sadaghiani C (2017) Prevention and Therapy of Type 2 Diabetes-
842 What Is the Potential of Daily Water Intake and Its Mineral Nutrients? *Nutrients* 9.:
843 <https://doi.org/10.3390/nu9080914>

844 Naumann J, Sadaghiani C (2014) Therapeutic benefit of balneotherapy and hydrotherapy in the
845 management of fibromyalgia syndrome: a qualitative systematic review and meta-analysis of
846 randomized controlled trials. *Arthritis Res Ther* 16:R141. <https://doi.org/10.1186/ar4603>

847 Pagourelas ED, Zorou PG, Tsaligopoulos M, et al (2011) Carbon dioxide balneotherapy and
848 cardiovascular disease. *Int J Biometeorol* 55:657–663. [https://doi.org/10.1007/s00484-010-0380-](https://doi.org/10.1007/s00484-010-0380-7)
849 7

850 Passali D, Gabelli G, Passali GC, et al (2017) Radon-enriched hot spring water therapy for upper and
851 lower respiratory tract inflammation. *Otolaryngol Pol* 71:8–13.
852 <https://doi.org/10.5604/01.3001.0010.2242>

853 Petticrew M, Egan M, Thomson H, et al (2008) Publication bias in qualitative research: what becomes
854 of qualitative research presented at conferences? *J Epidemiol Community Health* 62:552–554.
855 <https://doi.org/10.1136/jech.2006.059394>

856 Pittler MH, Karagülle MZ, Karagülle M, Ernst E (2006) Spa therapy and balneotherapy for treating
857 low back pain: meta-analysis of randomized trials. *Rheumatology* 45:880–884.
858 <https://doi.org/10.1093/rheumatology/kei018>

859 Quattrini S, Pampaloni B, Brandi ML (2016) Natural mineral waters: chemical characteristics and
860 health effects. *Clin Cases Miner Bone Metab* 13:173–180.
861 <https://doi.org/10.11138/ccmbm/2016.13.3.173>

862 Queirolo V, Galli D, Masselli E, et al (2016) PKC ϵ is a regulator of hypertrophic differentiation of
863 chondrocytes in osteoarthritis. *Osteoarthritis Cartilage* 24:1451–1460.
864 <https://doi.org/10.1016/j.joca.2016.04.003>

865 Queiroz LP (2013) Worldwide epidemiology of fibromyalgia. *Curr Pain Headache Rep* 17:356.
866 <https://doi.org/10.1007/s11916-013-0356-5>

867 Rachakonda TD, Schupp CW, Armstrong AW (2014) Psoriasis prevalence among adults in the United

- 868 States. *J Am Acad Dermatol* 70:512–516. <https://doi.org/10.1016/j.jaad.2013.11.013>
- 869 Raza HMA, Krutulyte G, Rimdeikiene I, Savickas R (2020) Efficacy of Balneotherapy and Mud
870 Therapy in Patients with Knee Osteoarthritis: A Systematic Literature Review. *Aktuelle*
871 *Rheumatologie*
- 872 Rinaldi L, Gobbi G, Pambianco M, et al (2006) Hydrogen sulfide prevents apoptosis of human PMN
873 via inhibition of p38 and caspase 3. *Lab Invest* 86:391–397.
874 <https://doi.org/10.1038/labinvest.3700391>
- 875 Roques C, -C. Mancret R, Tabone W (2012) Beneficial effect of balneotherapy: Achievements of the
876 French association of balneotherapy research and methodological difficulties. *Annals of Physical*
877 *and Rehabilitation Medicine* 55:e351
- 878 Roques CF (2014) Balneotherapy actual medical benefit. Data of evidence for the last twenty years.
879 *Ann Phys Rehabil Med* 57:e159. <https://doi.org/10.1016/j.rehab.2014.03.576>
- 880 Santos I, Cantista P, Vasconcelos C (2016) Balneotherapy in rheumatoid arthritis-a systematic review.
881 *Int J Biometeorol* 60:1287–1301. <https://doi.org/10.1007/s00484-015-1108-5>
- 882 Schuh A (2009) Die Evidenz der Klima- und Thalassotherapie. Ein Review. *Schweizerische*
883 *Zeitschrift für Ganzheitsmedizin / Swiss Journal of Integrative Medicine* 21:96–104
- 884 Shea BJ, Reeves BC, Wells G, et al (2017) AMSTAR 2: a critical appraisal tool for systematic
885 reviews that include randomised or non-randomised studies of healthcare interventions, or both.
886 *BMJ* 358:j4008. <https://doi.org/10.1136/bmj.j4008>
- 887 Stanhope J, Weinstein P, Cook A (2018) Do natural spring waters in Australia and New Zealand
888 affect health? A systematic review. *Journal of Water and Health* 16:1–13
- 889 Stier-Jarmer M, Kus S, Frisch D, et al (2015) Health resort medicine in non-musculoskeletal
890 disorders: is there evidence of its effectiveness? *Int J Biometeorol* 59:1523–1544.
891 <https://doi.org/10.1007/s00484-015-0953-6>
- 892 Sulaiman SK, Enakshee J, Traxer O, Somani BK (2020) Which Type of Water Is Recommended for
893 Patients with Stone Disease (Hard or Soft Water, Tap or Bottled Water): Evidence from a
894 Systematic Review over the Last 3 Decades. *Curr Urol Rep* 21:6. <https://doi.org/10.1007/s11934-020-0968-3>
- 896 Tenti S, Cheleschi S, Galeazzi M, Fioravanti A (2015) Spa therapy: can be a valid option for treating
897 knee osteoarthritis? *Int J Biometeorol* 59:1133–1143. <https://doi.org/10.1007/s00484-014-0913-6>
- 898 The Cochrane Group About Cochrane Reviews. <https://www.cochranelibrary.com/about/about-cochrane-reviews>. Accessed 25 Dec 2020
- 900 The European Parliament and the Council of the European Union (2009) Directive 2009/54/EC of the
901 European Parliament and of the Council of 18 June 2009 on the exploitation and marketing of
902 natural mineral waters. European Union law
- 903 van Tubergen A, van der Linden S (2002) A brief history of spa therapy. *Ann Rheum Dis* 61:273–
904 275. <https://doi.org/10.1136/ard.61.3.273>
- 905 Verhagen AP, Bierma-Zeinstra SMA, Boers M, et al (2007) Balneotherapy for osteoarthritis.
906 *Cochrane Database Syst Rev* CD006864. <https://doi.org/10.1002/14651858.CD006864>
- 907 Verhagen AP, Bierma-Zeinstra SMA, Boers M, et al (2015) Balneotherapy (or spa therapy) for
908 rheumatoid arthritis. *Cochrane Database Syst Rev* CD000518.

909 <https://doi.org/10.1002/14651858.CD000518.pub2>

910 Verhagen AP, de Vet HCW, de Bie RA, et al (1998) Balneotherapy and Quality Assessment:
 911 Interobserver Reliability of the Maastricht Criteria List and the Need for Blinded Quality
 912 Assessment. *J Clin Epidemiol* 51:335–341. [https://doi.org/10.1016/S0895-4356\(97\)00297-7](https://doi.org/10.1016/S0895-4356(97)00297-7)

913 Viegas J, Esteves AF, Cardoso EM, et al (2019) Biological Effects of Thermal Water-Associated
 914 Hydrogen Sulfide on Human Airways and Associated Immune Cells: Implications for
 915 Respiratory Diseases. *Front Public Health* 7:128. <https://doi.org/10.3389/fpubh.2019.00128>

916 Vina ER, Kwoh CK (2018) Epidemiology of osteoarthritis: literature update. *Curr Opin Rheumatol*
 917 30:160–167. <https://doi.org/10.1097/BOR.0000000000000479>

918 Vv.Aa. (2014) Hydroglobe. Definizione di un quadro di riferimento complessivo mondiale per
 919 l'idroterapia. Narcissus.me

920 Xiang J, Wu D, Li J 'an (2016) Clinical Efficacy of Mudpack Therapy in Treating Knee
 921 Osteoarthritis: A Meta-Analysis of Randomized Controlled Studies. *Am J Phys Med Rehabil*
 922 95:121–131. <https://doi.org/10.1097/PHM.0000000000000354>

923 Yuan D, Yu Z-X, Wang W, Chen Y (2019) Head-out immersion in natural thermal mineral water for
 924 the management of hypertension: a review of randomized controlled trials. *Int J Biometeorol*
 925 63:1707–1718. <https://doi.org/10.1007/s00484-019-01780-4>

926 Zhen-han D, Tuo Y, Hui L, et al (2014) A meta-analysis of mud therapy on knee osteoarthritis pain.
 927 *Journal of Clinical Rehabilitative Tissue Engineering Research* 2455–2460

928 洋晴上岡, 律雄黒柳, 泰喜小松, et al (2006) 温泉の治療と健康増進の効果に関する. *日温気物医誌* 69:155–166.
 929 <https://doi.org/10.11390/onki1962.69.155>




Figure 1. Flow-chart describing the study selection process.

The flow-chart was adapted from the model recommended by the PRISMA guidelines (Moher et al. 2009).




Figure 2. Number of included reviews for each health condition.

● Balneotherapy
 ● Mud therapy
 ● Spa therapy
 ● Hydroponic therapy
● Inhalation/irrigation therapy
● Any mineral water/mud-based therapy




Figure 3. Number of included reviews for each type of therapy.

TABLE 1. Main characteristics (PICOS, methodological quality and study authors' conclusions) of included systematic reviews.

Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
An et al. 2019 (An et al. 2019)	Both healthy and diseased subjects (212)	BT	Any type	Various (physiological and clinical)	SR (13 clinical studies of any type)	D	?	Demonstration of various cardiovascular and neuromuscular effects, but uncertain conclusions.
Brosseau et al. 2013 (Brosseau et al. 2002)	Patients with knee OA (172)	BT	Any type	Pain and functionality	SR (3 RCTs)	C	2	Only short-term effects were demonstrated, with significant benefits in terms of pain relief exclusively shown for baths in Dead Sea or hot sulphur water.
Cao et al. 2020 (Cao et al. 2020)	Patients with fibromyalgia (611)	BT	Any type	Pain, QoL and mood	SR + MA (10 RCTs) - CP	C	3	Pooled evidence from analyzed RCTs (follow-up: from 12 to 48 weeks) indicates that BT may reduce pain and improve QoL of patients with fibromyalgia.
Corvillo et al. 2019 (Corvillo et al. 2019)	Patients with neck pain of different origins (658)	BT	No treatment or standard rehabilitation	Pain, functionality, QoL and mood	SR (13 clinical studies of any type)	D	2	Beneficial effects for all studied algofunctional outcomes, and for psychophysical wellbeing.
Falagas et al. 2009 (Falagas et al. 2009)	Patients with any disease (1720 subjects with rheumatic conditions)	BT	Any type	Any relevant clinical outcome	SR (29 RCTs)	C	2	Clinical improvement of several rheumatic diseases (OA, fibromyalgia, ankylosing spondylitis, rheumatoid arthritis). Preliminary evidence of benefits for psoriasis and Parkinson's disease.
Harzy et al. 2009 (Harzy et al. 2009)	Patients with knee OA (493)	BT	Any type	Pain, functionality, drugs	SR (9 RCTs)	B	2	Beneficial effects on pain and joint functionality lasting up to 24 weeks from intervention.
Moraes Silva et al. 2019 (de Moraes Silva et al. 2019)	Patients with chronic venous insufficiency in leg veins (891)	BT	Any type	Any relevant clinical and QoL outcome	SR+MA (7 RCTs)	A	3	Low-to-moderate evidence indicates beneficial effects in terms of pain, QoL, and skin pigmentation due to chronic venous insufficiency.
Schuh 2009 (Schuh 2009)	Patients with any disease (?)	BT (TT)	Any type	Any relevant clinical outcome	SR (12 studies related to marine climatotherapy and TT)	D	?	A combination of thalassotherapy with marine climatotherapy can exert positive effects on psoriasis, atopic dermatitis, and bronchial asthma.
Yuan et al. (2019) (Yuan et al. 2019)	Patients with arterial hypertension (1122)	BT	Any type	Variations of blood pressure	SR (12 RCTs)	C	3	No worsening of blood pressure parameters.
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Beer et al. 2018 (Beer et al. 2018)	Patients with any disease (728)	MT	Any type	Any relevant clinical outcome	SR (35 clinical studies)	D	2	High-quality evidence indicates a beneficial effect for the symptomatic treatment of OA and fibromyalgia.
Espejo-Antunez et al. 2013 (Espejo-Antunez et al. 2013)	Patients with knee OA (2102)	MT	Any type	Symptoms and QoL	SR (20 studies of any type)	D	2	Significant improvement of pain, general symptoms and QoL.
Hou et al. 2020 (Hou et al. 2020)	Patients with knee OA (1106)	MT	Any type	Pain and functionality	SR + MA (11 RCTs)	B	3	Significant amelioration of knee pain and function.
Liu et al. 2013 (Liu et al. 2013)	Patients with knee OA (410)	MT	Any type	Pain and functionality	SR + MA (7 RCTs)	D	2	Significant improvement of OA-related pain.
Xiang et al. 2016 (Xiang et al. 2016)	Patients with knee OA (1010)	MT	Any type	Functionality	SR + MA (10 RCTs)	C	2	No significant improvement in joint functionality.
Zhen-han et al. 2014	Patients with knee OA (410)	MT	Any type	Pain	SR + MA (7 RCTs) - abstract	?	?	Mud therapy can significantly attenuate knee osteoarthritis pain.

(Zhen-han et al. 2014)					only			
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Antonelli & Donelli 2018 (Antonelli and Donelli 2018a)	Both healthy and diseased subjects (684)	SPA-T	Any type	Variations of salivary or serum cortisol levels	SR (15 clinical studies of any type)	C	2	The effect on cortisol levels suggests that intervention can have an anti-stress action and improve stress resilience.
Antonelli et al. 2018 (Antonelli et al. 2018)	Patients with knee OA (1599)	SPA-T	Any type	QoL, algofunctional indices, drugs	SR+MA (17 RCTs)	C	3	Significant improvement of QoL. Beneficial effects on algofunctional indices and painkiller intake.
Bai et al. 2019 (Bai et al. 2019)	Patients with chronic LBP (1038)	SPA-T	Any type	Back pain and mobility	SR+MA (12 RCTs)	C	2	Significant improvement for back pain and functionality.
Forestier & Francon 2008 (Forestier and Françon 2008)	Patients with OA of the limbs (1658)	SPA-T	Any type	Pain, functionality, QoL	SR+MA (19 RCTs)	D	2	Suggestive evidence of a possible beneficial effect, but analyzed studies have some limitations.
Forestier et al. 2016 (Forestier et al. 2016)	Patients with knee OA (2917)	SPA-T	Any type	Any relevant clinical outcome	SR (30 RCTs)	C	2	Evidence of relevant clinical improvements lasting, on average, from 3 to 6 months (and up to 9 months) after intervention.
Forestier et al. 2017 (Forestier et al. 2017)	Patients with chronic LBP (2146)	SPA-T	Any type	Pain, functionality, QoL, drugs	SR (18 RCTs)	C	2	Evidence of improvements in pain, disability, and QoL, lasting 3-6 months after intervention. Possible reduction of painkiller intake.
Fraioli et al. 2013 (Fraioli et al. 2013)	Patients with fibromyalgia (271)	SPA-T	Any type	Pain, symptoms and mood	SR (7 studies)	D	?	Evidence of symptomatic improvement, including pain and mood (depression).
Fraioli et al. 2018 (Fraioli et al. 2018)	Patients with knee OA (1649)	SPA-T	Any type	Pain, functionality, QoL, drugs	SR (12 clinical studies)	D	?	Improvement of pain, joint mobility, NSAID intake, and QoL.
Gravelier et al. 2020 (Gravelier et al. 2020)	Patients with burn scars (115)	SPA-T.	Any type	Pain, skin elasticity, QoL	SR (2 RCTs)	C	3	Potential beneficial effects for burn scar recovery, but further scientific evidence is needed.
Kamioka et al. 2006 (洋晴 et al. 2006)	Patients with any disease (1425)	SPA-T	Any type	Pain, QoL, drugs, occupational functionality and healthcare costs	SR (18 RCTs)	D	2	Amelioration of all analyzed outcomes with high-quality studies mostly supporting the efficacy of intervention for rheumatic disorders.
Karagülle M. & Karagülle M.Z. 2015 (Karagülle and Karagülle 2015)	Patients with chronic LBP (769)	SPA-T	Any type	Any relevant clinical outcome	SR (8 RCTs)	D	?	Evidence of an improvement in symptomatic management, but further studies are needed to confirm these results.
Langhorst et al. 2009 (Langhorst et al. 2009)	Patients with fibromyalgia (446)	SPA-T	Any type	Any clinical and QoL-related outcome	SR+MA (10 RCTs)	A	2	Moderate evidence of beneficial effects on health-related QoL.
Matsumoto et al. 2017 (Matsumoto et al. 2017)	Patients with knee OA (734)	SPA-T	Any type	Pain and functionality	SR+MA (8 RCTs)	B	3	Possible beneficial effects, but high heterogeneity is found across studies and evidence is not of sufficiently high quality.
Morer et al. 2017 (Morer et al. 2017)	Patients with any rheumatic condition (1118)	SPA-T	HT	Pain, functionality, drugs, QoL, laboratory parameters	SR (27 RCTs)	D	3	Possible beneficial effects, but high heterogeneity and potential risk of bias is reported within and across available studies.
Naumann &	Patients with	SPA-T	Any type	Pain, functionality,	SR+MA (12	B	2	Significant improvements of symptoms and

Sadaghiani 2014 (Naumann and Sadaghiani 2014)	fibromyalgia (553)			QoL, mood	RCTs about SPA-T and 12, excluded, about HT)			QoL with potentially long-lasting effects on pain. No significant effect was observed on depressive symptoms
Pittler et al. 2006 (Pittler et al. 2006)	Patients with chronic LBP (674)	SPA-T	Any type	Pain, functionality, QoL, drugs	SR+MA (5 RCTs)	B	2	Scant but promising evidence which suggests a possible clinical benefit.
Raza et al. 2020 (Raza et al. 2020)	Patients with knee OA (831)	SPA-T	Any type	Pain and functionality	SR+MA (10 RCTs)	C	2	Significant improvement in pain and functionality, as measured with the WOMAC scale.
Santos et al. 2016 (Santos et al. 2016)	Patients with rheumatoid arthritis (496)	SPA-T	Any type	Pain, functionality, QoL, drugs, laboratory parameters	SR (8 RCTs)	B	2	Evidence of a beneficial effect on studied outcomes even up to 3 months after intervention.
Verhagen et al. 2007 (Verhagen et al. 2007)	Patients with OA (498)	SPA-T	Any type	Pain, functionality, QoL	SR+ MA (7 RCTs)	A	3	A positive effect is found when intervention is compared to no treatment, but the quality of evidence is low.
Verhagen et al. 2015 (Verhagen et al. 2015)	Patients with rheumatoid arthritis (579)	SPA-T	Any type	Pain, functionality, QoL	SR+MA (9 RCTs)	A	3	Insufficient evidence to draw conclusions on the topic.
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Casale et al. 2018 (Casale et al. 2018)	Patients with rhinosinusitis (663)	IT/IR	Any type	Functionality of upper airways	SR (11 RCTs)	D	2	Nasal irrigations with mineral waters can be clinically beneficial for patients with rhinosinusitis in terms of endoscopic scores and mucociliary clearance if compared to isotonic solution.
Keller et al. 2014 (Keller et al. 2014)	Patients with rhinosinusitis (840)	IT/IR	Any type	Functionality of upper airways	SR + MA (13 clinical studies of any type)	D	2	Mineral waters (for example, those ones rich in sulphur) can have an integrative role in the management of chronic inflammatory diseases of the upper respiratory tract.
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Böhmer et al. 2000 (Bohmer et al. 2000)	Healthy subjects (54)	HPT (mineral waters with high calcium content)	Dairy products	Bioavailability of calcium	SR (4,controlled studies)	D	?	Bioavailability of calcium from calcium-rich waters is comparable to that one from dairy product consumption.
Naumann et al. 2017 (Naumann et al. 2017)	Patients with cardiovascular risk factors (1089)	HPT (various mineral waters)	Other drinks	Glycemic control	SR (15 RCTs)	C	3	Bicarbonate- and magnesium-rich waters may have a positive impact on glycemic control.
Sulaiman et al. 2020 (Sulaiman et al. 2020)	Both healthy and diseased subjects at risk of kidney stone formation (470 involved in interventional studies)	HPT (mineral waters with high calcium, bicarbonate or magnesium content)	Tap water	Kidney stone prevention, especially calcium stones	SR (10 clinical interventional studies and 5 observational studies)	C	2	Bicarbonate- and magnesium-rich waters may be useful against kidney stone formation. Consumption of waters with a high calcium content can lead to hypercalciuria.
Reference	Population (n)	Intervention	Control	Outcomes	Study design	RQ	TQ	Conclusions
Stier-Jarmer et al. 2015 (Stier-Jarmer et al. 2015)	Patients with any non-musculoskeletal disease (13782)	Any Hydrologic Therapy	Any type	Any improvement assessed clinically or with diagnostic devices	SR (41 clinical studies of any type)	B	3	Clinical benefits for various skin diseases (atopic dermatitis, psoriasis, ictiosis), respiratory and ENT illnesses (rhinosinusitis, COPD, catarrhal otitis), vascular problems (hypertension, chronic venous insufficiency), digestive complaints (dyspepsia, irritable bowel syndrome), and neuropsychiatric

								conditions (peripheral neuropathy, Parkinson's' disease, psychophysical stress).
--	--	--	--	--	--	--	--	--

Legends:

BT=Balneotherapy (only baths with mineral waters)

COPD=Chronic Obstructive Pulmonary Disease

CP=Conference Proceedings (review presented at a conference/lecture/symposium and then published in a refereed journal as an abstract)

HPT=Hydropinic therapy (drinking mineral waters)

HT=Hydrotherapy (use of tap non-mineral water)

IT/IR=Inhalation therapy/irrigations (aerosols, vapors, nebulizations, humages, and oral/nasal irrigations)

LBP=Low Back Pain

MA=Meta-analysis

MT=Mud therapy (use of therapeutic muds)

NSAID=NonSteroidal Anti-Inflammatory Drug

OA=Osteoarthritis

QoL=Quality of Life

RCT=Randomized Controlled Trial

RQ=Review Quality or overall quality assessed on the basis of review methods (A=high quality, B=moderate quality, C=low quality,

D=very low quality, ?=the review quality was not assessable because the full-text version was irretrievable)

SPA-T=Spa therapy (multicomponent spa-based treatments)

SR=Systematic Review

TQ=Trial Quality or average quality of clinical studies included in each analyzed review (1=high quality, 2=fair quality, 3=poor quality,

?=trial quality was not assessed by the authors of included reviews)

TT=Thalassotherapy (baths with sea water)

Caption:

Included reviews are grouped on the basis of analyzed intervention and alphabetically sorted according to the first author's surname.

TABLE 2. Main characteristics (PICOS and study authors' conclusions) of included high-quality narrative reviews.

Reference	Population (n)	Intervention	Control	Outcomes	Study design	Conclusions
Cacciapuoti et al. 2020 (Cacciapuoti et al. 2020)	Patients with chronic inflammatory skin diseases (?)	BT	Any type	Various health-related outcomes	NR (?)	Benefits for psoriasis and atopic dermatitis, but also for pruritus, prurigo, lichen ruber planus, acne vulgaris, and seborrheic dermatitis.
Bender 2016 (Bender 2016)	Patients with OA (?)	SPA-T	Any type	Pain and QoL	NR (50+ RCTs) - CP	Positive effect on pain and QoL of patients with OA of various joints.
Fortunati et al. 2016 (Fortunati et al. 2016)	Patients with hand OA (168)	SPA-T	Any type	Pain, functionality, QoL	NR (3 RCTs)	Suggestive evidence of a potential benefit, but further investigation is advised to draw conclusions.
Francon et al. 2015 (Françon et al. 2015)	Patients with chronic pain of rheumatic origin (2905)	SPA-T	Any type	Pain, functionality, QoL, drugs	NR (28 RCTs)	Improvement of pain, joint mobility, painkiller intake, and QoL.
Guidelli et al. 2012 (Guidelli et al. 2012)	Patients with fibromyalgia (314)	SPA-T	Any type	Pain, functionality, QoL, mood	NR (8 RCTs)	Evidence of benefits in terms of pain, functionality, QoL, and mood, lasting 3 to 9 months after intervention.
Roques 2014 (Roques 2014)	Patients with any disease (?)	SPA-T	Any type	Any relevant clinical outcome	NR (90 RCTs) - CP	Beneficial effects mostly for arthro-rheumatic diseases, but even for psoriasis, atopic dermatitis, vessel disorders of lower limbs, gynecological conditions, and psychosomatic diseases.
Tenti et al. 2015 (Tenti et al. 2015)	Patients with knee OA (1198)	SPA-T	Any type	Pain, functionality, QoL, drugs	NR (14 RCTs)	Evidence of a positive effect on pain, disability, and QoL, which may last up to 6-9 months.
Passali et al. 2017 (Passali et al. 2017)	Patients with chronic rhinosinusitis, allergic rhinitis or asthmatic bronchitis (110)	IT	Any type	Functionality of upper and lower airways	NR (4 clinical studies of any type)	A 2-week inhalation therapy with radon-enriched water may improve nasal function and respiratory obstruction in patients with allergic respiratory diseases.

Legends:

BT=Balneotherapy (only baths with mineral waters)

COPD=Chronic Obstructive Pulmonary Disease

CP=Conference Proceedings (review presented at a conference/lecture/symposium and then published in a refereed journal as an abstract)

HT=Hydrotherapy (use of tap non-mineral water)

IT/IR=Inhalation therapy/irrigations (aerosols, vapors, nebulizations, humages, and oral/nasal irrigations)

LBP=Low Back Pain

MT=Mud therapy (use of therapeutic muds)

NR=Narrative Review

NSAID=NonSteroidal Anti-Inflammatory Drug

OA=Osteoarthritis

QoL=Quality of Life

RCT=Randomized Controlled Trial

SPA-T=Spa therapy (multicomponent spa-based treatments)

Caption:

Included reviews are grouped on the basis of analyzed intervention and alphabetically sorted according to the first author's surname.

1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152

TABLE 3. Systematic reviews including only clinical studies conducted in a specific country or region of the world.

Reference	Country	Conclusions
Bender et al. 2014 (Bender et al. 2014)	Hungary	Some evidence of efficacy of Hungarian mud baths for several rheumatic conditions, mostly osteoarthritis.
Drobnik and Stebel 2020 (Drobnik and Stebel 2020)	Poland and Austria	Scientific evidence on the “Tolpa” peloid is not strong enough to formulate specific clinical indications.
Karagülle and Karagülle 2004 (Karagülle and Karagülle 2004)	Turkey	Efficacy of spa therapy with Turkish mineral waters and muds for various rheumatic conditions, including osteoarthritis, fibromyalgia, and rheumatoid arthritis.
Katz et al. 2012 (Katz et al. 2012)	Israel	Baths in Dead Sea water and mud can be useful for the treatment of rheumatic diseases and psoriasis.
Khalilzadeh et al. 2019 (Khalilzadeh et al. 2019)	Iran	Possible efficacy of balneotherapy with Persian mineral waters for the management of psoriasis.
Roques et al. 2012 (Roques et al. 2012)	France	Spa therapy with French mineral waters and muds can be effective for the complementary treatment of osteoarthritis, tendinopathies, benign chronic low back pain, and leg chronic venous insufficiency. Beneficial effects were found for anxiety. Preliminary studies also reported some improvements in patients with neurologic and metabolic conditions.
Stanhope et al. 2018 (Stanhope et al. 2018)	Australia and New Zealand	No evidence of efficacy due to complete lack of clinical studies on the topic.

1153

TABLE 4. Water types and clinical indications according to the “HydroGlobe” study (2014).

Affected apparatus	Example	Administration	Recommended water types
Ear-Nose-Throat and Respiratory tract	Chronic rhinosinusitis or bronchitis	Inhalations, irrigations	Sulphurous; Salt, bromine and iodine; Bicarbonate; Arsenical-ferruginous.
Cardiovascular system	Chronic venous insufficiency	Baths	Carbonic.
Gynecological apparatus	Chronic vaginitis	Irrigations	Sulphurous; Salt, bromine; Bicarbonate; Sulphate.
Urinary tract	Recurrent kidney stones	Oral intake	Oligomineral (low mineral content); Bicarbonate.
Gastrointestinal system	Irritable bowel disease with constipation	Oral intake	Bicarbonate; Sulphate; Salt.
Skin	Psoriasis, atopic dermatitis	Baths	Salt, bromine and iodine; Radioactive; Bicarbonate; Sulphurous.
Musculoskeletal system	Osteoarthritis, fibromyalgia	Baths	Sulphurous; Salt, bromine and iodine; Radioactive.

Legends:

According to relevant directives of the European Union and to traditional definitions, mineral waters are labelled as follows (Nappi 2001; Vv.Aa. 2014; Quattrini et al. 2016):

- “Arsenical-ferruginous” waters: when they contain both arsenic and iron, either as a ferrous or ferric ion. Waters are considered “ferrous” or “ferruginous” when iron content is > 1 mg/L.
- “Bicarbonate-rich” waters: when bicarbonate content is > 600 mg/L. If the calcium content of these waters is > 150 mg/L and magnesium content is > 50 mg/L, they are also defined as “calcium- and magnesium-rich” waters.
- “Carbonic” waters: they spring up naturally with a detectable content of free carbon dioxide. When CO₂ content is > 250 mg/L, they are defined as “acid waters” due to their low pH.
- “Oligomineral” waters: characterized by a low mineral content, with a fixed residue at 180°C inferior to 500 mg/L. On the contrary, when the fixed residue exceeds 1500 mg/L, waters are defined as “highly mineralized”.
- “Radioactive” waters: they have a radioactivity of at least 1 nCi/L, mostly due to their content of Radon.
- “Sulphate-rich” waters: when sulphate content is > 200 mg/L. If the calcium content of these waters is > 150 mg/L and magnesium content is > 50 mg/L, they are also defined as “calcium- and magnesium-rich” waters.
- “Sulphurous” waters: they have a high content of bivalent sulphur.
- “Waters rich in salt, bromine and iodine”: they have a high mineral content and, like seawater, they are rich in sodium (in the form of NaCl) and other minerals. If sodium content is > 200 mg/L, they are labelled as “sodium-rich” waters. They can also contain elements like bromine and iodine.



[Click here to access/download](#)

Electronic Supplementary Material
Electronic Supplementary Materials REVISED.docx

