

University of Parma Research Repository

Stabilization of Arthrospira platensis with high-pressure processing and thermal treatments: Effect on physicochemical and microbiological quality

This is the peer reviewd version of the followng article:

Original

Stabilization of Arthrospira platensis with high-pressure processing and thermal treatments: Effect on physico-chemical and microbiological quality / Alinovi, Marcello; Bancalari, Elena; Martelli, Francesco; Cirlini, Martina; Rinaldi, Massimiliano. - In: JOURNAL OF FOOD PROCESSING AND PRESERVATION. - ISSN 0145-8892. - (2021). [https://doi.org/10.1111/jfpp.15912]

Availability: This version is available at: 11381/2898408 since: 2022-02-18T12:18:06Z

Publisher: Wiley

Published DOI:https://doi.org/10.1111/jfpp.15912

Terms of use:

Anyone can freely access the full text of works made available as "Open Access". Works made available

Publisher copyright

note finali coverpage

(Article begins on next page)

1 2	Stabilisation of <i>Arthrospira platensis</i> with high-pressure processing and thermal treatments: effect on physico-chemical and microbiological quality	
3		
4	Running title: Thermal, non-thermal treatment on Spirulina	
5		
6	Marcello Alinovi ^{1a} , Elena Bancalari ^{1a*} , Francesco Martelli ¹ , Martina Cirlini ¹ , Massimiliano Rinaldi ¹	
7	¹ Department of Food and Drug, University of Parma, Parco Area delle Scienze 47-49/A, 43124, Parma, Italy	
8	*Corresponding: <u>elena.bancalari@unipr.it</u>	
9	^a These authors contributed equally to the work.	
10		

12 Abstract

Arthrospira platensis (Spirulina) is a cyanobacterium that has been recently studied for food applications due to its high 13 biological and nutritional value. When A. platensis is used as ingredient in food applications, proper treatments have to be 14 15 applied in order to reduce microbial contamination. This work compared the effect of thermal treatments (sterilization at 121°C and pasteurization at 90°C) and high-pressure processing (400, 600 MPa) on the chemical, physico-chemical and 16 17 microbial quality of 5% (w/v) A. platensis aqueous suspensions. Total antioxidant capacity, total polyphenols content, colour and pigments content were not strongly lowered/modified by the HPP treatments. HPP at 400 MPa even improved the release 18 19 of C-phycocyanin from the biomass because of the breakage of cell walls. HPP treatments were comparable to pasteurization 20 in reducing Yeasts, Coliforms, Staphylococci and total bacterial count. Conversely, sterilization was the only treatment that 21 guaranteed the inactivation of spore-forming species but affecting the final quality.

- 22
- 23

24 Keywords: non-thermal processing; emerging technologies; high pressure processing; Spirulina; C-phycocyanin

25

26 Novelty Impact Statement HPP treatments were found to be a good strategy to preserve or even improve some physical and 27 chemical properties of *Arthrospira platensis* (Spirulina), in particular antioxidant capacity, polyphenols, colour and pigments 28 content. Furthermore, HPP treatments were comparable to pasteurization in reducing microbial cell count, while sterilization 29 was the only treatment able to ensure the inhibition of spore-forming species.

- 30
- 31

32 1. Introduction

- Arthrospira platensis, a cyanobacterium commonly known as Spirulina, is widely cultivated and studied because of its
 richness in bioactive compounds with high nutritional value, that can be used in several food formulations (Niccolai, Chini
 Zittelli, et al., 2019; Golmakani et al., 2019; Batista et al., 2017; Barkallah et al., 2017).
- 36 It can be commercialized dehydrated as a food integrator, or as an ingredient for food formulations or, as recently reported,
- as a booster in the fermentation process or as natural antimicrobial (Martelli, Alinovi, Bernini et al., 2020; Zheng et al., 2020,

39 used as ingredients (-Martelli, Cirlini, Lazzi et al., 2020; Bancalari et al., 2021) 40 A. platensis is commonly cultivated in raceway ponds that is cheaper than cultivations in photobioreactors (Richardson et al., 41 2012) but one of its major drawbacks is the high level of microbial contamination. In fact, alterative and spoilage bacteria are 42 reported as contaminants in these cultivations and their presence could lead to safety issues for consumers (Wang et al., 2013). 43 For this reason, microalgal biomass cannot be marketed fresh but needs to be dehydrated in order to reduce its water activity 44 (aw) and avoid the potential bacterial proliferation. Otherwise, thermal or non-thermal treatments can be applied to inactivate 45 alterative and foodborne pathogens, but, on the other hand they may cause an unwanted loss of nutritional and functional 46 compounds (Martelli et al., 2014). One of the most promising emerging alternatives to thermal treatment is high pressure 47 processing (HPP), that can reduce the microbial load and stabilize food products, avoiding losses of nutritional and functional 48 compounds (Paciulli et al., 2019). Furthermore, high pressure processing could also improve the extraction of bioactive compounds by disrupting tissues, cell walls, membranes and organelles (Cox et al., 2014; Jun, 2013; Martelli, Favari, Mena 49 50 et al., 2020). Reducing bacterial contamination to an acceptable level maintaining unchanged or even enhance the composition 51 of A. platensis is a goal that must be reached to use it as a safe and functional ingredient in food formulations. Moreover, HPP 52 treatment showed to have a positive effect in stabilizing the C-phycocyanin mainly in presence of proteins or hydrocolloids 53 due to encapsulation induced by pressure (Zhang et al., 2021). In the last years, the extraction and stabilization of C-54 phycocyanin, an accessory photosynthetic blue-protein that is present in relatively high concentration in A. platensis, has been 55 widely investigated for potential food applications (Zhang et al., 2021; Martelli et al., 2014; Martinez et al., 2017). The 56 interest for this protein pigment has grown because the confectionary and drinks industry has a high demand in blue colorants, 57 that are however uncommon in nature and have to be synthesized (Martelli et al., 2014) and because of the high protein yield 58 and the relatively low cost of production of this pigment. Thus, the aims of the present study were to: i) investigate the effect 59 of the above-mentioned treatments on: phenolic content, antioxidant capacity, chlorophyll, carotenoids and C-phycocyanin 60 content, protein fraction and colour and ii) evaluate the efficacy of two thermal treatments (pasteurization and sterilization) 61 and two high pressure treatments (HPP at 400 and 600 MPa for 3 min) in reducing the microbial contamination of A. platensis 62 aqueous solution.

Bancalari et al., 2020). Moreover, several bioactive or functional compounds can be extracted from A. platensis biomass and

Commentato [MA1]: Martínez, J. M., Luengo, E., Saldaña, G., Álvarez, I., & Raso, J. (2017). Cphycocyanin extraction assisted by pulsed electric field from Artrosphira platensis. *Food Research International*, *99*, 1042-1047.

63

38

64 2. Materials and Methods

65 2.1 Materials

Gallic acid, sodium carbonate, Trolox ((±)-6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid), DPPH (2,2-diphenyl1-pirylhydrazyl free radical) and methanol were purchased from Sigma-Aldrich (St. Louis, MO, USA); Folin-Ciocalteu's
phenol reagent solution was obtained from VWR (Milan, Italy). Bi-distilled water was produced by a Millipore Alpha Q
purification system (Waters, Billerica, MA, USA).

Acrylamide, N,N'-methylenbisacrylamide, N,N,N,N, tetramethyl ethylene-diamine (TEMED), glycine, Sodium dodecyl
sulphate (SDS), Tris aminomethane (TRIZMA® BASE), ammonium persulfate, 1,4-ditioeritrol (DTE), Bromophenol blue,
glycerol, sigmamarker[™] wide range protein ladder (200-6.5 kDa), Coomassie brilliant blue G250, orto-phosphoric acid
(85%), aluminium sulphate, ethanol, and acetone were purchased from-Merck (Milan, Italy).

74

75 2.2 Experimental design and processing treatments

76 Dried *A. platensis* was kindly provided by S.A.Ba.R (Novellara, RE, Italy). One hundred grams of dried product were 77 rehydrated in 2 L of distilled water (5% w/v) under gentle stirring at 4°C overnight. After rehydration, the suspension was 78 divided into ten aliquots of 200 mL that were packed in sterilized bottles with an internal volume of 250 mL. The aliquots 79 were grouped in 5 different treatment (two 200-mL aliquots for each treatment): i) raw, non-treated sample (control); ii) 400 80 MPa for 3 min (HPP400), iii) 600 MPa for 3 min (HPP600), iv) pasteurized at 90°C for 10 min (PAST) and v) sterilized at 81 121°C for 20 min (STER).

Both HPP treatments were performed with a QUINTUS FOOD PRESS QFP 35 L (Avure Technologies Inc., Middletown,
OH, USA). Cold water (4 °C) was used as pressure medium; temperature during processing was not controlled but the
temperature increase due to compression was not higher than 2-3 °C/100 MPa.

Sterilization was performed by means of Tuttnauer autoclave (Cavallo s.r.l., Milan, Italy), at 121°C for 20 min. Pasteurization treatment was performed by placing the samples in a temperature-controlled laboratory bath (type M418-BM, MPM instruments, Bernareggio, Italy) at 90°C for 10 min; the treatment durations of sterilization and pasteurization were respectively calculated when the inner temperature of the autoclave reached $121 \pm 1^{\circ}$ C and when the water inside the bath reached $90\pm 1^{\circ}$ C after the immersion of the bottles. After both pasteurization and sterilization, the bottles were immediately cooled in a water bath and then, immediately analysed or stored at refrigerated temperature (4 ± 1°C),

ha formattato: Inglese (Stati Uniti)

91 Five out of the ten bottles subjected to different treatments were analysed immediately (0 d of storage) and the other five
92 after a short period of refrigerated storage at 4°C (6 d of storage). Two technical replicates for each analysis were performed
93 for each sample.

94

95 2.3 Total phenolic content determination

Total phenolic content was determined using the Folin-Ciocalteu's method, following the protocol reported by Wu et al. (2005) with some modifications. Briefly, 0.25 mL of water diluted sample (1/5, v/v) were transferred in the test tubes and added with 1 mL of Folin-Ciocalteu's phenol reagent solution (1/10, v/v in bi-distilled water) and with 2 mL of aqueous sodium carbonate (20 %, w/v). All the samples were incubated for 30 min, in the dark at room temperature. Absorbance was measured at 760 nm using a JASCO V-530 spectrophotometer (Easton, MD, USA), and water was used to adjust zero. A calibration curve was built analysing 5 different gallic acid standard solutions (10 – 100 mg GAE/L). Results were expressed as gallic acid equivalent (mg GAE/L).

103

115

104 2.4 Radical-scavenging activity by DPPH method

The antioxidant capacity test was performed applying the DPPH radical scavenging assay on the basis of the protocol reported by Dall'Asta et al. (2013) with slight modifications: an aliquot of 0.1 mL of the 1/5 diluted sample was putted in a test tube and added with 2.9 mL of a methanolic DPPH solution (0.05 mM). The resulting mixture was kept in the dark at room temperature for 30 minutes, then the absorbance at 517 nm was registered on a JASCO V-530 spectrophotometer (Easton, MD, USA), and methanol was used for setting the autozero. At the same time, a blank sample was prepared and submitted to the same procedure. After that, the inhibition of DPPH radical was determined on the basis of the following formula:

111
$$I\% = \frac{Abs_0 - Abs_1}{Abs_0} \cdot 100$$
 (2)

where Abs_0 was the absorbance of the blank and Abs_1 was the absorbance of the sample. A calibration curve was constructed using Trolox as reference, preparing 5 different standard solutions in a concentration range of 0.1 - 1 mM. Then, the antioxidant capacity of samples was expressed as TEAC value (Trolox Equivalent Antioxidant Capacity; mmol Trolox eq/kg). **Commentato [eb2]:** Forse ci diamo la zappa sui piedi ma è quello che abbiamo fatto, cosa ne pensate? Penso che il revisore volesse sapere questo

116 2.5 Quantification of chlorophylls, carotenoids and phycocyanin content

117 2.5.1 Sample preparation for spectrophotometric assays

118 To perform spectrophotometric analyses, the insoluble biomass of A. platensis was separated by centrifugation (6,500 g for 119 20 min at 4°C) from the supernatant containing water-soluble C-phycocyanin. Supernatant was extracted by pipetting and an 120 aliquot (100 µL) was diluted by adding 900 µL of ultrapure water for C-phycocyanin quantification. The pellet containing A. 121 platensis biomass was weighed and resuspended in ~900 µL of pure acetone (1:20 dilution). The suspension was vortexed at 122 2,000 rpm and sonicated for 15 min using a workbench bath (VWR ultrasonic Cleaner) (Hynstova et al. 2018). Extracted 123 pigments (chlorophyll and carotenoids) were separated by centrifuging at 13,700 g for 10 min at 4°C. Supernatant was filtered 124 by syringe filtration using 0.45 µm filters (VWR International, Milan, Italy) and 50 µL of permeate were diluted by adding 125 950 µL of pure acetone. Samples preparation was performed in dim light to avoid pigment degradation. Each sample was 126 prepared by means of two independent extractions.

127

128 2.5.2 Spectrophotometric determination of the pigments

For chlorophylls and carotenoids determination, methanol was used as the blank of analysis. Concentration of chlorophyll a
(Ca), chlorophyll b (Cb), total chlorophylls (Ca+b), total carotenoids (Cx+c), total chlorophylls and carotenoids (Ccc) and
pheophytins (Cph) were calculated according to Hynstova et al. (2018) and Lichtenthaler (1987) using equations 3-8:

132	$Ca(\mu g/mL) = 11.24A_{662} - 2.04A_{645}$	(3)
-----	---	-----

133	$Cb (\mu g/mL) = 20.13 A_{645} - 4.19 A_{662}$	(4)

134 $Ca + b (\mu g/mL) = 7.05 A_{662} + 18.09 A_{645}$ (5)

135
$$Cx + c (\mu g/mL) = \frac{(1,000 A_{470} - 1.90 Ca - 63.14Cb)}{214}$$
 (6)

136
$$Ccc (\mu g/mL) = (Ca + b) + (Cx + c)$$
 (7)

137
$$Cph(\mu g/mL) = 321.3 A_{653} - 208.4 A_{654}$$
 (8)

For C-phycocyanin, milliQ water was used as the blank of analysis. Concentration of water-soluble C-phycocyanin was
determined according to the method reported by de Marco Castro et al. (2019). Quantification of C-phycocyanin was

140 performed according to equation (9):

141
$$C - phycocyanin\left(\frac{mg}{mL}\right) = \frac{A_{615} - 0.474 A_{652}}{5.34}$$
 (9)

142 where A_{615} and A_{652} are absorbances of the sample measured at 615 nm and 652 nm.

143

144 2.6 Physical properties

145 Colorimetric parameters were evaluated with image analysis: samples were scanned by means of a desktop flatbed scanner 146 (Hewlett Packard Scanjet 8200, Palo Alto, CA, USA) at 236 pixels/cm (600 dpi of resolution; true colour - 24 bit), equipped 147 with a cold cathode lamp for reflective scanning. All images were scanned at the same conditions, during image acquisition, 148 the scanner was held in a black box to exclude surrounding light and external reflections. Flatbed scanner colour (R, G and 149 B) was corrected as previously reported by N'Dri et al. (2010) and conversed to L* (lightness), a* (redness at positive values, 150 greenness at negative values) and b* (yellowness at positive values, blueness at negative values). Total colour differences 151 (ΔE) between 0 and 6 days were also calculated. 152 Water holding capacity (WHC) of A. platensis suspensions was measured similarly to Barkallah et al. (2017). Samples aliquots

(~1 g) were placed in 1.5 mL tubes and centrifuged at 6,000 g per 10 min at 4°C with a benchtop centrifuge (mod. 5810R,
Eppendorf, Hamburg, Germany). WHC was calculated as follows (equation 1):

155
$$WHC\left(\%\frac{w}{w}\right) = \left(\frac{W_1}{W_2}\right) \cdot 100$$
 (1)

where W_1 is the weight of *A. platensis* centrifuged pellet and W_2 is the weight of the original sample.

157

158 2.7 Optical microscopy observation

An aliquot of 10 µL of each sample was observed by means of an optical light microscope Nikon 80i (Tokyo, Japan) equipped
with a 10X objective. Pictures of each sample were acquired by the Nis Elements software (ver. 2.10 Nikon, Tokyo, Japan).

162 2.8 Sodium Dodecyl Sulphate polyacrylamide gel electrophoresis (SDS PAGE)

163	Sodium dodecyl sulphate (SDS) PAGE was performed according to the method of Laemmi (1970). Stacking and separating
164	gels were casted by using a 4 and 15% acrylamide concentration, respectively. Diluted samples (proteins concentration of \sim 2 methods are casted by using a 4 and 15% acrylamide concentration, respectively.
165	$\mu g/\mu L) \mbox{ each sample were loaded in the gel wells (5 \mu L \mbox{ aliquots}). \mbox{ Samples were run in non-reducing conditions at 150 V on a loaded in the gel wells (5 \mu L \mbox{ aliquots}). \mbox{ Samples were run in non-reducing conditions at 150 V on a loaded in the gel wells (5 \mu L \mbox{ aliquots}). \mbox{ Samples were run in non-reducing conditions at 150 V on a loaded in the gel wells (5 \mu L \mbox{ aliquots}). \mbox{ Samples were run in non-reducing conditions at 150 V on a loaded in the gel wells (5 \mu L \mbox{ aliquots}). \mu = 100000000000000000000000000000000000$
166	Mini-Protean II cube (Bio-Rad, Hercules, CA, USA).

SDS PAGE gels were stained with Coomassie Brilliant Blue G250 according to Kang et al. (2002), destained in severalchanges of distilled water and scanned with a flatbed scanner.

169

170 2.9 Microbial counts

Each sample was ten-fold (1:10 v/v) diluted, up to 10⁻⁶ in Ringer's solution (Sigma Aldrich, Milan) and then inoculated in 171 172 specific culture media using the spread-plate technique. Total microbial count (TBC) was obtained by plating onto Plate Count 173 Agar (PCA, Oxoid, Basingstoke, Hampshire, UK) and incubating at 37 ± 1 °C for 48 h. Chromocult® Coliform Agar (Merck, 174 KGaA, Darmstadt, Germany) was used for the enumeration of Coliforms by incubating at 37 ± 1 °C for 24 h. Staphylococci 175 were counted by spreading the dilutions on Mannitol Salt Agar (Oxoid, UK) and incubated at 37 ± 1 °C for 48 h. For the 176 count of Listeria, Agar Listeria Ottaviani and Agosti medium (ALOA, Biolife Italiana, Milan) was used and plates incubated 177 for 48 h at 37 ± 1 °C. Yeast and molds were counted on Yeast Extract Glucose Chloramphenicol Agar (YGC, Merck KGaA, 178 Darmstadt, Germany) that was incubated at 25 ± 1 °C for 48–72 h.

For aerobic and anaerobic spore-forming bacteria, samples were heated to 80 °C for 10 min and then cooled before analysis. Then the samples were serially diluted in Ringer's solution, and plated on Tryptone Soy Agar (TSA, Merck KGaA, Darmstadt, Germany), followed by incubation for 24 h at 30 ± 1 °C under aerobic condition for the determination of aerobic spore-forming bacteria (Abdelmassih et al., 2011). For the count of anaerobic spore-forming bacteria, 1 mL of each dilution was inoculated in the medium using the pour-plate technique and then covered with a thin layer of the same medium in order to obtain a double-layer. The plates were anaerobically incubated into the jars with the AnaeroGen sachet (Oxoid, UK), at 30 ± 1 °C for 24 h.

186

187 2.10 Statistical analysis

Statistical analyses were performed using SPSS Statistics v. 25 (IBM, Armonk, NY, USA): one-way ANOVA and Tukey HSD post hoc test were applied to test significant differences among the treatments and storage times (P<0.05). Pearson correlation coefficients between total polyphenolic concentrations and antioxidant capacity values were also calculated considering a significant correlation for P<0.05.</p>

192

193 3. Results and Discussion

194 3.1 Total polyphenolic content (TPC)

Immediately after processing, HPP treatments did not affect the amount of polyphenols if compared to the control sample.
Conversely, pasteurized and sterilized samples (0 d of storage) showed significantly (P<0.05) lower concentrations of phenols in respect to the untreated suspension (Table 1). The loss of polyphenols in PAST and STER samples could be ascribed to the effect of the heating treatments.

The phenolic concentration of the aqueous suspension of *A. platensis* at 0 d (Table 1) was 149.49 ± 0.31 mg GAE/L,
significantly higher (P<0.05) than after the storage period (216.48 ± 2.07 mg GAE/L).

201 After 6 d of refrigerated storage, control and HPP400 samples showed a significant increase (P<0.05) of TPC. This trend 202 could be probably due to a release of phenolic compounds from cells; polyphenols are contained in cells and for their 203 completely extraction the lysis of cell walls results necessary (Parniakov et al., 2015). Conversely, HPP600 samples showed 204 a significantly lower value than HPP400 and control samples, but still a significantly higher (P<0.05) value than PAST and 205 STER (Table 1). Differences observed between the two HPP treatments after the storage could be ascribed to a degradation 206 of the compounds that react with Folin-Ciocalteau's reagent, such as C-phycocyanin (İlter et al., 2018). By increasing the 207 pressure applied in HPP processing, the degradation of C-phycocyanin may be observed, as reported in a previous work which 208 investigated the decomposition of this molecule at different pressure levels (50 - 600 MPa) (Li et al., 2020).

209

210 3.2 Antioxidant activity

A. platensis aqueous sample at 0 d presented an antioxidant activity of 0.472 ± 0.036 mmol Trolox eq/kg (Table 1). As already
 observed for TPC at 6 d, the antioxidant capacity of the control samples increased to 0.578 ± 0.027 mmol Trolox eq/kg
 (P<0.05). On the other hand, the other samples presented values comparable with those found in the control at 0 d, while after
 6 d of storage, all the samples presented a lower antioxidant activity in respect to the control. In particular, the pasteurized

215 sample showed the statistically lowest radical scavenging capacity (P<0.05) (Table 1). These results reflected those observed 216 for total polyphenolic content: a strongly positive relationship (r = 0.81) between the radical scavenging activity and the 217 phenolic concentration values was found. 218 Along with polyphenolic compounds, several other molecules naturally present in A. platensis showed antioxidant properties, 219 such as carotenoids, chlorophylls, phycocyanin and ascorbic acid (Chopra & Bishnoi, 2008). So, even if heating treatments 220 caused a reduction of TPC as in the case of sterilized samples, the antioxidant capacity remained almost comparable in respect 221 to the control, probably due to the presence of other compounds not belonging to the polyphenols class. The control sample, 222 prepared at a concentration of 5 % of A. platensis (50 μ g/mL), showed a concentration of antioxidant compounds of 0.472 \pm 223 0.036 mmol Trolox eq/kg, corresponding to an inhibition of 12.65 ± 0.97 %, resulting indeed higher as compared to 224 data already reported in literature. Also ethanolic suspension that contained 25 µg/mL of A. platensis, showed an inhibition 225 of 10.30 ± 0.80 % (Anbarasan et al., 2011). A. platensis is a well-known antioxidant source: some studies reported the 226 antioxidant properties of A. platensis water extracts in reducing cell death due to apoptosis caused by the presence of free 227 radicals (as DPPH), so the authors hypothesized that these products could be applied in food as natural preservatives (Chu et 228 al., 2010).

229

230 3.3 Physical properties of *A. platensis* aqueous dispersions

In general, all the treatments showed a significant effect on the colour of the samples, as depicted in Fig. 1. Regarding L*
values, all treatments caused a significant increase (P<0.05) (Table 2). Increase of L* values could be due to rapid oxidation
and/or pigment decomposition at high temperature, in accordance with Nouri et al. (2018).

234 Redness (positive a* scores) was significantly reduced (P<0.05) by all treatments apart from HPP400, that showed 235 an increase if compared to the control; the extent of reduction was in the following order: PAST>HPP600>STER. 236 Pasteurization (PAST) caused the highest decrease of a*; in particular, PAST was the only sample that showed a negative a* 237 value, indicating a green shade of the sample, that was also visually observable (Fig. 1). Interestingly, HPP400 showed a 238 HPP600. significantly higher (P<0.05) a* value than 239

Yellowness (<u>positive b* scores</u>) significantly increased with the thermal treatments (P<0.05), while HPP samples did not
show significant differences (P>0.05) compared to the control. The thermal degradation of C-phycocyanin in PAST and STER
probably caused a loss of blue hue towards a greener and browner colour (Fig.1).

Changes in b* and a* values were due to the effect of different processes that can cause pigments' degradation: e.g. for chlorophylls with the formation of various coloured compounds (Koca et al., 2007), and for C-phycocyanin which results in precipitation and fading of the blue pigment (increase of b*) (Chaiklahan et al. 2012). On the contrary, the HPP treatments compared to the control did not show significantly different values in terms of the b* parameter, despite the highest C-phycocyanin content was observed in HPP400 (Fig. 5) and apparently led to a bluer shade (Fig. 1), with this latter characteristic probably caused by the higher, positive a* score that led to a bluer hue.

250

251 <u>The highest total colour difference (ΔE) was higher for the thermally treated samples if compared to the HPP 252 ones (Table 2). Surprisingly, the highest total colour difference (ΔE), with an evident variation of colour compared to the 253 control, was observed for PAST and not for STER, that reported the second highest total colour difference with 254 the control sample. On the contrary, the lowest ΔE value was recorded for HPP600 and HPP400. These results 255 confirmed the strongest impact of the thermal treatment on colour characteristics, that still represent an important limitation 256 of *A. platensis* in food applications (Chaiklahan et al. 2012).</u>

During refrigerated storage, the thermally-treated samples showed a considerable stability, conversely, the control and the HPP samples showed significant variations (P<0.05), especially a decrease in L* and b* and an increase in a* values was observed, probably due to microbial activity and/or residual activity of enzymes.

As reported in Fig. 2, water holding capacity (WHC) of the control samples resulted 13.3 ± 0.1 %, confirming the characteristic of *A. platensis* to retain water thanks to the high protein (Benelhadj et al., 2016) and extracellular polysaccharides (EPS) content (Trabelsi et al., 2009). All the treatments caused a significant increase (P<0.05) in WHC values except for HPP600. The increase in WHC was probably related to gelation of *A. platensis* proteins (Chronakis, 2001) due to dissociation into subunits and unfolding of the protein molecules. As temperature increases, reactive groups are exposed favouring protein-protein or other electrostatic and hydrogen bonding interactions. Thermally treated samples (PAST and STER) presented

266 significant higher values (P<0.05) compared to high pressure treated ones. After 6 d of storage, WHC of the treatments

- 267 remained the same with no significant (P>0.05) variations, confirming the stability of the formed network.
- 268

269 3.4 Pigments concentration

270 Concentrations of chlorophyll a (Ca), carotenoids (Cx+c) and pheophytins (Cph) are reported in Table 3. The control sample 271 showed a relatively high Ca content (135 \pm 8 µg/mL at 0 d, corresponding to 2.71 \pm 0.15 mg/g dry weight), comparable to 272 values already reported in other studies (Hynstova et al., 2018; Park et al., 2018). Conversely, Cb content of raw *A. platensis* 273 was low (6 \pm 4 µg/mL) (Table 3).

- 274 Cph, the primary degradation product of chlorophylls due to thermal treatments, was found in a concentration of 779 ± 45 275 μ g/mL in the control. In particular, Cph content was found to be higher than Ca (Table 3) and this can be due to the initial
- 276 degradation of chlorophylls during the A. platensis industrial drying step (Hynstova et al., 2018).
- The concentration of C-phycocyanin (Fig. 3), the main phycobiliprotein present in *A. platensis*, was 760 \pm 20 µg/mL in the control, corresponding to 15.1 \pm 0.4 mg/g dry weight, in accordance with de Marco Castro et al. (2019).

Afterwards, the different treatments showed significant differences (P<0.05) in terms of pigments content (Table 3, Fig. 3). As expected, thermal treatments promoted a significant reduction of chlorophylls, pheophytin and C-phycocyanin. On the contrary, the reduction of carotenoids was only significant for STER (P<0.05). Interestingly, both HPP treatments did not cause a significant variation of total chlorophylls, carotenoids and pheophytin content, and this result was in line with the colorimetric analyses. This observation highlights the benefits of applying HPP treatments, compared to the thermal ones, which lead to significant denaturation of these thermolabile compounds that are known to have an important positive bioactive activity (Ariede et al., 2017; Kumar et al., 2015).

C-phycocyanin concentration was slightly, but significantly higher (P<0.05) for the HPP400 (Fig. 3) also if compared to the control, probably due to both an increase in extraction of this molecule from *A. platensis* cells and a limited denaturation.
Contrarily, Li et al. (2020) observed a significant reduction in the C-phycocyanin content in HPP-treated *A. platensis* at 400 MPa for 3.5 min and hypothesized that its decrease was related to the denaturation promoted by the HPP treatments. The microscopic observations allowed to observed that HPP treatment induced cell breakage, although not complete (Fig. 4). This can explain the increase in the C-phycocyanin concentration in sample treated at 400 MPa. On the contrary, stronger HPP

conditions (600 MPa x 3 min) caused a marked denaturation and a consequent reduction of C-phycocyanin concentration, in accordance with Li et al. (2020), that observed a heavier C-phycocyanin denaturation at higher pressure treatments. According to its low thermal stability (Martelli et al., 2014), C-phycocyanin was also heavily denatured by PAST and STER thermal treatments that showed about 87 and 97 % of reduction, respectively. It is quite well known that solutions containing Cphycocyanin are sensitive to heat treatment already at temperature above 47°C (Chaiklahan et al., 2012) with at least 50% of reduction after 30 min at 60°C. By considering first-order rate constants reported by (Chaiklahan et al., 2012) calculated residual activities agree with experimental data.

299 After 6 d of storage, a general increase of pigments, although not always significant, was observed; in particular, HPP400 and 300 pasteurized samples showed a significant increase (P<0.05) in the total chlorophyll and carotenoids concentration, while the 301 control, HPP400, HPP600 samples showed a significant increase (P<0.05) in the C-phycocyanin concentration. This result 302 was partly in accordance with the increase of antioxidant capacity of the control and of the polyphenols content of the control 303 and HPP400 (section 3.4). As stated before, an increase in the pigments' concentrations could be explained by the presence 304 of residual enzymatic activities that may cause the release of these compounds from algae cells. Previous studies demonstrated 305 that enzymes produced by bacteria can improve the extraction of C-phycocyanin (de Marco Castro et al., 2019; Zhu et al., 306 2007).

307

308 3.5 Optical microscopy observation

309 To further investigate whether each treatment affected the microstructure of A. platensis, the samples were observed using 310 light microscopy (Fig. 4). A. platensis is characterized by a particular morphological feature, such as the arrangement in an 311 open left-hand helix along the entire length (Ali & Saleh, 2012). The untreated A. platensis (Fig. 4a) already showed some 312 breakdown of the cell walls and the presence of cell's fragments, probably mainly related to the drying process of the biomass 313 (CIT). In the thermally treated samples, and particularly in STER (Fig. 4e), the appearance of some aggregates can be 314 observed. These are assumed to be caused by the sterilization process that may lead to aggregation of denatured proteins, that are the major constituent of the total biomass (Bernaerts et al., 2017). These aggregates are probably the main responsible for 315 316 the higher WHC (Fig. 2). Both HPP treatments (Fig. 4b, c), affected the structure of A. platensis to a lesser extent, by causing β17 a lower but observable breakdown of the cell walls, if compared to the more intact number of cells observed in the control (Li 318 et al., 2020).

Commentato [MA3]: Microalgae Nutraceuticals •August 2016 •Foods 5(3) • Follow journal •DOI: •10.3390/foods5030054

ha formattato: Evidenziato

320 3.6 Protein determination by SDS PAGE

321 Results of SDS PAGE are reported in Fig. 5. Several proteins bands were visible and the majority were present in the 322 medium molecular weight range (30-60 kDa) (Aiello et al., 2019). Biliproteins were the most intense bands present in the low 323 molecular weight range of the electrophoretogram (Li et al., 2020; de Marco Castro et al., 2019), which are mainly represented 324 by C-phycocyanin α and β subunits (Benelhadj et al., 2016). Chaiklahan et al. (2011) reported that C-phycocyanin α and β 325 subunits were characterized by a molecular weight of about 18.4 and 21.3 kDa, respectively; Rajakumar & Muthukumar, 326 (2018) reported a molecular weight range of 19-22 kDa. Our results were consistent with these observations, as C-phycocyanin 327 α and β subunits were found to have a molecular weight of about 18.5 and 20.5 kDa, respectively. 328 Other well-defined bands were observable at higher molecular weights (~44, 60, 102 kDa, corresponding to peaks 3, 4, 5

reported in Fig. 5). In particular, peak 3 (~44 kDa) is probably indicative of the presence of chlorophyll complexes (Kalaji et
 al., 2017) and peak 5 (~102 kDa) is probably related to a core protein linker of the phycobilisomes to thylakoid membrane

331 (Rajakumar & Muthukumar, 2018).

In general, protein bands showed a reduction because of both thermal and non-thermal treatments. The decrease of intensity was more evident for thermally-treated samples than HPP ones; this was expected as C-phycocyanin is denatured at temperatures above 40°C (Liu et al., 2016), and it was in accordance with pigments quantification (section 3.4). No observable protein bands were present in the sterilized sample.

- 336 Concerning HPP treatments, HPP400 showed a smaller reduction in protein bands than HPP600, if compared to the control\$37 sample.
- 338

339 3.7 Microbial counts

The result of total microbial count (TBC) on raw samples, showed a value of 5.96 ± 0.05 Log CFU/mL (Table 4), confirming
the high values of microbial contamination previously observed (Morais et al., 2009; Yu et al., 2019; Martelli, Cirlini, Lazzi
et al, 2021; Martelli, Marrella, Lazzi et al 2021). Regarding the treated samples, a different microbial reduction was observed
depending on the treatment: pasteurization, together with HPP400 and HPP600 allowed a reduction of respectively 4.30, 4.32

319

and 4.96 Log CFU/mL. Sterilization achieved a significantly higher reduction of microbial load, for all the species considered(Table 4).

346 If compared to both HPP treatments, pasteurization was more effective on Salmonella, allowing a reduction of 4 Log CFU/mL.
347 Despite no significant differences (P>0.05) were found between HPP600 and HPP400 on Salmonella, a greater reduction of
348 2.68 Log CFU/mL was observed in samples treated with HPP600 (Table 1). This could be due to the higher pressure, that
349 could induce a greater denaturation of enzymes and proteins, the breakdown of the cell membrane with the consequent loss
350 of internal substances, and a higher bacterial death (Abe, 2007).

351 Results on spores confirmed that neither pasteurization nor HPP treatments can completely inactivate endospore-forming 352 strains: their effect on those bacteria count was not consistent, allowing a small reduction of only about 1 Log CFU/mL. Our 353 results agreed with Reddy et al. (2006), who reported that bacterial spores cannot be inactivated by high pressure alone and 354 that they can survive at pressure treatments above 1000 MPa, probably due to the thickness structure of bacterial spore coat. 355 Under such conditions, the pressure tolerance of the endospore substantially exceeds that of the microorganism's original 356 vegetative state (Leggett et al., 2012). From these results, HPP effect on A. platensis microbial count, was comparable to the 357 effect of a high temperature pasteurization. Sterilization was the only treatment that can ensure the inhibition of spore-forming 358 species. Some authors reported that to achieve sterilization levels, HPP treatments can be used combined with an appropriate 359 temperature treatment or, in combination with other pressure cycle, to effectively inhibit endospore activity (Huang et al., 360 2014).

- Regarding the analysis after 6 d of refrigerated storage, the overall microbial count (TBC) increased for all the treated samples;
 probably due to the presence of heat-resistant, psychrotrophic gram-positive bacteria (Fromm & Boor, 2004). Significant
 (P<0.05) increments were also found in the yeast and Staphylococci enumeration.
- 364 For *Salmonella*, a significant reduction (P<0.05) of concentration was found after 6 d of storage for both HPP treatments
- 365 while spore forming bacteria, showed a small reduction in number for all the treatments applied.
- 366 Concerning the sterilized samples, no significative differences (P>0.05) were found during the storage.
- 367
- 368 4. Conclusions

369	In this study different thermal and HPP treatments were applied on 5% (w/v) A. platensis aqueous suspensions, to assess	
370	which treatments was the best compromise between the achievement of microbial safety and the preservation bioactive	
371	compounds. HPP treatments were found to be the best process in preserving the biological value and the physical properties	
372	of A. platensis. In particular, HPP400, even improved the release of C-phycocyanin from A. platensis probably because of the	
373	breakage of cell walls promoted by the mechanical effect of high pressures.	
374	From the microbiological analysis it was observed that HPP treatments were comparable to pasteurization in reducing	
375	microbial cell count, while sterilization was the only treatment able to ensure the inhibition of spore-forming species.	
376	These results suggest that HPP, despite being the best treatments in preserving the biological values of A. platensis, were not	
377	sufficient to achieve microbial stabilization. In conclusion, a combination of thermal and HPP treatments or a combination of	
378	HPP cycles, could be further investigated as a strategy to obtain a good compromise between microbiological and physico-	
379	chemical, nutritional quality. These results may be useful to design strategies to stabilize and transform food products	
380	containing A. platensis as ingredient.	

381

382 Declarations of Interest

- 383 None
- 384

385 References

Abdelmassih, M., Planchon, V., Anceau, C. & Mahillon, J. (2011). Development and validation of stable reference materials
 for food microbiology using *Bacillus cereus* and *Clostridium perfringens* spores. *Journal of Applied Microbiology*, *110*, 1524 1530. https://doi.org/10.1111/j.1365-2672.2011.05007.x

- Abe, F. (2007). Exploration of the effects of high hydrostatic pressure on microbial growth, physiology, and survival:
 perspectives from piezophysiology. *Bioscience, Biotechnology, and Biochemistry, 71,* 2347-2357.
 https://doi.org/10.1271/bbb.70015
- Aiello, G., Li, Y., Boschin, G., Bollati, C., Arnoldi, A., & Lammi, C. (2019). Chemical and biological characterization of
 spirulina protein hydrolysates: Focus on ACE and DPP-IV activities modulation. *Journal of Functional Foods*, 63, 103592.
 https://doi.org/10.1016/j.jff.2019.103592
- Ali, S. K., & Saleh, A. M. (2012). Spirulina an overview. *International Journal of Pharmacy and Pharmaceutical Sciences*,
 4(3), 9-15.

- Anbarasan, V., Kumar, V. K., Kumar, P. S., & Venkatachalam, T. (2011). In vitro evaluation of antioxidant activity of blue
 green algae *Spirulina platensis*. *Journal of Pharmaceutical Science and Research*, *2*, 2616-2618.
- Antelo, F. S., Costa, J. A. V., & Kalil, S. J. (2008). Thermal degradation kinetics of the phycocyanin from *Spirulina platensis*.
 Biochemical Engineering Journal, 41, 43–47. <u>https://doi.org/10.1016/j.bej.2008.03.012</u>
- Ariede, M. B., Candido, T. M., Jacome, A. L. M., Velasco, M. V. R., de Carvalho, J. C. M., & Baby, A. R. (2017). Cosmetic
 attributes of algae—A review. *Algal Research*, *25*, 483–487. <u>https://doi.org/10.1016/j.algal.2017.05.019</u>
- Bancalari, E., Martelli, F., Berinini, V., Neviani, E., & Gatti, M. (2020). Bacteriostatic or bactericidal? Impedometric
 measurements to test the antimicrobial activity of *Arthrospira platensis* extract. *Food Control, 118, 107380.*https://doi.org/10.1016/j.foodcont.2020.107380
- Bancalari, E., Martelli, F., Bottari, B., V., Neviani, E., & Gatti, M (2021). *Arthrospira platensis* Extract: A Non-Invasive
 Strategy to Obtain Adjunct Attenuated Cultures. *Foods*, *10*, 588. <u>https://doi.org/10.3390/foods10030588</u>
- Barkallah, M., Dammak, M., Louati, I., Hentati, F., Hadrich, B., Mechichi, T., Ayadi, M. A., Fendri, I., Attia, H., & Abdelkafi,
 S. (2017). Effect of *Spirulina platensis* fortification on physicochemical, textural, antioxidant and sensory properties of yogurt
 during fermentation and storage. *LWT Food Science and Technology*, *84*, 323–330.
- 410 during fermentation and storage. LWT Food Science and Technology, 84,
 411 <u>https://doi.org/10.1016/j.lwt.2017.05.071</u>
- Bartlett, D. H. (2002). Pressure effects on in vivo microbial processes. *Biochimica et Biophysica Acta (BBA)-Protein Structure* and Molecular Enzymology, 1595, 367-381. <u>https://doi.org/10.1016/S0167-4838(01)00357-0</u>
- Batista, A. P., Niccolai, A., Fradinho, P., Fragoso, S., Bursic, I., Rodolfi, L., Biondi, N., Tredici, M. R., Sousa, I., &
 Raymundo, A. (2017). Microalgae biomass as an alternative ingredient in cookies: Sensory, physical and chemical properties,
 antioxidant activity and in vitro digestibility. *Algal Research*, *26*, 161–171. <u>https://doi.org/10.1016/j.algal.2017.07.017</u>
- 417 Benelhadj, S., Gharsallaoui, A., Degraeve, P., Attia, H., & Ghorbel, D. (2016). Effect of pH on the functional properties of
 418 Arthrospira (Spirulina) platensis protein isolate. Food Chemistry, 194, 1056-1063.
 419 https://doi.org/10.1016/j.foodchem.2015.08.133
- Bernaerts, T. M. M., Panozzo, A., Doumen, V., Foubert, I., Gheysen, L., Goiris, K., Moldenaers, P., Hendrickxa, M.E., &
 Van Loey, A. M. (2017). Microalgal biomass as a (multi)functional ingredient in food products: Rheological properties of
 microalgal suspensions as affected by mechanical and thermal processing. *Algal Research*; *25*, 452–463.
 https://doi.org/10.1016/j.algal.2017.05.014
- Chaiklahan, R., Chirasuwan, N., & Bunnag, B. (2012). Stability of phycocyanin extracted from Spirulina sp.: influence of
 temperature, pH and preservatives. *Process Biochemistry*, 47, 659-664. https://doi.org/10.1016/j.procbio.2012.01.010
- Chaiklahan, R., Chirasuwan, N., Loha, V., Tia, S., & Bunnag, B. (2011). Separation and purification of phycocyanin from *Spirulina sp.* using a membrane process. *Bioresource Technology*, 102(14), 7159-7164.
 https://doi.org/10.1016/j.biortech.2011.04.067

- Chopra, K., & Bishnoi, M. (2008). Spirulina in Human Nutrition and Health. In M. E. Gershwin, A. Belay (Eds) Antioxidant
 profile of Spirulina: a blue-green microalga. (pp. 101-116) CRC Press, Boca Raton, FL, USA,
- 431 Chronakis, I. S. (2001). Gelation of edible blue-green algae protein isolate (Spirulina platensis strain pacifica): thermal
- transitions, rheological properties, and molecular forces involved. *Journal of Agricultural and Food Chemistry*, 49, 888-898.
 <u>https://doi.org/10.1021/jf0005059</u>
- Chu, W-L., Lim, Y-W., Radhakrishnan, A.K., & Lim, P-E. (2010). Protective effect of aqueous extract from *Spirulina platensis* against cell death induced by free radicals. *BMC Complementary and Alternative Medicine*, 10, 53.
 https://doi.org/10.1186/1472-6882-10-53
- 437 Cox, S., Hamilton Turley, G., Rajauria, G., Abu-Ghannam, N., & Jaiswal, A. K. (2014). Antioxidant potential and
 438 antimicrobial efficacy of seaweed (*Himanthalia elongata*) extract in model food systems. *Journal of Applied Phycology*, 26,
- 439 1823–1831. <u>https://doi.org/10.1007/s10811-013-0215-0</u>
- da Silva, M. F., Casazza, A. A., Ferrari, P. F., Aliakbarian, B., Converti, A., Bezerra, R. P., Porto, A. F. L., & Perego, P.
 (2017). Recovery of phenolic compounds of food concern from *Arthrospira platensis* by green extraction techniques. *Algal Research*, *25*, 391–401. https://doi.org/10.1016/j.algal.2017.05.027
- Dall'Asta, C., Cirlini, M., Morini, E., Rinaldi, M., Ganino, T., & Chiavaro, E. (2013). Effect of chestnut flour supplementation
 on physico-chemical properties and volatiles in bread making. *LWT Food Science and Technology*, *53*, 233-239.
 https://doi.org/10.1016/j.lwt.2013.02.025
- de Marco Castro, E., Shannon, E., & Abu-Ghannam, N. (2019). Effect of Fermentation on Enhancing the Nutraceutical
 Properties of *Arthrospira platensis* (Spirulina). *Fermentation*, 5, 28. https://doi.org/10.3390/fermentation5010028
- Fromm, H. I., & Boor, K. J. (2004). Characterization of pasteurized fluid milk shelf-life attributes. *Journal of Food Science*,
 69, 207-214. <u>https://doi.org/10.1111/j.1365-2621.2004.tb09889.x</u>
- Fromm, H. I., & Boor, K. J. (2004). Characterization of pasteurized fluid milk shelf-life attributes. *Journal of Food Science*,
 69, 207-214. <u>https://doi.org/10.1111/j.1365-2621.2004.tb09889.x</u>
- Golmakani, M.-T., Soleimanian-Zad, S., Alavi, N., Nazari, E., & Eskandari, M. H. (2019). Effect of Spirulina (*Arthrospira platensis*) powder on probiotic bacteriologically acidified feta-type cheese. *Journal of Applied Phycology*, *31*, 1085–1094.
 https://doi.org/10.1007/s10811-018-1611-2
- Huang, H-W., Lung, H-M., Yang, B.B., & Wang, C-Y. (2014). Responses of microorganisms to high hydrostatic pressure
 processing, a Review. *Food Control*, 40, 250-259. https://doi.org/10.1016/j.foodcont.2013.12.007
- 457 Hynstova, V., Sterbova, D., Klejdus, B., Hedbavny, J., Huska, D., & Adam, V. (2018). Separation, identification and
- 458 quantification of carotenoids and chlorophylls in dietary supplements containing Chlorella vulgaris and Spirulina platensis
- using High Performance Thin Layer Chromatography. *Journal of Pharmaceutical and Biomedical Analysis*, *148*, 108–118.
 https://doi.org/10.1016/j.jpba.2017.09.018

- 461 İlter, I., Akyıl, S., Demirel, Z., Koç, M., Conk-Dalay, M., & Kaymak-Ertekin, F. (2018). Optimization of phycocyanin
 462 extraction from *Spirulina platensis* using different techniques. *Journal of Food Composition and Analysis, 70, 78–88.*463 https://doi.org/10.1016/j.jfca.2018.04.007
- Jun, X. (2013). High-Pressure Processing as Emergent Technology for the Extraction of Bioactive Ingredients From Plant
 Materials. *Critical Reviews in Food Science and Nutrition*, 53, 837–852. <u>https://doi.org/10.1080/10408398.2011.561380</u>
- Kalaji, M. H., Goltsev, V. N., Żuk-Gołaszewska, K., Zivcak, M., & Brestic, M. (2017). Chlorophyll fluorescence:
 understanding crop performance—basics and applications. CRC Press, Boca Raton, FL, USA.
- Kang, D., Gho, Y. S., Suh, M., & Kang, C. (2002). Highly sensitive and fast protein detection with Coomassie brilliant blue
 in sodium dodecyl sulfate-polyacrylamide gel electrophoresis. *Bulletin of the Korean Chemical Society*, 23, 1511–1512.
- 470 <u>https://doi.org/10.5012/bkcs.2002.23.11.1511</u>
- Koca, N., Karadeniz, F., & Burdurlu, H. S. (2007). Effect of pH on chlorophyll degradation and colour loss in blanched green
 peas. *Food Chemistry*, 100, 609-615. <u>https://doi.org/10.1016/j.foodchem.2005.09.079</u>
- Kumar, K., Mishra, S. K., Shrivastav, A., Park, M. S., & Yang, J.-W. (2015). Recent trends in the mass cultivation of algae
 in raceway ponds. *Renewable and Sustainable Energy Reviews*, *51*, 875–885. https://doi.org/10.1016/j.rser.2015.06.033
- Laemmi, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, 227,
 680-685. <u>https://doi.org/10.1038/227680a0</u>
- Leggett, M. J., McDonnell, G., Denyer, S. P., Setlow, P. & Maillard, J.-Y. (2012). Bacterial spore structures and their
 protective role in biocide resistance. *Journal of Applied Microbiology*, *113*, 485-498. <u>https://doi.org/10.1111/j.1365-</u>
 2672.2012.05336.x
- Li, Y., Zhang, Z., Paciulli, M., & Abbaspourrad, A. (2020). Extraction of phycocyanin—A natural blue colorant from dried
 spirulina biomass: Influence of processing parameters and extraction techniques. *Journal of Food Science*, *85*, 727 735.
 https://doi.org/10.1111/1750-3841.14842
- Lichtenthaler, H. K. (1987). Chlorophylls and Carotenoids: Pigments of Photosynthetic Biomembranes. *Methods in Enzymology*, 148, 350–382. <u>https://doi.org/10.1016/0076-6879(87)48036-1</u>
- Liu, Q., Huang, Y., Zhang, R., Cai, T., & Cai, Y. (2016). Medical application of *Spirulina platensis* derived C-phycocyanin.
 Evidence-Based Complementary and Alternative Medicine, 2016, 7803846. <u>https://doi.org/10.1155/2016/7803846</u>
- Machu, L., Misurcova, L., Vavra Ambrozova, J., Orsavova, J., Mlcek, J., Sochor, J., & Jurikova, T. (2015). Phenolic Content
 and Antioxidant Capacity in Algal Food Products. *Molecules*, 20, 1118-1133. <u>https://doi.org/10.3390/molecules20011118</u>
- Martelli, F., Alinovi, M., Bernini, V., Gatti, M., & Bancalari, E. (2020). *Arthrospira platensis* as Natural Fermentation Booster
 for Milk and Soy Fermented Beverages. *Foods*, 9, 350. https://doi.org/doi:10.3390/foods9030350

- Martelli, F., Cirlini, M., Lazzi, C., Neviani, E., & Bernini, V. (2020). Edible Seaweeds and Spirulina Extracts for Food
 Application: In Vitro and In Situ Evaluation of Antimicrobial Activity towards Foodborne Pathogenic Bacteria. *Foods*, *9*,
 1442. https://doi.org/10.3390/foods9101442
- 494 Martelli, F., Favari, C., Mena, P., Guazzetti, S., Ricci, A., Del Rio, D., Lazzi, C., et al. (2020). Antimicrobial and Fermentation
 495 Potential of *Himanthalia elongata* in Food Applications. Microorganisms, 8, 248.
 496 http://doi.org/10.3390/microorganisms8020248
- Martelli, F., Cirlini, M., Lazzi, C., Neviani, E., & Bernini, V. (2021). Solid-State Fermentation of *Arthrospira platensis* to
 Implement New Food Products: Evaluation of Stabilization Treatments and Bacterial Growth on the Volatile Fraction. *Foods*,
 10, 67. https://doi.org/10.3390/foods10010067
- 500 Martelli, F., Marrella, M., Lazzi, C., Neviani, E., & Bernini, V. (2021). Microbiological contamination of ready to eat algae 501 and evaluation of *Bacillus cereus* behavior by microbiological challenge test. *Journal of food protection*, 10, 4315.
- Martelli, G., Folli, C., Visai, L., Daglia, M., & Ferrari, D. (2014). Thermal stability improvement of blue colorant C Phycocyanin from *Spirulina platensis* for food industry applications. *Process Biochemistry*, 49, 154–159.
 https://doi.org/10.1016/j.procbio.2013.10.008
- Morais, M. G., Radmann, E. M., Andrade, M. R., Teixeira G. G., Brusch, L. R. F., & Costa J. A. V. (2009). Pilot scale
 semicontinuous production of Spirulina biomass in southern Brazil. *Aquaculture 294*, 60–64.
 https://doi.org/10.1016/j.aquaculture.2009.05.009
- N'Dri, D., Calani, L., Mazzeo, T., Scazzina, F., Rinaldi, M., Del Rio, D., Pellegrini, N., & Brighenti, F. (2010). Effects of
 different maturity stages on antioxidant content of Ivorian Gnagnan (*Solanum indicum* L.) berries. *Molecules*, *15*, 7125-7138.
- 510 <u>https://doi.org/10.3390/molecules15107125</u>
- Niccolai, A., Chini Zittelli, G., Rodolfi, L., Biondi, N., & Tredici, M. R. (2019). Microalgae of interest as food source:
 Biochemical composition and digestibility. *Algal Research*, *42*, 101617. <u>https://doi.org/10.1016/j.algal.2019.101617</u>
- Nouri, E., Abbasi, H., & Rahimi, E. (2018). Effects of processing on stability of water-and fat-soluble vitamins, pigments (C phycocyanin, carotenoids, chlorophylls) and colour characteristics of *Spirulina platensis*. *Quality Assurance and Safety of*
- 515 Crops & Foods, 10, 335-349. <u>https://doi.org/10.3920/QAS2018.1304</u>
- Paciulli, M., Palermo, M., Chiavaro, E., & Pellegrini, N. (2017). Chlorophylls and colour changes in cooked vegetables. In
 E. M. Yahia (Ed.) *Fruit and Vegetable Phytochemicals: Chemistry and Human Health: Second Edition*, 703–719.
- 518 <u>https://doi.org/10.1002/9781119158042.ch31</u>
- 519 Paciulli, M., Rinaldi, M., Rodolfi, M., Ganino, T., Morbarigazzi, M., & Chiavaro, E. (2019). Effects of high hydrostatic
- pressure on physico-chemical and structural properties of two pumpkin species. *Food Chemistry*, 274, 281–290.
 https://doi.org/10.1016/j.foodchem.2018.09.021

- Park, W. S., Kim, H. J., Li, M., Lim, D. H., Kim, J., Kwak, S. S., Kang, C.-M., Ferruzzi, M. G., & Ahn, M. J. (2018). Two
 classes of pigments, carotenoids and C-phycocyanin, in Spirulina powder and their antioxidant activities. *Molecules*, 23, 2065.
 https://doi.org/10.3390/molecules23082065
- 525 Parniakov, O., Apicella, E., Koubaa, M., Barba, F. J., Grimi, N., Lebovka, N., Pataro, G., Ferrari, G., & Vorobiev, E. (2015).
- 526 Ultrasound-assisted green solvent extraction of high-added value compounds from microalgae *Nannochloropsis* spp.
 527 *Bioresource Technology*, 198, 262–267, <u>https://doi.org/10.1016/j.biortech.2015.09.020</u>
- Rajakumar, M. S., & Muthukumar, K. (2018). Influence of pre-soaking conditions on ultrasonic extraction of *Spirulina platensis* proteins and its recovery using aqueous biphasic system. *Separation Science and Technology*, *53*, 2034-2043.
 https://doi.org/10.1080/01496395.2018.1442860
- Reddy, N. R., Tetzloff, R. C., Solomon, H. M., & Larkin, J. W. (2006). Inactivation of *Clostridium botulinum* non proteolytic
 type B spores by high pressure processing at moderate to elevated high temperatures. *Innovative Food Science & Emerging Technologies*, 7, 169–175. https://doi.org/10.1016/j.ifset.2006.03.002
- Richardson, J. W., Johnson, M. D., & Outlaw, J. L. (2012). Economic comparison of open pond raceways to photo bioreactors for profitable production of algae for transportation fuels in the Southwest. *Algal Research*, *1*, 93–100.
 https://doi.org/10.1016/j.algal.2012.04.001
- Trabelsi, L., M'sakni, N. H., Ouada, H. B., Bacha, H., & Roudesli, S. (2009). Partial characterization of extracellular
 polysaccharides produced by cyanobacterium *Arthrospira platensis*. *Biotechnology and Bioprocess Engineering*, *14*, 27-31.
 https://doi.org/10.1007/s12257-008-0102-8
- Wang, H., Zhang, W., Chen, L., Wang, J., & Liu, T. (2013). The contamination and control of biological pollutants in mass
 cultivation of microalgae. *Bioresource Technology*, *128*, 745–750. <u>https://doi.org/10.1016/j.biortech.2012.10.158</u>
- Wu, L-C., Ho, J-A. A., Shien, M-C., & Lu, I-W. (2005). Antioxidant and Antiproliferative Activities of Spirulina and
 Chlorella Water Extracts. *Journal of Agricultural and Food Chemistry*, 53, 4207-4212. https://doi.org/10.1021/jf0479517
- Yu, J., Hu, H., Wu, X., Wang, C., Zhou, T., Liu, Y., Ruan, R., & Zheng H. (2019). Continuous cultivation of *Arthrospira platensis* for phycocyanin production in large-scale outdoor raceway ponds using microfiltered culture medium. *Bioresource Technology*, 287, 121420. https://doi.org/10.1016/j.biortech.2019.121420
- Zarbà, C., Chinnici, G., & D'Amico, M. (2020). Novel Food: The Impact of Innovation on the Paths of the Traditional Food
 Chain. *Sustainability*, *12*, 555. <u>https://doi.org/10.3390/su12020555</u>
- Zhang, Z., Cho, S., Dadmohammadi, Y., Li, Y., & Abbaspourrad, A. (2021). Improvement of the storage stability of Cphycocyanin in beverages by high-pressure processing. *Food Hydrocolloids*, *110*, 106055.
 <u>https://doi.org/10.1016/j.foodhyd.2020.106055</u>
- Zheng, J. X., Yin, H., Shen, C. C., Zhang, L., Ren, D. F., & Lu, J. (2020). Functional and structural properties of spirulina
 phycocyanin modified by ultra-high-pressure composite glycation. *Food Chemistry*, *306*, 125615.
 <u>https://doi.org/10.1016/j.foodchem.2019.125615</u>

- 555 Zhu, Y., Chen, X. B., Wang, K. B., Li, Y. X., Bai, K. Z., Kuang, T. Y., & Ji, H. B. (2007). A simple method for extracting C-
- 556 phycocyanin from Spirulina platensis using Klebsiella pneumoniae. Applied Microbiology and Biotechnology, 74, 244-248.
- 557 <u>https://doi.org/10.1007/s00253-006-0636-7</u>
- 558

559 Table 1. Total phenolic content and antioxidant capacity of aqueous suspension of A. platensis analysed after different

560 treatment at time zero and after 6 days of storage

Storage time (d)	Treatment	Total phenolic content $(mg/L GAF)$	Antioxidant capacity	Antioxidant capacity	Tabella formattata
	Control	$149.49^{aB} \pm 0.31$	$0.472^{aB} + 0.036$	$12.65^{aAr} + 0.07$	
	HPP400	$144.02^{aB}\pm 5.53$	0.472 ± 0.030 $0.424^{aA} \pm 0.026$	$\frac{12.03}{11.36^{84} \pm 0.70}$	
0	HPP600	$146.58^{\mathrm{aA}}\pm2.72$	$0.424^{\circ} \pm 0.020^{\circ}$ $0.437^{aA} \pm 0.015^{\circ}$	$11.30^{\circ} \pm 0.70^{\circ}$	
	PAST	$95.80^{bA}\pm6.45$	$0.372^{aA} \pm 0.028$	$9.97^{aA} \pm 0.74$	
	STER	$112.62^{bA}\pm 6.42$	$0.450^{\mathrm{aA}} \pm 0.021$	$\frac{12.05^{aA} \pm 0.56}{2}$	
	Control	$216.48^{aA}\pm 2.07$	$0.578^{\rm aA}\pm 0.027$	$\frac{15.50^{aA} \pm 0.72}{2}$	
	HPP400	$205.94^{\mathrm{aA}}\pm5.67$	$0.461^{bA}\pm 0.011$	$\frac{12.35^{bA} \pm 0.29}{2}$	
6	HPP600	$142.68^{bA}\pm 0.82$	$0.427^{bA} \pm 0.004$	$\frac{11.45^{bA} \pm 0.12}{2}$	
	PAST	$113.89^{\mathrm{cA}}\pm6.31$	$0.348^{cA} \pm 0.008$	$9.33^{b,cA} \pm 0.22$	
	STER	$110.31^{cA}\pm 1.97$	$0.413^{bA}\pm 0.016$	$\frac{11.06^{bA} \pm 0.44}{2}$	
ower case superscrip	ts highlight differe	ences and/or analogies among dif	fferent treatments within a colum	nn.	

562 A-B upper case superscripts indicate a significant difference (within a column) between the same treatment analysed at 0 d and 6 d of storage

563 time.

Abbreviations: raw, non-treated sample (control); 400 MPa for 3 min (HPP400), 600 MPa for 3 min (HPP600), pasteurized

565 (PAST) and sterilized (STER).

566

567

568

Table 2. Colorimetric parameters (L*, a*, b*, ΔE) of *A. platensis* aqueous suspensions (5% w/v). Treatments of *A. platensis*

570 aqueous dispersion were: fresh, non-treated A. platensis (control); high pressure processed (HPP) sample at 400MPa

571 (HPP400); HPP sample at 600MPa (HPP600); 90°C x 10 min pasteurized sample (PAST); 121°C x 20 min sterilized sample

572 (STER). Treatments were analysed at time zero (0 d) and after 6 days of refrigerated storage.

Storage time (d) Treatment		L*	a*	b*	ΔΕ
	Control	$4.06^{\text{bA}}\pm0.92$	$6.03^{\mathrm{bB}}\pm1.00$	$1.25^{\text{bA}}\pm0.28$	-
	HPP400	$9.04^{\mathrm{aA}}\pm0.77$	$8.67^{aB}\pm2.07$	$1.34^{\mathrm{bA}}\pm0.36$	$5.85^{\text{cA}}\pm1.36$
0	HPP600	$10.31^{\mathrm{aA}}\pm1.09$	$0.61^{\text{dB}}\pm0.99$	$1.73^{bA}\pm0.29$	$8.37^{\mathrm{cA}}\pm0.63$
	PAST	$11.85^{\mathrm{aA}}\pm2.32$	$\textbf{-}11.25^{eA}\pm1.58$	$13.11^{\mathrm{aA}}\pm3.11$	$22.45^{aB}\pm3.54$
	STER	$11.66^{aA}\pm2.04$	$3.39^{\mathrm{cA}}\pm0.35$	$12.19^{\mathrm{aA}}\pm1.97$	$13.60 \ ^{\rm bA} \pm 2.73$
	Control	$6.23^{bcA}\pm1.35$	$12.75^{bA} \pm 1.47$	$0.32^{bB}{\pm}0.50$	-
	HPP400	$5.68^{\mathrm{cB}}\pm0.84$	$17.08^{aA} {\pm}~0.88$	$\text{-}1.46^{\text{bB}} \pm 0.54$	$4.81^{\text{cA}}\pm0.78$
6	HPP600	$6.30^{bcB}{\pm}1.17$	$6.80^{cA} \!\pm\! 0.87$	$0.97^{\mathrm{bB}}\pm0.15$	$6.08^{\mathrm{cB}}\pm0.87$
	PAST	$11.36^{\mathrm{aA}}\pm2.83$	$\textbf{-}11.50^{eA}\pm1.94$	$12.23^{\mathrm{aA}}\pm3.35$	$27.66^{\rm aA} {\pm}~ 3.55$
	STER	$9.67^{abA}\pm1.60$	$1.14^{\text{dA}}\pm0.47$	$10.67^{\mathrm{aA}}\pm1.29$	$15.86^{bA}\pm1.20$

573 a-c lower case superscripts highlight differences and/or analogies among different treatments within a column.

574 A-B upper case superscripts indicate a significant difference (within a column) between the same treatment analysed at 0 d and 6 d of storage

575 time.

577

578

576 Abbreviations: lightness (L*), Redness (a*), Yellowness (b*) and total colour differences (Δ E) between 0 and 6 days.

ha formattato: Inglese (Stati Uniti)

579	Table 3. Total content of chlorophyll a (Ca), chlorophyll b (Cb), total chlorophylls (Ca + Cb), total carotenoids (Cx+c), total
580	chlorophylls and carotenoids (Ccc), pheophytins (Cph) of different treatments of A. platensis in aqueous suspension (5% w/v).
581	Treatments of A. platensis aqueous dispersion were: fresh, non-treated A. platensis (control); high pressure processed (HPP)
582	sample at 400MPa (HPP400); HPP sample at 600MPa (HPP600); 90°C x 10 min pasteurized sample (PAST); 121°C x 20
583	min sterilized sample (STER). Treatments were analysed at time zero (0 d) and after 6 days of shelf life.

Storage time (d)	Treatment	Ca (µg/mL)	Cb (µg/mL)	Ca+Cb (µg/mL)	Cx+c (µg/mL)	$Ccc (\mu g/mL)$	Cph (µg/mL)
	Control	$135^{aA} \pm 8$	5.9 ^{aA} ± 4.1	186 ^{aA} ± 6	$46^{aA} \pm 4$	233 ^{aA} ± 10	779 ^{aA} ± 45
	HPP400	148 ^{aB} ± 5	$3.5^{abA} \pm 0.4$	201 ^{aB} ± 6	47 ^{aA} ± 1	249 ^{aB} ± 7	$820^{aB} \pm 23$
0	HPP600	142 ^{aA} ±13	$2.8^{abA} \pm 0.3$	192ª ^A ± 18	45 ^{aB} ± 2	$237^{aA} \pm 20$	784 ^{aA} ± 73
	PAST	92 ^{bB} ± 8	$2.1^{abA} \pm 0.6$	125 ^{bB} ± 11	43 ^{aB} ± 5	$168^{bB} \pm 16$	476 ^{bB} ± 43
	STER	43 ^{cA} ± 3	$0.6^{bA} \pm 0.2$	57 ^{cA} ± 4	26 ^{bA} ± 1	83 ^{cA} ± 5	184 ^{cA} ± 11
	Control	192ª ^A ± 21	5.0 ^{aA} ± 2.3	260 ^{aA} ± 30	60 ^{aA} ± 6	320 ^{aA} ± 36	1062 ^{aA} ± 123
	HPP400	189 ^{aA} ± 10	$5.0^{aA} \pm 2.1$	257ª ^A ± 11	$58^{aA} \pm 4$	315 ^{aA} ± 14	1017 ^{aA} ± 49
6	HPP600	185 ^{aA} ± 10	$4.6^{abA} \pm 1.8$	251 ^{aA} ± 11	59 ^{aA} ± 3	309 ^{abA} ± 15	1034 ^{aA} ± 43
	PAST	138 ^{bA} ± 7	$1.2^{abA} \pm 0.6$	185 ^{bA} ± 9	$60^{aA} \pm 3$	245 ^{bcA} ± 12	$702^{abA} \pm 44$
	STER	105 ^{bA} ± 21	$0.7^{bA} \pm 0.5$	141 ^{bA} ± 29	53 ^{aA} ± 12	194 ^{cA} ± 41	484 ^{bA} ± 113

584 a-e lower case superscripts highlight differences and/or analogies among different treatments within a column.

A-B upper case superscripts indicate a significant difference (within a column) between the same treatment analysed at 0 d and 6 d of storage
time.

Table 4. Microbial concentration at 0 d and after 6 d of storage at refrigerated temperatures. Microbial concentration was
 expressed as Log CFU/ml.

		Microbial species						
Storage time (d)	Treatment	Aerobic spor.	Anaerobic spor.	TBC	Yeast	Salmonella	Coliform	Staphylococci
	Control	2.84 ± 0.12^{aA}	2.66 ± 0.19^{aA}	5.96 ± 0.05^{aA}	4.44 ± 0.41^{aA}	$4.53\pm0.15^{\mathrm{aA}}$	$\begin{array}{c} 5.06 \pm \\ 0.08^{aA} \end{array}$	4.94 ± 0.19^{aA}
	HPP400	1.88 ± 0.29^{bA}	1.83 ± 0.27^{bA}	1.66 ± 0.08^{bA}	<1 ^{bA}	2.70 ± 0.17^{bA}	nd ^{bA}	1.70 ± 0.20^{bA}
0	HPP600	1.81 ± 0.31^{bA}	$1.70\pm0.20^{b\rm A}$	$1.00\pm0.17^{b\rm A}$	<1 ^{bA}	1.85 ± 0.08^{bA}	nd ^{bA}	1.70 ± 0.10^{bA}
	PAST	$1.86\pm0.32^{b\rm A}$	1.66 ± 0.25^{bA}	1.66 ± 0.19^{bA}	nd ^{bA}	ndcA	nd ^{bA}	<1 ^{bA}
	STER	nd ^{cA}	nd ^{cA}	nd ^{cA}	nd ^{bA}	ndcA	nd ^{bA}	nd ^{bA}
	Control	1.44 ± 0.29^{aB}	2.65 ± 0.17^{aA}	6.90 ± 0.07^{aB}	5.14 ± 0.12^{aB}	2.10 ± 0.10^{aB}	$\begin{array}{c} 6.83 \pm \\ 0.15^{aB} \end{array}$	6.76 ± 0.21^{aB}
	HPP400	1.86 ± 0.26^{bA}	1.85 ± 0.19^{bA}	$3.40\pm0.09^{\rm bB}$	1.18 ± 0.17^{bA}	nd ^{bB}	nd ^{bA}	1.83 ± 0.08^{bB}
6	HPP600	1.71 ± 0.21^{bB}	1.56 ± 0.17^{bB}	2.35 ± 0.12^{bB}	1.30 ± 0.08^{bB}	nd ^{bB}	nd ^{bA}	1.81 ± 0.13^{bB}
	PAST	1.48 ± 0.26^{bB}	1.50 ± 0.31^{bB}	2.00 ± 0.08^{bB}	nd ^{cA}	nd ^{bA}	nd ^{bA}	1.78 ± 0.31^{bB}
	STER	nd ^{cA}	nd ^{cA}	nd ^{cA}	nd ^{cA}	nd ^{bA}	nd ^{bA}	nd ^{cA}

599 Results are shown as the mean \pm standard deviation. nd, not detected

600 a- c Different lower-case superscript letters highlight significant differences (P<0.05) among treatments within each column

601 A-B Different upper-case superscript letters highlight significant differences (P<0.05) between the same treatments at 0 and 6

602 days of storage

Abbreviations: aerobic spore-forming bacteria (Aerobic spor), anaerobic spore-forming bacteria (Anaerobic spor.), Total

604 <u>Bacteria Count (TBC).</u>

605

606 Figure captions

Fig.1 Visual appearance of *A. platensis* in aqueous suspensions (5% w/v) colour characteristics at 0 and 6 d of storage time.
Samples were: fresh, non-treated *A. platensis* (control); high pressure processed (HPP) sample at 400 MPa (HPP400); HPP
sample at 600 MPa (HPP600); 90°C x 10 min pasteurized sample (PAST); 121°C x 20 min sterilized sample (STER).
Treatments were analysed at time zero (0 d) and after 6 days of refrigerated storage.

611

Fig.2 Water holding capacity (WHC) of *A. platensis* in aqueous suspensions (5% w/v) at 0 and 6 d of storage time. Samples
were: fresh, non-treated *A. platensis* (control); high pressure processed (HPP) sample at 400 MPa (HPP400); HPP sample at
600 MPa (HPP600); 90°C x 10 min pasteurized sample (PAST); 121°C x 20 min sterilized sample (STER). Treatments were
analysed at time zero (0 d) and after 6 days of refrigerated storage.

616

Fig.3 Concentration of phycocyanin (μg/mL) of different treatments of *A. platensis* in aqueous suspension (5% w/v).
Treatments of *A. platensis* aqueous dispersion were: fresh, non-treated *A. platensis* (control); high pressure processed (HPP)
sample at 400MPa (HPP400); HPP sample at 600MPa (HPP600); 90°C x 10 min pasteurized sample (PAST); 121°C x 20
min sterilized sample (STER). Treatments were analysed at time zero (0 d) and after 6 days of shelf life.

621

622	Fig.4 Light microscopy observations (100x magnification) of A. platensis in aqueous suspensions (5% w/v) processed by
623	applying different treatments. (a) raw, control suspension; (b) HPP sample processed at 400 MPa x 3 min; (c) HPP sample
624	processed at 600 MPa x 3 min; (d) sample pasteurized at 90 °C x 10 min; (b) sample sterilized at 121 °C x 20 min. <u>CF: cell's</u>
625	fragments: CB: cell's breakdown; AG: aggregates.

626

Fig.5 Sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS PAGE) results of *A. platensis* aqueous suspensions
(5 w/v) at 0 d of storage. Treatments of *A. platensis* aqueous dispersion were: fresh, non-treated *A. platensis* (control); high
pressure processed (HPP) sample at 400MPa (HPP400); HPP sample at 600MPa (HPP600); 90°C x 10 min pasteurized sample
(PAST); 121°C x 20 min sterilized sample (STER). Treatments were analysed at time zero (0 d) and after 6 days of shelf life.
Band 1: subunit of C-phycocyanin; band 2: subunit of C-phycocyanin; band 3: chlorophyll a complex; band 4: chlorophyll a

ha formattato: Tipo di carattere: Grassetto